

High frequency of new particle formation events driven by summer monsoon in the central Tibetan Plateau, China

Lizi Tang¹, Min Hu^{*1,2}, Dongjie Shang¹, Xin Fang¹, Jianjiong Mao², Wanyun Xu³, Jiacheng Zhou⁴, Weixiong Zhao⁴, Yaru Wang¹, Chong Zhang¹, Yingjie Zhang⁵, Jianlin Hu², Limin Zeng^{1,2}, Chunxiang Ye¹, Song Guo^{1,2}, Zhijun Wu^{1,2}

¹State Key Joint Laboratory of Environmental Simulation and Pollution Control, International Joint Laboratory for Regional Pollution Control, Ministry of Education (IJRC), College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

²Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control. Nanjing University of Information Science & Technology, Nanjing 210044, China

³State Key Laboratory of Severe Weather & Key Laboratory for Atmospheric Chemistry of CMA, Institute of Atmospheric Composition, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

⁴Laboratory of Atmospheric Physico-Chemistry, Anhui Institute of Optics and Fine Mechanisms, HFIPS, Chinese Academy of Science, Hefei 230031, Anhui, China

⁵School of Ecology and Nature Conservation, Beijing Forestry University, Beijing 100083, China

Correspondence to: Min Hu (minhu@pku.edu.cn).

Abstract

New particle formation (NPF) is an important source of cloud condensation nuclei (CCN), which affects Earth's radiative balance and global climate. The mechanism and CCN contribution of NPF at the high-altitude mountains, especially in the Tibetan Plateau (TP), was unclear due to lack of measurements. In this study, intensive measurements were conducted at Nam Co station (4730 m a.s.l) in the central TP during both pre-monsoon and summer monsoon seasons. The frequencies of NPF events exhibited extreme distinction with 15% in pre-monsoon season and 80% in monsoon season. were favorable for NPFThe comprehensive analysis of the measured CS, gaseous precursors and meteorological conditions, supplemented by the model simulations of SO₂ and VOC, points to the organic involved nucleation as the dominant mechanism. Condensation sink and gaseous sulfuric acid could have no significant effect on the occurrence of NPF events. The frequent NPF events in summer monsoon season may result from the higher frequency of southerly and southwesterly air masses, which brought the organic precursors to participate in the NPF process. The level of organic vapours governed

带格式的: 下标

~~the occurrence of NPF events, while condensation sink and gaseous sulfuric acid had no effect. The frequent NPF events in summer monsoon season resulted from the higher concentration of organic vapours, which was brought from northeast India by the strong southerly monsoon.~~ It had increased the aerosol number concentrations and CCN at supersaturation of 1.2% by more than 2 and 0.6 times compared with those in pre-monsoon season, respectively. Considering that the smaller particles formed by NPF may further grow and reach CCN size during the following days due to the low-level coagulation sink, the amount of potential CCN in monsoon season could be much larger than our local measurement results. Our results emphasized the importance of considering the seasonal effect of NPF when simulating the amounts of aerosols and CCN in the high-altitude atmosphere. Long-term investigations with a full set of instrumentation are required for deeper scientific understanding of NPF process and its role in the global budget in the TP. organics to NPF in high-altitude atmosphere, and the seasonal effect of NPF at the high-altitude sites should be carefully considered in model simulations to reduce the uncertainty of global CCN budget.

1 Introduction

Atmospheric aerosols can affect Earth's radiative balance through direct interaction with solar radiation and by serving as cloud condensation nuclei (CCN), and thus global climate. There is considerable uncertainty about the total radiative forcing contributed by aerosol particles, mainly from the number concentrations and sizes of CCN (IPCC, 2021). New particle formation (NPF) is a physical and chemical process, comprising nucleation of gaseous precursors and the subsequent growth of the nucleated clusters into aerosol particles. NPF has been considered to be an important source of aerosols and contributes a major fraction to the global CCN budget. Model simulations found that up to 45% of global CCN were secondary aerosols derived from NPF process, with a large fraction created in the free troposphere (Merikanto et al., 2009). Furthermore, NPF may be more important to the total CCN budget in the pristine atmosphere than that in the industrial atmosphere (Gordon et al., 2017). Similarly, the field observations in diverse atmospheric environments showed the more significant CCN contribution from NPF events in pristine atmosphere, especially in high-altitude mountains (Kerminen et al., 2018).

~~With the development of technology and instrumentation, NPF events have been observed in diverse atmospheric environments around the world, including urban, rural, mountain, boreal, and polar regions (Nieminen et al., 2018; Kerminen et al., 2018; Wang et al., 2017; Peng et al., 2014). High altitude mountains are particularly interesting environment for atmospheric NPF studies due to near absence of primary particle sources and long aerosol particle lifetimes in such environment (Kerminen et al., 2018). Furthermore, the NPF events at the high-altitude sites have been considered as~~

60 a potentially important source of particles injected into the free troposphere (Venzac et al., 2008). Unfortunately, there were
few NPF studies at pristine high-altitude sites, mainly due to the challenges in measuring particle numbers and precursors
under the adverse meteorological and topographic conditions such as the extremely low temperature, the thin air and the
steep mountains. Moreover, the parameters and the exact mechanisms of NPF in the limited high-altitude measurements
showed large temporal and spatial variations. For example, variable frequencies of NPF event days were found: 20% during
one year at Jungfraujoch, Switzerland (3580 m a.s.l.) (Tröstl et al., 2016a), 56% in spring and 43% in summer at Storm Peak
Laboratory, USA (3210 m a.s.l.) (Hallar et al., 2011), 67% during one year in Maïdo observatory, Réunion (2165 m a.s.l.)
(Rose et al., 2019), and 80% in post-monsoon season at Mt. Daban, China (3295 m a.s.l.) (Du et al., 2015). In Jungfraujoch,
65 NPF was limited to the availability of highly oxidized organic species, which was restricted by the contact time in the air
masses with the boundary layer (Bianchi et al., 2016). In Storm Peak Laboratory, NPF events were correlated with UV
irradiance, but not with O₃ concentration and existing particle surface area (i.e. condensation sink) (Hallar et al., 2011). But
at Chacaltaya Station in Bolivia, Maïdo observatory, and Jungfraujoch, the NPF frequency was found to be positively
correlated with CS, which was mainly because the increase of transmitted condensable precursors was often accompanied
70 by the increase of CS (Rose et al., 2015; Rose et al., 2019; Boulon et al., 2010). These results indicated the location and
seasonal effects of NPF at the high-altitude sites, and more measurements and studies were needed in high-altitude
atmosphere.

75 _____ The Tibetan Plateau (TP) is the largest plateau in China and the highest plateau in the world, with an average
altitude of over 4000 m a.s.l. It is known as the "roof of the world" and "the third pole" of the Earth and plays a fundamental
role in the regional climate and environment through various dynamic and thermal effects (Yanai and Wu, 2006). Due to the
rare anthropogenic activities, the TP is considered one of the most pristine locations around the world and an ideal location
for characterizing for observations of free-tropospheric air masses and NPF events remote and regional background aerosols.
At the same time, NPF is an assumed key source of CCN production in the TP and has large impact on cloud processes, due
80 to the negligible primary particle emissions. However, Unfortunately, there were few NPF studies at pristine high-altitude
sites in the TP, mainly due to the challenges in measuring particle numbers and precursors under the adverse meteorological
and topographic conditions such as the extremely low temperature, the thin air and the steep mountains. research about NPF
in the TP is rather limited. Several long-term and short-term observations have been conducted in different regions of the TP
in recent years. It restrained the understanding of mechanisms and CCN contribution of NPF in the TP, and brought
85 considerable uncertainty about the radiative forcing. The significant seasonal variation of NPF frequency was observed in the

90 ~~TP. For example, Some scientists have conducted~~ the long-term NPF measurement at Himalayan Nepal Climate Observatory at Pyramid (NCO-P) site on the southern TP ~~showed, and found that~~ the distinct NPF frequency ~~varied in each~~ different seasons with the highest in the monsoon period (Venzac et al., 2008). The occurrence of NPF events in NCO-P was initiated by the up-valley winds which sent the biogenic condensable vapours from Khumbu valley to the high-altitude location (Bianchi et al., 2021). ~~The occurrence of NPF events in NCO-P was initiated by the up-valley winds which sent the biogenic condensable vapours from Khumbu valley to the high-altitude location (Bianchi et al., 2021).~~ At Mt. Yulong on the southeastern TP, the NPF frequency was only 14% during pre-monsoon season and the occurrence of NPF events was related to an elevated boundary layer or transported biomass burning pollutants from southern Asia, but not biogenic condensable vapours (Shang et al., 2018; Du et al., 2015). For Mt. Daban on the northeastern TP, the high frequency of NPF events was observed during post-monsoon season (80%) (Du et al., 2015). These results indicated that the frequency and mechanism of NPF may associate with air mass origins and monsoon shift in the southern, southeastern and northeastern TP. However, there were few NPF studies conducted in the central TP especially for the reaction of NPF on monsoon season shifting in this region.

95 ~~Overall, few NPF studies were conducted in the central TP which can better represent the regional characteristics of TP compared with the southern and southeastern TP. The reason for the variation of NPF frequencies, the mechanisms for nucleation and growth of nanoparticles, and CCN contribution of NPF in various seasons in the TP remained ambiguous.~~

100 In this study, intensive measurements during pre-monsoon and monsoon seasons were conducted at Nam Co station (4730 m a.s.l) in the central TP, which can better represent the regional characteristics of TP during pre-monsoon and monsoon seasons. Collocated measurements including particle number size distributions (PNSD), trace gases and meteorological parameters, and assisted Weather Research and Forecasting (WRF) and Community Multiscale Air Quality (CMAQ) models were employed to investigate the characteristics of PNSD and NPF. This study aimed to (1) characterize NPF events in pre-monsoon and summer monsoon season in the central TP, (2) investigate the source of the NPF events occurrence in pre-monsoon and summer monsoon season in the central TP, and (3) quantify the CCN contribution of NPF in pre-monsoon and summer monsoon season in the central TP.

110 **2 Material and methods**

2.1 Measurement site

An intensive field campaign was carried out at a high mountain observatory at the central Tibetan Plateau, i.e., Nam Co

station (30.8°N, 91.0°E; 4730 m a.s.l) during the in-depth study of the atmospheric chemistry over the Tibetan Plateau in the year of 2019, referred as @Tibet 2019 field campaign (Fig. 1). The Nam Co station is located near Nam Co Lake (area: 1920 km²), the highest and largest saltwater lake in the world which is backed by the Nyainqentanglha mountains in the South. The ecology of the surrounding area is semi-arid land dominated by alpine meadow and barren areas. The capital city (Lhasa) of the Tibet Autonomous Region is about 100 km southeast of the station. The closest town, Dangxiong, is about 70 km southeast of the station, between which are the Nyainqentanglha mountains. Overall, Nam Co station is located in a typical pristine environment and there are almost no local anthropogenic source emissions in this area. The measurement was conducted from 26 April to 22 May, 2019 and 15 June to 25 June, 2019, and can be representative corresponding to the of the pre-monsoon season and the summer monsoon season, respectively (Text S1) (Bonasoni et al., 2010; Cong et al., 2015). –

2.2 Instruments and methods

The sampling was conducted at the observatory field of Nam Co station (Fig. 1c). O₃ was detected using ultraviolet (UV) absorption by a “Model 49C Ozone Analyzer” (Thermo Scientific, USA). CO and water content (H₂O) were measured by Picarro G2401 based on cavity ring-down spectroscopy. Meteorological parameters including wind speed (WS), wind direction (WD), temperature (T) and relative humidity (RH) were measured by the automatic meteorological station (Met one Instrument Inc). The photolysis frequencies of O₃ (JO¹D) were monitored using a spectral radiometer (ultra-fast CCD-detector spectrometer, METCON GmbH, Germany). 99 types of volatile organic compounds (VOC) were measured by an online gas chromatograph coupled with a mass spectrometer and flame ionization detectors (GC-MS/FID) (TH-PKU 300B, Wuhan Tianhong Instrument Co. Ltd., China) in pre-monsoon season (Wang et al., 2014). Black carbon (BC) was measured by Aethalometer (Magee Scientific, model AE33), and the concentration of BC at 880 nm was used in this study to reduce the influence of Brown carbon (Kirchstetter et al., 2004).

PNSD in the stokes size range of 4nm and 700 nm were obtained by integrating two sets of scanning mobility particle spectrometers (SMPS). The first SMPS measured particles with the size of 4-45 nm, consisting of a TSI Model 3085 DMA and a TSI Model 3776 CPC (with a flow rate of 1.5 L min⁻¹). The second SMPS measured particles with the size of 45-700 nm, consisting of a TSI Model 3081 DMA and a TSI Model 3775 CPC (with a flow rate of 0.3 L min⁻¹). A silicon diffusion tube was placed before the SMPS, which kept the relative humidity (RH) of the sampling air under 40%. PNSD were corrected for particle losses in the SMPS and the sampling tube, following the method of “equivalent pipe length” as described in Wiedensohler et al. (2012).

145 ~~Considering that the concentration of VOC in monsoon season has not been measured in this observation, Weather Research and Forecasting (WRF) (version 4.2.1) and The Community Multiscale Air Quality version 5.3.2 (CMAQv5.3.2) models have been adopted to simulate the VOC levels in the surrounding area to assist in the analysis of the role of VOC in NPF events. Similarly, SO₂ during pre-monsoon and monsoon was also simulated to help analyze the role of sulfuric acid in NPF events. Detailed information about the model setting and evaluation is provided in Text S1.~~

To reveal the transport pathway of air masses that arrive at the site, the 48h backward trajectories of the air mass at 5600m above the ground (52330 m a.s.l) were computed using the HYSPLIT (Hybrid Single Particle Lagrangian Integrated Trajectory) model and Global Data Assimilation System (GDAS) data.

2.3 Model simulation

150 ~~Considering that the limited measurements on concentration of SO₂, VOC and VOC in monsoon season has not been measured in this observation (only VOC during pre-monsoon season), Weather Research and Forecasting/Community Multiscale Air Quality (WRF/CMAQ) modeling system Weather Research and Forecasting (WRF) (version 4.2.1) and The Community Multiscale Air Quality version 5.3.2 (CMAQv5.3.2) models have been was adopted to simulate the VOC levels of SO₂ and VOC in the whole observation period in the surrounding area, to assist in the analysis of the role of sulfuric acid VOC and organics in NPF events. Similarly, SO₂ during pre-monsoon and monsoon was also simulated to help analyze the role of sulfuric acid in NPF events.~~

160 ~~Weather Research and Forecasting (WRF) (version 4.2.1) model was used to simulate the meteorological conditions with the FNL reanalysis dataset. The 6 h FNL data were obtained from the U.S. National Centre for Atmospheric Research (NCAR), with a spatial resolution of 1.0° × 1.0° (<http://rda.ucar.edu/datasets/ds083.2/>, last accessed on 28 April 2022). The Community Multiscale Air Quality version 5.3.2 (CMAQv5.3.2) model, being one of the three-dimensional chemical transport models (CTMs) (Appel et al., 2021), configured with the gas-phase mechanism of SAPRC07tic and the aerosol module of AERO6i, was employed in this study to simulate the air quality over Tibet in the observation period (26 April to 22 May and 15 June to 25 June in 2019). Air quality simulations were performed with a horizontal resolution of 12 km. The corresponding domain covered Tibet and the surrounding countries and regions with 166 × 166 grids (Fig. S2), with the 18 layers in vertical resolution. Detailed information about the model setting is provided in Text S2.~~

165 ~~The Multi-resolution Emission Inventory for China version 1.3 (MEICv1.3) (<http://www.meicmodel.org>) and Regional Emission inventory in ASia (REASv3.2) (<https://www.nies.go.jp/REAS/>) were used to provide the anthropogenic emissions from China and neighboring countries and regions, respectively. The MEICv1.3 emissions of the year 2019 were~~

170 used. For REAS, the emission inventory in the year 2015 was used for 2019 as no emission inventory was released for the
years after 2015. Although emission inventories are usually released 3 years behind, we acknowledge that this may cause
additional uncertainties in the simulation. Biogenic emissions were generated using the Model for Emissions of Gases and
Aerosols from Nature (MEGANv2.1) (Guenther et al., 2012). The open biomass burning emissions were processed using
the Fire Inventory for NCAR (FINN) during the entire study period (Wiedinmyer et al., 2011).

175 The model evaluation is introduced in Text S2. The WRF and CMAQ models successfully reproduced the
meteorological fields and air pollutants including PM and O₃ with model performance indices meeting the suggested
benchmarks. For VOC, the observed VOC and predicted VOC in pre-monsoon season were compared to examine the model
performance. The benchmarks for VOC had not been reported, but the statistical metrics of MFB (mean fractional bias,
-0.47) and MFE (mean fractional error, 0.49) in this study are within the range reported in previous VOC modelling result
(Hu et al., 2017). The correlation coefficient (R) between simulated and observed VOC is 0.41, which reflected that the
180 model can fairly simulate the variation of VOC concentration. It should be noted that VOC was underpredicted on the
whole, which may due to the uncertainty of the emission inventory as mentioned before. For SO₂, the WRF/CMAQ models
have been successfully reproduced SO₂ in major regions in China with R of 0.25-0.79 (Mao et al., 2022). The simulated
SO₂ level in the model domain is comparable with that measured at Mt. Yulong (Shang et al., 2018), with average values of
0.03±0.02 ppbv and 0.06±0.05 ppbv. At the same time, considering that both BC and SO₂ are mainly emitted from coal
combustion and biomass burning, BC could be a good indicator for SO₂ especially for pristine environment without local
185 anthropogenic source emissions. As shown in Fig.S6, a good correlation between SO₂ and BC measured at Mt. Yulong was
found with correlation coefficient (R) of 0.79 (Shang et al., 2018). In this study, the modelled SO₂ and measured BC also
showed good correlation with R of 0.58 (Fig. S6). In general, the results of model simulation showed good performance in
statistical parameters and correlation analysis with other tracers. The modelled VOC and SO₂ may be helpful for the NPF
190 analysis.

Detailed information about the model setting and evaluation is provided in Text S1.

2.4.3 Parameterization of NPF

195 In this study, a typical NPF event was defined by that there is the criteria that a burst in the 3-10 nm particles number
concentration (PN₃₋₁₀, PN₃₋₁₀ (particle number concentration in the diameter of 3-10 nm) increased obviously, and lasted for
more than 2 hours subsequent growth of these newly formed particles (Fang et al., 2020; Dal Maso et al., 2005). The days

带格式的: 下标

without newly particle formation were defined as non-event days. Days Other days in which the increase of PN_{3-10} was observed but without the particles growing to the larger size, or days in which the later phase of a mode growing in the Aitken mode size range can be observed lasted for less than 2 hours were treated as undefined days (Dal Maso et al., 2005).

The examples of the classification of NPF events are shown in Fig.S8 (Dal Maso et al., 2005)

During the NPF events, the formation rate was calculated using the following formula (Cai and Jiang, 2017):

$$J_{d_k} = \frac{dN_{[d_k, d_u]}}{dt} + \sum_{d_g=d_k}^{d_u-1} \sum_{d_i=d_{min}}^{+\infty} \beta_{(i,g)} N_{[d_i, d_{i+1}]} N_{[d_g, d_{g+1}]} - \frac{1}{2} \sum_{d_g=d_{min}}^{d_u-1} \sum_{d_i^2 = \max(d_{min}^2, d_k^2 - d_{min}^2)}^{d_{i+1}^2 + d_{g+1}^2 \leq d_u^2} \beta_{(i,g)} N_{[d_i, d_{i+1}]} N_{[d_g, d_{g+1}]} + n_u \cdot GR_u$$

(1)

Where J_{d_k} is the formation rate of particles at size d_k , and d_u is the upper size bound of the target size range. Here d_k and d_u are selected to be 4 and 25 nm, respectively. $N_{[d_k, d_u]}$ is the total number concentration of particles in the diameters of $[d_k, d_u]$. d_i is the lower bound of each measured size bin, and d_{min} is lowest size limit detected by measuring instrument. $\beta_{(i,g)}$ is the coagulation coefficient for the collision between the particle at size of d_i and the particle at size of d_g . n_u is the particle size distribution function (dN/dd_p). GR_u is the growth rate at size of d_u .

The growth rate (GR) was obtained by the mode-fitting method described in Dal Maso et al. (2005). In short, the PNSD during NPF event days were fitted as the sum of three-mode lognormal distribution. GR was calculated as the variation of the geometric mean diameter D_m of newly formed mode (4-25 nm) in unit internal (Dal Maso et al., 2005):

$$GR = \frac{\Delta D_m}{\Delta t}$$

(2)

To evaluate the scavenging effects of preexisting particles on condensable vapours, the condensation sink (CS) was calculated as follow (Dal Maso et al., 2005):

$$CS = 2\pi D \sum \beta_m(D_{p,i}) D_{p,i} N_i$$

(3)

where D is the diffusion coefficient of the condensing vapor, β_m is the transition regime correction factor, and $D_{p,i}$ and N_i are the diameter and number concentration in the size class i , respectively.

2.54 Calculation of CCN concentration

The CCN number concentration was calculated based on the assumption that particles larger than a certain diameter could act as CCN. The critical diameter (D_c) of the CCN activation at the supersaturation (S_c) can be calculated based on

κ – Köhler theory (Petters and Kreidenweis, 2007):

$$\kappa = \frac{4A^3}{27D_c^3 m^2 S_c}$$

(4)

$$A = \frac{4\sigma_{s/a}M_w}{RT\rho_w}$$

(5)

Where κ is the hygroscopicity parameter about the composition-dependence of the solution water activity. $\sigma_{s/a}$ is the water surface tension (0.0728 N m⁻¹). M_w and ρ_w are the molecular weight and density of water, respectively. R is the universal gas constant (J mol⁻¹ K⁻¹), and T is the absolute temperature (K).

3 Results and discussions

3.1 Characteristics of meteorology and atmospheric pollutants

The distinct meteorological characterizations were exhibited in the two seasons. As shown in Fig. 2 and Fig. S97, temperature behavior was characterized by higher value in monsoon season (10.4±4.1 °C) and lower value in pre-monsoon season (3.1±3.6 °C) with an average value of 5.3±5.1 °C. The relative humidity (RH) seems very similar in the two seasons (50%±21% in pre-monsoon season vs 48%±19% in monsoon season), but the water content (H₂O) in the air during monsoon season (1.0%±0.2%) was clearly higher than that in pre-monsoon season (0.6%±0.2%), which reflected the wetter environment in the monsoon period. Wind speed (WS) was comparable during the two seasons, which was 4.2±2.7 m s⁻¹ in pre-monsoon season and 4.5±2.7 m s⁻¹ in monsoon season, respectively. Wind direction (WD) showed a clear divergence, with westerly and southwesterly winds prevailing in pre-monsoon season, and southerly winds prevailing in monsoon season (Fig. S108). The frequencies of air masses arriving at the observation station from various directions during the two seasons were shown in Fig. S119. In pre-monsoon season, strong westerlies pass through western Nepal, northwest India and Pakistan (i.e., southern Himalayas). In monsoon season, air masses were more derived from Bangladesh and northeast India and brought moisture that originated in the Bay of Bengal. The seasonality of meteorology was generally in agreement with the previous studies in the TP, which is strongly influenced by the large-scale Asian monsoon circulation (Cong et al., 2015; Bonasoni et al., 2010).

As a background high altitude site on the TP, the Nam Co station displayed a low particle concentration. On average PM_{0.8} was 1.8±1.0 µg m⁻³, which was similar with PM₁ (2 µg m⁻³) measured by a high-resolution time-of-flight aerosol mass spectrometer at Nam Co station in 2015 (Xu et al., 2018a). The particulate matter concentration at Nam Co station was

lower than the PM₁ observed at Qomolangma Station (4.4 μg m⁻³) on the southern TP and Waliguan Observatory (9.1 μg m⁻³) on the northeastern TP, which resulted from a much longer transport distance of anthropogenic emissions compared with Qomolangma Station and Waliguan Observatory (Zhang et al., 2021). The average concentration of BC was 223±135 ng m⁻³ during the whole measurement period, which was comparable with previous observations (Xu et al., 2018a; Xu et al., 2020; Wang et al., 2016; Xu et al., 2018b), and represented the background level in the TP region. The average O₃ concentration was 58±15 ppbv, which was similar to the results from some high-elevation sites in the TP (Bonasoni et al., 2010; Shang et al., 2018) and higher than the results from Beijing during spring (Chen et al., 2020). Consistent with previous research (Xu et al., 2018a; Cong et al., 2015; Yin et al., 2021), higher PM, BC and O₃ concentrations were found during pre-monsoon season (Fig. S12⁹). The higher BC and PM may result from the active biomass-burning emissions in pre-monsoon season on the southern TP (Cong et al., 2015). The higher O₃ in pre-monsoon season were primarily attributed to stratospheric intrusion of ozone (Yin et al., 2021). In contrast, there was no noticeable difference in CO between the two seasons, and the average concentration was 0.12±0.14 ppmv.

3.2 Higher frequency of new particle formation in monsoon season

Fig. 2g shows the evolution of PNSD for the entire study. It can be seen that NPF events in the monsoon period were observed almost every day (8 in 10 days NPF days and 2 undefined days, 80%), while the frequency of NPF events was extremely low during the pre-monsoon period (4 in 27 days, 15%). As shown in Table 1, the event frequency at Nam Co station during monsoon season was higher than the result reported at NCO-P site on the southern TP in monsoon season (57%) (Venzac et al., 2008), and similar to the frequency reported at Mt. Daban on the northeastern TP in post-monsoon season (80%) (Du et al., 2015). The frequency at Nam Co station during pre-monsoon season was lower than NCO-P (38%) (Venzac et al., 2008) and comparable with Mt. Yulong on the southeastern TP (14%) (Shang et al., 2018) in the same season. In addition, the NPF frequencies of other high-altitude sites, such as Mt. Tai (Lv et al., 2018), Jungfrauoch (Tröstl et al., 2016a), Maïdo observatory (Rose et al., 2019), Storm Peak Laboratory (Hallar et al., 2011), were basically between those in pre-monsoon and monsoon seasons of Nam Co station. In general, the frequencies of the two seasons at Nam Co station can represent the highest and lowest values of NPF frequencies observed in the TP and other high-altitude sites, and even various environments (such as urban, rural, etc.) (Nieminen et al., 2018).

Table 1 summarizes the formation rate (J), growth rate (GR), and condensation sink (CS) at Nam Co station and other high-altitude sites. The J_4 at Nam Co station varied from 0.38 to 2.43 cm⁻³ s⁻¹, with an average value of 1.15 ± 0.58 cm⁻³ s⁻¹, which was comparable with the results at Mt. Yulong (J_3 , 1.18 cm⁻³ s⁻¹) (Shang et al., 2018) and Jungfrauoch ($J_{3,2}$, 0.2-7.5

cm³ s⁻¹) (Tröstl et al., 2016a). However, the J at Nam Co station was higher than the values at NCO-P site (J_{I0} , 0.14-0.19 cm³ s⁻¹) (Venzac et al., 2008) and Storm Peak Laboratory (J_8 , 0.39-1.19 cm³ s⁻¹) (Hallar et al., 2011), but lower than that at Mt. Tai (J_3 , 1.33-52.54 cm³ s⁻¹) (Lv et al., 2018) and Maïdo observatory (J_2 , 0.5-30 cm³ s⁻¹) (Rose et al., 2019), which may be due to the distinction of particle size range used in the calculation and precursor concentrations between the sites. For example, SO₂ at Mt. Tai (3.2 ppb) (Lv et al., 2018) was much higher than that at Mt. Yulong (0.06 ppb) (Shang et al., 2018). The average GR in the size range of 4-25 nm at Nam Co station varied from 1.5 to 5.6 nm h⁻¹, with the average value of 4.0 ± 1.2 nm h⁻¹, which was comparable with those at Mt. Yulong (3.2 nm h⁻¹) (Shang et al., 2018) and Jungfraujoch (4.0 nm h⁻¹) (Tröstl et al., 2016a), and higher than those at NCO-P site (1.8 ± 0.7 nm h⁻¹) (Venzac et al., 2008) and Mt. Daban (2.0 nm h⁻¹) (Du et al., 2015), but lower than that at Maïdo observatory (8-45 nm h⁻¹) (Rose et al., 2019). The average CS was $0.15 \times 10^{-2} \pm 0.07 \times 10^{-2} \text{ s}^{-1}$ at Nam Co station, which was comparable with those at Mt. Yulong ($0.2 \times 10^{-2} \text{ s}^{-1}$) (Shang et al., 2018), Maïdo observatory ($0.02 \times 10^{-2} - 2 \times 10^{-2} \text{ s}^{-1}$) (Rose et al., 2019) and Storm Peak Laboratory ($0.12 \times 10^{-2} \text{ s}^{-1}$) (Hallar et al., 2011), and much lower than that at Mt. Tai ($1.1 \times 10^{-2} - 4.8 \times 10^{-2} \text{ s}^{-1}$) (Lv et al., 2018). In a word, the J , GR, and CS at Nam Co station were within average levels in the high-altitude sites. Meanwhile, there was no significant variation in J , GR and daily CS between pre-monsoon and monsoon seasons, although the NPF frequencies of the two seasons were quite different. The key factors determining the occurrence of NPF events need to be further studied.

3.3 Frequent NPF events driven ~~by organic matter brought by summer monsoon~~

Whether an NPF event can occur is mainly related to 1) the CS, which mainly referred to the scavenging rate of precursors, clusters, and newly formed particles by background aerosols. High CS can lead to the continual reduction in newly formed particle number concentration, and inhibit the occurrence of NPF; 2) the gaseous precursors that can participate in nucleation and growth, including sulfuric acid (Kulmala et al., 2013), dimethylamine (Yao et al., 2018), ammonia (Xiao et al., 2015) and VOC (Tröstl et al., 2016b; Fang et al., 2020; Qiao et al., 2021). A sufficiently high concentration of low volatility vapors (precursors) can contribute to persistent nucleation and generating new atmospheric particles; 3) ~~meteorological factors including air mass origins and meteorological factors including~~ WD, RH, temperature, ~~the intensity of solar radiation~~, etc, which can influence the occurrence and intensity of NPF events by directly or indirectly affecting the source and sink parameters.

In the following, ~~those potential reasons will be examined. the discussion will focus on the key factors for the occurrence of NPF events described above.~~ For clarity, NPF events days occurring in pre-monsoon and monsoon seasons were named NPF-pre days and NPF-monsoon days, respectively. Non-events days in pre-monsoon season were termed

non-event days, as non-event days in monsoon season were scarce.

3.3.1 Condensation sink

CS is the key factor controlling the occurrence of NPF events especially in urban environment (Yan et al., 2021). At some high-altitude observations at a larger scale, the important role of the transported pre-existing particles in NPF events was also emphasized (Rose et al., 2019; Boulon et al., 2010). Here we analyzed the CS levels in NPF days and non-event days at Nam Co station. As shown in Fig. 34a, the levels of CS in NPF-pre days, NPF-monsoon days and non-event days were ~~approximatesimilar~~ during 11:00–18:00 (the occurrence time of NPF events), although the CS in the early morning of NPF-pre days seems to be slightly lower. The CS was mainly in the range of 0.1×10^{-2} – $0.15 \times 10^{-2} \text{ s}^{-1}$ during the NPF occurrence time, which was much lower than that at most locations in China, such as $\sim 0.01 \text{ s}^{-1}$ in urban Beijing (Deng et al., 2021), and 0.1×10^{-2} to $28.4 \times 10^{-2} \text{ s}^{-1}$ at Mt. Tai (Lv et al., 2018), and comparable with that at Mt. Yulong ($\sim 0.2 \times 10^{-2} \text{ s}^{-1}$) (Shang et al., 2018). The result varied from previous studies which reported much lower CS during NPF days than that in non-event days (Zhou et al., 2021; Lv et al., 2018). It indicated that ~~the occurrence of NPF events at Nam Co station was not influenced by the lower CS. The similar phenomenon was found at Mt. Yulong (Shang et al., 2018). In addition, previous studies found that CS was on average positively correlated with the occurrence of NPF events in some high-altitude observations at a larger scale, such as Maïdo observatory and Jungfraujoeh (Rose et al., 2019; Boulon et al., 2010). When the gaseous precursors were transported to the observation site to trigger NPF events, the pre-existing particles were transported simultaneously and caused the increase in CS. Although the topography and the rapidly changing conditions related to complex atmospheric dynamics at Nam Co station were similar with Maïdo observatory and Jungfraujoeh, the relationship between CS and the occurrence of NPF events seemed different. Whatever, the result indicated that~~ CS was not the decisive factor controlling the occurrence of NPF events at Nam Co station.

3.3.2 Gas precursors

Gaseous sulfuric acid is identified as the key precursor for nucleation and initial growth due to its low volatility (Kulmala et al., 2013; Qiao et al., 2021). ~~Nucleation based on sulfuric acid and stabilizing species, such as ammonia, amines, is regarded as the dominant NPF mechanism in urban and rural environment. However, the role of gaseous sulfuric acid in NPF events may not be significant at Nam Co station. The concentration of sulfuric acid in the atmosphere is related to the degree of SO₂, photochemical rate and CS. On the one hand, the level of SO₂ was considered to be comparable between NPF days and non-event days or maybe higher in non-event days. The SO₂ concentration was not measured~~

带格式的：下标

带格式的：下标

directly in the observation periods, and the WRF/CMAQ modelled SO₂ concentration was employed for analysis. It can be found that the modelled SO₂ concentration in the area was at a very low level, with the average value of 0.03±0.02 ppb (Fig. S13), which was comparable with that measured at Mt. Yulong (Shang et al., 2018). For NPF and non-event days, slightly higher modelled SO₂ concentration was observed in non-event days and NPF-pre days compared with that in NPF-monsoon days. It indicates that the level of SO₂ may be related to the season influence, and there is no significant difference between NPF days and non-event days. On the other hand, the photochemical oxidation intensity of SO₂ and CS were comparable between NPF days and non-event days. The photochemical oxidation intensity of SO₂ was indicated by J (O¹D). As shown in Fig. 34b, NPF-pre days, NPF-monsoon days and non-event days shared the same level of J (O¹D) ($3 \times 10^{-2} - 6 \times 10^{-2} \text{ s}^{-1}$) during the occurrence time of NPF events. The CS levels were close among those three types of days as introduced in Sec. 3.3.1. With speculated comparable/lower SO₂ concentration and similar CS and J (O¹D) level, the abundance of gaseous sulfuric acid in NPF days would be approaching, or little lower than that in non-event days. Whatever, it indicated that sulfuric acid may not govern the occurrence of NPF events at Nam Co station, and the photochemical formation of SO₂ was the major source for sulfuric acid in the atmosphere. However, the level of SO₂, the photochemical oxidation intensity of SO₂, and CS were comparable between NPF days and non-event days, which indicated that sulfuric acid may not govern the occurrence of NPF events at Nam Co station. Here, the low levels of SO₂ (0.03±0.03 ppb) in NPF days and non-event days were present in Fig. S8. The photochemical oxidation intensity of SO₂ was indicated by J (O¹D). As shown in Fig. 3b, NPF-pre days, NPF-monsoon days and non-event days shared the same level of J (O¹D) ($3 \times 10^{-2} - 6 \times 10^{-2} \text{ s}^{-1}$) during the occurrence time of NPF events.

In addition to sulfuric acid, organics are also considered to be an important factor of NPF events. Observations and laboratory experiments have found that organics may participate in or even dominate the nucleation and growth process in NPF events in pristine environments and the preindustrial atmosphere. For example, CLOUD (Cosmics Leaving Outdoor Droplets) experiments observed obvious NPF events from highly oxidized organics without the involvement of sulfuric acid (Kirkby et al., 2016). At the high-altitude sites of Jungfraujoch and Himalaya, NPF events occurred mainly through the condensation of highly oxygenated molecules (HOMs) (Bianchi et al., 2016; Bianchi et al., 2021). Due to instrument status, VOC measurement was only available in pre-monsoon season. The concentration of TVOC (total VOC) showed a noticeably higher value (20%) during 11:00-18:00 on NPF-pre days compared with non-event days (Fig. 34c). Aromatics, which can be used as the indicator of anthropogenic emissions, also exhibited a higher level (20%) during NPF-pre days (Fig. 3-4d). This suggested that VOC may be the key factor in determining the occurrence of NPF events. In order to further evaluate the role of VOC, we used WRF/CMAQ models to simulate the spatial distribution of VOC concentration in

带格式的：下标

带格式的：下标

带格式的：下标

带格式的：下标

带格式的：上标

带格式的：上标

带格式的：上标

带格式的：下标

pre-monsoon and monsoon seasons. ~~The detailed information about the model setting and evaluation can be found in Text S2 (Hu, 2017 #180), 62 (Hu et al., 2017)~~ As shown in Fig. 45, the simulated VOC levels in NPF days including NPF-pre and NPF-monsoon days were higher than those in non-event days. The average modelled VOC concentrations in NPF-pre days and NPF-monsoon days were 25% and 88% higher than those in non-event days, respectively. Therefore, we further considered that the organic matter could be the key factor to determine whether NPF event occurred. Nucleation of pure organics or organics involved could be the dominant mechanism at Nam Co station. In addition, higher organic concentrations were observed in monsoon season (NPF-monsoon days) compared with that in pre-monsoon season (NPF-pre days and non-event days). The result is consistent with one recent research which has found that the concentration of monoterpene-derived HOMs in East Asia was higher in summer (June-August) than that in Spring (March-May) by using GEOS-Chem global chemical transport model (Xu et al., 2022). This means that the frequent NPF events in monsoon season could be caused by the higher organic matters. ~~It can be found that the average VOC concentration of NPF-monsoon days was about 2 ppb higher than that of NPF-pre days, and about 4 ppb higher than that of non-event days at Nam Co station (Fig. 4). In addition, one recent research has found that the concentration of monoterpene derived HOMs in East Asia was higher in summer (June-August) than that in Spring (March-May) by using GEOS-Chem global chemical transport model (Xu et al., 2022). All the results indicated that the frequent NPF events in monsoon season were caused by the higher concentration of organic precursors. And pure organic nucleation may be the dominant NPF mechanism at Nam Co station.~~

3.3.3 ~~Air mass origins and m~~ Meteorology

While the concentrations of organic precursors ~~have explained~~ could be the most possible reason for ~~the notable distinction in NPF frequencies in pre-monsoon season and monsoon season~~ occurrence of NPF events, the external factors driving the ~~apparent~~ difference in VOC levels between the ~~NPF and non-event days two seasons~~ and other conditions that may affect the characteristics of NPF were still unknown. This indicated that a further investigation into other NPF-related variables was still required.

~~WD can reflect the local situation for air mass and the source of air pollutants. The wind rose plots of non event days, NPF pre days and NPF monsoon days were present in Fig. 5. It can be found that the dominant WD on non event days was westerly and the WS was mostly below 10 m/s. In comparison, WD on NPF pre days and NPF monsoon days mainly come from the South, mostly above 8 m/s. Similar patterns can be found from the time series (Fig. S2), that was, the WD from 3 May to 5 May (NPF pre days)~~

resembled that during summer monsoon, mainly from the South, and the overall WS was relatively high. The example of the spatial distribution of wind in non-event days, NPF-pre days and NPF-monsoon days showed the same phenomenon on a larger scale (Fig. S9). These results suggested that when the southerly wind with high WD occurred, it may bring organic precursors from the southern region (northeast India) to Nam Co station, driving the occurrence of NPF events in this area. This explained the phenomenon of the higher VOC concentration in NPF days than non-event days. The frequent NPF events in monsoon season resulted from the higher concentration of organic vapours brought by the frequent southerly wind (summer monsoon). This conclusion was confirmed by the more frequent air masses from the South during monsoon season (Fig. S6). Similar result was found in the recent study which showed that the Indian summer monsoon acted as a facilitator for transporting gaseous pollutants (Yin et al., 2021).

There are almost no local anthropogenic source emissions at Nam Co station. The air pollutants at the observation site mainly brought by air mass transmission. Backward trajectories were utilized to identify the air mass origins associated with NPF events. The frequencies of the 48 h back trajectories of air masses arriving at Nam Co station during the occurrence time of nucleation (11:00-18:00) in non-event days, NPF-pre days and NPF-monsoon days are present in Fig. 5. It can be found that the dominant air masses in non-event days were from the west (almost 100%) and passed by western Nepal, northwest India and Pakistan. In comparison, air masses in NPF-pre days and NPF-monsoon days mainly come from the south and southwest (57% and 75%) and originated in the northeast India. WD, which can reflect the local situation for air masses and the source of air pollutants, showed a high frequency of strong southerly wind in NPF-pre days and NPF-monsoon days and westerly wind in non-event days (Fig. S14). The example of the spatial distribution of wind field in non-event days, NPF-pre days and NPF-monsoon days displayed the similar phenomenon on a larger scale (Fig. S15). These results suggested that the occurrence of NPF events was related to the southerly and southwesterly air masses. When the southerly air mass occurred, it may bring organic precursors from the southern region (northeast India) to Nam Co station, driving the occurrence of NPF events in this area. As for the significantly higher NPF frequency in monsoon season, it resulted from the more frequent southerly air masses (summer monsoon) in monsoon season in comparison with pre-monsoon season as introduced in Sec. 3.1. The summer monsoon can bring the higher organic concentrations in monsoon season (NPF-monsoon days) compared with those in pre-monsoon season (NPF-pre and non-event days) (Fig.4), thus triggered almost daily NPF events. Similar result was found in the recent study which showed that the Indian summer monsoon acted as a facilitator for transporting gaseous pollutants (Yin et al., 2021).

(Yin et al., 2021) In Fig. 6, we show the diurnal variations of meteorological factors during NPF-pre days,

- 带格式的: 字体颜色: 自动设置
- 带格式的: 字体颜色: 自动设置
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号

420 NPF-monsoon days and non-event days at Nam Co station. ~~First, the~~ temperature in NPF-pre days and non-event days were generally close, but the temperature of NPF-monsoon days was obviously higher. The similar temperature in NPF-pre days and non-event days suggests that temperature is not a crucial factor for NPF event occurrence. Previous studies have found that higher temperature can increase the formation rate of HOMs, but reduce the volatility of HOMs in the meanwhile. The effect of temperature on the nucleation rate may not be important in a limited ambient temperature range (Stolzenburg et al., 2018). But the higher temperature during NPF-monsoon days may promote biogenic VOC monoterpene-emissions, which favored particle nucleation and growth (Andreae et al., 2022). ~~This~~ This may partly explain the phenomenon for the higher modelled VOC concentration in NPF-pre-monsoon days than NPF-premonsoon days. ~~Second, WS shared the same level during 11:00-18:00 on NPF days and non-event days, while WS before the occurrence of NPF events in NPF days was higher than that of non-event days.~~ ~~Third, the~~ The average RH was similar in both NPF days and non-event days, although 430 NPF-pre days had a wide range of RH changes. Laboratory studies have shown that water vapour can suppress the formation of extremely low volatility organic compounds (ELVOC) from monoterpenes (Bonn et al., 2002). However, our result was the opposite. During 11:00-18:00, the water content (H₂O) in the air during NPF-monsoon days was the highest, NPF-pre days was the second, and non-event days was the lowest. This indicated the inhibition effect of water vapour was not important at Nam Co station. The similar finding was also observed at remote sites in the subboreal forest of North America (Andreae et al., 2022).

435 Overall, our results were most consistent with the organics involved nucleation mechanism. Condensation sink and gaseous sulfuric acid could have no significant effect on the occurrence of NPF events. The frequent NPF events in summer monsoon season may result from the higher frequency of southerly and southwesterly air masses, which brought the organic precursors to participate in the NPF process.

440 ~~In all, the level of organic vapours governed the occurrence of NPF events at Nam Co station, while condensation sink and gaseous sulfuric acid had no effect. The frequent NPF events in summer monsoon season resulted from the higher concentration of organic vapours, which was brought from northeast India by the strong southerly monsoon.~~

3.4 Significant increase of CCN in monsoon season

445 The CCN concentration was estimated following the method introduced in Sect. 2.4. Considering the similar proportions of chemical components between Nam Co station and Mt. Yulong with around 70% fraction of organics, and the stability of the fractions of chemical components, ~~The~~ the hygroscopicity parameter κ was assumed to be a constant

value of 0.12 throughout the measurement period, according to the previous measurement [at Mt. Yulong in the TP of chemical composition in TP](#) (Shang et al., 2018). As a result, the D_c at S_c levels of 0.6% and 1.2% were 73.4 ± 1.3 and 46.3 ± 0.8 nm, respectively. No noticeable difference in D_c was found between pre-monsoon and monsoon seasons ($S_c = 0.6\%$: 74.4 ± 1.0 vs 72.4 ± 1.1 nm, $S_c = 1.2\%$: 46.9 ± 0.6 vs 45.7 ± 0.7 nm). There could be uncertainties in the values of κ and D_c , but they had little impact on the final result of CCN concentration (Shang et al., 2018).

The high-frequency NPF events in the summer monsoon period markedly increased the number concentration of atmospheric aerosols. The average PNSD during pre-monsoon and monsoon seasons are plotted in Fig. 7a with much higher number concentrations observed during monsoon season. The mean total particle number concentration (PN_{4-700}) in monsoon season was 3647 ± 2671 cm^{-3} , which was more than 2 times higher than that of pre-monsoon season (1163 ± 1026 cm^{-3}). Although the measured particle size ranges were not the same with other studies, the results can still be comparable as the background particles were mainly distributed in tens to hundreds of nanometers. As shown in Table 1, PN_{4-700} at Nam Co station in pre-monsoon season was comparable with other high-altitude sites around the world, while PN_{4-700} in monsoon season was much higher due to the frequent NPF events. The atmospheric particles contributed by new particle nucleation and growth in monsoon season were mainly concentrated below 100 nm. Among them, the concentration of nucleation mode particles (4-25 nm) in monsoon season was about 2 times higher than that in pre-monsoon season, and the concentration of Aitken mode particles (25-100 nm) was 3.5 times higher than that in pre-monsoon season. In contrast, accumulation-mode particles (>100 nm) which were related to the secondary formation process (Wang et al., 2013), were nearly at the same level in the two seasons (around 200 cm^{-3}). As for CCN, the average number concentrations of CCN at S_c of 0.6% and 1.2% in monsoon season were 434 ± 242 and 863 ± 628 cm^{-3} , respectively (Fig. 8). The results were about 0.1 and 0.6 times higher than those in pre-monsoon season at S_c of 0.6% (396 ± 177 cm^{-3}) and 1.2% (552 ± 261 cm^{-3}).

In addition to the average particle number concentration in the two seasons, the important impact of NPF events is more reflected in the increased number concentration of aerosol and CCN in a short time, that is, the aerosol and CCN production. The aerosol production and CCN production during one day can be obtained by comparing the particle number concentration at the beginning of the increase of the target particles (N_{init}) with the maximum number concentration (N_{max}), as introduced by Rose et al. (2017). The N_{init} and N_{max} are hourly average concentrations as shown in Fig. S16. Fig. 9 showed the average daily variation of PNSD, three modes particles and CCN in the whole periods in pre-monsoon season and monsoon season. Higher daily aerosol production and CCN production were observed during monsoon season compared with those in pre-monsoon season (Fig. 7). In monsoon season, nucleation-mode particles (4-25 nm) started to increase quickly at around 11:00 when nucleation occurred. The freshly nucleated particles grew to larger sizes due to

condensation and coagulation of the pre-existing particles within several hours, which contributed to the increase of Aitken-mode particles (25-100 nm) from 15:00. The average daily aerosol production of nNucleation-mode particles and Aitken-mode particles in monsoon season was around 3400 cm⁻³ and 1200 cm⁻³, respectively, and the enhancement factors (EFs, $N_{\text{max}}/N_{\text{min}}$) were 9.2 and 3.6, respectively. As for CCN, the average production of CCN at S_c of 0.6% and 1.2% in monsoon season was 180 and 518 cm⁻³, respectively, and the EFs were 1.7 and 2.2, respectively. The production EFs of CCN were lower than previous studies because they only considered NPF days and their environment was relatively polluted (Rose et al., 2017; Shen et al., 2016). Although the particles and CCN at around 11:00 in pre-monsoon season were comparable with monsoon season, the production of aerosols and CCN was much lower in pre-monsoon season, although the particles and CCN at around 11:00 were comparable with monsoon season. The average daily aerosol production of nNucleation-mode particles and Aitken-mode particles in pre-monsoon season were around 500 cm⁻³ and 300 cm⁻³, respectively. And the CCN production at S_c of 0.6% and 1.2% was 160 and 286 cm⁻³, respectively. The average daily production of nNucleation-mode particles, Aitken-mode particles and CCN at S_c of 1.2% in monsoon season was 5.8, 3 and 0.8 times higher than that in pre-monsoon season, respectively.

The high frequency NPF events in the summer monsoon period markedly increased the number concentration of atmospheric aerosols. The average PNSD during pre-monsoon and monsoon seasons were plotted in Fig. 8a with much higher number concentrations occurring during monsoon season. The mean total particle number concentration (PN_{4-700}) in monsoon season was 3647 ± 2671 cm⁻³, which was more than 2 times higher than that of pre-monsoon season (1163 ± 1026 cm⁻³). Although the measured particle size ranges were not the same with other studies, the results can still be comparable as the background particles were mainly distributed in tens to hundreds of nanometers. As shown in Table 1, PN_{4-700} at Nam Co station in pre-monsoon season was comparable with other high-altitude sites around the world, while PN_{4-700} in monsoon season was much higher due to the frequent NPF events. The atmospheric particles contributed by new particle nucleation and growth in monsoon season were mainly concentrated below 100 nm. Among them, the concentration of Nucleation-mode particles in monsoon season was about 2 times higher than that in pre-monsoon season, and the concentration of Aitken-mode particles was 3.5 times higher than that in pre-monsoon season. In contrast, Accumulation-mode particles (>100 nm) which were related to the secondary formation process and long-range transport (Wang et al., 2013; Vu et al., 2015), were nearly at the same level in the two seasons (around 200 cm⁻³). As shown in Fig. 9, the average number concentrations of CCN at S_c of 0.6% and 1.2% in monsoon season were 434 ± 242 and 863 ± 628 cm⁻³, respectively. The results were 10% and 56% higher than those in pre-monsoon season at S_c of 0.6% (396 ± 177 cm⁻³) and 1.2% (552 ± 261 cm⁻³). It should be noted that the CCN concentration and production we estimated above were local ground

510 levels, which can be considered as the minimum values. That is because those smaller particles (<40 nm) formed by NPF may further grow and reach CCN size in the following days during transportation, as the coagulation sink affecting the lifetime of aerosols was at a low level. This mechanism was verified at different high-altitude stations such as NCO-P site, Chacaltaya and Jungfraujoch (Venzac et al., 2008; Rose et al., 2015; Tröstl et al., 2016a). The Nucleation-mode and Aitken-mode particle bands in the morning reflected the continuation of NPF events. As a result, the amount of potential CCN in monsoon season could be much larger than the result measured here, due to the high level of the smaller particles. Therefore, the climate effect from aerosols may have a large variance between pre-monsoon and monsoon season.

4 Summary and conclusions

515 The PNSD, trace gases and meteorological parameters were measured at Nam Co station (43747309 m a.s.l) in the central Tibetan Plateau (TP) during pre-monsoon and summer monsoon seasons. Firstly, meteorological conditions between pre-monsoon and monsoon seasons were distinct with higher temperature and water content in monsoon season. Strong westerlies pass through western Nepal, northwest India and Pakistan in pre-monsoon season, while air masses were mainly derived from the South (Bangladesh and northeast India) in the monsoon period. Secondly, as the pristine high-altitude site, low levels of particles and gaseous pollutants were displayed at Nam Co station. Relatively higher pollutant concentration was found during pre-monsoon season due to less favorable atmospheric circulation.

520 The most important finding of this study was that there was an evident distinction in the frequencies of NPF events at Nam Co station with 15% in pre-monsoon season and 80% in monsoon season, which can represent the lowest and highest values of NPF frequencies observed in the TP and other high-altitude sites, even around the world. In addition, there was no noticeable variation in *J*, GR, and CS between pre-monsoon and monsoon seasons at Nam Co station. ~~—were favorable for NPF~~ Although the measured precursors were limited at our sites, the comprehensive analysis of the measured CS, gaseous precursors and meteorological conditions, supplemented by the modelled SO₂ and VOC, points to the organic involved nucleation as the dominant mechanism. Condensation sink and gaseous sulfuric acid could have no significant effect on the occurrence of NPF events. The frequent NPF events in summer monsoon season may result from the higher frequency of southerly and southwesterly air masses, which brought the organic precursors to participate in the NPF process. ~~Through the comprehensive analysis of the measured CS, gaseous precursors and meteorological conditions, supplemented by the model simulations, we found that the level of organic vapours governed the occurrence of NPF events, while condensation sink and gaseous sulfuric acid had no effect. The frequent NPF events in monsoon season resulted from the higher concentration of organic vapours, which was brought from northeast India by the strong southerly monsoon.~~

带格式的：下标

The frequent NPF events in the summer monsoon season had elevated the number concentration of atmospheric aerosols and CCN at Nam Co station. The average total particle number concentration in monsoon season was more than 2 times higher than that in pre-monsoon season, mainly contributed by nucleation mode and Aitken mode particles. And the average number concentrations of CCN at S_c of 0.6% and 1.2% in monsoon season were 0.1 and 0.6 times higher than those in pre-monsoon season. As for aerosol and CCN production, the average daily production of nucleation mode particles, Aitken mode particles and CCN at S_c of 1.2% in monsoon season was 5.8, 3 and 0.8 times higher than that in pre-monsoon season, respectively. The average daily production of Nucleation-mode particles, Aitken-mode particles and CCN at S_c of 1.2% in monsoon season was 5.8, 3 and 0.8 times higher than that in pre-monsoon season, respectively. The average total particle number concentration in monsoon season was more than 2 times higher than that in pre-monsoon season, mainly contributed by Nucleation-mode and Aitken-mode particles. And the average number concentration of CCN in monsoon season was 10%-56% higher than that in pre-monsoon season. Considering that the smaller particles (< 40 nm) formed by NPF may further grow and reach CCN size during the following days due to the low-level coagulation sink, the amount of potential CCN in monsoon season could be much larger than our local measurement results. It may markedly affect the earth's radiation balance and global climate. Our results emphasized that seasonal effect of NPF should be considered in model simulations when calculating the amounts of aerosols and CCN in high-altitude atmosphere. Long-term measurement campaigns with a more comprehensive set of instrumentation to investigate the role of NPF in high-altitude atmosphere are required for a deeper scientific understanding of NPF process and its role in the global aerosol budget. Our result emphasized the role of organic in the occurrence of NPF events in the TP. Seasonal effect of NPF should be considered in model simulations when calculating the amount of aerosols and CCN in high-altitude atmosphere.

Data availability. The data provided in this paper can be obtained from the author upon request (minhu@pku.edu.cn).

Supplement. An independent supplement document is available.

Authorship contributions. Lizi Tang: Investigation, Data curation, Methodology, Formal analysis, Writing - original draft, Writing - review & editing. Min Hu: Project administration, Supervision, Funding acquisition, Writing - review & editing. Dongjie Shang: Investigation, Data curation, Methodology, Formal analysis. Xin Fang: Investigation, Data curation, Methodology, Formal analysis. Jianjiong Mao: Investigation, Data curation. Wanyun Xu: Data curation. Jiacheng Zhou: Data curation. Weixiong Zhao: Data curation. Yaru Wang: Data curation. Chong Zhang: Data curation. Yingjie Zhang: Data curation. Jianlin Hu: Data curation. Limin Zeng: Data curation. Chunxiang Ye: Project administration, Funding acquisition,

Data curation. Song Guo: Writing - review & editing. Zhijun Wu: Writing - review & editing.

565 *Competing interests.* The authors declare that they have no conflict of interest.

570 *Acknowledgements.* The research has been supported by the National Natural Science Foundation of China (91844301 and 91544214), National Research Program for Key Issues in Air Pollution Control (DQGG0103), National Key Research and Development Program of China (No. 2016YFC0202000: Task 3), and the second Tibetan Plateau Scientific Expedition and Research Program (STEP, 2019QZKK0606).

575

580

585

590 **References**

- 595 [Andreae, M. O., Andreae, T. W., Ditas, F., and Pöhlker, C.: Frequent new particle formation at remote sites in the subboreal forest of North America, *Atmos. Chem. Phys.*, 22, 2487-2505, 10.5194/acp-22-2487-2022, 2022.](#)
- Appel, K. W., Bash, J. O., Fahey, K. M., Foley, K. M., Gilliam, R. C., Hogrefe, C., Hutzell, W. T., Kang, D., Mathur, R., Murphy, B. N., Napelenok, S. L., Nolte, C. G., Pleim, J. E., Pouliot, G. A., Pye, H. O. T., Ran, L., Roselle, S. J., Sarwar, G., Schwede, D. B., Sidi, F. I., Spero, T. L., and Wong, D. C.: The Community Multiscale Air Quality (CMAQ) model versions 5.3 and 5.3.1: system updates and evaluation, *Geosci. Model Dev.*, 14, 2867-2897, 10.5194/gmd-14-2867-2021, 2021.
- 600 Bianchi, F., Junninen, H., Bigi, A., Sinclair, V. A., Dada, L., Hoyle, C. R., Zha, Q., Yao, L., Ahonen, L. R., Bonasoni, P., Buenrostro Mazon, S., Hutterli, M., Laj, P., Lehtipalo, K., Kangasluoma, J., Kerminen, V. M., Kontkanen, J., Marinoni, A., Mirme, S., Molteni, U., Petäjä, T., Riva, M., Rose, C., Sellegri, K., Yan, C., Worsnop, D. R., Kulmala, M., Baltensperger, U., and Dommen, J.: Biogenic particles formed in the Himalaya as an important source of free tropospheric aerosols, *Nature Geoscience*, 14, 4-9, 10.1038/s41561-020-00661-5, 2021.
- Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H. E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J., and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing, *Science*, 352, 1109-1112, doi:10.1126/science.aad5456, 2016.
- 605 Bonasoni, P., Laj, P., Marinoni, A., Sprenger, M., Angelini, F., Arduini, J., Bonafè, U., Calzolari, F., Colombo, T., Decesari, S., Di Biagio, C., Di Sarra, A., Evangelisti, F., Duchi, R., Facchini, M. C., Fuzzi, S., Gobbi, G. P., Maione, M., Panday, A., Roccatto, F., Sellegri, K., Venzac, H., Verza, G., Villani, P., Vuillermoz, E., and Cristofanelli, P.: Atmospheric Brown Clouds in the Himalayas: first two years of continuous observations at the Nepal Climate Observatory-Pyramid (5079 m), *Atmospheric Chemistry and Physics*, 10, 7515-7531, 10.5194/acp-10-7515-2010, 2010.
- Bonn, B., Schuster, G., and Moortgat, G. K.: Influence of Water Vapor on the Process of New Particle Formation during Monoterpene Ozonolysis, *The Journal of Physical Chemistry A*, 106, 2869-2881, 10.1021/jp012713p, 2002.
- 610 Boulon, J., Sellegri, K., Venzac, H., Picard, D., Weingartner, E., Wehrle, G., Collaud Coen, M., Bütikofer, R., Flückiger, E., Baltensperger, U., and Laj, P.: New particle formation and ultrafine charged aerosol climatology at a high altitude site in the Alps (Jungfrauoch, 3580 m a.s.l., Switzerland), *Atmos. Chem. Phys.*, 10, 9333-9349, 10.5194/acp-10-9333-2010, 2010.
- Cai, R. and Jiang, J.: A new balance formula to estimate new particle formation rate: reevaluating the effect of coagulation scavenging, *Atmos. Chem. Phys.*, 17, 12659-12675, 10.5194/acp-17-12659-2017, 2017.
- Chen, S., Wang, H., Lu, K., Zeng, L., Hu, M., and Zhang, Y.: The trend of surface ozone in Beijing from 2013 to 2019: Indications of the persisting strong atmospheric oxidation capacity, *Atmospheric Environment*, 242, 117801, <https://doi.org/10.1016/j.atmosenv.2020.117801>, 2020.
- 615 Cong, Z., Kang, S., Kawamura, K., Liu, B., Wan, X., Wang, Z., Gao, S., and Fu, P.: Carbonaceous aerosols on the south edge of the Tibetan Plateau: concentrations, seasonality and sources, *Atmos. Chem. Phys.*, 15, 1573-1584, 10.5194/acp-15-1573-2015, 2015.
- 620 Dal Maso, M., Kulmala, M., Riipinen, I., and Wagner, R.: Formation and growth of fresh atmospheric aerosols: Eight years of aerosol size distribution data from SMEAR II, Hyttiälä, Finland, *Boreal Environment Research*, 10, 323-336, 2005.
- Deng, C., Cai, R., Yan, C., Zheng, J., and Jiang, J.: Formation and growth of sub-3 nm particles in megacities: impact of background aerosols, *Faraday Discussions*, 226, 348-363, 10.1039/D0FD00083C, 2021.
- 625 Du, W., Sun, Y. L., Xu, Y. S., Jiang, Q., Wang, Q. Q., Yang, W., Wang, F., Bai, Z. P., Zhao, X. D., and Yang, Y. C.: Chemical characterization of submicron aerosol and particle growth events at a national background site (3295 m a.s.l.) on the Tibetan Plateau, *Atmos. Chem. Phys.*, 15, 10811-10824, 10.5194/acp-15-10811-2015, 2015.

带格式的: 字体: 五号

带格式的: 字体: 五号

带格式的: 字体: 五号

Fang, X., Hu, M., Shang, D., Tang, R., Shi, L., Olenius, T., Wang, Y., Wang, H., Zhang, Z., Chen, S., Yu, X., Zhu, W., Lou, S., Ma, Y., Li, X., Zeng, L., Wu, Z., Zheng, J., and Guo, S.: Observational Evidence for the Involvement of Dicarboxylic Acids in Particle Nucleation, *Environmental Science & Technology Letters*, 7, 388-394, 10.1021/acs.estlett.0c00270, 2020.

Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozhkov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R., Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *Journal of Geophysical Research: Atmospheres*, 122, 8739-8760, <https://doi.org/10.1002/2017JD026844>, 2017.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.

Hallar, A. G., Lowenthal, D. H., Chirokova, G., Borys, R. D., and Wiedinmyer, C.: Persistent daily new particle formation at a mountain-top location, *Atmospheric Environment*, 45, 4111-4115, <https://doi.org/10.1016/j.atmosenv.2011.04.044>, 2011.

Hu, J., Wang, P., Ying, Q., Zhang, H., Chen, J., Ge, X., Li, X., Jiang, J., Wang, S., Zhang, J., Zhao, Y., and Zhang, Y.: Modeling biogenic and anthropogenic secondary organic aerosol in China, *Atmos. Chem. Phys.*, 17, 77-92, 10.5194/acp-17-77-2017, 2017.

IPCC, 2021: *Climate Change 2021: The Physical Science Basis*. (eds [Masson-Delmotte, V. et al.]) (Cambridge Univ. Press, 2021).

Kerminen, V.-M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi, F.: Atmospheric new particle formation and growth: review of field observations, *Environmental Research Letters*, 13, 103003, 10.1088/1748-9326/aadf3c, 2018.

Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *Journal of Geophysical Research (Atmospheres)*, 109, D21208, 10.1029/2004jd004999, 2004.

Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521-526, 10.1038/nature17953, 2016.

Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P., Paasonen, P., Mikkilä, J., and Worsnop, D.: Direct Observations of Atmospheric Aerosol Nucleation, *Science (New York, N.Y.)*, 339, 943-946, 10.1126/science.1227385, 2013.

Lv, G., Sui, X., Chen, J., Jayaratne, R., and Mellouki, A.: Investigation of new particle formation at the summit of Mt. Tai, China, *Atmos. Chem. Phys.*, 18, 2243-2258, 10.5194/acp-18-2243-2018, 2018.

Mao, J., Li, L., Li, J., Sulaymon, I. D., Xiong, K., Wang, K., Zhu, J., Chen, G., Ye, F., Zhang, N., Qin, Y., Qin, M., and Hu, J.: Evaluation of Long-Term Modeling Fine Particulate Matter and Ozone in China During 2013–2019, *Frontiers in Environmental Science*, 10, 10.3389/fenvs.2022.872249, 2022.

Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN,

带格式的: 字体: 五号

带格式的: 字体: 五号

带格式的: 字体: 五号

带格式的: 字体: 五号

675 Atmos. Chem. Phys., 9, 8601-8616, 10.5194/acp-9-8601-2009, 2009.

Nieminen, T., Kerminen, V. M., Petäjä, T., Aalto, P. P., Arshinov, M., Asmi, E., Baltensperger, U., Beddows, D. C. S., Beukes, J. P., Collins, D., Ding, A., Harrison, R. M., Henzing, B., Hooda, R., Hu, M., Hörrak, U., Kivekäs, N., Komsaare, K., Krejci, R., Kristensson, A., Laakso, L., Laaksonen, A., Leitch, W. R., Lihavainen, H., Mihalopoulos, N., Németh, Z., Nie, W., O'Dowd, C., Salma, I., Sellegri, K., Svenningsson, B., Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler, A., Wu, Z., Virtanen, A., and Kulmala, M.: Global analysis of continental boundary layer new particle formation based on long-term measurements, *Atmos. Chem. Phys.*, 18, 14737-14756, 10.5194/acp-18-14737-2018, 2018.

Peters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, 10.5194/acp-7-1961-2007, 2007.

685 Qiao, X., Yan, C., Li, X., Guo, Y., Yin, R., Deng, C., Li, C., Nie, W., Wang, M., Cai, R., Huang, D., Wang, Z., Yao, L., Worsnop, D. R., Bianchi, F., Liu, Y., Donahue, N. M., Kulmala, M., and Jiang, J.: Contribution of Atmospheric Oxygenated Organic Compounds to Particle Growth in an Urban Environment, *Environmental Science & Technology*, 55, 13646-13656, 10.1021/acs.est.1c02095, 2021.

Rose, C., Foucart, B., Picard, D., Colomb, A., Metzger, J. M., Tulet, P., and Sellegri, K.: New particle formation in the volcanic eruption plume of the Piton de la Fournaise: specific features from a long-term dataset, *Atmos. Chem. Phys.*, 19, 13243-13265, 10.5194/acp-19-13243-2019, 2019.

690 Rose, C., Sellegri, K., Moreno, I., Velarde, F., Ramonet, M., Weinhold, K., Krejci, R., Andrade, M., Wiedensohler, A., Ginot, P., and Laj, P.: CCN production by new particle formation in the free troposphere, *Atmos. Chem. Phys.*, 17, 1529-1541, 10.5194/acp-17-1529-2017, 2017.

Rose, C., Sellegri, K., Velarde, F., Moreno, I., Ramonet, M., Weinhold, K., Krejci, R., Ginot, P., Andrade, M., Wiedensohler, A., and Laj, P.: Frequent nucleation events at the high altitude station of Chacaltaya (5240 m a.s.l.), Bolivia, *Atmospheric Environment*, 102, 18-29, <https://doi.org/10.1016/j.atmosenv.2014.11.015>, 2015.

695 Shang, D., Hu, M., Zheng, J., Qin, Y., Du, Z., Li, M., Fang, J., Peng, J., Wu, Y., Lu, S., and Guo, S.: Particle number size distribution and new particle formation under the influence of biomass burning at a high altitude background site at Mt. Yulong (3410&thinspm), China, *Atmos. Chem. Phys.*, 18, 15687-15703, 10.5194/acp-18-15687-2018, 2018.

700 Shen, X., Sun, J., Zhang, X., Zhang, Y., Zhang, L., and Fan, R.: Key features of new particle formation events at background sites in China and their influence on cloud condensation nuclei, *Frontiers of Environmental Science & Engineering*, 10, 5, 10.1007/s11783-016-0833-2, 2016.

Stolzenburg, D., Fischer, L., Vogel, A. L., Heinritzi, M., Schervish, M., Simon, M., Wagner, A. C., Dada, L., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bianchi, F., Breitenlechner, M., Brilke, S., Buenrostro Mazon, S., Chen, D., Dias, A., Draper, D. C., Duplissy, J., El Haddad, I., Finkenzeller, H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., He, X., Helm, J., Hofbauer, V., Hoyle, C. R., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lampilahti, J., Lawler, M., Lehtipalo, K., Leiminger, M., Mai, H., Mathot, S., Mentler, B., Molteni, U., Nie, W., Nieminen, T., Nowak, J. B., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Quéléver, L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S., Tauber, C., Tomé, A., Wagner, R., Wang, M., Weitz, L., Wimmer, D., Xiao, M., Yan, C., Ye, P., Zha, Q., Baltensperger, U., Curtius, J., Dommen, J., Flagan, R. C., Kulmala, M., Smith, J. N., Worsnop, D. R., Hansel, A., Donahue, N. M., and Winkler, P. M.: Rapid growth of organic aerosol nanoparticles over a wide tropospheric temperature range, *Proceedings of the National Academy of Sciences of the United States of America*, 115, 9122-9127, 10.1073/pnas.1807604115, 2018.

705 Tröstl, J., Herrmann, E., Frege, C., Bianchi, F., Molteni, U., Bukowiecki, N., Hoyle, C. R., Steinbacher, M., Weingartner, E., Dommen, J., Gysel, M., and Baltensperger, U.: Contribution of new particle formation to the total aerosol concentration at the high-altitude site Jungfrauoch (3580 m asl, Switzerland), *Journal of Geophysical Research: Atmospheres*, 121, 11,692-611,711, <https://doi.org/10.1002/2015JD024637>, 2016a.

710 Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R.,

带格式的: 字体: 五号

带格式的: 字体: 五号

带格式的: 字体: 五号

带格式的: 字体: 五号

Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Möhler, O., Nieminen, T., Onnela, A., Petäjä, T., Piel, F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipilä, M., Smith, J. N., Steiner, G., Tomè, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop, D. R., Donahue, N. M., and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533, 527-531, 10.1038/nature18271, 2016b.

Venzac, H., Sellegri, K., Laj, P., Villani, P., Bonasoni, P., Marinoni, A., Cristofanelli, P., Calzolari, F., Fuzzi, S., Decesari, S., Facchini, M.-C., Vuillermoz, E., and Verza, G. P.: High frequency new particle formation in the Himalayas, *Proceedings of the National Academy of Sciences*, 105, 15666-15671, doi:10.1073/pnas.0801355105, 2008.

Wang, M., Xu, B., Wang, N., Cao, J., Tie, X., Wang, H., Zhu, C., and Yang, W.: Two distinct patterns of seasonal variation of airborne black carbon over Tibetan Plateau, *Science of The Total Environment*, 573, 1041-1052, <https://doi.org/10.1016/j.scitotenv.2016.08.184>, 2016.

Wang, M., Zeng, L., Lu, S., Shao, M., Liu, X., Yu, X., Chen, W., Yuan, B., Zhang, Q., Hu, M., and Zhang, Z.: Development and validation of a cryogen-free automatic gas chromatograph system (GC-MS/FID) for online measurements of volatile organic compounds, *Analytical Methods*, 6, 9424-9434, 10.1039/C4AY01855A, 2014.

Wang, Z. B., Hu, M., Wu, Z. J., Yue, D. L., He, L. Y., Huang, X. F., Liu, X. G., and Wiedensohler, A.: Long-term measurements of particle number size distributions and the relationships with air mass history and source apportionment in the summer of Beijing, *Atmos. Chem. Phys.*, 13, 10159-10170, 10.5194/acp-13-10159-2013, 2013.

Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Grünig, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H. G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Löschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term observations of atmospheric particle number size distributions, *Atmos. Meas. Tech.*, 5, 657-685, 10.5194/amt-5-657-2012, 2012.

Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625-641, 10.5194/gmd-4-625-2011, 2011.

Xiao, S., Wang, M. Y., Yao, L., Kulmala, M., Zhou, B., Yang, X., Chen, J. M., Wang, D. F., Fu, Q. Y., Worsnop, D. R., and Wang, L.: Strong atmospheric new particle formation in winter in urban Shanghai, China, *Atmos. Chem. Phys.*, 15, 1769-1781, 10.5194/acp-15-1769-2015, 2015.

Xu, J., Hettiyadura, A. P. S., Liu, Y., Zhang, X., Kang, S., and Laskin, A.: Regional Differences of Chemical Composition and Optical Properties of Aerosols in the Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031226, <https://doi.org/10.1029/2019JD031226>, 2020.

Xu, J., Zhang, Q., Shi, J., Ge, X., Xie, C., Wang, J., Kang, S., Zhang, R., and Wang, Y.: Chemical characteristics of submicron particles at the central Tibetan Plateau: insights from aerosol mass spectrometry, *Atmos. Chem. Phys.*, 18, 427-443, 10.5194/acp-18-427-2018, 2018a.

Xu, R., Tie, X., Li, G., Zhao, S., Cao, J., Feng, T., and Long, X.: Effect of biomass burning on black carbon (BC) in South Asia and Tibetan Plateau: The analysis of WRF-Chem modeling, *Science of The Total Environment*, 645, 901-912, <https://doi.org/10.1016/j.scitotenv.2018.07.165>, 2018b.

带格式的: 字体: 五号

765 Xu, R., Thornton, J. A., Lee, B. H., Zhang, Y., Jaeglé, L., Lopez-Hilfiker, F. D., Rantala, P., and Petäjä, T.: Global simulations of monoterpene-derived peroxy radical fates and the distributions of highly oxygenated organic molecules (HOMs) and accretion products, *Atmos. Chem. Phys.*, 22, 5477-5494, 10.5194/acp-22-5477-2022, 2022.

770 Yan, C., Yin, R., Lu, Y., Dada, L., Yang, D., Fu, Y., Kontkanen, J., Deng, C., Garmash, O., Ruan, J., Baalbaki, R., Schervish, M., Cai, R., Bloss, M., Chan, T., Chen, T., Chen, Q., Chen, X., Chen, Y., Chu, B., Dällenbach, K., Foreback, B., He, X., Heikkinen, L., Jokinen, T., Junninen, H., Kangasluoma, J., Kokkonen, T., Kurppa, M., Lehtipalo, K., Li, H., Li, H., Li, X., Liu, Y., Ma, Q., Paasonen, P., Rantala, P., Pileci, R. E., Rusanen, A., Samela, N., Simonen, P., Wang, S., Wang, W., Wang, Y., Xue, M., Yang, G., Yao, L., Zhou, Y., Kujansuu, J., Petäjä, T., Nie, W., Ma, Y., Ge, M., He, H., Donahue, N. M., Worsnop, D. R., Kerminen, V.-M., Wang, L., Liu, Y., Zheng, J., Kulmala, M., Jiang, J., and Bianchi, F.: The Synergistic Role of Sulfuric Acid, Bases, and Oxidized Organics Governing New-Particle Formation in Beijing, *Geophysical Research Letters*, 48, e2020GL091944, <https://doi.org/10.1029/2020GL091944>, 2021.

Yanai, M. and Wu, G.-X.: Effects of the Tibetan Plateau, in: *The Asian Monsoon*, edited by: Wang, B., Springer Berlin Heidelberg, Berlin, Heidelberg, 513-549, 10.1007/3-540-37722-0_13, 2006.

775 Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S., Ehn, M., Paasonen, P., Sipilä, M., Wang, M., Wang, X., Xiao, S., Chen, H., Lu, Y., Zhang, B., Wang, D., Fu, Q., and Wang, L.: Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278-281, 10.1126/science.aao4839, 2018.

Yin, X., Kang, S., Foy, B. d., Rupakheti, D., Rupakheti, M., Cong, Z., Wan, X., Zhang, G., and Zhang, Q.: Impacts of Indian summer monsoon and stratospheric intrusion on air pollutants in the inland Tibetan Plateau, *Geoscience Frontiers*, 12, 101255, <https://doi.org/10.1016/j.gsf.2021.101255>, 2021.

780 Zhang, X., Xu, J., Kang, S., Sun, J., Shi, J., Gong, C., Sun, X., Du, H., Ge, X., and Zhang, Q.: Regional Differences in the Light Absorption Properties of Fine Particulate Matter Over the Tibetan Plateau: Insights From HR-ToF-AMS and Aethalometer Measurements, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035562, <https://doi.org/10.1029/2021JD035562>, 2021.

785 Zhou, Y., Hakala, S., Yan, C., Gao, Y., Yao, X., Chu, B., Chan, T., Kangasluoma, J., Gani, S., Kontkanen, J., Paasonen, P., Liu, Y., Petäjä, T., Kulmala, M., and Dada, L.: Measurement report: New particle formation characteristics at an urban and a mountain station in northern China, *Atmos. Chem. Phys.*, 21, 17885-17906, 10.5194/acp-21-17885-2021, 2021.

带格式的: 字体: 五号




Figure 1. Location map for (a) the Tibetan Plateau, (b) Nam Co station, colored according to altitude. (c) The appearance drawing of atmospheric environment observatory field of Nam Co station (Yin et al., 2020).

800

805








Figure 2. Time series during pre-monsoon and monsoon: (a) the wind speed and wind direction, (b) the ambient temperature and the relative humidity (RH), (c) the pressure and water content (H_2O) in air, (d) CO concentration and the black carbon (BC) concentration, (e) O_3 concentration and the photolysis frequencies of O_3 (JO^1D), (f) the $PM_{0.8}$ mass concentration and the number concentration of particles in size of 4-700 nm (PN_{4-700}), (g) the particle number size distribution (PNSD).






Figure 3. Diurnal variations of (a) condensation sink (CS), (b) JO^1D , the total concentration of (c) VOC (TVOC) and (d) aromatics in NPF-pre days, NPF-monsoon days and non-event days. The upper and lower bars indicate the 75th and 25th percentiles, the markers are the average values.




Figure 4. Model spatial distribution of TVOC during (a) non-event days, (b) NPF-pre days, and (c) NPF-monsoon days. The star is Nam Co station.




Figure 5. The frequencies of the 48 h back trajectories of air masses arriving at Nam Co station from different directions during the occurrence time of nucleation (11:00-18:00) in (a) non-event days, (b) NPF-pre days and (c) NPF-monsoon days.




Figure 5. Wind rose plots of (a) non-event days, (b) NPF pre days and (c) NPF monsoon days. The length of each spoke on the circle represents the probability of wind coming from a particular direction at a certain range of wind speed.





Figure 6. Diurnal variations of (a) temperature, (b) wind speed, (c) RH and (d) the water content (H_2O) in the air in NPF-pre days, NPF-monsoon days and non-event days. The upper and lower bars indicate the 75th and 25th percentiles, the markers are the average values.



865 **Figure 7.** Diurnal variations of particle number size distributions (PNSD) and the number concentrations of Nucleation-mode particles (PN_{Nuc}), Aitken-mode particles (PN_{Ait}), Accumulation-mode particles (PN_{Acc}), the total particles (PN_{Tot}), CCN at S_g of 0.6% ($CCN_{0.6\%}$) and 1.2% ($CCN_{1.2\%}$) during pre-monsoon (a and c) and monsoon (b and d) seasons.




Figure 78. Particle number size distribution (PNSD) at Nam Co station. (a) Mean PNSD in pre-monsoon and monsoon; the contribution of different size ranges (4-25 nm, 25-100 nm, 100-700 nm) to the total particle number concentration (b) in pre-monsoon and (c) in monsoon.




Figure 89. Mean number concentration of CCN at S_c of 0.6% and 1.2% in pre-monsoon and monsoon seasons.




Figure 9.7. Diurnal variations of particle number size distributions (PNSD) and the number concentrations of nucleation-mode particles (PN_{Nuc}), Aitken-mode particles (PN_{Air}), accumulation-mode particles (PN_{Acc}), the total particles (PN_{Tot}), CCN at S_c of 0.6% (CCN_{0.6%}) and 1.2% (CCN_{1.2%}) during pre-monsoon (a and c) and monsoon (b and d) seasons.

Table 1. Comparison of particle number concentration (PN) and NPF parameters (NPF frequency, CS, J, GR) with other high-altitude sites around the world

Site	Altitude (m)	Observation Date and season ^a	PN (cm ⁻³)	NPF frequency	CS×10 ⁻² (s ⁻¹)	J (cm ⁻³ s ⁻¹)	GR (nm h ⁻¹)	Reference
Nam Co station, China	4379	May 2019, pre-monsoon	PN ₃₋₇₀₀ = 1163±1026	15%	0.14±0.07	$J_4 = 1.11±0.79$	GR ₄₋₂₅ = 4.2±0.9	This study
		June 2019 monsoon	PN ₃₋₇₀₀ = 3647±2671	80%	0.15±0.05	$J_4 = 1.08±0.21$	GR ₄₋₂₅ = 3.8±0.8	
NCO-P ^b , Nepal	5079	April-June 2007, pre-monsoon	NA	38%	NA	$J_{10} = 0.14$	GR = 1.8±0.7	Venzac et al. (2008)
		July-September 2007, monsoon	NA	57%	NA	$J_{10} = 0.19$	NA	
Mt. Daban, China	3295	September-October 2013, post-monsoon	PN ₁₂₋₄₇₈ = 2400	80%	NA	NA	GR = 2.0	Du et al. (2015)
Mt. Yulong, China	3410	May–April 2015, pre-monsoon	PN ₃₋₁₀₀₀₀ = 1600±1290	14%	0.2	$J_3 = 1.18$	3.2	Shang et al. (2018)

Mt. Tai, China	1534	July-August 2014, summer	NA	21%	1.1-4.8	$J_3 = 1.33-52.54$	$GR_{3-20} = 1.15-7.76$	Lv et al. (2018)
Jungfrauoch, Switzerland	3580	July 2013-June 2014, one year	$PN_{10-600} = 200-1000$	20%	NA	$J_{3,2} = 0.2-7.5$	$GR_{5-15} = 4\pm 2.3$	Tröstl et al. (2016a)
Maïdo observator, Réunion	2165	2015, one year	NA	67%	0.02-2	$J_2 = 0.5-30$	$GR_{12-19} = 8-45$	Rose et al. (2019)
Storm Peak Laboratory, USA	3210	March-May 2001-2009, spring June-July 2001-2009, summer	NA	56%	0.12 ± 0.05	$J_8 = 0.39\pm 0.05$	$GR = 7.5\pm 4.5$	Hallar et al. (2011)
			NA	43%	0.13 ± 0.06	$J_8 = 1.19\pm 0.34$	$GR = 9.1\pm 6.9$	

^a The seasons in Tibet sites were divided into pre-monsoon, monsoon and post-monsoon. The seasons not in the Tibet sites were divided into spring, summer, autumn and winter.

^b Nepal Climate Observatory at Pyramid site

NA: Data not found in the literature