

1 **An assessment of land energy balance over East Asia from**
2 **multiple lines of evidence and the roles of Tibet Plateau,**
3 **aerosols, and clouds**

4
5 Qiuyan Wang^{1,2,6}, Hua Zhang^{1,2}, Su Yang³, Qi Chen², Xixun Zhou², Bing Xie⁴, Yuying
6 Wang¹, Guangyu Shi^{1,5}, Martin Wild⁶

7
8 ¹ Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University
9 of Information Science and Technology, Nanjing 210044, China

10 ² State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, Beijing 100081,
11 China

12 ³ National Meteorological Information Center, China Meteorological Administration, Beijing 100081, China

13 ⁴ Laboratory for Climate Studies of China Meteorological Administration, National Climate Center, Beijing
14 100081, China

15 ⁵ State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics,
16 Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

17 ⁶ Institute for Atmospheric and Climate Science, ETH Zurich, 8092 Zurich, Switzerland

18

19 *Corresponding to:* Hua Zhang (huazhang@cma.gov.cn)

20

21 **Abstract.** With high emissions of aerosols and the known world’s “Third Pole” of the Tibet Plateau (TP) in
22 East Asia, knowledge on the energy budget over this region has been widely concerned. This study first
23 attempts to estimate the present-day land energy balance over East Asia by combining surface and satellite
24 observations, as well as the atmospheric reanalysis and Coupled Model Intercomparison Project phase 6
25 (CMIP6) simulations. Compared to the global land budget, a substantially larger fraction of atmospheric
26 shortwave radiation of 5.2% is reflected, highly associated with the higher aerosol loadings and more clouds
27 over East Asian land. While a slightly smaller fraction of atmospheric shortwave absorption of 0.6% is
28 unexpectedly estimated, possibly related to the lower water vapor content effects due to the thinner air over
29 the TP to overcompensate for the aerosol and cloud effects over East Asian land. The weaker greenhouse
30 effect and fewer low clouds due to the TP are very likely the causes for the smaller fraction of East Asian-
31 land surface downward longwave radiation. Hence, high aerosol loadings, clouds, and the TP over East Asia
32 play vital roles in the shortwave budgets, while the TP is responsible for the longwave budgets during this
33 regional energy budget assessment. The further obtained cloud radiative effects suggest that the presence of
34 clouds results in a larger cooling effect on the climate system over East Asian land than that over globe. This
35 study provides a perspective to understand fully the roles of potential factors in influencing the different
36 energy budget assessments over regions.

37

38 **1. Introduction**

39 Current patterns of Earth’s weather and climate are largely determined by the spatiotemporal
40 distributions of energy exchanges between the surface, atmosphere, and space. Theoretically, the outgoing
41 longwave radiation (OLR) is balanced by the incoming and reflected solar radiation at the top of the
42 atmosphere (TOA) to produce an equilibrium climate. The incoming solar radiation can be scattered by
43 clouds and aerosols or absorbed by the intermediary atmosphere, thereby contributing to the diverse energy
44 transformation at the surface (Trenberth et al., 2009; Wild et al., 2013a). The Earth’s surface energy balance
45 is of particular significance because it is the key driver of atmospheric and oceanic circulations, hydrological
46 cycles, and various surface processes (Wild et al., 2008; Mercado et al., 2009; Wild et al., 2013a; L’Ecuyer
47 et al., 2015). Anthropogenic influences on climate change are driven by the uneven distribution of the TOA
48 net radiation caused by forcings perturbed by variations of the atmospheric composition of greenhouse gases
49 and aerosols as well as aerosol-cloud interactions (Trenberth et al., 2009; Stephens et al., 2012; Wild et al.,
50 2013a; Trenberth et al., 2014; L’Ecuyer et al., 2015; Wild et al., 2019; Z. Wang et al., 2021).

51 Many efforts have been made to quantify the magnitudes of different radiative components or energy
52 budgets in the climate system over a range of time-space scales, such as on global scales (Lin et al., 2008;
53 Trenberth et al., 2009; Stephens et al., 2012; Wild et al., 2013b; Wild et al., 2015; L’Ecuyer et al., 2015;
54 Wild et al., 2019; Wild, 2020), over land and ocean domains or the energy transport between them (Fasullo
55 and Trenberth, 2008a, b; Trenberth et al., 2009; Wild et al., 2015; L’Ecuyer et al., 2015), over the Arctic
56 (Previdi et al., 2015; Christensen et al., 2016), and over individual continents and ocean basins (L’Ecuyer et
57 al., 2015; Kim and Lee, 2018; Thomas et al., 2020). The energy balance at the TOA can be accurately

58 monitored by satellites from the most advanced Clouds and the Earth's Radiant Energy System (CERES)
59 Energy Balanced and Filled (EBAF) data product (Loeb et al., 2018), while considerably larger uncertainties
60 appear at the surface fluxes owing to weaker observational constraints (Raschke et al., 2016; Kato et al.,
61 2018; Huang et al., 2019). These assessments mostly build upon complementary approaches from a
62 combination of space and surface observations, climate models, and reanalyses. To date, the discrepancies
63 of independent global mean surface radiative fluxes have estimated to be within a few W m^{-2} (Wild, 2017a,
64 b), enabling the accurate quantification of global surface budgets. In addition, the surface radiative
65 components simulated by various climate models vary substantially in a range of around 10–20 W m^{-2} on
66 global scales, but exhibit greater inter-model discrepancies on regional scales (Li et al., 2013; Wild et al.,
67 2013a; Boeke and Taylor, 2016; Wild et al., 2015; Wild, 2017a, b, 2020). Existing challenges on the surface
68 energy estimates include considerable uncertainties from surface albedo and skin temperature, as well as the
69 partitioning of surface net radiation into sensible and latent heat (SH; LH) (Wild, 2017a, b).

70 Due to the large population and the largest emission source of aerosols and their precursors, East Asia,
71 especially China, has long been a hotspot in climate change research. Aerosols can interact with radiation
72 directly by scattering and absorbing solar/thermal radiation (Ghan et al., 2012) and indirectly by modifying
73 cloud microphysical properties and lifetimes (Li et al., 2011), thereby influencing Earth's radiation balance.
74 As the world's largest and highest plateau, the Tibet Plateau (TP) covers nearly one fifth of the East Asian
75 land area, significantly affecting the atmospheric circulation, energy budget, and water cycles of climate
76 system through its orographic and thermal effects (Liu et al., 2007; Xu et al., 2008a, b; Wu et al., 2015).
77 Deeper insights into the energy budget differences over East Asian and global land under the background of
78 high aerosol emissions and the role of the TP in East Asia are of the meaningful and essential attempts.
79 Moreover, clouds play a key role in modulating global and regional energy budgets and hydrological cycles
80 through increasing the reflected solar radiation and also the downward thermal radiation, leading to a cooling
81 and warming of climate system (Stephens, 2005; Wild et al., 2013a; Li et al., 2015; H. Wang et al., 2021).
82 Therefore, our emphasis in this study is on the regional characterization of the East Asian energy balance
83 under both all-sky and clear-sky conditions based on a combination of surface observations, satellite-derived
84 products, reanalysis, and Coupled Model Intercomparison Project phase 6 (CMIP6) models. The cloud
85 influence on the radiative energy budgets at the TOA, within the atmosphere, and at the surface is further
86 quantified over this region. Section 2 introduces the different data sources used in this study, including
87 surface and satellite observations, climate models, and reanalysis. Sections 3 and 4 provide detailed analyses
88 of the all-sky and clear-sky estimates of the energy balance components. The inferred cloud radiative effects
89 (CREs) at the TOA, within the atmosphere, and at the surface are presented in Section 5. Summary and
90 conclusions are given in Section 6. The present-day in this study represents years of 2010–2014, which
91 corresponds to the last five years of the historical simulations in CMIP6 climate models. East Asian land as
92 considered in this study consists of five countries, including China, Japan, South and North Korea, as well
93 as Mongolia.

94

95 2. Data sources

96 2.1. Surface observations

97 Considering the efforts to diminish the inhomogeneities in the measurement of ground-based surface
98 (downward) solar radiation (SSR) (Tang et al., 2011; Wang, 2014; Wang et al., 2015; Wang and Wild, 2016;
99 He et al., 2018; Yang et al., 2018, 2019) and the large amount of observational stations over China, the
100 homogenized monthly all-sky and clear-sky SSR datasets from the China Meteorological Administration
101 (CMA) National Meteorological Information Center (NMIC) are used in this study (<http://data.cma.cn/enl>)
102 (Yang et al., 2018, 2019). In this dataset, the clear-sky condition at observational sites is defined based on
103 the measured cloud fraction per day of no more than 15% (Yang et al., 2018). Taking clear-sky data (with
104 relatively complex missing months compared to the all-sky dataset) as an example, sites with more than one
105 year of > 2 missing months were deleted to ensure ≥ 4 years of available data during the period 2010-2014,
106 then the spline interpolation was performed on the missing months of the selected sites. As a consequence,
107 99 and 76 sites are available for the all-sky and clear-sky studies, respectively. Besides, to further explore
108 the anthropogenic influence on SSR, 84 (62) urban and 15 (14) rural stations for all-sky (clear-sky)
109 conditions are defined according to the administrative divisions of China (Wang et al., 2017).

110 For the remaining East Asian sites, we use the monthly Global Energy Balance Archive (GEBA) dataset
111 (<http://www.geba.ethz.ch>) (Wild et al., 2017), which contains a worldwide widespread distribution of
112 monthly data from many sources, e.g., from the World Radiation Data Center (WRDC), the Baseline Surface
113 Radiation Network (BSRN), etc. Among these data sources, the BSRN dataset has a much higher precision
114 and temporal resolution (up to 1 min) compared to the GEBA, but its site number is very limited over East
115 Asia (only a few sites located in Japan and one site in Xianghe, China, but with no data available during this
116 study period). Moreover, the relative random error of the monthly SSR from the GEBA data evaluated by
117 Gilgen et al. (1998) is 5%.

118 In order to retain as many sites as possible during the study period, we widen the selection criterion of
119 the GEBA data, i.e., sites with data ≥ 4 years and missing months ≤ 3 . Eventually, 8, 2, 4, and 14 sites are
120 selected from GEBA in China, Mongolia, South and North Korea, and Japan, respectively. Especially,
121 among the 14 sites in Japan, five pairs of the duplicate sites are obtained from the WRDC and BSRN sources,
122 respectively, and the left 4 sites are only from the WRDC (9 sites available). For China, only one site from
123 Hongkong out of 8 GEBA sites is not repetitive from the above-mentioned CMA sites (1 site available).
124 Therefore, 16 out of 28 GEBA sites are available under all-sky conditions (including 15 sites over regions
125 outside China and 1 site over Hongkong, China) by taking the average of these duplicate sites in Japan
126 instead, while the clear-sky reference sites are obtained from the interpolated CERES EBAF clear-sky
127 estimates at the GEBA sites (also 16 sites) due to the limited numbers of observational sites over these
128 regions. Additionally, we regard four island sites in Japan as rural stations (not shown in the figures), while
129 the sites in Mongolia as well as South and North Korea are all urban sites.

130 As shown in Fig. S1, there are 99 (rural/total: 15/99) and 16 (rural/total: 4/16) sites from the CMA and
131 GEBA available under all-sky conditions, respectively, whereas 76 (rural/total: 14/99) and 16 (from the
132 CERES-interpolated data at the 16 GEBA sites) sites are considered for clear-sky conditions, respectively.
133 More detailed station information is given in Table S1.

134

135 2.2. Satellite observation

136 Owing to the excellent temporal and spatial coverage of satellite instruments, CERES data products are
137 widely used to track variations of Earth's energy budgets. The newly released CERES EBAF Edition 4.1
138 with a monthly $1^\circ \times 1^\circ$ latitude-longitude resolution is used in this study (<https://ceres.larc.nasa.gov/data/>). In
139 this dataset, the TOA radiation components are adjusted within their uncertainty ranges based on the
140 independent observational ocean estimates of global heating rate (Loeb et al., 2018). Unlike the directly
141 measured TOA energy budget, the EBAF-surface energy fluxes are calculated by the cloud and aerosol
142 properties from satellite-derived products as well as the atmospheric profiles from reanalysis, with a lower
143 accuracy than their TOA counterparts (Kato et al., 2018). The uncertainty ranges in $1^\circ \times 1^\circ$ regional monthly
144 all-sky and clear-sky longwave (LW) and shortwave (SW) radiation fluxes at the TOA are also documented
145 by Loeb et al. (2018).

146

147 2.3. Climate models and reanalysis

148 Data from 40 CMIP6 climate models are used for the analyses in this study with their model
149 abbreviations, modeling groups, and resolutions in Table S2. A detailed description of the modeling groups
150 participating in CMIP6 is provided at <https://pcmdi.llnl.gov/CMIP6/>. The CMIP6 model-calculated radiation
151 fluxes under investigation for this study include energy budgets under both all-sky and clear-sky conditions
152 from 'historical all forcings' experiments covering the period 2010-2014. In these historical simulations,
153 both natural (e.g., solar variability and volcanic aerosols) and anthropogenic (e.g., greenhouse gases, aerosols,
154 and land use) forcings are considered to reproduce the climate change and evolution since preindustrial times
155 as accurately as possible (Eyring et al., 2016). Only the first ensemble member of each model is selected for
156 the analysis and the model numbers vary slightly among different available energy components.

157 In the long history of the European Center for Medium-range Weather Forecast (ECMWF), ERA5 is
158 the fifth generation product. It is a comprehensive reanalysis from 1959 to near real time, which assimilates
159 as many observations as possible in the upper air and near surface
160 ([https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-means?tab=form)
161 [means?tab=form](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-means?tab=form)). Monthly means of the radiative components from ERA5 are used in this study with a
162 resolution of $0.25^\circ \times 0.25^\circ$ (regridDED to $1^\circ \times 1^\circ$). Compared to previous reanalyses (such as ERA-Interim), a
163 major strength of ERA5 is the much higher temporal and spatial resolutions, as well as a higher vertical
164 resolution with 137 levels (Hersbach et al., 2020). Several independent studies have evaluated the
165 performance of ERA5 since its release. For example, excellent closure of the Arctic energy budget based on

166 ERA5 atmospheric data has been assessed by Mayer et al. (2019). The representation of surface irradiance
 167 of ERA5 has been compared with other reanalyses and with ground and satellite observations (Trollet et al.,
 168 2018; Urraca et al., 2018). Specifically, Trollet et al. (2018) found that the surface solar irradiance over the
 169 tropical Atlantic Ocean from ERA5 exhibits fewer biases than the second version of the Modern-Era
 170 Retrospective Analysis for Research and Applications (MERRA-2). Urraca et al. (2018) reported that ERA5
 171 can be a valid alternative for satellite-derived products in terms of surface irradiance in most inland stations
 172 compared to ERA-Interim or MERRA-2. Furthermore, based on BSRN station data, Tang et al. (2021)
 173 pointed out that the accuracy of the ERA5 over land in terms of surface downward longwave radiation is
 174 higher than CERES-derived product on average both at hourly and monthly times scales.


175

176 **3. Assessment of land energy balance budgets under all-sky conditions**

177 3.1. Shortwave components

178 Under all-sky conditions, the present-day annual land-mean anomalies of TOA incident solar radiation
 179 as well as the SW net radiation at the TOA, within the atmosphere, and at the surface regarding to their
 180 respective multi-model means as simulated by various CMIP6 models over East Asia are shown in Fig. 1a.
 181 A summary of the CMIP6 model statistics (such as available model number, model spread, and the standard
 182 deviation (SD)), along with the corresponding multi-model mean, ERA5-, and CERES-derived estimates of
 183 different energy balance components are listed in Table 1. As shown in Fig. 1a, with the exception of the
 184 BCC-CSM2-MR and BCC-CESM1 models, all models give an estimate around 334 W m^{-2} for TOA
 185 incoming solar radiation with a very small SD of 0.2, closely matching the multi-model mean as well as the
 186 CERES and ERA5 estimates (Table 1). The multi-model means of solar absorption at the TOA, within the
 187 atmosphere, and at the surface are 217, 73, and 144 W m^{-2} , respectively, all within 2 W m^{-2} of the biases
 188 against the CERES-derived estimates, while they are $3\text{--}4 \text{ W m}^{-2}$ larger for those from ERA5 at the TOA and
 189 within the atmosphere, yielding 1 W m^{-2} of bias against the CERES-based estimate at the surface (Table 1).
 190 However, the individual models vary significantly in their simulated annual East Asian land-mean solar
 191 absorption both at the TOA and surface (Fig. 1a), with SDs of around 6 W m^{-2} and inter-model spreads of
 192 more than 20 W m^{-2} (Table 1). Considering the smaller absolute amount of atmospheric and surface solar
 193 absorption compared to the TOA counterpart (73 and 144 vs. 217 W m^{-2} ; Table 1), the relative (percentage)
 194 differences relative to their respective multi-model means (relative (percentage) difference =
 195 $\frac{\text{range}}{\text{multi-model mean}} \times 100\%$) indicate that the uncertainties within the atmosphere and at the surface are larger
 196 than that at the TOA (i.e., TOA: $\frac{22}{217} \times 100\% = 10\%$; Atmosphere: $\frac{19}{73} \times 100\% = 26\%$; Surface:
 197 $\frac{23}{144} \times 100\% = 16\%$).

198



199

200 **Figure 1.** Annual land mean anomalies of (a, b) shortwave (SW) and (c, d) longwave (LW) budgets
 201 (Units: $W m^{-2}$) with regard to their respective multi-model means for present-day climate under (a, c)
 202 all-sky and (b, d) clear-sky conditions over East Asia as simulated by various CMIP6 models. The black, red,
 203 blue, and green lines represent the TOA incoming solar radiation, as well as the net SW/LW radiation at
 204 the TOA, within the atmosphere, and at the surface, respectively.

205

206 **Table 1.** Annual land mean estimates (Units: $W m^{-2}$) of the magnitudes of various energy balance
 207 components and cloud radiative effects (CREs) over East Asia under all-sky and clear-sky conditions at the
 208 TOA, within the atmosphere, and at the surface, respectively. The CMIP6 model statistics (e.g., available
 209 model number, spread, standard deviation (SD)), as well as the corresponding multi-model mean, ERA5-,
 210 and CERES-derived estimates are also given in the Table.


Component ($W m^{-2}$)	CMIP6				ERA5	CERES
	models	spread	SD	mean		
TOA						
Solar down	39	4	0.2	334	334	334
Solar up all-sky	39	23	6	-117	-115	-118

Solar net all-sky	39	22	6.1	217	219	216
Solar up clear-sky	39	24	7	-76	-78	-72
Solar net clear-sky	39	24	6.9	258	256	262
SW CRE	39	26	6.5	-41	-37	-46
Thermal up all-sky	39	12	3.5	-224	-225	-226
Thermal up clear-sky	39	15	3.2	-247	-246	-250
LW CRE	39	12	2.4	23	21	24
Net CRE	39	24	5.8	-18	-16	-22
Atmosphere						
SW absorption all-sky	39	19	3.8	73	78	74
SW absorption clear-sky	35	19	3.8	69	77	71
SW CRE	32	33	6.9	4	2	3
LW net all-sky	39	22	5.1	-152	-150	-157
LW net clear-sky	35	16	3.6	-151	-151	-154
LW CRE	32	14	3.3	-2	1	-3
Net CRE	32	35	7.8	1	2	0
Surface						
SW down all-sky	39	33	7.6	186	191	178
SW up all-sky	39	24	6.5	-43	-50	-36
SW absorbed all-sky	39	23	6.1	144	141	142
SW down clear-sky	35	25	4.6	242	238	236
SW up clear-sky	35	27	6.8	-53	-59	-45
SW absorbed clear-sky	32	36	7.8	189	179	191
SW CRE	35	28	6.6	-46	-38	-49
LW down all-sky	39	27	7.9	280	273	285
LW up all-sky	39	23	7.1	-352	-347	-354
LW net all-sky	39	23	5.7	-71	-74	-69
LW down clear-sky	35	26	6.8	256	253	256
LW up clear-sky	35	23	7.1	-351	-347	-353
LW net clear-sky	35	18	4.1	-95	-94	-97
LW CRE	35	12	3.5	24	20	27
net CRE	32	31	6	-21	-18	-22
net radiation	39	20	5.3	72	67	73
LH	40	26	4.7	-43	-38	—
SH	40	21	5.2	-31	-29	—

211

212 The simulated SSR, however, shows the largest spread of more than 30 W m⁻² (ranging from 172–205
213 W m⁻²) among all the substantially differing all-sky surface radiation components, with a large SD of 7.6 W
214 m⁻² (Fig. 2a; Table 1). The multi-model mean SSR is estimated to be 186 W m⁻², suggesting positive and
215 negative deviations of 8 and 5 W m⁻² from the CERES- and ERA5- derived estimates, respectively (Table
216 1). Interestingly, although the discrepancy between them is very large (8 or 5 W m⁻²), both the resulting
217 surface solar absorption differences are very small (within 3 W m⁻²), indicating that a higher SSR goes
218 together with a higher surface albedo (Table 1), which agrees well with that on a global mean level (Wild et
219 al., 2015).

220



221


222 **Figure 2.** Annual land mean surface downward (a) SW and (b) LW radiation (Units: $W m^{-2}$) under both
 223 all-sky (orange bars) and clear-sky (green bars) conditions over East Asia as calculated by various CMIP6
 224 models.

225

226 3.2. Best estimates for the surface downward SW radiation

227 As a major component of Earth's energy balance, the solar radiation reaching the Earth's surface
 228 governs a wide range of surface physical and chemical processes. The spatial distributions of the site-based
 229 annual mean SSR from the CMA and GEBA (Section 2.1) over East Asia under all-sky conditions are
 230 presented in Fig. 3a, together with the classified rural and urban sites. In short, the high values are mainly
 231 located at the high elevation stations over western China and a few island sites in Japan (e.g.,
 232 Minamitorishima, Japan; not shown in the figure), especially over the TP, with the largest value reaching
 233 $263 W m^{-2}$ (Geer, Tibet), which is associated with the high atmospheric transparency over these regions.
 234 However, the low annual mean values are primary over southwestern China, with the smallest value of 103
 235 $W m^{-2}$ (Shapingba, Chongqing), which is possibly caused by the higher aerosol loadings (Liao et al., 2015;
 236 de Leeuw et al., 2018) and more clouds (Li et al., 2017; You et al., 2019; Lei et al., 2020; Zhang et al., 2020)
 237 over these regions. This distribution pattern is highly consistent with that over China documented by Q.
 238 Wang et al. (2021).

239




240

241 **Figure 3.** Spatial distributions of annual mean surface downward solar radiation (SSR) (Units: $W m^{-2}$) under
 242 (a) all-sky and (b) clear-sky conditions over East Asia. The all-sky sites are available from 99 CMA and 16
 243 GEBA stations, while there are 76 CMA and 16 CERES-interpolated sites for clear-sky conditions. The
 244 cross and circle symbols indicate rural (19 vs. 18 for all-sky and clear-sky conditions) and urban stations (96
 245 vs. 74), respectively.

246

247 Figure 4 shows the distributions of annual mean SSR biases derived from the CERES, CMIP6 multi-
 248 model mean, and ERA5 against the surface observations, as well as the comparisons of their respective
 249 annual land means at the surface sites with their observed counterparts. The corresponding quantifications
 250 of the magnitudes of station-mean biases are also given in Table 2. According to the comparisons, they all
 251 correlate well with the ground-based observations, with their respective high correlation coefficients of 0.93,
 252 0.87, and 0.89, indicative of the highest accuracy in the CERES-derived estimate (Figs. 4b, d, and f). To
 253 quantify their SSR mean biases against the corresponding observed counterparts, the CERES-based bias at
 254 all sites is the smallest, with a station-mean bias of $3.8 W m^{-2}$, followed by the CMIP6 multi-model mean
 255 and the ERA5 reanalysis (with respective station-mean biases of 13.8 and $16.5 W m^{-2}$) (Table 2).
 256 Additionally, among all the aforementioned SSR estimates, the East Asian urban sites are in general more
 257 significantly overestimated than the rural sites on average compared to the surface observations (Figs. 4b, d,
 258 and f; Table 2). This further supports the argument that rural stations might be more representative for larger
 259 scale comparisons (e.g., the general circulation model grid scales) than the urban stations (which are
 260 vulnerable to local pollution) (Wang et al., 2018). The overestimations are mainly located in the high-latitude
 261 regions over East Asia for CERES-derived estimates (among them the underestimations mostly from rural
 262 sites), while the underestimates are primarily located in lower-latitude and eastern coastal regions (Figs. 4a
 263 and b). The CMIP6 multi-model mean and ERA5-derived SSR generally greatly overestimate the surface-
 264 based observations both at urban and rural sites, except for the regions over northern and northeastern Inner
 265 Mongolia, northwestern Heilongjiang (located in the northeastern China), and some individual sites over
 266 southwestern China (Figs. 4c-f). The annual land-mean area-weighted average SSR over East Asia derived
 267 from CERES is estimated to be $178 W m^{-2}$, which is closest to the surface observational estimate of $174 W$
 268 m^{-2} , compared to the much higher overestimations of both the CMIP6 multi-model mean and ERA5 (186
 269 and $191 W m^{-2}$) against the surface observations (Table 3), which shows a high consistency with their bias
 270 distributions and the collocated quantifications (Fig. 4; Table 2).



272

273 **Figure 4.** Spatial distributions of annual mean SSR biases (Units: $W m^{-2}$) derived from (a) CERES-EBAF,
 274 (c) CMIP6 multi-model mean, and (e) ERA5 reanalysis at a combination of the CMA and GEBA sites under
 275 all-sky conditions over East Asia. The corresponding comparisons of their respective annual means at the
 276 surface sites with their observed counterparts are displayed in (b), (d), and (f), respectively. The cross and
 277 circle symbols in Figs. a, c, e as well as the orange and green stars in Figs. b, d, f indicate rural and urban
 278 stations, respectively.

279

280 **Table 2.** Annual station-mean SSR biases (Units: $W m^{-2}$) derived from CERES-EBAF, CMIP6 multi-model
 281 mean, and ERA5 compared to the surface observational sites under all-sky and clear-sky conditions during
 282 2010-2014 over East Asian land, together with the separate station averages of biases at urban and rural sites.
 283 The values in parentheses represent the percentages of SSR biases relative to their respective station-mean
 284 averages with the largest percentages around 10% and 4% for all-sky and clear-sky conditions.

Station-mean SSR biases	All-sky		Clear-sky	
	Urban	Rural	Urban	Rural

(Unit: W m^{-2})	all	urban	rural	all	urban	rural
CERES-EBAF	3.8 (2.3%)	4.2 (2.6%)	1.7 (0.9%)	0.4 (0.2%)	0.5 (0.2%)	-0.3 (-0.1%)
CMIP6	13.8 (8.3%)	15 (9.2%)	7.4 (4.1%)	9.1 (4%)	9.7 (4.3%)	6.4 (2.8%)
ERA5	16.5 (10%)	17.2 (10.5%)	12.7 (7%)	5.7 (2.5%)	6.2 (2.7%)	3.6 (1.5%)

285


286 **Table 3.** Annual land mean area-weighted average SSR (Units: W m^{-2}) from a combination of the CMA and
 287 GEBA (CERES-interpolated) site observations under all-sky (clear-sky) conditions during the period 2010-
 288 2014 over East Asia, together with the corresponding estimates from the CERES-EBAF, CMIP6 multi-
 289 model means, and ERA5, respectively.

Average annual mean SSR during 2010-2014 over East Asia (Units: W m^{-2})	Surface observations	CERES-EBAF	CMIP6	ERA5
All-sky	174	178	186	191
Clear-sky	230	236	242	238

290

291 However, the ground-based observations are spatially limited with sparse stations in some remote
 292 regions and are thus inadequate for many applications, as they may be not representative for real situations.
 293 To better constrain the large spread in the model-based SSR outlined above, we combine the ground-based
 294 observations to obtain the best estimate referring to the approach introduced in Wild et al. (2013a). Figure
 295 5a gives various CMIP6 model biases of all-sky SSR at all the surface sites and their respective East Asian
 296 land means. The higher overestimations relative to surface observations generally correspond to higher
 297 model-based East Asian land means, with a much higher correlation coefficient of 0.96 than that of 0.88 on
 298 the global scale (Wild et al., 2015). Thus, the best estimate of the annual East Asian land-mean SSR is
 299 deduced to be $174.2 \pm 1.3 \text{ W m}^{-2}$ (2σ uncertainty) in light of the linear regression analysis. The corresponding
 300 estimates from CERES and ERA5 are also labeled in the figure, at 178 and 191 W m^{-2} , respectively, implying
 301 a slight and substantial overestimation for CERES and ERA5 estimates. There is an overall tendency that
 302 most models overestimate the surface downward SW fluxes (36 out of 39 sites) compared to the ground-
 303 based observations, with a multi-model mean overestimation relative to site observations of 13.8 W m^{-2} ,
 304 which is also a longstanding issue in climate modelling (Wild et al., 1995; Wild et al., 2015).

305



306

307 **Figure 5.** Annual land mean SSR (Units: W m^{-2}) of various CMIP6 models as well as their respective model
308 biases relative to an average over surface sites (99 CMA and 16 GEBA for all-sky; 76 CMA and 16 CERES-
309 interpolated sites for clear-sky) under (a) all-sky and (b) clear-sky conditions during 2010-2014 over East
310 Asia. Green stars represent various CMIP6 models. Best estimate here (orange circle) can be inferred from
311 the intersection between the linear regression line (green solid lines) and the zero-bias line (blue dotted lines).
312 Furthermore, the corresponding estimates from CERES-EBAF and ERA5 are also given by red triangle and
313 blue square, respectively.

314

315 3.3. Longwave components

316 Similar to the all-sky SW counterparts, obvious discrepancies can still be noted in the annual land-mean
317 LW radiation over East Asia among models, especially for those within the atmosphere and at the surface
318 (Fig. 1c). Correspondingly, the simulated TOA OLR varies in a range of 12 W m^{-2} , which is almost 10 W
319 m^{-2} lower than that within the atmosphere (22 W m^{-2}) and at the surface (23 W m^{-2}) (Table 1). The estimated
320 annual East Asian land-mean TOA OLR from the CMIP6 multi-model mean is -224 W m^{-2} , within 2 W m^{-2}
321 of the deviations from the CERES- and ERA5-inferred estimates. The model spread of the simulated annual
322 land-mean net LW radiation becomes larger from the TOA to the surface, with SDs of 3.5, 5.1, and 5.7 W
323 m^{-2} , respectively, which shows the same tendency as the relative (percentages) differences with respect to
324 their multi-model means (5.4%, 14.5%, and 32.4%).

325 These large discrepancies in surface net LW radiation between models are particularly evident in the
326 surface downward LW radiation (Fig. 2b; Table 1), with a range up to 27 W m^{-2} (from 267 to 294 W m^{-2})
327 and a SD of 7.9 W m^{-2} , which is also the largest deviation among all components under all-sky conditions.
328 Compared to the CERES estimates, the slightly lower surface upward LW radiation (-352 vs. -354 W m^{-2})
329 and much lower surface downward LW radiation (280 vs. 285 W m^{-2}) from the multi-model means are the
330 major reason for the small deviation (within 2 W m^{-2}) of the surface net LW radiation between them (Table
331 1). It's interesting to note that the annual East Asian land-mean surface upward LW radiation estimated from
332 the ERA5 is the lowest among all these estimates, at -347 W m^{-2} , suggesting the lowest surface skin
333 temperature of the ERA5 product according to the Stefan-Boltzmann law, followed by the estimates from
334 the multi-model mean and CERES (Table 1). In addition, the annual land-mean surface downward LW
335 radiation estimated by ERA5 is 273 W m^{-2} , approximately 7 and 12 W m^{-2} lower than the estimates by the
336 CMIP6 multi-model mean and CERES, respectively (Table 1). Therefore, both the lower surface upward
337 and downward LW radiation fluxes result in the small deviation in the estimated surface net LW radiation
338 from ERA5 compared to those from the multi-model mean and CERES (Table 1). Since the reanalysis
339 products take as many observed atmospheric parameters with global coverage as possible into consideration
340 during the radiative transfer calculations, they are widely used to obtain more accurate surface LW radiation
341 (Simmons et al., 2004; Wild et al., 2015). We also examined the corresponding surface LW fluxes from
342 another reanalysis, namely MERRA-2, and found much lower annual land means than those from ERA5, in
343 particular for the surface downward LW radiation (not shown), which arrives at the similar conclusions with
344 that documented by Urraca et al. (2018). Thus, considering the limited observational surface LW radiation
345 data over East Asia, ERA5 might be the best reference for the estimates of the annual land-mean surface
346 upward and downward LW radiation, at -347 and 273 W m^{-2} , respectively (Table 1).


347

348 3.4. Discussion of land energy balance over East Asia under all-sky conditions


349 3.4.1. Radiative components

350 Figure 6a displays the schematic diagram of the all-sky land mean energy balance over East Asia,
351 including the above-mentioned SW and LW radiation budgets and other radiative components discussed in
352 the following. The estimated annual East Asian land-mean incoming, reflected, and net SW radiation as well
353 as the OLR at the TOA are therefore 334, -118, 216, and -226 W m^{-2} (Table 1), respectively, based on the
354 CERES EBAF dataset. The corresponding uncertainties are obtained from the uncertainty of 2.5 (1σ)
355 uncertainty) W m^{-2} for both SW and LW fluxes given by (Loeb et al., 2018). The annual East Asian land-
356 mean TOA OLR in CERES-EBAF is estimated to be 10 W m^{-2} larger than the TOA absorbed SW radiation,
357 implying an energy loss of 10 W m^{-2} at the TOA under all-sky conditions, which should be compensated by
358 the LH and SH transported from regions outside East Asia (Fig. 6a).

359



360



361

362 **Figure 6.** Diagrams of the annual land mean energy balance (Units: W m^{-2}) over East Asia under (a) all-
 363 sky and (b) clear-sky conditions for present-day climate. The uncertainty ranges are also given in
 364 parentheses.

365

366 For the SSR, the annual East Asian land-mean best estimate based on the CMIP6 multi-model
 367 simulations and surface observations is 174.2 W m^{-2} (Fig. 5a and Fig. 6a). Considering the abnormally high
 368 overestimation by ERA5 compared to surface observation, the high value of the uncertainty range is given
 369 by the estimate from CERES EBAF (178 W m^{-2}), while its low value is from the lowest model estimate (172
 370 W m^{-2} ; Fig. 2a) (Fig. 6a). The all-sky surface albedo information is derived from the ratio between the
 371 CERES-derived surface upward and downward solar radiation, with a radiation weighted average of around
 372 0.2 ($36.4/178.3$) over East Asian land. However, the corresponding surface albedos estimated by the CMIP6
 373 multi-model mean and ERA5 are substantially higher than that from the CERES, with respective averages
 374 of around 0.23 ($42.7/186.4$) and 0.26 ($49.6/191$). Considering the large spatial coverage of remote sensing
 375 measurement to map albedo globally, the CERES-derived annual East Asian land-mean surface albedo is
 376 adopted as the best estimate in this study. Therefore, considering the rounded best SSR estimate of 174 W
 377 m^{-2} , the calculated surface reflected and absorbed SW radiation fluxes are around -35 and 139 W m^{-2} ,
 378 respectively. As shown in Table 1, the uncertainty range of the surface absorbed SW radiation is $132\text{--}144$
 379 W m^{-2} according to the lowest value of CMIP6 models and the highest estimate among the aforementioned
 380 estimates, which gives rise to an uncertainty range of the surface reflected solar radiation of $34\text{--}40 \text{ W m}^{-2}$.
 381 Together with the annual East Asian land-mean SW absorption at the TOA and surface of 216 and 139 W
 382 m^{-2} , the best estimate for the atmospheric SW absorption is therefore to be 77 W m^{-2} , which is within 4 W
 383 m^{-2} of the differences between those estimated from the CMIP6 multi-model mean and CERES and closes

384 to the ERA5-derived estimate of 78 W m^{-2} (Table 1). The uncertainty range of the atmospheric SW
385 absorption is also determined by the estimates from different data sources as shown in Fig. 6a.

386 The downward LW radiation emitted by the atmosphere is mainly sensitive to the near-surface
387 temperature, water vapor, and cloud properties, while the surface emission is in proportion to the skin
388 temperature according to the Stefan-Boltzmann law. As analyzed in section 3.3, the best estimates of the
389 East Asian annual land-mean surface upward and downward LW radiation amount to -347 and 273 W m^{-2} ,
390 respectively, with uncertainty ranges coming also from the above-discussed different data sources (Fig. 6a).
391 The surface net LW radiation is then estimated to be -74 W m^{-2} based on the surface upward and downward
392 LW radiation outlined above. Combined with TOA outgoing thermal radiation of -226 W m^{-2} , the estimated
393 atmospheric net LW radiation is -152 W m^{-2} , which is close to the collocated estimates from the multi-model
394 mean (-152 W m^{-2}) and ERA5 (-150 W m^{-2}) but deviates substantially from the CERES-derived estimate of
395 -157 W m^{-2} (Table 1). Considering the surface absorbed SW radiation of 139 W m^{-2} , a best estimate for
396 surface net radiation is 65 W m^{-2} , suggesting that around 65 W m^{-2} of energy is available for the non-radiative
397 SH and LH. Besides, the ERA5 estimate of 67 W m^{-2} is very close to the best estimate of 65 W m^{-2} , while
398 much higher estimates of 72 and 73 W m^{-2} are obtained from the multi-model mean and CERES (Table 1),
399 respectively.

400

401 3.4.2. Nonradiative components

402 The surface net radiation is mainly balanced by the non-radiative components of SH and LH in addition
403 to a very small proportion of ground heat flux and melt (less than 1%) (Ohmura, 2004). However, due to the
404 lack of constraints from in-situ and space observations, this partitioning of the surface net radiation into SH
405 and LH is still subject to considerable uncertainties. As shown in Fig. S2, the simulated annual East Asian
406 land-mean LH and SH vary greatly between different models, with a range of 26 and 21 W m^{-2} , respectively,
407 as well as the relative discrepancies relative to their respective multi-model means of 60% ($\frac{26}{43} \times 100\%$) and
408 68% ($\frac{21}{31} \times 100\%$), respectively, showing larger discrepancies between models with larger uncertainties in SH
409 (Table 1). The best SH estimate can therefore be obtained from the residual of the LH. To obtain a more
410 accurate surface LH from available datasets of the multi-model mean and ERA5, we take an average of them
411 as the best estimate, namely -40 W m^{-2} , the uncertainty ranges of which are also given according to these
412 estimates (Fig. 6a). Note that all the values in this study are calculated on the basis of one decimal point,
413 which may result in 1 W m^{-2} of bias during the rounding process. Combined with the surface net radiation
414 and LH of 65 and -40 W m^{-2} , respectively, the surface SH is estimated to be -25 W m^{-2} , the uncertainty range
415 of which is also given by the existing estimates from various CMIP6 models and ERA5 (Fig. 6a). In addition,
416 although the annual land-mean SH estimated from the MERRA-2 is much higher than the estimates from
417 multi-model mean and ERA5 (not shown), the estimated LH is around -39 W m^{-2} (not shown), very close to
418 the best estimate of -40 W m^{-2} , which increases our confidence in the estimation of this quantity.

419

420 3.4.3. Comparisons with global annual land-mean estimates

421 Notable discrepancies exist in the global land-mean energy budgets reported by Wild et al. (2015) and
 422 the regional land-mean estimates over East Asia in this study (Fig. S3; Table 4). For the SW budgets, the
 423 estimated annual land-mean TOA incident solar radiation over East Asia is 9 W m^{-2} higher than that over
 424 global land (334 vs. 325 W m^{-2}), implying a slightly lower land-mean solar zenith angle over East Asia.
 425 Comparisons also show a slightly higher relative percentage of TOA reflected solar radiation of 0.8% despite
 426 of the much lower surface reflected SW radiation of 4.3% over East Asian land compared to global land with
 427 respect to their respective TOA incident solar radiation (thereafter call ‘relative percentage’ for short). This
 428 suggests much more relative atmospheric SW reflection of 5.2% over East Asian land, which agrees fairly
 429 well with more aerosols (Wei et al., 2019) and more clouds (King et al., 2013; Fan et al., 2018; also see Fig.
 430 S4) over this region compared to global land. However, the annual land-mean solar radiation reaching the
 431 East Asian surface is around 10 W m^{-2} lower than that over global land (174 vs. 184 W m^{-2}), approximately
 432 accounting for 52.1% and 56.6% of their respective incident solar radiation at the TOA, respectively,
 433 indicating lower fraction of solar energy arriving at the East Asian surface compared to global land. Together
 434 with the lower annual land-mean surface albedo over East Asian land compared to global land (20% vs.
 435 26%), this leads to the similar relative percentages of surface absorptions (41.6% vs. 41.9%). Although the
 436 magnitude of the atmospheric SW absorptions over East Asian and global land are nearly the same (both
 437 around 77 W m^{-2}), the corresponding relative percentage over East Asian land is a little bit lower than that
 438 over global land (around 0.6%). This is somewhat unexpected due to the fact of more clouds and aerosol
 439 loadings over East Asian land, which is possibly offset by the lower water vapor contents caused by the
 440 higher altitudes and thinner air over the TP.

441

442 **Table 4.** Comparisons of the annual mean SW/LW energy balance components (Units: W m^{-2}) over East
 443 Asian land (this study) and global land (Wild et al., 2015) as well as the corresponding relative percentages
 444 with regard to their respective TOA incident solar radiation/surface LW emissions, along with the relative
 445 percentage differences between them.

Component	East Asian land		Global land		Percentage difference
	Annual mean	Relative percentage	Annual mean	Relative percentage	
SW budget					
TOA solar down	334	1	325	1	—
TOA solar up	-118	35.3%	-112	34.5%	0.8%
Atmospheric SW absorption	77	23.1%	77	23.7%	-0.6%
Atmospheric SW reflection	-83	24.9%	-64	19.7%	5.2%
Surface solar down	174	52.1%	184	56.6%	-4.5%
Surface solar up	-35	10.5%	-48	14.8%	-4.3%
Surface solar absorption	139	41.6%	136	41.9%	-0.3%
LW budget					
TOA LW up	-226	65.1%	-232	62.4%	2.7%
Atmospheric LW absorption	-152	43.8%	-166	44.6%	-0.8%
surface LW down	273	78.7%	306	82.3%	-3.6%
Surface LW up	-347	1	-372	1	—

446

447 For the LW budgets, the regional surface LW emission over East Asia is estimated to be much lower
448 than the global land-mean estimates in Wild et al. (2015) (Fig. S3), which mainly results from the lower
449 temperature over the TP induced by high altitudes. The relative percentage of land mean surface downward
450 LW radiation with respect to the surface emission over East Asia is about 78.7 %, which is lower than the
451 global estimate of 82.3%, corresponding well to a reduction in greenhouse effect and fewer low clouds due
452 to the TP (Fig. S4) considering its coverage over East Asian land. Ultimately, a higher percentage of LW
453 radiation is emitted to space over East Asian land compared to global land (65.1% vs. 62.4%). Our estimates
454 also indicate approximately similar amounts of LH (40 vs. 38 W m⁻²) and much lower SH (25 vs. 32 W m⁻²)
455 over East Asia compared to the global land-mean estimates (Fig. S3), which is possibly related to the
456 lower East Asian-land surface temperature.

457 In general, as can be concluded from Table 4, although much less surface SW radiation of 4.3% is
458 reflected over East Asian land compared to global land, a slightly more SW reflection of 0.8% is estimated
459 at the TOA, indicating much larger atmospheric SW reflection of 5.2% due to the stronger scattering from
460 aerosols and clouds over East Asian land than global land. However, the SW absorption within the
461 atmosphere over East Asian land is 0.6% lower than that over global land despite of the more absorption
462 from clouds and aerosols, which is possibly offset by the lower water vapor contents caused by the thinner
463 air over the TP. The lower surface temperature, weaker greenhouse effect and fewer low clouds due to the
464 high altitudes and the thinner air over the TP in East Asian land are the major reasons for the relative lower
465 surface LW emission, less and more fractions of surface downward LW radiation of 3.6% and the OLR of
466 2.7% over East Asian land compared to global land, respectively.


467

468 **4. Assessment of land energy balance budgets under clear-sky conditions**

469 The clear-sky land energy balance budgets over East Asia are similarly evaluated as all-sky conditions.
470 Detailed analyses are given in Supplemental material if interested. The annual land-mean SW clear-sky
471 absorptions at the TOA and surface over East Asia show larger variations among different models than that
472 under all-sky conditions (Fig. 1a and b; Table 1), which is consistent with that reported by Wild et al. (2019)
473 but is amazingly in contrast to the recognition that the representation of clouds is the largest uncertainties in
474 climate models (Dolinar et al., 2015). Specially, the surface SW clear-sky absorptions simulated by various
475 models still exhibit a larger uncertainty than the TOA counterparts despite of the lower absolute values (Fig.
476 1b; Table 1). Contrary to the all-sky counterparts, the simulated clear-sky SSR among different models,
477 shows notably smaller inter-model spread and SD than the surface SW absorptions (Table 1), with much
478 smaller model discrepancy compared to the all-sky conditions (Fig. 2a; Table 1).

479 To further constrain the outlined inter-model discrepancy of the simulated clear-sky SSR, surface
480 observations from the CMA and CERES-interpolated estimates at the GEBA sites are utilized in this study.
481 The high values of the station-based clear-sky SSR are mainly located in the TP, but with an abnormally
482 high value located at the southern China (Fig. 3b). All the East Asian land-mean clear-sky SSR estimates
483 from CERES, CMIP6 multi-model mean, and ERA5 agree reasonably well with the surface observations,

484 but with smaller correlation coefficients ranging from 0.78 to 0.82 compared to the all-sky conditions (Figs.
485 7 b, d, and f). The CERES-derived clear-sky SSR is mainly overestimated in central and western China, but
486 with slight underestimations mainly located in northeastern, eastern, and southern China (Fig. 7a). Similar
487 bias patterns can also be found in the clear-sky SSR from the CMIP6 multi-model mean and ERA5 compared
488 to the surface observations, except for some individual sites over northeastern Inner Mongolia, eastern China,
489 western Mongolia, and Japan (Figs. 7c and e), but with relatively smaller overestimations than the all-sky
490 counterparts (Figs. 4c and e; Table 2). Specifically, the smallest station mean bias in CERES-derived SSR
491 compared to the multi-model mean and ERA5 (Table 2) can be attributed to its even distributed surface sites
492 of overestimations and underestimations (Figs. 7b, d, f). Again, among all the aforementioned clear-sky SSR
493 biases, more overestimations exist in urban stations than the rural stations (b, d, f in Figs. 4 and 7; Table 2).
494 Consequently, all East Asian land-mean area-weighted averages of clear-sky SSR from CERES, CMIP6
495 multi-model mean, and ERA5 show higher overestimations of around 6, 12, and 8 W m^{-2} , respectively,
496 compared to the surface observed counterpart of 230 W m^{-2} (Table 3). Based on the similar method
497 introduced in Wild et al. (2015), the best estimate for the East Asian land-mean clear-sky SSR is determined
498 to be $234 \pm 1.1 \text{ W m}^{-2}$ (2σ uncertainty), with a slightly smaller correlation coefficient of 0.94 and smaller
499 deviations from the CERES and ERA5 estimates compared to the all-sky counterparts (Fig. 5b; Table 3).
500 Besides, the overestimations still exist in the observed land-mean clear-sky SSR for most climate models
501 over East Asia, with a smaller multi-model mean overestimation of 9.1 W m^{-2} than the all-sky counterparts.
502



503

504 **Figure 7.** Spatial distributions of annual mean SSR biases derived from (a) CERES-EBAF, (b) CMIP6
 505 multi-model mean, and (c) ERA5 reanalysis against surface observations from a combination of the CMA
 506 and CERES-interpolated sites under clear-sky conditions over East Asia. The corresponding comparisons of
 507 their respective annual land means at the surface sites with their observed counterparts are displayed in (b),
 508 (d), and (f), respectively. The cross and circle symbols in Figs. a, c, e as well as the orange and green stars
 509 in Figs. b, d, f indicate rural and urban stations, respectively.

510

511 This clear-sky energy budget only represents the removal of cloud but maintains the same atmospheric
 512 conditions as the all-sky conditions, which is not balanced because it is not the equilibrium state the Earth
 513 would achieve when no clouds could form. Ultimately, the clear-sky East Asian land-mean energy budget is
 514 not closed and with no quantifications of SH and LH as displayed in Fig. 6b. In addition to the analyses
 515 above, the clear-sky TOA energy budgets are derived from CERES-derived product, with uncertainty ranges
 516 referred to Loeb et al. (2018), while the surface LW budgets are again from ERA5 reanalysis. Also,

517 additional clear-sky radiation weighted surface albedo of 0.19 from CERES is obtained to estimate the
 518 surface reflected and absorbed SW radiation. Apart from the TOA budget, all the rest uncertainty ranges are
 519 given by different data sources from various CMIP6 models, as well as the multi-model mean, CERES-, and
 520 ERA5-derived estimates.

521 We doublecheck the energy balance components evaluated in this study by referring to the uncertainty
 522 ranges from CERES-derived product given by Kato et al. (2018) (Table 5), which indicates that all estimated
 523 energy components fall within these uncertainty ranges, except for the all-sky surface downward LW
 524 radiation, with about 3 W m⁻² lower than the corresponding lowest CERES range. This is in line with its
 525 much higher CERES-derived estimate compared to that of the ERA5 (285 vs. 273 W m⁻²) (Table 1).

526

527 **Table 5.** Uncertainties (Units: W m⁻²) in 1°×1° regional monthly surface SW, LW, and net (SW + LW)
 528 fluxes under all-sky and clear-sky conditions for the CERES-EBAF Edition 4.1 product (referring to Kato
 529 et al. (2018)), as well as its corresponding estimates of various surface fluxes.

Uncertainties(1 σ)	All-sky	Clear-sky
SW down	178±14	236±6
SW up	36±11	45±11
SW net	142±13	191±13
LW down	285±9	256±8
LW up	354±15	353±15
LW net	69±17	97±17
SW + LW net	73±20	95±20

530

531 Overall, around 21.6% and 56.9% of the TOA incoming solar radiation are absorbed by the atmosphere
 532 and surface, respectively, for clear-sky conditions, while these absorptions are 23.1% and 41.6% for all-sky
 533 conditions. This implies that the existence of clouds results in more atmospheric SW absorption of around
 534 1.5% and much less surface solar absorption of around 15.3% with respect to the TOA incoming solar
 535 radiation.

536

537 5. The cloud radiative effects (CREs)

538 According to the annual land-mean best estimates of radiative components over East Asia under all-sky
 539 and clear-sky conditions obtained in previous sections, the present-day CREs can be inferred quantitatively
 540 over this region. The calculated SW, LW, and net CREs at the TOA, within the atmosphere, and at the
 541 surface are therefore presented in Fig. 8. Moreover, the corresponding calculation formulas are also given in
 542 the followings:

543

$$544 \text{ TOA SW CRE} = \text{TOA outgoing SW}_{\text{all-sky}} - \text{TOA outgoing SW}_{\text{clear-sky}}$$

$$545 \text{ TOA LW CRE} = \text{TOA outgoing LW}_{\text{all-sky}} - \text{TOA outgoing LW}_{\text{clear-sky}}$$

546
$$\text{TOA Net CRE} = \text{TOA SW CRE} + \text{TOA LW CRE}$$

547

548
$$\text{Surface Net SW CRE} = \text{Surface Net SW}_{\text{all-sky}} - \text{Surface Net SW}_{\text{clear-sky}}$$

549
$$\text{Surface Net LW CRE} = \text{Surface Net LW}_{\text{all-sky}} - \text{Surface Net LW}_{\text{clear-sky}}$$


550
$$\text{Surface Net total CRE} = \text{Surface Net SW CRE} + \text{Surface Net LW CRE}$$

551

552
$$\text{Atmospheric SW CRE} = \text{TOA SW CRE} - \text{Surface Net SW CRE}$$

553
$$\text{Atmospheric LW CRE} = \text{TOA LW CRE} - \text{Surface Net LW CRE}$$

554



555

556 **Figure 8.** Diagram of the annual land mean SW, LW, and net (SW + LW) cloud radiative effects (CREs)
 557 (Units: W m^{-2}) at the TOA, within the atmosphere, and at the surface over East Asia, calculated by the
 558 differences between all-sky and clear-sky radiation budgets as given in Fig. 7.

559


560 Best estimates for the annual East Asian land-mean reflected solar radiation at the TOA under all-sky
 561 and clear-sky conditions are -118 and -72 W m^{-2} , respectively, differing by -46 W m^{-2} , indicating that the
 562 clouds give rise to an extra 46 W m^{-2} solar reflection at the TOA, thus cooling the Earth-atmosphere system.
 563 Similarly, the TOA LW CRE, obtained as the difference between the TOA thermal radiation under all-sky

564 and clear-sky conditions, is 24 W m^{-2} , suggesting a warming effect of clouds on the system. Thus, the
565 estimated TOA net CRE is -22 W m^{-2} , pointing out that the overall effects of clouds result in an energy loss
566 and net cooling to the system, not only in the global mean, but also over East Asian land.

567 At the Earth's surface, the shading effects of clouds are estimated to reduce the surface solar radiation
568 by 60 W m^{-2} , from 234 to 174 W m^{-2} , while the surface solar absorption differs by 51 W m^{-2} , from 190 to
569 139 W m^{-2} , namely the surface net SW CRE is -51 W m^{-2} . On cloudy skies, the estimated surface downward
570 LW radiation increases from 253 to 273 W m^{-2} , with an increase of 20 W m^{-2} , illustrating that the surface
571 net LW CRE is 20 W m^{-2} and therefore leads to a surface warming. Thus, the surface net CRE, i.e., the sum
572 of the surface net SW and LW CRE, is then -31 W m^{-2} , indicating that clouds contribute more to the SW
573 energy budgets. Eventually, the clouds lead to the enhancement of the SW and LW absorption within the
574 atmosphere of around 5 and 4 W m^{-2} , respectively, thus resulting in an atmospheric net CRE of 9 W m^{-2} over
575 East Asian land.

576 The above CRE best estimates are compared with the corresponding estimates from different data
577 sources (Fig. 9; Table 1). Generally, compared to the LW CREs (Fig. 9b), the simulated SW CREs show
578 larger spreads and SDs amongst models (Fig. 9a; Table 1). For the SW CREs at the TOA, within the
579 atmosphere, and at the surface, the CERES-derived estimates match perfectly with the best estimates
580 mentioned above, within 2 W m^{-2} of the biases, followed by the estimates from the multi-model means and
581 ERA5 (Table 1). For the LW CREs, the calculated TOA LW CREs from the CMIP6 multi-model mean and
582 CERES differ by no more than 1 W m^{-2} compared to the best estimate, while large differences are noted at
583 the surface LW CREs, thereby leading to their opposite signs in the atmospheric LW CREs (Fig. 9b; Table
584 1). Specifically, since the ERA5-based TOA LW CRE deviates by no more than 3 W m^{-2} with the best
585 estimate of 24 W m^{-2} with nearly the same surface LW CRE, the estimated atmospheric LW CRE is therefore
586 the closest to the best estimate (Table 1). This is owing to the fact that we make use of the ERA5 data as the
587 reference to estimate the surface LW radiation. Thus, the major reason for the large discrepancies in the
588 atmospheric and surface LW CREs estimated from different data sources with respect to the best estimates
589 in this study is the determination of the surface downward and upward LW radiation, which is also the reason
590 for the large deviations in their net CREs (Fig. 9c).

591



592

593 **Figure 9.** Annual land mean anomalies of (a) SW, (b) LW, and (c) net (SW + LW) CREs (Units: $W m^{-2}$) at
 594 the TOA (red line), within the atmosphere (blue line), and at the surface (green line) with regard to their
 595 respective multi-model means over East Asia, respectively, as represented by various CMIP6 models. The
 596 numbers in the parentheses indicate the available CMIP6 climate models for the corresponding radiation
 597 components.

598

599 A better comparison with the global annual mean best estimates of CREs by Wild et al. (2019) is given
 600 in Fig. S5. At the TOA, a slightly lower and much lower East Asian land-mean SW and LW CREs of $1 W m^{-2}$
 601 and $4 W m^{-2}$ result in $3 W m^{-2}$ more energy loss at the TOA compared to the globe. At the surface, much
 602 lower annual East Asian land-mean SW and LW CREs by $3 W m^{-2}$ and $8 W m^{-2}$ are estimated compared to
 603 the values over the globe, leading to a net CRE deviation of $5 W m^{-2}$, indicative of $5 W m^{-2}$ more energy loss
 604 at the surface. However, lower and higher annual East Asian land-mean SW and LW CREs of 2 and $4 W m^{-2}$
 605 within the atmosphere contribute to the nearly close net CRE with a deviation of no more than $2 W m^{-2}$
 606 compared to the global mean estimates. On the whole, lower annual East Asian land-mean best estimates in
 607 the absolute values of surface SW and LW CREs as well as the TOA LW CRE compared to their global
 608 mean counterparts give rise to the CRE differences between them.

609 **6. Summary and conclusions**

610 This study aims to explore how the energy budgets are interrupted by the complex orographic and
611 thermal effects of the TP, as well as the high anthropogenic aerosol emissions over East Asian land compared
612 to global land, based on complementary data sources from space and surface observations, as well as the
613 CMIP6 climate models and ERA5 reanalysis. A further quantitative investigation of CREs at the TOA,
614 within the atmosphere, and at the surface is also conducted.

615 Comparisons between all-sky and clear-sky energy budgets indicate that the overall effects of clouds
616 greatly reduce the surface solar absorption by about 15.3% and enhance that within the atmosphere by 1.5%.
617 Compared to the global land energy budget estimates from Wild et al. (2015), for the SW budgets, notably
618 more atmospheric SW reflection of 5.2% but with a slightly less atmospheric SW absorption of 0.6% with
619 respect to their respective TOA incident solar radiation are estimated over East Asian land, possibly
620 indicating that the lower water vapor content effects due to TP overcompensate for the aerosol and cloud
621 effects over East Asian land. For the LW budgets, a substantially lower surface LW emission of around 25
622 W m^{-2} and smaller relative surface downward LW radiation of around 3.6% with respect to their respective
623 surface emissions can be noticed over East Asian land compared to global land, which possibly result from
624 the lower regional surface skin temperature, as well as the weaker greenhouse effect and fewer low clouds
625 mainly induced by the high altitude and thinner air over TP, thus leading to a higher percentage of regional
626 OLR of 2.7%.

627 The CREs over East Asian land are inferred through the energy budget differences between all-sky and
628 clear-sky conditions. The clouds reduce the solar absorption at the TOA by 46 W m^{-2} and enhance the TOA
629 thermal radiation by 24 W m^{-2} , respectively, leading to a TOA net CRE of -22 W m^{-2} , a more cooling effect
630 on the regional climate system than that over globe (-19 W m^{-2}). At the surface, the net CRE is estimated to
631 be -31 W m^{-2} according to less solar absorption of 51 W m^{-2} and more downward thermal radiation of 20 W
632 m^{-2} , indicative of larger cloud impacts on SW radiation. Within the atmosphere, the estimated net CRE is 9
633 W m^{-2} due to an increase of 5 W m^{-2} of solar absorption and 4 W m^{-2} of the net thermal radiation, respectively.
634 Compared to the global mean best estimates of CREs as introduced by Wild et al. (2019), relatively lower
635 East Asian land-mean best estimates of surface SW and LW CREs as well as the TOA LW CRE contribute
636 to the CRE differences between them.

637 On the whole, all the estimated land-mean energy balance components over East Asia in this study fall
638 within the uncertainty ranges of the CERES-derived assessments, except for the all-sky surface downward
639 LW radiation. More accurate and reliable datasets should be utilized to reduce the substantial uncertainties
640 in the regional energy balance estimates, particularly in the surface budgets, and more widespread temporal
641 and spatial representations of energy budget research are recommended for more comprehensive
642 comparisons in future. For example, newly published surface radiation products with high resolutions based
643 on satellite datasets (e.g., Letu et al., 2022; Xu et al., 2022) are expected to make sense in improving the
644 accuracy of the regional/global surface radiation budget studies.

645 *Acknowledgments.* This research was funded by the National Key Research and Development Program of
646 China (2017YFA0603502) and the Science and Technology Development Fund of CAMS
647 (2021KJ004&2022KJ019). The Global Energy Balance Archive (GEBA) is co-funded by the Federal
648 Office of Meteorology and Climatology Meteo Swiss within the framework of GCOS Switzerland.

649

650 *Data Availability.* The CERES SYN1deg data is available at [https://ceres-tool.larc.nasa.gov/ord-](https://ceres-tool.larc.nasa.gov/ord-tool/jsp/SYN1degEd41Selection.jsp)
651 [tool/jsp/SYN1degEd41Selection.jsp](https://ceres-tool.larc.nasa.gov/ord-tool/jsp/SYN1degEd41Selection.jsp); The AIRS data is accessible from
652 https://disc.gsfc.nasa.gov/datasets/AIRS3STM_006/summary?keywords=AIRS; The MODIS data is from
653 [https://ladsweb.modaps.eosdis.nasa.gov/archive/allData/61/MYD08_M3/?process=ftpAsHttp&path=allDat](https://ladsweb.modaps.eosdis.nasa.gov/archive/allData/61/MYD08_M3/?process=ftpAsHttp&path=allData%2f61%2fMYD08_M3)
654 [a%2f61%2fMYD08_M3](https://ladsweb.modaps.eosdis.nasa.gov/archive/allData/61/MYD08_M3/?process=ftpAsHttp&path=allData%2f61%2fMYD08_M3); The CloudSat data is from [http://www.cloudsat.cira.colostate.edu/data-](http://www.cloudsat.cira.colostate.edu/data-products/level-2b/2b-cwc-ro)
655 [products/level-2b/2b-cwc-ro](http://www.cloudsat.cira.colostate.edu/data-products/level-2b/2b-cwc-ro); The MERRA-2 dataset is obtained at
656 https://disc.gsfc.nasa.gov/datasets/M2IMNPANA_5.12.4/summary?keywords=merra-2. The ERA-Interim
657 is from <https://apps.ecmwf.int/datasets/data/interim-full-moda/levtype=sfc>.

658

659 *Author contributions.* HZ, MW, and QW proposed the main ideas of this study. QW designed and wrote the
660 manuscript. SY provided the homogenized ground-based surface solar radiation data. QC, XZ, and GS
661 contributed to the interpretation of the results. BX and YW assisted with the figures. All co-authors
662 participated in discussions and provided constructive suggestions.

663

664 *Competing interests.* The authors declare that they have no conflict of interest.

665

666 **References**

- 667 Boeke, R. C., and Taylor, P. C.: Evaluation of the Arctic surface radiation budget in CMIP5 models, *J. Geophys.*
668 *Res.-Atmos.*, 121, 8525–8548, <https://doi.org/10.1002/2016JD025099>, 2016.
- 669 Christensen, M. W., Behrangi, A., L’Ecuyer, T. S., Wood, N. B., Lebsock, M. D., and Stephens, G. L.: Arctic
670 observation and reanalysis integrated system: A new data product for validation and climate study, *B. Am.*
671 *Meteorol. Soc.*, 97, 907–916, <https://doi.org/10.1175/BAMS-D-14-00273.1>, 2016.
- 672 de Leeuw, G., Sogacheva, L., Rodriguez, E., Kourtidis, K., Georgoulias, A. K., Alexandri, G., Amiridis, V.,
673 Proestakis, E., Marinou, E., Xue, Y., and van der A, R.: Two decades of satellite observations of AOD over
674 mainland China using ATSR-2, AATSR and MODIS/Terra: data set evaluation and large-scale patterns, *Atmos.*
675 *Chem. Phys.*, 18, 1573–1592, <https://doi.org/10.5194/acp-18-1573-2018>, 2018.
- 676 Dolinar, E. K., Dong, X., Xi, B., Jiang, J. H., and Su, H.: Evaluation of CMIP5 simulated clouds and TOA
677 radiation budgets using NASA satellite observations, *Clim. Dynam.*, 44, 2229–2247,
678 <https://doi.org/10.1007/s00382-014-2158-9>, 2015.
- 679 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the
680 Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci.*
681 *Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 682 Fan, T., Zhao, C., Dong, X., Liu, X., Yang, X., Zhang, F., Shi, C., Wang, Y., and Wu, F.: Quantify contribution
683 of aerosol errors to cloud fraction biases in CMIP5 Atmospheric Model Intercomparison Project simulations,
684 *Int. J. Climatol.*, 38, 3140–3156, <https://doi.org/10.1002/joc.5490>, 2018.
- 685 Fasullo, J. T., and Trenberth, K. E.: The annual cycle of the energy budget. Part I: Global mean and land-ocean
686 exchanges, *J. Climate*, 21, 2297–2312, <https://doi.org/10.1175/2007JCLI1935.1>, 2008a.
- 687 Fasullo, J. T., and Trenberth, K. E.: The annual cycle of the energy budget. Part II: Meridional structures and
688 poleward transports, *J. Climate*, 21, 2313–2325, <https://doi.org/10.1175/2007JCLI1936.1>, 2008b.
- 689 Ghan, S. J., Liu, X., Easter, R. C., Zaveri, R., Rasch, P. J., Yoon, J.-H., and Eaton, B.: Toward a minimal

690 representation of aerosols in climate models: Comparative decomposition of aerosol direct, semidirect, and
691 indirect radiative forcing, *J. Climate*, 25, 6461–6476, <https://doi.org/10.1175/JCLI-D-11-00650.1>, 2012.

692 Gilgen, H., Wild, M., and Ohmura, A.: Means and trends of shortwave irradiance at the surface estimated from
693 global energy balance archive data, *J. Climate*, 11, 2042–2061, [https://doi.org/10.1175/1520-0442\(1998\)011<2042:MATOSI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2042:MATOSI>2.0.CO;2), 1998.

694 He, Y., Wang, K., Zhou, C., and Wild, M.: A revisit of global dimming and brightening based on the sunshine
695 duration, *Geophys. Res. Lett.*, 45, 4281–4289, <https://doi.org/10.1029/2018GL077424>, 2018.

696 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu,
697 R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot,
698 J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes,
699 R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S.,
700 Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and
701 Thépaut, J.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049,
702 <https://doi.org/10.1002/qj.3803>, 2020.

703 Huang, G., Li, Z., Li, X., Liang, S., Yang, K., Wang, D., and Zhang, Y.: Estimating surface solar irradiance from
704 satellites: Past, present, and future perspectives, *Remote Sens. Environ.*, 233, 111371,
705 <https://doi.org/10.1016/j.rse.2019.111371>, 2019.

706 Kato, S., Rose, F. G., Rutan, D. A., Thorsen, T. J., Loeb, N. G., Doelling, D. R., Huang, X., Smith, W. L., Su, W.,
707 and Ham, S.: Surface Irradiances of Edition 4.0 Clouds and the Earth’s Radiant Energy System (CERES)
708 Energy Balanced and Filled (EBAF) data product, *J. Climate*, 31, 4501–4527, <https://doi.org/10.1175/JCLI-D-17-0523.1>, 2018.

709 Kim, B., and Lee, K.: Radiation component calculation and energy budget analysis for the Korean Peninsula
710 region, *Remote Sens.*, 10, 1147, <https://doi.org/10.3390/rs10071147>, 2018.

711 King, M. D., Platnick, S., Menzel, W. P., Ackerman, S. A., and Hubanks, P. A.: Spatial and temporal distribution
712 of clouds observed by MODIS onboard the Terra and Aqua satellites, *IEEE T. Geosci. Remote Sens.*, 51, 3826–
713 3852, <https://doi.org/10.1109/TGRS.2012.2227333>, 2013.

714 L’Ecuyer, T. S., Beaudoin, H. K., Rodell, M., Olson, W., Lin, B., Kato, S., Clayson, C. A., Wood, E., Sheffield,
715 J., Adler, R., Huffman, G., Bosilovich, M., Gu, G., Robertson, F., Houser, P. R., Chambers, D., Famiglietti, J.
716 S., Fetzer, E., Liu, W. T., Gao, X., Schlosser, C. A., Clark, E., Lettenmaier, D. P., and Hilburn, K.: The observed
717 state of the energy budget in the early twenty-first century, *J. Climate*, 28, 8319–8346,
718 <https://doi.org/10.1175/JCLI-D-14-00556.1>, 2015.

719 Lei, Y., Letu, H., Shang, H., and Shi, J.: Cloud cover over the Tibetan Plateau and eastern China: a comparison
720 of ERA5 and ERA-Interim with satellite observations, *Clim. Dynam.*, 54, 2941–2957,
721 <https://doi.org/10.1007/s00382-020-05149-x>, 2020.

722 Letu, H., Nakajima, T. Y., Wang, T., Shang, H., Ma, R., Yang, K., Baran, A. J., Riedi, J., Ishimoto, H., and Yoshida,
723 M.: A new benchmark for surface radiation products over the East Asia–Pacific region retrieved from the
724 Himawari-8/AHI next-generation geostationary satellite, *B. Am. Meteorol. Soc.*, 103, E873–E888,
725 <https://doi.org/10.1175/BAMS-D-20-0148.1>, 2022.

726 Li, J., Waliser, D. E., Stephens, G., Lee, S., L’Ecuyer, T., Kato, S., Loeb, N., and Ma, H.: Characterizing
727 and understanding radiation budget biases in CMIP3/CMIP5 GCMs, contemporary GCM, and reanalysis, *J.*
728 *Geophys. Res.-Atmos.*, 118, 8166–8184, <https://doi.org/10.1002/jgrd.50378>, 2013.

729 Li, J., and Mao, J.: A preliminary evaluation of global and East Asian cloud radiative effects in reanalyses, *Atmos.*
730 *and Ocean. Sci. Lett.*, 8, 100–106, <https://doi.org/10.3878/AOSL20140093>, 2015.

731 Li, J., Mao, J., and Wang, F.: Comparative study of five current reanalyses in characterizing total cloud fraction
732 and top-of-the-atmosphere cloud radiative effects over the Asian monsoon region, *Int. J. Climatol.*, 37, 5047–
733 5067, <https://doi.org/10.1002/joc.5143>, 2017.

734 Li, Z., Niu, F., Fan, J., Liu, Y., Rosenfeld, D., and Ding, Y.: Long-term impacts of aerosols on the vertical
735 development of clouds and precipitation, *Nat. Geosci.*, 4, 888–894, <https://doi.org/10.1038/ngeo1313>, 2011.

736 Liao, H., Chang, W., and Yang, Y.: Climatic effects of air pollutants over china: A review, *Adv. Atmos. Sci.*, 32,
737 115–139, <https://doi.org/10.1007/s00376-014-0013-x>, 2015.

738 Lin, B., Stackhouse Jr., P. W., Minnis, P., Wielicki, B. A., Hu, Y., Sun, W., Fan, T., and Hinkelman, L. M.:
739 Assessment of global annual atmospheric energy balance from satellite observations, *J. Geophys. Res.-Atmos.*,
740 113, <https://doi.org/10.1029/2008JD009869>, 2008.

741 Liu, Y., Bao, Q., Duan, A., Qian, Z. A., and Wu, G.: Recent progress in the impact of the Tibetan Plateau on
742 climate in China, *Adv. Atmos. Sci.*, 24, 1060–1076, <https://doi.org/10.1007/s00376-007-1060-3>, 2007.

743 Loeb, N. G., Doelling, D. R., Wang, H., Su, W., Nguyen, C., Corbett, J. G., Liang, L., Mitrescu, C., Rose, F. G.,
744 and Kato, S.: Clouds and the Earth’s Radiant Energy System (CERES) Energy Balanced and Filled (EBAF)
745 Top-of-Atmosphere (TOA) Edition-4.0 data product, *J. Climate*, 31, 895–918, <https://doi.org/10.1175/JCLI-D-17-0208.1>, 2018.

749 Mayer, M., Tietsche, S., Haimberger, L., Tsubouchi, T., Mayer, J., and Zuo, H.: An improved estimate of the
750 coupled Arctic energy budget, *J. Climate*, 32, 7915–7934, <https://doi.org/10.1175/JCLI-D-19-0233.1>, 2019.

751 Mercado, L. M., Bellouin, N., Sitch, S., Boucher, O., Huntingford, C., Wild, M., and Cox, P. M.: Impact of
752 changes in diffuse radiation on the global land carbon sink, *Nature*, 458, 1014–1017,
753 <https://doi.org/10.1038/nature07949>, 2009.

754 Ohmura, A.: Cryosphere During the Twentieth Century, *The state of the planet: frontiers and challenges in
755 geophysics*, *Geophys. Monogr. Ser.*, 150, 239–257, <https://doi.org/10.1029/150gm19>, 2004.

756 Previdi, M., Smith, K. L., and Polvani, L. M.: How well do the CMIP5 models simulate the Antarctic atmospheric
757 energy budget? *J. Climate*, 28, 7933–7942, <https://doi.org/10.1175/JCLI-D-15-0027.1>, 2015.

758 Raschke, E., Kinne, S., Rossow, W. B., Stackhouse, P. W., and Wild, M.: Comparison of radiative energy flows
759 in observational datasets and climate modeling, *J. Appl. Meteorol. Clim.*, 55, 93–117,
760 <https://doi.org/10.1175/JAMC-D-14-0281.1>, 2016.

761 Simmons, A. J., Jones, P. D., Da Costa Bechtold, V., Beljaars, A. C. M., Källberg, P. W., Saarinen, S., Uppala, S.
762 M., Viterbo, P., and Wedi, N.: Comparison of trends and low-frequency variability in CRU, ERA-40, and
763 NCEP/NCAR analyses of surface air temperature, *J. Geophys. Res.-Atmos.*, 109,
764 <https://doi.org/10.1029/2004JD005306>, 2004.

765 Stephens, G. L.: Cloud feedbacks in the climate system: A critical review, *J. Climate*, 18, 237–273,
766 <https://doi.org/10.1175/JCLI-3243.1>, 2005.

767 Stephens, G. L., Li, J., Wild, M., Clayson, C. A., Loeb, N., Kato, S., L'Ecuyer, T., Stackhouse, P. W., Lebsock,
768 M., and Andrews, T.: An update on Earth's energy balance in light of the latest global observations, *Nat. Geosci.*,
769 5, 691–696, <https://doi.org/10.1038/ngeo1580>, 2012.

770 Tang, W. J., Qin, J., Yang, K., Zhu, F., and Zhou, X.: Does ERA5 outperform satellite products in estimating
771 atmospheric downward longwave radiation at the surface? *Atmos. Res.*, 252, 105453,
772 <https://doi.org/10.1016/j.atmosres.2021.105453>, 2021.

773 Tang, W. J., Yang, K., Qin, J., Cheng, C. C. K., and He, J.: Solar radiation trend across China in recent decades:
774 a revisit with quality-controlled data, *Atmos. Chem. Phys.*, 11, 393–406, <https://doi.org/10.5194/acp-11-393-2011>, 2011.

775 Thomas, C. M., Dong, B., and Haines, K.: Inverse modeling of global and regional energy and water cycle fluxes
776 using earth observation data, *J. Climate*, 33, 1707–1723, <https://doi.org/10.1175/JCLI-D-19-0343.1>, 2020.

777 Trenberth, K. E., Fasullo, J. T., and Balmaseda, M. A.: Earth's energy imbalance, *J. Climate*, 27, 3129–3144,
778 <https://doi.org/10.1175/JCLI-D-13-00294.1>, 2014.

779 Trenberth, K. E., Fasullo, J. T., and Kiehl, J.: Earth's global energy budget, *B. Am. Meteorol. Soc.*, 90, 311–324,
780 <https://doi.org/10.1175/2008BAMS2634.1>, 2009.

781 Trolliet, M., Walawender, J. P., Bourlès, B., Boilley, A., Trentmann, J., Blanc, P., Lefèvre, M., and Wald, L.:
782 Downwelling surface solar irradiance in the tropical Atlantic Ocean: a comparison of re-analyses and satellite-
783 derived data sets to PIRATA measurements, *Ocean Sci.*, 14, 1021–1056, <https://doi.org/10.5194/os-14-1021-2018>, 2018.

784 Urraca, R., Huld, T., Gracia-Amillo, A., Martinez-de-Pison, F. J., Kaspar, F., and Sanz-Garcia, A.: Evaluation of
785 global horizontal irradiance estimates from ERA5 and COSMO-REA6 reanalyses using ground and satellite-
786 based data, *Sol. Energy*, 164, 339–354, <https://doi.org/10.1016/j.solener.2018.02.059>, 2018.

787 Wang, H., Zhang, H., Xie, B., Jing, X., He, J., and Liu, Y.: Evaluating the Impacts of Cloud Microphysical and
788 Overlap Parameters on Simulated Clouds in Global Climate Models, *Adv. Atmos. Sci.*,
789 <https://doi.org/10.1007/s00376-021-0369-7>, 2021.

790 Wang, K.: Measurement biases explain discrepancies between the observed and simulated decadal variability of
791 surface incident solar radiation, *Sci. Rep.*, 4, 6144, <https://doi.org/10.1038/srep06144>, 2014.

792 Wang, K., Ma, Q., Li, Z., and Wang, J.: Decadal variability of surface incident solar radiation over China:
793 Observations, satellite retrievals, and reanalyses, *J. Geophys. Res.-Atmos.*, 120, 6500–6514,
794 <https://doi.org/10.1002/2015JD023420>, 2015.

795 Wang, Q., Zhang, H., Yang, S., Chen, Q., Zhou, X., Shi, G., Cheng, Y., and Wild, M.: Potential driving factors
796 on surface solar radiation trends over China in recent years, *Remote Sens.*, 13, 704,
797 <https://doi.org/10.3390/rs13040704>, 2021.

798 Wang, Y., Trentmann, J., Yuan, W., and Wild, M.: Validation of CM SAF CLARA-A2 and SARA-E surface
799 solar radiation datasets over China, *Remote Sens.*, 10, 1977, <https://doi.org/10.3390/rs10121977>, 2018.

800 Wang, Y., Wild, M., Sanchez-Lorenzo, A., and Manara, V.: Urbanization effect on trends in sunshine duration in
801 China, *Ann. Geophys.*, 35, 839–851, <https://doi.org/10.5194/angeo-35-839-2017>, 2017.

802 Wang, Y., and Wild, M.: A new look at solar dimming and brightening in China, *Geophys. Res. Lett.*, 43, 11,
803 711–777, 785, <https://doi.org/10.1002/2016GL071009>, 2016.

804 Wang, Z., Lin, L., Xu, Y., Che, H., Zhang, X., Zhang, H., Dong, W., Wang, C., Gui, K., and Xie, B.: Incorrect
805 Asian aerosols affecting the attribution and projection of regional climate change in CMIP6 models, *npj Clim.*
806 807

808 Atmos. Sci., 4, 1–8, <https://doi.org/10.1038/s41612-020-00159-2>, 2021.

809 Wei, J., Peng, Y., Guo, J., and Sun, L.: Performance of MODIS Collection 6.1 Level 3 aerosol products in spatial-
810 temporal variations over land, *Atmos. Environ.*, 206, 30–44, <https://doi.org/10.1016/j.atmosenv.2019.03.001>,
811 2019.

812 Wild, M.: Progress and challenges in the estimation of the global energy balance, *AIP Conference Proceedings*,
813 1810, 20004, <https://doi.org/10.1063/1.4975500>, 2017a.

814 Wild, M.: Towards global estimates of the surface energy budget, *Curr. Clim. Change Rep.*, 3, 87–97,
815 <https://doi.org/10.1007/s40641-017-0058-x>, 2017b.

816 Wild, M.: The global energy balance as represented in CMIP6 climate models, *Clim. Dynam.*, 55, 553–577,
817 <https://doi.org/10.1007/s00382-020-05282-7>, 2020.

818 Wild, M., Folini, D., Hakuba, M. Z., Schär, C., Seneviratne, S. I., Kato, S., Rutan, D., Ammann, C., Wood, E. F.,
819 and König-Langlo, G.: The energy balance over land and oceans: an assessment based on direct observations
820 and CMIP5 climate models, *Clim. Dynam.*, 44, 3393–3429, <https://doi.org/10.1007/s00382-014-2430-z>, 2015.

821 Wild, M., Folini, D., Schär, C., Loeb, N., Dutton, E. G., and König-Langlo, G.: The global energy balance from
822 a surface perspective, *Clim. Dynam.*, 40, 3107–3134, <https://doi.org/10.1007/s00382-012-1569-8>, 2013a.

823 Wild, M., Folini, D., Schär, C., Loeb, N., Dutton, E. G., and König-Langlo, G.: A new diagram of the global
824 energy balance, *AIP Conference Proceedings*, 1531, 628–631, <https://doi.org/10.1063/1.4804848>, 2013b.

825 Wild, M., Grieser, J., and Schär, C.: Combined surface solar brightening and increasing greenhouse effect support
826 recent intensification of the global land-based hydrological cycle, *Geophys. Res. Lett.*, 35,
827 <https://doi.org/10.1029/2008GL034842>, 2008.

828 Wild, M., Hakuba, M. Z., Folini, D., Dörig-Ott, P., Schär, C., Kato, S., and Long, C. N.: The cloud-free global
829 energy balance and inferred cloud radiative effects: an assessment based on direct observations and climate
830 models, *Clim. Dynam.*, 52, 4787–4812, <https://doi.org/10.1007/s00382-018-4413-y>, 2019.

831 Wild, M., Hakuba, M. Z., Folini, D., Schär, C., and Long, C.: New estimates of the Earth radiation budget under
832 cloud-free conditions and cloud radiative effects, *AIP Conference Proceedings*, 1810, 90012,
833 <https://doi.org/10.1063/1.4975552>, 2017.

834 Wild, M., Ohmura, A., Gilgen, H., and Roeckner, E.: Validation of general circulation model radiative fluxes
835 using surface observations, *J. Climate*, 8, 1309–1324, [https://doi.org/10.1175/1520-0442\(1995\)008<1309:VOGCMR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<1309:VOGCMR>2.0.CO;2), 1995.

836 Wu, G., Duan, A., Liu, Y., Mao, J., Ren, R., Bao, Q., He, B., Liu, B., and Hu, W.: Tibetan Plateau climate
837 dynamics: recent research progress and outlook, *Natl. Sci. Rev.*, 2, 100–116,
838 <https://doi.org/10.1093/nsr/nwu045>, 2015.

839 Xu, J., Liang, S., and Jiang, B.: A global long-term (1981–2019) daily land surface radiation budget product from
840 AVHRR satellite data using a residual convolutional neural network, *Earth Syst. Sci. Data*, 14, 2315–2341,
841 <https://doi.org/10.5194/essd-14-2315-2022>, 2022.

842 Xu, X., Lu, C., Shi, X., and Gao, S.: World water tower: An atmospheric perspective, *Geophys. Res. Lett.*, 35,
843 <https://doi.org/10.1029/2008GL035867>, 2008a.

844 Xu, X., Zhang, R., Koike, T., Lu, C., Shi, X., Zhang, S., Bian, L., Cheng, X., Li, P., and Ding, G.: A new integrated
845 observational system over the Tibetan Plateau, *B. Am. Meteorol. Soc.*, 89, 1492–1496, 2008b.

846 Yang, S., Wang, X. L., and Wild, M.: Homogenization and trend analysis of the 1958–2016 in situ surface solar
847 radiation records in China, *J. Climate*, 31, 4529–4541, <https://doi.org/10.1175/JCLI-D-17-0891.1>, 2018.

848 Yang, S., Wang, X. L., and Wild, M.: Causes of dimming and brightening in China inferred from homogenized
849 daily clear-sky and all-sky in situ surface solar radiation records (1958–2016), *J. Climate*, 32, 5901–5913,
850 <https://doi.org/10.1175/JCLI-D-18-0666.1>, 2019.

851 You, Q., Liu, J., and Pepin, N.: Changes of summer cloud water content in China from ERA-Interim reanalysis,
852 *Global Planet. Change*, 175, 201–210, <https://doi.org/10.1016/j.gloplacha.2019.02.014>, 2019.

853 Zhang, H., Zhao, M., Chen, Q., Wang, Q., Zhao, S., Zhou, X., and Peng, J.: Water and ice cloud optical thickness
854 changes and radiative effects in East Asia, *J. Quant. Spectrosc. Radiat. Transf.*, 254, 107213,
855 <https://doi.org/10.1016/j.jqsrt.2020.107213>, 2020.