Simulating organic aerosol in Delhi with WRF-Chem using the VBS 1

approach: Exploring model uncertainty with a Gaussian Process 2

emulator 3

- Ernesto Reyes-Villegas^{1,a}, Douglas Lowe^{1,b}, Jill S. Johnson^{2,c}, Kenneth S. Carslaw², Eoghan Darbyshire^{1,d}, 4
- Michael Flynn¹, James D. Allan^{1,e}, Hugh Coe¹, Ying Chen³, Oliver Wild³, Scott Archer-Nicholls⁴, Alex 5
- Archibald⁴, Siddhartha Singh⁵, Manish Shrivastava⁶, Rahul A, Zaveri⁶, Vikas Singh⁷, Gufran Beig⁸, 6
- Ranjeet Sokhi⁹, Gordon McFiggans¹ 7
- 8 ¹Department of Earth and Environmental Sciences, University of Manchester, Manchester, M13 9PL, UK
- 9 ²Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK
- 10 ³Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK
- 11 ⁴NCAS-Climate, Department of Chemistry, University of Cambridge, Cambridge, CB2 1EW, UK
- 12 ⁵Ozone Unit, India Meteorology Department, New Delhi, 110003, India
- 13 ⁶Atmospheric Sciences & Global Change Division, Pacific Northwest National Laboratory, Richland, Washington 993522, 14 United States
- 15 ⁷National Atmospheric Research Laboratory, Gadanki, AP, India
- 16 ⁸Indian Institute of Tropical Meteorology, Pune, 411008, India
- 17 ⁹Centre for Atmospheric and Climate Physics Research, University of Hertfordshire, Hertfordshire, AL10 9AB, UK
- 18 ^a now at: Tecnologico de Monterrey, Escuela de Ingeniería y Ciencias, Ay, General Ramon Corona 2514, Nuevo Mexico, 19 Zapopan CP 45201, Jalisco, Mexico. School of Engineering and Sciences, Tecnologico de Monterrey, Guadalajara, 45201, 20 Mexico
- 21
- 22 ^b now at: IT Services, University of Manchester, Manchester, M13 9PL, UK
- 23 ^c now at University of Sheffield, Sheffield, S10 2TN, UK
- 24 ^d now at: The Conflict and Environment Observatory, Hebden Bridge, HX7 5HZ, UK
- 25 ^e National Centre for Atmospheric Science, The University of Manchester, Manchester, M13 9PL, UK
- 26

27 Correspondence to: Gordon McFiggans (g.mcfiggans@manchester.ac.uk)

28 Abstract. The nature and origin of organic aerosol in the atmosphere remain unclear. The gas-particle partitioning of semi-29 volatile organic compounds (SVOC) that constitute primary organic aerosols (POA) and the multigenerational chemical aging 30 of SVOCs are particularly poorly understood. The volatility basis set (VBS) approach, implemented in air quality models such 31 as WRF-Chem, can be a useful tool to describe emissions of POA and its chemical evolution POA production and aging. 32 However, the main disadvantage is its complexity, making the evaluation of model uncertainty and the optimal model 33 parameterisation maybe expensive to probe using only WRF-Chem simulations. Gaussian process emulators, trained on 34 simulations from relatively few WRF-Chem simulations, are capable of reproducing model results and estimating the sources 35 of model uncertainty within a defined range of model parameters. In this study, a WRF-Chem VBS parameterisation is 36 proposed; we then generate a perturbed parameter ensemble of 111 model runs, perturbing ten parameters of the WRF-Chem 37 model relating to organic aerosol emissions and the VBS oxidation reactions. This allowed us to cover the model's uncertainty 38 space and compare output from each run to aerosol mass spectrometer observations of organic aerosol concentrations and O:C 39 ratios measured in New Delhi, India. The simulations spanned the organic aerosol concentrations measured with the AMS. 40 However, they also highlighted potential structural errors in the model that may be related to unsuitable diurnal cycles in the 41 emissions and/or failure to adequately represent the dynamics of the planetary boundary layer. While the structural errors 42 prevented us from clearly identifying an optimised VBS approach in WRF-Chem, we were able to apply the emulator in two 43 periods: the full period (1st -29th May) and the <u>a subperiod</u> period 14:00- 16:00 hrs local time, 1st-29th May. The combination 44 of emulator analysis and model evaluation metrics allowed us to identify plausible parameter combinations for the analysed 45 periods. We demonstrate that the methodology presented in this study can be used to determine the model uncertainty and 46 identify the appropriate parameter combination for the VBS approach, and hence provide valuable information to improve our 47 understanding on <u>SOA</u> production.

48 1 Introduction

49 Over the last decades, India has been facing air pollution problems and is ranked fifth in the 2020 world air quality ranking 50 (IQair, 2021) and Delhi ranked as one of the most polluted cities in the world with related health burden of about 10,000 51 premature deaths annually (Chen et al., 2020a), based on $PM_{2.5}$ measurements (particulate matter lower than 2.5 micrometers 52 in diameter). This situation has a remarkable impact on Indian citizens due to India having a population that is larger than one 53 billion inhabitants.

54 Organic aerosols (OA) are one of the main constituents of submicron particulate matter, accounting for between 20% - 90%55 of the total aerosol mass concentration globally-in urban environments (Kanakidou et al., 2005;Zhang et al., 2007). Various 56 studies have been performed in India looking at the particulate matter composition and source identification of OA using 57 receptor modelling tools (Kompalli et al., 2020; Jain et al., 2020; Cash et al., 2021; Reves-Villegas et al., 2021) along with 58 investigating the health risks associated with aerosols (Shivani et al., 2019;Gadi et al., 2019). However, one limitation of 59 receptor models is that they do not involve chemical processing. The use of regional atmospheric models allows the study of 60 the temporal and spatial behaviour of various chemical species of OA. The Weather Research and Forecasting model coupled 61 with Chemistry (WRF-Chem) is a regional 3-D atmospheric model that simulates the emissions and dispersion of gaseous and 62 particulate species, including the chemical processes and their interaction with meteorology. There have been recent WRF-63 Chem studies investigating PM_{2.5} concentrations (Bran and Srivastava, 2017;Chen et al., 2020b;Jat et al., 2021; Ghosh et al., 64 2021) and volatile organic compounds (VOC) compounds (Chutia et al., 2019) over India.

65 Despite the recent studies on aerosol sources and processes involving both observations and modelling, there is still a gap 66 between observations and modelling studies, for example with particulate organic matter being generally underestimated by 67 models (Bergström et al., 2012; Tsigaridis et al., 2014), mainly attributed to the lack of understanding of the emission sources, 68 and the POA processes and SOA mechanisms. Hence, we need to understand the capability of organic matter to produce and 69 retain fine particulate mass in order to fully understand their processes and impacts on air quality and climate (Carlton et al., 70 2010; von Schneidemesser et al., 2015). It is here where the volatility basis set (VBS) scheme can be valuable when 71 implemented in chemical transport models. The VBS scheme describes the chemical ageing of particulate organic matter, its 72 chemical processing and associated volatility (Donahue et al., 2006;Shrivastava et al., 2011; Bianchi et al., 2019). It treats 73 POA emissions as semi volatile and distributes particulate organic matter by its volatility. This distribution, based on their 74 saturation concentration (C^{*}), includes low volatility (LVOC), semivolatile (SVOC) and intermediate volatility (IVOC) 75 organic compounds (Tsimpidi et al., 2016). POA constitutes emissions from anthropogenic combustion processes and open 76 biomass burning (Stewart et al., 2021a; Stewart et al., 2021b) and by being considered to be semivolatile, the initial particulate 77 organic matter partially evaporates due to atmospheric dilution followed by the oxidation of evaporated semi-volatile organic 78 vapors. The resulting low volatility oxidized organic vapors can condense to produce oxidized primary organic aerosols 79 (oPOA)secondary organic aerosol (SOA) (Shrivastava et al., 2008). This favours the formation of IVOCs and SVOCs in the 80 gas phase. Previous studies have found that IVOCs and SVOCs can act as a reservoir of organic species that are able to 81 repartition to the particle phase after suffering chemical processing (Robinson et al., 2007;Lane et al., 2008).

Regional (Li et al., 2016; Akherati et al., 2019) and global models (Tsigaridis et al., 2014; Tilmes et al., 2019) have been
 successfully used to simulate aerosol dispersion and chemical processing to some extent. However, they can be highly

uncertain (Bellouin et al., 2016;Johnson et al., 2020), particularly when comparing with on-site observations in a high time
resolution. This uncertainty can be due to a wide range of parameter settings, emission sources or missing processes, and is
challenging to comprehensively evaluate by only running direct model simulations, due to the computing time and expense
needed. Statistical analysis to evaluate model performance over parameter uncertainty can be made tractable through the use
of a statistical emulator (Carslaw et al., 2018). With a trained emulator, it is possible to study thousands or millions of model
variants (parameter combinations) and estimate the sources of uncertainty (Lee et al., 2011;Johnson et al., 2018;Wild et al.,
2020)

91 The VBS approach is often tuned to the environment of interest (Bergström et al., 2012; Shrivastava et al., 2013; Tilmes et al., 92 2019;Shrivastava et al., 2019;Shrivastava et al., 2022) and, as mentioned before, doing this only with WRF-Chem runs is 93 particularly challenging and time consuming. The aim of this study is to determine an effective way of tuning the VBS scheme 94 using observations, and also to learn about the processes controlling OA in Delhi. Hence, we need to explore the combination 95 of different techniques, i.e., observations, WRF-Chem modelling with VBS implementation and statistical emulators, to better 96 understand the partitioning of matter between gaseous and particulate phases, of organic matter and the chemical 97 agingevolution of POA. In this study, a WRF-Chem parameterisation is proposed to simulate organic mass concentrations and 98 organic to carbon (O:C) ratios over the region of New Delhi, India, that includes detailed primary and ageing parameters in 99 the VBS and secondary organic aerosol (SOA) formation schemes. In this parameterisation we explore the perturbation to the 100 chosen anthropogenic POA and biomass burning POA parameters that would be needed to give the best fit to the observed 101 OA, without perturbing the SOA parameters from the base case. The model performance is evaluated over a multi-dimensional 102 parameter uncertainty space that explores parameter uncertainty in these schemes. We generate a perturbed parameter 103 ensemble (PPE) of 111 model runs that cover the model's uncertainty space and compare output from each run to AMS 104 observations of OA concentrations and O:C ratios measured at New Delhi, India. The PPE is then used to construct statistical 105 emulators and sample densely over the uncertainty for a more detailed comparison over a specific time-period of the 106 observations. The evaluation over specific time-periods will allow to study the behaviour of the model setup under different 107 conditions, i.e., high vs low mass concentrations, and analyse the impact the different parameter setups have on the organic 108 mass concentrations.

109 2 Methodology

110 2.1 WRF-Chem parameterisation and setup

111 The Weather Research and Forecasting model coupled with Chemistry (WRF-Chem) is used to simulate the emission, 112 transport, mixing, and chemical transformation of trace gases and aerosols concurrently with meteorology data (Grell et al., 113 2005; Fast et al., 2006). Here, WRF-Chem version 3.8.1 is run with a 15 km domain, 12755 x 12755 grid cells, (Figure 1) and 114 a simulation period from 19th April - 29th May 2018, with substantial modification, details in below. This period was selected 115 in order to compare with aerosol measurements performed at New Delhi (Reyes-Villegas et al., 2021). Table 1 lists the 116 components that contribute to our model set-up, including the chemistry and aerosol schemes, emissions inventories and 117 boundary condition specifications. Gas-phase chemistry is simulated with the Common Representative Intermediates (CRI) 118 mechanism which permits a reasonably detailed representation of volatile organic compound oxidation. The aerosol chemistry 119 is simulated using the sectional MOSAIC module (Zaveri et al., 2008), including N₂O₅ heterogeneous chemical reactions 120 (Archer-Nicholls et al., 2014;Bertram and Thornton, 2009) and is coupled to the aqueous phase, which allows aerosols to act 121 as cloud condensation nuclei, as well as the removal of aerosols through wet deposition processes. The aerosol size distribution 122 in MOSAIC is described by eight size bins spanning a dry particle diameter range of 39nm to 10µm (Zaveri et al., 2008).

123 Table 1: WRF-Chem setup

Parameter Set up

Gas phase mechanism	CRI-v2R5 (Watson et al., 2008;Archer-Nicholls et al., 2014)
Aerosol module	MOSAIC (Zaveri et al., 2008;Fast et al., 2006)
	with VBS (Shrivastava et al., 2011)
	with SOA (Tsimpidi et al., 2010)
Anthropogenic emissions	EDGAR-HTAP and SAFAR-India (CRI-v2R5 speciation)
Fire emissions	FINN 1.5 (Wiedinmyer et al., 2011)
Biogenic emissions	MEGAN <u>2.04</u> (Guenther et al., 2006)
Chemical Boundaries	CESM2/WACCM (Danabasoglu et al., 2020)
Meteorological Boundaries	ECMWF Reanalysis (Hersbach et al., 2018)

125 Our main modifications are focused on the treatment of the organic aerosol (OA) components. Primary organic aerosol (POA) 126 is treated as semi-volatile, using the Volatility Basis Set (VBS) treatment of Shrivastava et al. (2011). Their 9 volatility bin 127 VBS scheme has been adapted for use in the 8 size bin version of MOSAIC. Secondary organic aerosol (SOA) has been 128 included based on the scheme described in Tsimpidi et al. (2010), providing 'anthropogenic' (ARO1 and ARO2 in the original 129 scheme, SAPRC99) and 'biogenic' (iIsoprene and monoterpenes) SOA components, each covering 4 volatility bins with C* values (at 298 K) of 1, 10, 100, and 1000 μ g.m-3. ARO1 represents the aromatics with OH reaction rates less than $2x10^4$ ppm⁻¹ 130 131 min⁻¹, and ARO2 the aromatics with OH reaction rates greater than $2x10^4$ ppm⁻¹ min⁻¹. In mapping these to the CRI-v2R5 132 scheme we have used TOLUENE and BENZENE as the precursors for the ARO1 reactions, OXYL (xylene and other 133 aromatics) for the ARO2 reactions, and APINENE for the monoterpenes. Indicative SOA yields are given in Table S1 in the 134 supplementary material. Co-condensation of water has been added for these semi-volatile organics, and they have been coupled 135 to the aqueous phase in the same manner as other aerosol compounds in MOSAIC. Previous studies have demonstrated that 136 the condensation of semivolatile organic material onto aerosol particles substantially increases the soluble mass of particles, 137 their chemical composition and eventually their effective dry size (Topping et al., 2013; Crooks et al., 2018). The mapping of 138 CESM2/WACCM compounds to CRI-v2R5 and MOSAIC components, for the chemical boundaries, is detailed in Table S2 139 in the supplementary material. A spin-up period of 11 days was used, from 19th April to 1st May. The meteorological driving 40 fields were taken from ERA5 reanalysis data. Spectral nudging of the uv wind parameters, temperature and geopotential height variables to these, above model level 18 and for wavelengths greater than 950km, was used. The domain is conformed of 38 41 142 model layers, variable height and terrain following, model levels, up to a pressure of 50 hPa. The first model layer has a mean 143 height of 59m over Delhi (and a mean height of 56m over the whole model domain).

Previous studies using the VBS have used scaling factors from POA to derive SVOC emissions in each volatility bin based on equilibrium partitioning calculations, as well as volatility distributions based on laboratory studies and assumed oxygenation and chemical reaction rates (Shrivastava et al., 2011;Fountoukis et al., 2014). To investigate the impact of these assumptions on the model predictions, we have modified the model code so that the VBS emissions, the oxygenation rates and VBS reaction rates, can be directly controlled via namelist options. The parameters which are perturbed in this way for this study are described in more detail in Section 2.3.

150 The volatility distribution for of open biomass burning emissions is taken from May et al. (2013), and multiplied by a scaling 151 factor of 3 (based on equilibrium partitioning calculations) to ensure reasonably similar condensed mass at emission as that 152 reported in the FINN 1.5 emission dataset. Similar calculations have been made in previous studies, giving roughly the same 153 scaling factor (Shrivastava et al., 2011; Fountoukis et al., 2014; Denier Van Der Gon et al., 2015; Ciarelli et al., 2017). The 154 volatility distribution for anthropogenic emissions is also multiplied by a scaling factor of 3 for the same reasons as above. It 155 is worth mentioning that the perturbed space explored here is embedded in the parent VBS scheme that has been adopted. 156 There have been a large number of developments in, and variants of, the VBS aiming to address particular questions related 157 to SOA formation at various levels of complexity; for example, the mechanistic measurement-constrained radical 2D-VBS 158 examining the role of ELVOC and ULVOC in new particle formation; (Zhao et al., 2020; Zhao et al., 2021). In the current study, our implementation has been developed from the VBS version available in the distribution version of WRF-Chem and 159 160 our results should be interpreted in the context of the structural capabilities and limitations therein. More information about 161 the VBS distributions and parameter space setup is in section S1 in the supplementary material.

Anthropogenic emissions are derived from the EDGAR-HTAP, SAFAR-India (CRI-v2R5 speciation) and NMVOC global
 emission datasets, with NMVOC emissions speciated for the CRI-v2R5 chemical scheme, and applying diurnal activity cycles

- to the emissions based on emission sectors in Europe (Olivier et al., 2003). We used these diurnal activity cycles (Figure S1
- in supplement) as there were no data available for activity behavior in Delhi. Biogenic emissions are calculated online using
- the MEGAN model (Guenther et al., 2006). Biomass burning emissions are taken from the FINNv1.5 global inventory
- **167** (Wiedinmyer et al., 2011).



Figure 1. WRF-Chem model domain with PM₁-concentrationstopography data. White The red marker highlights the location of IMD New Delhi, where the AMS observations were taken and the red rectangle shows the area that covers the model results.

<u>resul</u>

169

173 2.2 Observations

174 Aerosol observations were made at the Indian Meteorology Department (IMD) at Lodhi road in New Delhi, India (Lat 28.588, 175 Lon 77.217) from 26th April to 30th May 2018 as part of the PROMOTE campaign (Reves-Villegas et al., 2021). A High-176 Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-TOF-AMS, Aerodyne Research Inc.), hereafter referred to as 177 AMS, was used to measure mass spectra of non-refractory particulate matter with an aerodynamic diameter equal or lower 178 than 1 µm (PM₁), including organic aerosols (OA), sulphate (SO₄²⁻), nitrate (NO₃⁻), ammonium (NH₄⁺) and chloride (Cl⁻), in a 179 5-minute time resolution. The AMS operation principle has been previously described by DeCarlo et al. (2006). The AMS was 180 calibrated during the campaign for the ionisation efficiency of nitrate (IE) and the relative ionisation efficiency (RIE) of other 181 inorganic compounds using nebulised ammonium nitrate and ammonium sulphate with a diameter of 300 nm. The data were 182 analysed using the IGOR Pro (WaveMetrics, Inc., Portland, OR, USA) based software SQUIRREL (Sequential Igor data 183 Retrieval) v.1.63I and PIKA (Peak Integration by Key Analysis) v.1.23I. The organic to carbon (O:C) ratios were calculated 184 with PIKA using the improved-Ambient elemental analysis method for AMS spectra measured in air (Canagaratna et al., 185 2015). The AMS data, OA mass concentrations and O:C ratios, are used to compare with the WRF-Chem model outputs: total 186 organic matter mass concentration (Total OM) and organic to carbon ratios (OC ratio).

There were no Planetary boundary layer height (PBLH) measurements available at IMD Lodhi road, hence, PBLH data were sourced from ECMWF ERA5 with 0.25 deg. results in 1-hour resolution for the coordinates closest to the IMD site.
Meteorology data was downloaded from https://ncdc.noaa.gov/ (last access: 05/01/2019) for the Indira Gandhi International Airport, India meteorology station.

191 The meteorology data were used to interpret the diurnal behaviour of the chemical species and to compare with meteorology 192 outputs from WRF-Chem. A dataset of meteorology was not available at IMD. The use of meteorology from airports has been 193 previously used and is considered to be representative of regional meteorology without being affected by surrounding buildings

194 (Reyes-Villegas et al., 2016).

195 2.3 Perturbed Parameter Ensemble

To evaluate the sensitivity to variations in the VBS emission and processing parameters of our WRF-Chem model of the simulated OA over the New Delhi region, we generated a perturbed parameter ensemble (PPE). We choose a set of simulations with optimal space-filling properties that provide effective coverage across the multi-dimensional space of the uncertain model parameters. Here, we perturb ten parameters of the WRF-Chem model that relate to <u>semi-volatile P</u>OA emissions and the aging of these VBS compounds. The parameters correspond to five processes in the model, which are perturbed with respect to both anthropogenic emissions and biomass burning emissions. These process parameters are:

- 2021.VBS ageing rate: The reaction rates of VBS compounds with OH each reaction reduces the volatility of the
compound by a factor of 10 (1 decade in saturation concentration, Ci*, position), and adds between 7.5% and 40%
oxygen (determined by the SVOC oxidation rate parameter, below). Ci* is the condensed mass loading at which half
of the organic material in that volatility bin will be in the condensed phase, and half will be in gas phase (Donahue et
al., 2006).
- 207 2. **SVOC volatility distribution:** This parameter is expressed in terms of an "equivalent age", determined using a simple 208 ageing model. At time = 0 all VBS molecules will be highly volatile, with a $Ci^*=4$. These compounds are processed 209 at a fixed reaction rate (at each step 0.1% of the gaseous mass in a volatility bin is moved to the next volatility bin), 210 with simple equilibrium partitioning of the VBS components between the gas and condensed phases (to roughly 211 simulate the manner in which VBS compounds are partitioned and aged within the WRF-Chem scheme). This 212 processing reduces the overall volatility of the VBS compounds, first providing a spread of mass across the volatility 213 range, before accumulating the mass in the lowest volatility bins until 90% of the VBS mass is in the Ci^{*}=-2 volatility 214 bin ("time" = 1). This parameter is a scalar variable (between 0-1), that indicates the dimensionless position between 215 these two points, and has an associated volatility distribution. After examining the range of volatility distributions 216 given by this simple ageing model, we have chosen to use distributions within the range of 0.05 to 0.4. Using values 217 above 0.05 ensures there will always be some lower volatility compounds to condense. Above 0.4 almost everything 218 is condensed, so we have excluded values above this so that our PPE does not become too heavily weighted towards 219 these scenarios. Example volatility distributions across the chosen range are shown in supplementary figure S2.
- 3. SVOC oxidation rate: This parameter represents the degree of oxidation that occurs with (or is induced by) each reaction with an OH molecule. Previous studies have used values between 0.075 (7.5%) extra oxygen (or one oxygen atom) (Robinson et al., 2007) and 0.40 (40%) extra oxygen (or five extra oxygen atoms per reaction) (Grieshop et al., 2009). Grieshop et al. (2009) stated that with 7.5%, there is not enough addition of oxygen to the organic mass, while with the 40% there is a noticeable improvement to the OA oxygen content with little effect on the predicted organic mass production. In our study, the lowest level is 0.075 extra oxygen (or one oxygen atom) and the uppermost level is 0.45 (or six extra oxygen atoms per reaction).
- 227 4. **IVOC scaling:** IVOC compounds bridge the gap from SVOC to VOC (log10(C*) 4-6). Including the IVOC 228 independently to parameter (2) (based on our simple ageing model) enables us to still include these within the 229 volatility distribution (this does restrict the impact of parameter (2) to influencing the shape of the volatility 230 distribution for the lower C* values only. These IVOC emissions are calculated using a fixed volatility distribution 231 which scales from the non-volatile OA mass in the emissions inventory. The fractional emitted masses are: 0.2 for 232 $Ci^* = 4$; 0.5 for $Ci^* = 5$; and 0.8 for $Ci^* = 6$ (as shown in supplementary Figure S2) (0.2+0.5+0.8=1.5), this is the 233 initial emission amount that then will be scaled by another factor, between 0-3, to probe the sensitivity of the model 234 to the abundance of IVOCs.. These fractions are then adjusted by this scaling factor, in the range 0.3.

235 5. **SVOC scaling:** This parameter is the scaling factor of the SVOC emissions, (which have been given a volatility 236 distribution by parameter 2). Traditionally such scaling has been used: to ensure that the condensed mass of the 237 emitted SVOC is the same as the non-volatile OA mass in the emissions inventory; however, this scaling could also 238 be used to off-set errors in the emission inventory estimates of OA emissions. The scaling needed to ensure that the 239 emitted condensed mass is the same will never be less than 1, but could go to x20 (or more) for the "younger" SVOC 240 volatility ranges (as estimated using the equilibrium partitioning tool for parameter 2). However, in order to 241 accommodate potential over-estimates of the emission inventories, and to avoid too much OA being generated after 242 aging of any highly-volatile emissions, we chose an SVOC scaling range 0.5 to 4.

243 Table 2 shows the uncertainty ranges applied to each of the parameters, that we explore with the PPE, and Table S34 in the 244 supplementary information shows an example of a 'namelist input' file with the parameters to control the VBS scheme, that 245 was used to create the model simulation. A total of 111 model simulations make up the ensemble. Following the statistical 246 methodology outlined in Lee et al. (2011), the combinations of input parameters used for the simulations in the PPE were 247 selected using an optimal Latin hypercube statistical design algorithm (Stocki, 2005), providing a good coverage of the multi-248 dimensional parameter space. The selection of combinations was performed in three subsets, for use in building statistical 249 emulators to densely sample key outputs from the model over its uncertainties. First, a single design of 61 runs was generated 250 for training the emulators (subset 1), and then a second set of 20 runs was made that 'augmented' into the larger gaps of the 251 first design, for use in validating the emulators (subset 2). On an initial comparison to observations, the observations were 252 found to be outside the range of the PPE's output, and following an investigation into this, the lower bound of the anthropogenic 253 SVOC scaling parameter (parameter 5) was extended from 0.5x down to 0.1x. Hence, an extra, third, set of 30 runs were 254 designed and simulated to cover the extended parameter space (subset 3), leading to a total of 111 runs in the final PPE. Table 255 S42 in supplementary information provides a list of the model runs that make up the PPE with their respective values.

256 Table 2: Range of the parameter space used for SVOCs co-emitted within anthropogenic POAsemi volatile POA emissions 257 and processing in the PPE with 111 model variants.

Parameter number	Parameter name	min	Max
1	Anthropogenic VBS ageing rate (cm ³ molec ⁻¹ s ⁻¹)	1.00E-	1.00E-
		13	11
2	Anthropogenic SVOC volatility distribution	0.05	0.4
3	Added oxygen per generation of ageing	0.075	0.45
4	Anthropogenic IVOC scaling	0	3
5	Anthropogenic SVOC scaling	* 0.1	4
6	Biomass Burning VBS ageing rate (cm ³ molec ⁻¹ s ⁻¹)	1.00E-	1.00E-
		13	11
7	Biomass Burning SVOC volatility distribution	0.05	0.4
8	Added oxygen per generation of ageing	0.075	0.45
9	Biomass Burning IVOC scaling	0	3
10	Biomass Burning SVOC scaling	0.5	4

258 * 81 runs were performed with an anthropogenic SVOC scaling min = 0.5 and max = 4 and 30 runs were performed with an 259 anthropogenic SVOC scaling min = 0.1 and max = 0.5. This due to a min = 0.5 and max = 4 giving high Org mass 260 concentrations, when compared with AMS.

261 2.4 Emulation

262 For each PPE member, a time series of the OC ratio and Total OM from the WRF-Chem model run was extracted at the 263 closest coordinates to the IMD site (Lat 28.628, Lon 77.209) in the model output. Gaussian process emulators (O'Hagan,

264 2006; Lee et al, 2011) were built using the PPE. Similarly to the approach described in Johnson et al. (2018), initial emulators were constructed using only training simulations (subsets 1 and 3) and these were validated using the validation runs (subset 266 2). Once validated, a further new emulator was then constructed using both the training and validation simulations of the PPE 267 together as training data, to obtain a final emulator based on all of the information that the PPE contains. An additional 268 verification of the quality of each final emulator was obtained via a 'leave-one-out' validation procedure (where each 269 simulation in turn is removed from the full set of 111 runs and a new emulator is built and used to predict that removed 270 simulation).

Monte Carlo sampling of the emulators enabled dense samples of model output to be generated over the 10-dimensional parameter uncertainty of the model. We produced output samples for a set of 0.5 million input parameter combinations across the uncertainty space, hereafter called 'model variants', to explore the model's uncertainty.

274 2.5 Model evaluation

Alongside the emulation, outputs from the 111 model runs (OC_ratio and Total_OM) were additionally evaluated, against the AMS observations (O:C and OA), using various model evaluation tools, including the fraction of predictions within a factor of two (FAC2), mean bias (MB) and the index of agreement (IOA). Section S²³ of the supplementary information provides a detailed explanation of the calculations for each evaluation metric and information on how to interpret the values.

279 3 Results and discussion

280 3.1 Model outputs and observational analysis

281 The model outputs of the central WRF-Chem run, from the original parameter space (Subsets 1 and 2), are used to compare 282 with observations in order to analyse the model performance. As mentioned in the methods section, the VBS setup will affect 283 OA concentrations and PM, with no implications to inorganic aerosols or gaseous species. As mentioned in the methods 284 section, the VBS setup will directly affect OA concentrations and PM. The oxidative budget for inorganic chemistry is not 285 directly affected, however, by changing the aerosol size distribution there are some indirect effects on inorganic aerosol and 286 gaseous species through changes in aerosol water content, cloud fields, and aerosol-radiation interactions. Figure 2 shows the comparison for the full dataset (1st – 29th May 2018) between model outputs and observations performed at IMD Lodhi road, 287 288 where we see higher PM_{25} and NO_x concentrations in the model simulation. The high NOx concentrations in the model seem 289 to be related to high NO₂ concentrations as the NO concentrations are in line with the range of the observations of NO. Looking 290 at the meteorological parameters, we can see similar temperatures and wind speeds between the model and observations, with 291 lower RH and higher PBLH in the model.





Figure 2. Comparison of observations (At Lodhi Road for air quality and IGI Airport for meteorology parameters) and model
 outputs of various parameters. May 2018. Bars highlight medians, quartiles and 95%, triangles highlight the mean.

296 3.2 Model runs and AMS observations

297 Here, we analyse and compare the mean values of Total OM (modelled particle phase) and OC ratio for the full period, 1st – 298 29th May 2018, of the 111 WRF-Chem model runs (Table S42 in supplement) with the AMS observations (OA and O:C). The 299 top panel in figure 3 shows a bar plot of the mean OC ratio for the model runs coloured by the mean total OM concentrations. 300 The bottom panel shows the mean total OM concentrations for the model runs coloured by the mean OC ratio. The model 301 runs are sorted from low to high values of the y-variable. The continuous and dashed red lines show the mean \pm one standard 302 deviation (SD) of the O:C ratio (top) and OA (bottom) measured with the AMS. In general, compared to mean values measured 303 with the AMS, a large number of WRF-Chem runs had a low O:C_ratio and high mean Total_OM concentrations. The bottom 304 panel shows the mean total OM concentrations of 47 runs lay within one SD of the mean OA concentration of 21.77 µg.m⁻³ 305 measured with the AMS. Moreover, the model runs with mean Total OM concentrations near the mean OA concentrations 306 have OC ratio mean values near the O:C mean AMS value (0.5), with a cyan colour. This analysis shows a number of model 307 runs with mean Total OM and OC ratio values near the mean values measured with an AMS.



293

309

310





Figure 3. Analysis of the 111 model runs for the full period. Mean OC_ratio coloured by mean Total_OM (top plot) and mean Total_OM coloured by mean OC_ratio (bottom plot). The red lines highlight the mean \pm SD of AMS observations (O:C top and OA bottom). The mean AMS values are O:C = 0.5 and OA = 21.77 µg.m⁻³.

316 3.3 Diurnal analysis to WRF-Chem runs

317 The high time resolution data collected with the AMS provides the opportunity of analysing the WRF-Chem outputs in more 318 detail, for example by looking at the diurnal cycles. Figure 4 shows the diurnal cycles of chosen WRF-Chem runs with 319 Total OM concentrations and OC ratio close to the AMS observations. In the model runs, we were able to span high and low 320 Total OM and OC ratio. However, in the case of OC ratio, we were not able to span the range of the O:C from AMS 321 observations with mean values of 0.3 at night and 0.7 during the day. Looking at the Total OM concentrations, we identified 322 two potential structural errors in the WRF-Chem outputs, the early morning peak and the late evening low concentrations. This 323 could be due to application of unsuitable diurnal activity cycles to the emissions or WRF-Chem not being able to capture 324 completely the dynamics of the planetary boundary layer. With no activity data available for Delhi, we used diurnal cycles of \$25 activities based on emission sectors in Europe (Olivier et al., 2003) (Figure S1 in supplement).-We can observe in figure S5 a 326 slightly better comparison in CO model vs observations, with flatter CO concentrations when looking at the observations. For 327 the diurnal cycles of meteorology (Figure S4), we can see that the model agrees with the PBLH- ERA5 in the early morning 328 and until 14:00 h, time when PBLH- ERA5 starts dropping and PBLH-Model remains high, perhaps preventing concentrations 329 to accumulate. This makes building and testing the emulator challenging as we may get the correct concentrations for the 330 wrong reasons. The emulator can be built over a specific time-period and be compared with the observations. Hence, the 331 emulator was built over two periods of interest: the full period (1st-29th May) and a period where no potential structural errors 332 were identified from 14:00- 16:00 hrs for 1st-29th May (2-4 pm period). Emulator analysis involving the filtering of model 333 results to avoid structural errors has been successfully performed previously in constraining a climate model (Johnson et al., 334 2020). Looking at the mean OC ratio and Total OM of the model runs for the 2 4 pm period (Figure S6), 34 runs lay within 335 one SD of the OA mean concentration (12.20 ug.m⁻³) measured with the AMS, compared with the 47 runs identified from \$36 figure 3. This means that even by analysing the 2-4 pm period we still have model runs that cover the AMS observations.



Figure 4. Diurnal cycle of selected WRF-Chem runs with values near the AMS observations (black line).

340 3.4 Model evaluation

338

341 There are various tools that can be used to compare the model outputs with the observations. In this study, we use a number 342 of statistical metrics (see Section S3 in the supplementary information for a detailed description of each metric we consider) 343 to evaluate the ensemble of 111 model runs for the 2-4 pm period and the full period. The fraction of predictions within a 344 factor of two (FAC2) represents the fraction of data where predictions are within a factor of two of observations. The Mean 345 Bias (MB) gives an indication of the mean over- or under-estimation of predictions. The Index of Agreement (IOA) is a 346 commonly used metric in model evaluation (Willmott et al., 2012), ranging between -1 and +1, with values close to +1347 representing a better model performance. Table S3 shows the results of the model evaluation for the 2-4 pm period and table 348 S4 the results for the full period. When comparing the performance of the two periods; the model runs of the 2-4 pm period 349 have a better performance with 103 runs for O:C and 29 runs for OA with FAC2 > 0.6 compared to 94 runs for O:C and 4 runs 350 for OA with FAC2 > 0.6 for the full period. The negative MB in O:C suggests the models are underestimating the O:C ratios 351 (between -0.01 to -0.15) measured with the AMS. However, the FAC2 values of 0.96 and higher indicate that the models are 352 doing a good job overall at simulating the O:C ratios. This is not the same for OA concentrations, where the models show an 353 over-estimate of the concentration compared to observations, and where only 0.56 -0.62 of predictions were within a factor of 354 two of the OA observations.

355 The IOA provides similar results with a better model performance in the 2-4 pm period, with 10 model runs for the 2-4 pm 356 period and only two runs for the full period with IOA values equal or higher than 0.45. It is interesting to see that while FAC2 357 was higher, for OA and O:C, in the 2-4 period runs compared to the full period, IOA values in 2-4 period were high with OA 358 but low with O:C, which reached IOA values of 0.53 in the 2-4 period and 0.56 in the full period. Previous studies performing 359 modelling evaluation determined similar IOA values using various models (Ciarelli et al., 2017; Fanourgakis et al., 2019). For 360 instance, Chen et al. (2021), modelling SOA formation, obtained IOA between 0.39 - 0.49. Huang et al. (2021) published 361 recommendations on model evaluation and identified IOA of around 0.5 for organic carbon. Lee et al. (2020) performed a 362 sensitivity analysis to two different SOA modules and obtained IOA values of 0.46 - 0.52.

The model evaluation metrics, along with the parameter setup for each ensemble member, allow us to analyse the model setup that gives a better performance. Figure 5 shows the relative variation (%) of the five anthropogenic parameters of the PPE (1 - 5) for the 2-4 pm period (Figure S7 in supplementary material shows the analysis for the full period). Each pentagon represents the 5-D parameter space and the positions of the dots connected with lines show the position of each parameter

367 within its range for that specific ensemble member. The filled area within the dots represents the explored parameter space in 368 each ensemble member. We are analysing the five anthropogenic PPE only since the five parameters related to biomass burning 369 represented a low contribution to the Total OM concentrations. We are looking for blue, light blue or green colours in the 370 lines and dots (high FAC2 values from the O:C analysis) and blue, light blue or green colours in the filled area (high FAC2 371 values from the OA analysis) to identify the model runs with a good evaluation. In figure 5, we can see that the best runs 372 according to the O:C and OA model evaluation are TRAIN127 and TRAIN121 with other TRAIN runs also with good 373 performance such as (126, 036, 117, 104, 115, 119 and 058). In general, these model runs have low SVOC volatility distribution \$74 (emitted VBS compounds are more volatile) and SVOC scaling. TRAIN127 and TRAIN121 have low VBS ageing rate, SVOC 375 volatility distribution and SVOC scaling and with either high SVOC Oxidation rate or high IVOC scaling.



379 Figure 5. Relative variation (%) of the 5 anthropogenic PPE (1-5) for the 2-4 pm period. Each pentagon represents the 5-D 380 parameter space and the positions of the dots connected with lines show the position of each parameter within its range for 381 that specific ensemble member. The filled area within the dots represents the explored parameter space in each ensemble 382 member. Anticlockwise from top there are the five anthropogenic parameters: VBS ageing rate (P1), SVOC volatility 383 distribution (P2), SVOC Oxidation rate (P3), IVOC scaling (P4) and SVOC scaling (P5). The values of the 5 parameters have 384 been normalised dividing by their respective maximum values, hence their values in this plot range from 0 - 1. Example of 385 interpretation in bottom right: the five parameters are towards their high values = 1.0. The colour in the lines and dots represents 386 the FAC2 values from the O:C analysis and the fill colour represents the FAC2 values from the OA analysis. Red = 0 - 0.2. 387 orange = 0.2 - 0.4, vellow = 0.4 - 0.6, green = 0.6 - 0.8, light blue/cyan = 0.8 - 0.9 and blue = 0.9 - 1.0

388 3.5 Emulator analysis

389 3.5.1 Emulator building and testing

Once we confirm that the ensemble of 111 model runs span the AMS observations we can use it to build the emulator. The emulators are tested using the leave-one-out validation approach (Johnson et al., 2018). In this analysis, each ensemble run is first excluded from the emulator build, and then the emulator is used to predict the output at the parameter setting of the excluded run. Figure 6 shows plots of the emulator predictions (with 95% credible intervals from the emulator model) vs the model outputs of the 111 runs from the leave-one-out validation for OA. Predictions from a perfect emulator would follow exactly along the 1:1 line on the plots.

396 We built and tested the emulator for the full period $(1^{st} - 29^{th} \text{ May})$ to have an overview of the emulator performance. The 397 emulator can be built over a specific time-period to compare with the observations. This allows to study the model performance 398 under different conditions, i.e., high/low aerosol concentrations, day/night, etc. We selected four period time-slots to build and 399 test the emulator under high and low Total OM concentrations and two time-slots. These four emulators showed a good 400 validation analysis (Refer to section S5.1.1 in the supplementary material). However, due to the potential structural errors 401 identified from the diurnal analysis (Section 3.3), we will focus on the selected period without structural errors, 2-4 pm period. 402 Figures S11 and S12 in supplementary material show the spread of Total OM and OC ratio respectively, for the ensemble of 403 111 model runs vs the 10 parameters.

We see in Figure 6 that overall, the emulators built for the two periods; full period (6.a and 6.b) and 2-4 pm period (6.c and 6.d) show a good performance; For the 2-4 pm period, Total_OM with only nine runs that are not within the 95% CI from prediction (red markers) and OC_ratio with ten runs that are not within the 95% CI from prediction. With the new 30 runs (error bars in blue) we managed to reduce the Total_OM concentrations with good prediction on the emulator. However, there is a compromise in the OC_ratio with eight runs with high OC_ratio values that at not within the 95% of the prediction interval of the emulator.



412 Figure 6. Validation of the full (a and b) and 2-4 pm (c and d) periods for O:C ratio and Total OM. Circles are the original 81 413 runs. Squares with error bars in blue are the new 30 runs with low settings of the anthropogenic SVOC scaling parameter 414 (which has led to low aerosol mass). Runs where the actual model output lies outside the 95% prediction interval of the 415 emulator are shown in red.

416 3.5.2 Emulator sensitivity analysis

417 We use a variance-based sensitivity analysis (Lee et al., 2011; Johnson et al., 2018) to decompose the overall variance in the 418 model output for key variables of interest into percentage fractions for the 10 parameters. This analysis was performed to the 419 full period and the 2-4 pm period (Figure 7). Looking at the parameters for the two periods, the anthropogenic SVOC scaling 420 has the highest contribution to the variance, which suggests that constraining this parameter would lead to a reduction in the 421 uncertainty in these outputs from the model. Anthropogenic SVOC volatility distribution has some impact on O:C ratios with 422 a fraction of variance of around 15%.





426 3.5.3 Impact of constraint on uncertainty

427 The emulator was used to predict model outputs for a sample of size 0.5 million, for the full period and the 2-4 pm period. 428 Figure 8 shows the probability distribution of OC ratio and Total OM predicted over the full parameter uncertainty. The 429 AMS mean \pm 1SD are shown in red. We can see the higher density (lower values) of the Total_OM show a good agreement 430 with the AMS-OA concentrations. However, in the case of O:C, the higher density lies on the low O:C ratios compared to the 431 O:C-AMS observations which lie in the upper tail of the predicted distribution. The OC_ratio varies within the two periods, 432 with a wider density range for the full period, 0.25-0.55, which represents the variability of the OC_ratio over the full day. In 433 the case of the 2-4 pm period, we can see more narrow density, 0.3-0.5, which, while lower than the mean O:C ratio measured 434 with the AMS (0.65), may be representative of the O:C ratios estimated with the WRF-Chem runs. This suggests that when 435 analysing diurnal behaviour of WRF-Chem outputs without structural errors, we would be able to analyse more into detail the 436 WRF-Chem performance over different hours of the day.



Figure 8. 0.5 million emulator sample, before constraint, covering the full parameter uncertainty space of the model for the full period (a and b) and for the 2-4 pm period (c and d). Red highlights the AMS mean +/- SD observations.

439 3.5.4 Constraint effect.

440 The AMS observations, OA concentrations and O:C ratios, are used to constrain the emulation, applying an observation 441 uncertainty as mean \pm SD. With mean as the emulator prediction and 1 SD uncertainty, we apply the constraint when 442 accounting for emulator prediction uncertainty, by retaining the variant if the range mean \pm SD overlaps with the observation 443 uncertainty range.

Figure 9 is a 2-d histogram for joint constraint (Total_OM and OC_ratio) for the 2-4pm period, with colour showing frequency of variants in a pixel of an underlying grid arranged as a pairwise (shown by the label box on each axis (above/to right). Each 2-d pairwise space has been split into a 25x25 uniform grid to calculate the frequencies. Where the plots show yellow to red, more variants are retained than in the green / blue areas, highlighting the most likely (higher probability) area of space. This analysis shows that when constraining both Total_OM and O:C ratios, the emulator retains 52310 variants from 0.5 million, which is approximately a 10.46% of the original variants generated. Figure S13 shows the histogram

White areas indicate no variants at all retained in that pixel, so that 2-d space is ruled out with respect to all 10 dimensions. (probability=0). Where the colour is uniform, e.g., biomass burning parameter plots in figure 9, the parameter is essentially un-constrained, and all parts of parameter space with respect to those 2 parameters are equally likely/covered by variants (as it was before the constraint was applied). These plots show where in parameter space is most likely given the comparison to observation. These are the variants that we cannot rule out (are plausible) given the uncertainty – it does not mean they are all 'good'. It is worth mentioning that with this analysis we do not locate the exact 'best' run, we provide a range of potential combinations to test the WRF-Chem set-up.

- 457 These results agree with the analysis in the model evaluation (Section 3.4). Figure 9 shows, in red colour, the higher probability
- that with low SVOC volatility distribution and low SVOC scaling would give a good model performance. However, there is no clear pattern with the other parameters.





464

461

465 **3.5.5 Marginal parameter constraints.**

These plotsFigure 10 shows the marginal constraint (1-d projection) on the parameters over their ranges. The unconstrained
 sample (black) has even coverage (is sampled uniformly) across all parameter ranges and the parameter space. The
 unconstrained sample covers the full 10-d space.

469 Where the probability density function (pdf) of the constrained sample is above the black unconstrained pdf, this means the 470 likelihood of the parameter taking a value at that point of its range is increased on constraint (more probability). Where it is 471 below, it is now less likely on constraint. (less probability). The more 'squashed' the unconstrained distribution is – the more

the likelihood of the parameter taking values in the range with higher density is. This analysis is a useful tool to identify the more likely values of the 10 parameters over all the parameter space. Here, we can see that low SVOC volatility distribution and low SVOC are clear parameter values that we can use to improve the WRF-Chem model setup. Other parameters that we can start testing on WRF-Chem are; high BB VBS ageing rate (6) and BB IVOC scaling (9). It is worth highlighting the similarity of the effects on the anthropogenic and biomass burning parameters.



477



479

480

482 **3.6** Analysis of model evaluation and emulator runs.

483 Table 4 shows the WRF-Chem runs with both mean Org and mean O:C values close to AMS observations for the two periods 484 and also selected runs from the 2-d histograms (Figure 9). Here we can see a couple of interesting findings. First, the O:C 485 ratios presented a better performance with the model evaluation metrics; FAC2 values higher than 0.9 compared with FAC2 486 values up to 0.73 for the Total OM. Looking at the Total OM, there are higher FAC2 values in the 2-4 pm period, which might 487 be related to the structural errors impacting the model performance int eh full period. The MB provides an estimation of the 488 over prediction of the Total OM. In this study, WRF-Chem runs were in general overpredicting the Total OM concentrations. 489 Hence, MB is an important metric. In both periods, there are runs where the overprediction was 5 µg.m⁻³ or lower, i.e. 490 TRAIN110, TRANI121, TRAIN117, etc. This highlights the use of all the analysis presented in this study where we are able 491 to identify probable values for the VBS model parameters and be able to model Total OM and O:C ratios.

492 Table <u>34</u>. Analysis of model evaluation metrics and comparison with observations for the full and 2-4 pm periods. The

FAC2 ranking is based on high FAC2 values of the Total_OM analysis. Mean AMS values for the full period: OA = 21.77494 µg.m⁻³ and O:C = 0.5. Mean AMS values for 2-4 pm period: OA = 12.20 µg.m⁻³ and O:C = 0.67.

Full period	Total_OM					O:C ratio						
model	FAC2 ranking	FAC2	MB	IOA	Total_OM mean	Total_OM SD	FAC2 ranking	FAC2	MB	IOA	O:C ratio mean	O:C ratio SD
TRAIN110	1	0.62	2.23	0.45	23.75	16.58	27	0.94	-0.04	0.48	0.46	0.12
TRAIN126	2	0.61	5.13	0.38	26.42	19.83	20	0.95	-0.04	0.51	0.46	0.11
TRAIN119	5	0.60	9.54	0.31	30.83	22.05	7	0.97	-0.04	0.54	0.47	0.10
TRAIN117	6	0.59	3.18	0.41	24.56	16.93	10	0.97	-0.01	0.53	0.49	0.11
TRAIN009	8	0.59	10.54	0.30	68.50	36.13	15	0.96	-0.08	0.51	0.42	0.11
TRAIN121	9	0.59	2.87	0.41	24.17	18.59	21	0.95	-0.05	0.50	0.45	0.11
TRAIN104	11	0.58	5.77	0.39	24.17	18.59	8	0.97	-0.01	0.56	0.45	0.11
VALID002	12	0.58	13.27	0.24	34.49	24.15	2	0.98	-0.08	0.52	0.43	0.09
TRAIN003	13	0.57	12.65	0.24	33.73	23.56	6	0.97	0.00	0.55	0.50	0.12
TRAIN127	16	0.56	4.78	0.37	26.12	20.02	5	0.97	-0.02	0.55	0.48	0.10
	Total_OM											
2-4 pm period				Total_C	M				0:	C ratio		_
2-4 pm period model	FAC2 ranking	FAC2	MB	Total_C	DM Total_OM mean	Total_OM SD	FAC2 ranking	FAC2	О: МВ	C ratio IOA	O:C ratio mean	O:C ratio SD
2-4 pm period model TRAIN127	FAC2 ranking 1	FAC2 0.73	MB 4.37	Total_C IOA 0.44	DM Total_OM mean 15.64	Total_OM SD 10.72	FAC2 ranking 3	FAC2 0.99	0: MB 0.02	C ratio IOA 0.51	O:C ratio mean 0.50	O:C ratio SD 0.06
2-4 pm period model TRAIN127 TRAIN121	FAC2 ranking 1 3	FAC2 0.73 0.72	MB 4.37 1.02	Total_(IOA 0.44 0.48	DM Total_OM mean 15.64 14.48	Total_OM SD 10.72 11.67	FAC2 ranking 3 7	FAC2 0.99 0.98	O: MB 0.02 0.00	C ratio IOA 0.51 0.52	O:C ratio mean 0.50 0.44	O:C ratio SD 0.06 0.08
2-4 pm period model TRAIN127 TRAIN121 TRAIN126	FAC2 ranking 1 3 4	FAC2 0.73 0.72 0.72	MB 4.37 1.02 4.35	Total_0 IOA 0.44 0.48 0.43	DM Total_OM mean 15.64 14.48 15.77	Total_OM SD 10.72 11.67 9.35	FAC2 ranking 3 7 12	FAC2 0.99 0.98 0.98	O: MB 0.02 0.00 0.01	C ratio IOA 0.51 0.52 0.50	O:C ratio mean 0.50 0.44 0.46	O:C ratio SD 0.06 0.08 0.08
2-4 pm period model TRAIN127 TRAIN121 TRAIN126 TRAIN110	FAC2 ranking 1 3 4 5	FAC2 0.73 0.72 0.72 0.70	MB 4.37 1.02 4.35 2.03	Total_C IOA 0.44 0.48 0.43 0.53	DM Total_OM mean 15.64 14.48 15.77 13.45	Total_OM SD 10.72 11.67 9.35 9.42	FAC2 ranking 3 7 12 23	FAC2 0.99 0.98 0.98 0.96	O: MB 0.02 0.00 0.01 0.02	C ratio IOA 0.51 0.52 0.50 0.47	O:C ratio mean 0.50 0.44 0.46 0.45	O:C ratio SD 0.06 0.08 0.08 0.09
2-4 pm period model TRAIN127 TRAIN121 TRAIN126 TRAIN110 TRAIN036	FAC2 ranking 1 3 4 5 11	FAC2 0.73 0.72 0.72 0.70 0.69	MB 4.37 1.02 4.35 2.03 5.13	Total_C IOA 0.44 0.48 0.43 0.53 0.40	DM Total_OM mean 15.64 14.48 15.77 13.45 17.23	Total_OM SD 10.72 11.67 9.35 9.42 12.85	FAC2 ranking 3 7 12 23 1	FAC2 0.99 0.98 0.98 0.96 1.00	O: MB 0.02 0.00 0.01 0.02 0.03	C ratio IOA 0.51 0.52 0.50 0.47 0.51	O:C ratio mean 0.50 0.44 0.46 0.45 0.52	O:C ratio SD 0.06 0.08 0.08 0.09 0.05
2-4 pm period model TRAIN127 TRAIN121 TRAIN126 TRAIN110 TRAIN036 TRAIN117	FAC2 ranking 1 3 4 5 11 13	FAC2 0.73 0.72 0.72 0.70 0.69 0.68	MB 4.37 1.02 4.35 2.03 5.13 1.27	Total_C IOA 0.44 0.48 0.43 0.53 0.40 0.47	DM Total_OM mean 15.64 14.48 15.77 13.45 17.23 16.66	Total_OM SD 10.72 11.67 9.35 9.42 12.85 14.80	FAC2 ranking 3 7 12 23 1 5	FAC2 0.99 0.98 0.98 0.96 1.00 0.99	O: MB 0.02 0.00 0.01 0.02 0.03 0.04	C ratio IOA 0.51 0.52 0.50 0.47 0.51 0.48	O:C ratio mean 0.50 0.44 0.46 0.45 0.52 0.51	O:C ratio SD 0.06 0.08 0.08 0.09 0.05 0.08
2-4 pm period model TRAIN127 TRAIN121 TRAIN126 TRAIN10 TRAIN036 TRAIN117 TRAIN104	FAC2 ranking 1 3 4 5 11 13 14	FAC2 0.73 0.72 0.72 0.70 0.69 0.68	MB 4.37 1.02 4.35 2.03 5.13 1.27 5.50	Total_C IOA 0.44 0.48 0.43 0.53 0.40 0.47 0.47	DM Total_OM mean 15.64 14.48 15.77 13.45 17.23 16.66 16.41	Total_OM SD 10.72 11.67 9.35 9.42 12.85 14.80 11.18	FAC2 ranking 3 7 12 23 1 5 14	FAC2 0.99 0.98 0.98 0.96 1.00 0.99 0.98	O: MB 0.02 0.00 0.01 0.02 0.03 0.04 0.03	C ratio IOA 0.51 0.52 0.50 0.47 0.51 0.48 0.47	O:C ratio mean 0.50 0.44 0.46 0.45 0.52 0.51 0.54	O:C ratio SD 0.06 0.08 0.08 0.09 0.05 0.08 0.06
2-4 pm period model TRAIN127 TRAIN121 TRAIN126 TRAIN100 TRAIN036 TRAIN117 TRAIN104 TRAIN115	FAC2 ranking 1 3 4 5 11 13 13 14 16	FAC2 0.73 0.72 0.72 0.70 0.69 0.68 0.68	MB 4.37 1.02 4.35 2.03 5.13 1.27 5.50 3.27	Total_C IOA 0.44 0.48 0.43 0.53 0.40 0.47 0.47 0.39	DM Total_OM mean 15.64 14.48 15.77 13.45 17.23 16.66 16.41 18.17	Total_OM SD 10.72 11.67 9.35 9.42 12.85 14.80 11.18 15.48	FAC2 ranking 3 7 12 23 1 5 14 6	FAC2 0.99 0.98 0.98 0.96 1.00 0.99 0.98 0.99	O: MB 0.02 0.00 0.01 0.02 0.03 0.04 0.03 0.05	C ratio IOA 0.51 0.52 0.50 0.47 0.51 0.48 0.47 0.46	O:C ratio mean 0.50 0.44 0.46 0.45 0.52 0.51 0.54 0.51	O:C ratio SD 0.06 0.08 0.08 0.09 0.05 0.08 0.06 0.06
2-4 pm period model TRAIN127 TRAIN121 TRAIN126 TRAIN10 TRAIN036 TRAIN117 TRAIN104 TRAIN115 TRAIN119	FAC2 ranking 1 3 4 5 11 13 14 16 19	FAC2 0.73 0.72 0.72 0.70 0.69 0.68 0.68 0.68 0.67	MB 4.37 1.02 4.35 2.03 5.13 1.27 5.50 3.27 7.12	Total_C IOA 0.44 0.48 0.43 0.53 0.40 0.47 0.47 0.47 0.39 0.35	DM Total_OM mean 15.64 14.48 15.77 13.45 17.23 16.66 16.41 18.17 18.96	Total_OM SD 10.72 11.67 9.35 9.42 12.85 14.80 11.18 15.48 11.26	FAC2 ranking 3 7 12 23 1 5 14 6 10	FAC2 0.99 0.98 0.98 0.96 1.00 0.99 0.98 0.99 0.98	O: MB 0.02 0.01 0.02 0.03 0.04 0.03 0.05 0.01	C ratio IOA 0.51 0.52 0.50 0.47 0.51 0.48 0.47 0.46 0.51	O:C ratio mean 0.50 0.44 0.46 0.45 0.52 0.51 0.54 0.51 0.49	O:C ratio SD 0.06 0.08 0.09 0.05 0.08 0.06 0.06 0.06

495

496 5 Conclusions

In this study we aimed to determine an effective way of tuning the VBS scheme using observations, and also to learn about the processes controlling OA in Delhi. WRF-Chem model runs with the VBS setup that successfully span the OA concentrations and O:C ratios from AMS observations can be identified, with many model runs overestimating organic mass concentrations and underestimating the O:C ratios compared with AMS observations. However, we identified two structural errors in the model related to a combination of unsuitable diurnal activity cycles applied to the emissions and/or WRF-Chem

- not being able to capture completely the dynamics of the planetary boundary layer. It is worth mentioning that these structural errors might also be related to representation of other organic aerosol processes not represented by the VBS approach. Recent studies, for example, have examined particle-phase and multiphase chemistry in aqueous aerosols and clouds (Shrivastava et al., 2022), and reactions of SOA precursors with other radicals like chlorine relevant to Indian conditions (Gunthe et al., 2021).
- 506 The structural errors prevented us from providing an optimised VBS approach in WRF-Chem. However, we were able to apply 507 the emulator in two periods: the full period (1st -29th May) and the 2-4 pm period (14:00- 16:00 hrs, 1st-29th May) to present 508 a methodology to evaluate a model performance using Gaussian emulators and metrics such as FAC2, IOA and MB. 509 Optimization is a stage-by-stage process, future analysis would imply to do an emulation study to address diurnal activity and 510 PBL directly, perhaps using NOx or total PM.
- The performance of the two emulators, the full period and the 2-4 pm period, was similar, with the two emulators performing
 a good prediction of the model outputs and presenting a similar high variance of the anthropogenic SVOC scaling (Parameter
 5). The model performance would highly improve if we are able to constrain the input values for the parameter 5.
- 514 When looking at the emulator sensibility analysis, we identified that the parameter anthropogenic SVOC scaling has the highest 515 contribution to the variance, with fractions higher than 70%. This suggests that constraining this parameter would lead to a 516 reduction in the uncertainty in these outputs from the model. Anthropogenic SVOC volatility distribution has little impact on 517 the fraction of variance to O:C ratios with a fraction of variance of around 15%. None of the parameters show a clear variance 518 to improve the model performance.
- The model evaluation analysis based on FAC2, IOA and MB agreed with the emulator analysis in identifying that using low SVOC volatility distribution and low SVOC scaling would give improved model performance. Based on the MB analysis, for both the full and the 2-4 pm periods, there are runs where the Total OM overprediction was 5 ug.m⁻³ or lower, i.e, TRAIN110, TRANI121, TRAIN117, etc. This overprediction is considered low compared to the mean Total_OM concentrations of ~20 – 30 μ g.m⁻³. Hence, we are able to identify probable values for the VBS model parameters and are able to model Total OM and O:C ratios in the range of the AMS observations.
- The combination of the emulator analysis and the model evaluation metrics (FAC2, IOA and mean bias) allowed us to identify the plausible parameter combinations for the analysed periods. The more plausible combinations were found to be with a low SVOC volatility distribution and low SVOC scaling, which means a more volatile distribution. The methodology presented in this study is shown to be a useful approach to determine the model uncertainty and determine the optimal parameterisation to the WRF-Chem VBS setup. This information is valuable to increase our understanding on secondary organic aerosol formation, which in turn will help to improve regional and global model simulations, emission inventories as well as making informed decisions towards the improvement of air quality in urban environments.
- 532 Data availability
- 533 Emission generation scripts: <u>https://github.com/douglowe/WRF_UoM_EMIT</u>
- 534 Scripts for running WRF-Chem (and reducing the outputs to key diagnostics):
- 535 <u>https://github.com/douglowe/promote_wrfchem_scripts</u>
- 536
- 537 Scenario configuration files, and python script for calculating the "pseudo-age" of the emitted VBS:
- 538 <u>https://github.com/douglowe/PROMOTE_VBS_scenarios</u>
- 539 Scenario chemistry input files
- 540 <u>https://github.com/douglowe/PROMOTE_VBS_scenarios/tree/master/Scenario_Configurations/scenario_chemistry_files_A</u>
- 541 <u>ug2020</u>
- 542

544 Financial support

545

- 546 This research has been supported by the UK NERC and MoES, India through the PROMOTE project
- 547 under the Newton Bhabha Fund programme "Air Pollution and Human Health in a Developing
- 548 Megacity (APHH-India)", NERC grant numbers NE/P016480/1 and, NE/P016405. M.S. was supported
- 549 by the U.S. Department of Energy (DOE) Office of Science, Office of Biological and Environmental
- 550 Research (BER) through the Early Career Research Program. R.A.Z. acknowledges support from the
- 551 Office of Science of the U.S. DOE through the Atmospheric System Research (ASR) program at Pacific
- 552 Northwest National Laboratory (PNNL). PNNL is operated for DOE by Battelle Memorial Institute
- under contract DE-AC06-76RLO 1830. This paper is based on interpretation of scientific results and in
- no way reflects the viewpoint of the funding agencies.

555 <u>Acknowledgements</u>

We acknowledge use of the WRF-Chem preprocessor tools mozbc, fire_emiss, bio_emiss and anthro_emiss, provided by the
 Atmospheric Chemistry Observations and Modeling Lab (ACOM) of NCAR.

558 References

Akherati, A., Cappa, C. D., Kleeman, M. J., Docherty, K. S., Jimenez, J. L., Griffith, S. M., Dusanter, S., Stevens, P.

- 560 S., and Jathar, S. H.: Simulating secondary organic aerosol in a regional air quality model using the statistical
- oxidation model Part 3: Assessing the influence of semi-volatile and intermediate-volatility organic
- 562 compounds and NOx, Atmos. Chem. Phys., 19, 4561-4594, 10.5194/acp-19-4561-2019, 2019.
- Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., Hodnebrog, Ø., Denier van der Gon,
- 564 H., and McFiggans, G.: Gaseous chemistry and aerosol mechanism developments for version 3.5.1 of the online 565 regional model, WRF-Chem, Geosci. Model Dev., 7, 2557-2579, 10.5194/gmd-7-2557-2014, 2014.
- 566 Bellouin, N., Baker, L., Hodnebrog, Ø., Olivié, D., Cherian, R., Macintosh, C., Samset, B., Esteve, A., Aamaas, B.,
- 567 Quaas, J., and Myhre, G.: Regional and seasonal radiative forcing by perturbations to aerosol and ozone
- 568 precursor emissions, Atmos. Chem. Phys., 16, 13885-13910, 10.5194/acp-16-13885-2016, 2016.
- 569 Bergström, R., Denier van der Gon, H. A. C., Prévôt, A. S. H., Yttri, K. E., and Simpson, D.: Modelling of organic
- aerosols over Europe (2002–2007) using a volatility basis set (VBS) framework: application of different
- assumptions regarding the formation of secondary organic aerosol, Atmos. Chem. Phys., 12, 8499-8527,
 10.5194/acp-12-8499-2012, 2012.
- 573 Bertram, T. H., and Thornton, J. A.: Toward a general parameterization of N₂O₅
- 574 reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, Atmos.
- 575 Chem. Phys., 9, 8351-8363, 10.5194/acp-9-8351-2009, 2009.
- 576 Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crounse, J. D., Wennberg, P. O.,
- 577 Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A., Donahue, N.,
- 578 Kjaergaard, H. G., and Ehn, M.: Highly Oxygenated Organic Molecules (HOM) from Gas-Phase Autoxidation

579 Involving Peroxy Radicals: A Key Contributor to Atmospheric Aerosol, Chemical Reviews, 119, 3472-3509,

580 10.1021/acs.chemrev.8b00395, 2019.

- Bran, S. H., and Srivastava, R.: Investigation of PM2.5 mass concentration over India using a regional climate
 model, Environmental Pollution, 224, 484-493, <u>https://doi.org/10.1016/j.envpol.2017.02.030</u>, 2017.
- 584 Canagaratna, M. R., Jimenez, J. L., Kroll, J. H., Chen, Q., Kessler, S. H., Massoli, P., Hildebrandt Ruiz, L., Fortner,
- 585 E., Williams, L. R., Wilson, K. R., Surratt, J. D., Donahue, N. M., Jayne, J. T., and Worsnop, D. R.: Elemental ratio
- measurements of organic compounds using aerosol mass spectrometry: characterization, improved calibration,
 and implications, Atmos. Chem. Phys., 15, 253-272, 10.5194/acp-15-253-2015, 2015.
- 588 Carlton, A. G., Bhave, P. V., Napelenok, S. L., Edney, E. O., Sarwar, G., Pinder, R. W., Pouliot, G. A., and Houyoux,
- 589 M.: Model Representation of Secondary Organic Aerosol in CMAQv4.7, Environmental Science & Technology, 590 44, 8553-8560, 10.1021/es100636g, 2010.
- 591 Carslaw, K. S., Lee, L. A., Regayre, L. A., and Johnson, J. S.: Climate models are uncertain, but we can do 592 something about it, Eos, 99, <u>https://doi.org/10.1029/2018E0093757C</u>, 2018.
- Cash, J. M., Langford, B., Di Marco, C., Mullinger, N. J., Allan, J., Reyes-Villegas, E., Joshi, R., Heal, M. R., Acton,
- 594 W. J. F., Hewitt, C. N., Misztal, P. K., Drysdale, W., Mandal, T. K., Shivani, Gadi, R., Gurjar, B. R., and Nemitz, E.:
- Seasonal analysis of submicron aerosol in Old Delhi using high-resolution aerosol mass spectrometry: chemical
 characterisation, source apportionment and new marker identification, Atmos. Chem. Phys., 21, 10133-10158,
- 597 10.5194/acp-21-10133-2021, 2021.
- Chen, X., Zhang, Y., Zhao, J., Liu, Y., Shen, C., Wu, L., Wang, X., Fan, Q., Zhou, S., and Hang, J.: Regional modeling
 of secondary organic aerosol formation over eastern China: The impact of uptake coefficients of dicarbonyls and
- 600 semivolatile process of primary organic aerosol, Science of The Total Environment, 793, 148176,
- 601 <u>https://doi.org/10.1016/j.scitotenv.2021.148176</u>, 2021.
- 602 Chen, Y., Wild, O., Conibear, L., Ran, L., He, J., Wang, L., and Wang, Y.: Local characteristics of and exposure to
- fine particulate matter (PM2.5) in four indian megacities, Atmospheric Environment: X, 5, 100052,
 https://doi.org/10.1016/j.aeaoa.2019.100052, 2020a.
- 605 Chen, Y., Wild, O., Ryan, E., Sahu, S. K., Lowe, D., Archer-Nicholls, S., Wang, Y., McFiggans, G., Ansari, T., Singh,
- 606 V., Sokhi, R. S., Archibald, A., and Beig, G.: Mitigation of PM2.5 and ozone pollution in Delhi: a sensitivity study
- 607 during the pre-monsoon period, Atmos. Chem. Phys., 20, 499-514, 10.5194/acp-20-499-2020, 2020b.
- 608 Chutia, L., Ojha, N., Girach, I. A., Sahu, L. K., Alvarado, L. M. A., Burrows, J. P., Pathak, B., and Bhuyan, P. K.:
- 609 Distribution of volatile organic compounds over Indian subcontinent during winter: WRF-chem simulation
- versus observations, Environmental Pollution, 252, 256-269, <u>https://doi.org/10.1016/j.envpol.2019.05.097</u>,
 2019.
- 612 Ciarelli, G., Aksoyoglu, S., El Haddad, I., Bruns, E. A., Crippa, M., Poulain, L., Äijälä, M., Carbone, S., Freney, E.,
- 613 O'Dowd, C., Baltensperger, U., and Prévôt, A. S. H.: Modelling winter organic aerosol at the European scale with
- 614 CAMx: evaluation and source apportionment with a VBS parameterization based on novel wood burning smog
- 615 chamber experiments, Atmos. Chem. Phys., 17, 7653-7669, 10.5194/acp-17-7653-2017, 2017.
- 616 Crooks, M., Connolly, P., and McFiggans, G.: A parameterisation for the co-condensation of semi-volatile
- organics into multiple aerosol particle modes, Geosci. Model Dev., 11, 3261-3278, 10.5194/gmd-11-3261-2018,
 2018.
- 619 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo,
- 620 J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M.,

- 621 Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A.
- 622 S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-
- 623 Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K.,
- Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2
- 625 (CESM2), Journal of Advances in Modeling Earth Systems, 12, e2019MS001916,
- 626 <u>https://doi.org/10.1029/2019MS001916</u>, 2020.
- 627 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K.,
- Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight
 aerosol mass spectrometer, Anal Chem, 78, 8281-8289, Doi 10.1021/Ac061249n, 2006.
- 630 Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D., and
- 631 Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe revised estimates and
- 632 an evaluation, Atmos. Chem. Phys., 15, 6503-6519, 10.5194/acp-15-6503-2015, 2015.
- 633 Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled Partitioning, Dilution, and Chemical
- Aging of Semivolatile Organics, Environmental Science & Technology, 40, 2635-2643, 10.1021/es052297c, 2006.
- Fanourgakis, G. S., Kanakidou, M., Nenes, A., Bauer, S. E., Bergman, T., Carslaw, K. S., Grini, A., Hamilton, D. S.,
- Johnson, J. S., Karydis, V. A., Kirkevåg, A., Kodros, J. K., Lohmann, U., Luo, G., Makkonen, R., Matsui, H.,
- 637 Neubauer, D., Pierce, J. R., Schmale, J., Stier, P., Tsigaridis, K., van Noije, T., Wang, H., Watson-Parris, D.,
- 638 Westervelt, D. M., Yang, Y., Yoshioka, M., Daskalakis, N., Decesari, S., Gysel-Beer, M., Kalivitis, N., Liu, X.,
- Mahowald, N. M., Myriokefalitakis, S., Schrödner, R., Sfakianaki, M., Tsimpidi, A. P., Wu, M., and Yu, F.:
- Evaluation of global simulations of aerosol particle and cloud condensation nuclei number, with implications for cloud droplet formation, Atmos. Chem. Phys., 19, 8591-8617, 10.5194/acp-19-8591-2019, 2019.
- Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and
- 643 Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston
- using a fully coupled meteorology-chemistry-aerosol model, Journal of Geophysical Research: Atmospheres,
 111, https://doi.org/10.1029/2005JD006721, 2006.
- Fountoukis, C., Megaritis, A. G., Skyllakou, K., Charalampidis, P. E., Pilinis, C., Denier van der Gon, H. A. C.,
- 647 Crippa, M., Canonaco, F., Mohr, C., Prévôt, A. S. H., Allan, J. D., Poulain, L., Petäjä, T., Tiitta, P., Carbone, S.,
- 648 Kiendler-Scharr, A., Nemitz, E., O'Dowd, C., Swietlicki, E., and Pandis, S. N.: Organic aerosol concentration and
- 649 composition over Europe: insights from comparison of regional model predictions with aerosol mass
- 650 spectrometer factor analysis, Atmos. Chem. Phys., 14, 9061-9076, 10.5194/acp-14-9061-2014, 2014.
- 651 Gadi, R., Shivani, Sharma, S. K., and Mandal, T. K.: Source apportionment and health risk assessment of organic
- 652 constituents in fine ambient aerosols (PM2.5): A complete year study over National Capital Region of India,
- 653 Chemosphere, 221, 583-596, <u>https://doi.org/10.1016/j.chemosphere.2019.01.067</u>, 2019.
- 654 Ghosh, S., Verma, S., Kuttippurath, J., and Menut, L.: Wintertime direct radiative effects due to black carbon
- 655 (BC) over the Indo-Gangetic Plain as modelled with new BC emission inventories in CHIMERE, Atmos. Chem.
- 656 Phys., 21, 7671–7694, https://doi.org/10.5194/acp-21-7671-2021, 2021.
- 657 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled 658 "online" chemistry within the WRF model, Atmos Environ, 39, 6957-6975,
- 659 <u>https://doi.org/10.1016/j.atmosenv.2005.04.027</u>, 2005.
- Grieshop, A. P., Logue, J. M., Donahue, N. M., and Robinson, A. L.: Laboratory investigation of photochemical
- oxidation of organic aerosol from wood fires 1: measurement and simulation of organic aerosol evolution,
- 662 Atmos. Chem. Phys., 9, 1263-1277, 10.5194/acp-9-1263-2009, 2009.

- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial
 isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmos. Chem. Phys.,
 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.
- 666 Gunthe, S. S., Liu, P., Panda, U., Raj, S. S., Sharma, A., Darbyshire, E., Reyes-Villegas, E., Allan, J., Chen, Y., Wang,
- K., Song, S., Pöhlker, M. L., Shi, L., Wang, Y., Kommula, S. M., Liu, T., Ravikrishna, R., McFiggans, G., Mickley, L. J.,
- 668 Martin, S. T., Pöschl, U., Andreae, M. O., and Coe, H.: Enhanced aerosol particle growth sustained by high 669 continental chlorine emission in India, Nat Geosci, 14, 77-84, 10.1038/s41561-020-00677-x, 2021.
- 670 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R.,
- 671 Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels
- 672 from 1979 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). Copernicus,
- 673 10.24381/cds.bd0915c6, 2018.
- Huang, L., Zhu, Y., Zhai, H., Xue, S., Zhu, T., Shao, Y., Liu, Z., Emery, C., Yarwood, G., Wang, Y., Fu, J., Zhang, K.,
- and Li, L.: Recommendations on benchmarks for numerical air quality model applications in China Part 1:
- 676 PM2.5 and chemical species, Atmos. Chem. Phys., 21, 2725-2743, 10.5194/acp-21-2725-2021, 2021.
- 677 World Air Quality: <u>https://www.iqair.com/</u>, access: 05/08/2021, 2021.
- Jain, S., Sharma, S. K., Vijayan, N., and Mandal, T. K.: Seasonal characteristics of aerosols (PM2.5 and PM10) and
- their source apportionment using PMF: A four year study over Delhi, India, Environmental Pollution, 262,
 114337, https://doi.org/10.1016/j.envpol.2020.114337, 2020.
- Jat, R., Gurjar, B. R., and Lowe, D.: Regional pollution loading in winter months over India using high resolution
- 682 WRF-Chem simulation, Atmos Res, 249, 105326, https://doi.org/10.1016/j.atmosres.2020.105326, 2021.
- Johnson, J. S., Regayre, L. A., Yoshioka, M., Pringle, K. J., Lee, L. A., Sexton, D. M. H., Rostron, J. W., Booth, B. B.
- B., and Carslaw, K. S.: The importance of comprehensive parameter sampling and multiple observations for
 robust constraint of aerosol radiative forcing, Atmos. Chem. Phys., 18, 13031-13053, 10.5194/acp-18-130312018, 2018.
- Johnson, J. S., Regayre, L. A., Yoshioka, M., Pringle, K. J., Turnock, S. T., Browse, J., Sexton, D. M. H., Rostron, J.
 W., Schutgens, N. A. J., Partridge, D. G., Liu, D., Allan, J. D., Coe, H., Ding, A., Cohen, D. D., Atanacio, A., Vakkari,
- 689 V., Asmi, E., and Carslaw, K. S.: Robust observational constraint of uncertain aerosol processes and emissions in
- a climate model and the effect on aerosol radiative forcing, Atmos. Chem. Phys., 20, 9491-9524, 10.5194/acp-20-9491-2020, 2020.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens,
 B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K.,
- 694 Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and
- 695 global climate modelling: a review, Atmos. Chem. Phys., 5, 1053-1123, 10.5194/acp-5-1053-2005, 2005.
- 696 Kompalli, S. K., Suresh Babu, S. N., Satheesh, S. K., Krishna Moorthy, K., Das, T., Boopathy, R., Liu, D., Darbyshire,
- 697 E., Allan, J. D., Brooks, J., Flynn, M. J., and Coe, H.: Seasonal contrast in size distributions and mixing state of
- black carbon and its association with PM1.0 chemical composition from the eastern coast of India, Atmos.
 Chem. Phys., 20, 3965-3985, 10.5194/acp-20-3965-2020, 2020.
- Lane, T. E., Donahue, N. M., and Pandis, S. N.: Simulating secondary organic aerosol formation using the
- 701 volatility basis-set approach in a chemical transport model. Atmos Environ. 42, 7439-7451.
- 702 https://doi.org/10.1016/j.atmosenv.2008.06.026, 2008.
- Lee, H.-J., Jo, H.-Y., Song, C.-K., Jo, Y.-J., Park, S.-Y., and Kim, C.-H.: Sensitivity of Simulated PM2.5 Concentrations
- 704 over Northeast Asia to Different Secondary Organic Aerosol Modules during the KORUS-AQ Campaign,
- 705 Atmosphere, 11, 1004, 2020.

- Lee, L. A., Carslaw, K. S., Pringle, K. J., Mann, G. W., and Spracklen, D. V.: Emulation of a complex global aerosol
 model to quantify sensitivity to uncertain parameters, Atmos. Chem. Phys., 11, 12253-12273, 10.5194/acp-1112253-2011, 2011.
- Li, J.-L., Zhang, M.-G., Gao, Y., and Chen, L.: Model analysis of secondary organic aerosol over China with a
- regional air quality modeling system (RAMS-CMAQ), Atmospheric and Oceanic Science Letters, 9, 443-450,
 10.1080/16742834.2016.1233798, 2016.
- May, A. A., Levin, E. J. T., Hennigan, C. J., Riipinen, I., Lee, T., Collett Jr., J. L., Jimenez, J. L., Kreidenweis, S. M.,
- and Robinson, A. L.: Gas-particle partitioning of primary organic aerosol emissions: 3. Biomass burning, Journal
- of Geophysical Research: Atmospheres, 118, 11,327-311,338, <u>https://doi.org/10.1002/jgrd.50828</u>, 2013.
- Olivier, J., Peters, J., Granier, C., Pétron, G., Muller, J. F., and Wallens, S.: POET inventory Metadata, GEIA-
- 716 ACCENT emission data portal, France, 2003.
- 717 Reyes-Villegas, E., Green, D. C., Priestman, M., Canonaco, F., Coe, H., Prévôt, A. S. H., and Allan, J. D.: Organic
- aerosol source apportionment in London 2013 with ME-2: exploring the solution space with annual and
- 719 seasonal analysis, Atmos. Chem. Phys., 16, 15545-15559, 10.5194/acp-16-15545-2016, 2016.
- Reyes-Villegas, E., Panda, U., Darbyshire, E., Cash, J. M., Joshi, R., Langford, B., Di Marco, C. F., Mullinger, N. J.,
- Alam, M. S., Crilley, L. R., Rooney, D. J., Acton, W. J. F., Drysdale, W., Nemitz, E., Flynn, M., Voliotis, A.,
- 722 McFiggans, G., Coe, H., Lee, J., Hewitt, C. N., Heal, M. R., Gunthe, S. S., Mandal, T. K., Gurjar, B. R., Shivani, Gadi,
- R., Singh, S., Soni, V., and Allan, J. D.: PM1 composition and source apportionment at two sites in Delhi, India,
- 724 across multiple seasons, Atmos. Chem. Phys., 21, 11655-11667, 10.5194/acp-21-11655-2021, 2021.
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E.,
- Pierce, J. R., and Pandis, S. N.: Rethinking Organic Aerosols: Semivolatile Emissions and Photochemical Aging,
 Science, 315, 1259-1262, doi:10.1126/science.1133061, 2007.
- 728 Shivani, Gadi, R., Sharma, S. K., and Mandal, T. K.: Seasonal variation, source apportionment and source
- attributed health risk of fine carbonaceous aerosols over National Capital Region, India, Chemosphere, 237,
 124500, https://doi.org/10.1016/j.chemosphere.2019.124500, 2019.
- 731 Shrivastava, M., Fast, J., Easter, R., Gustafson Jr, W. I., Zaveri, R. A., Jimenez, J. L., Saide, P., and Hodzic, A.:
- Modeling organic aerosols in a megacity: comparison of simple and complex representations of the volatility
 basis set approach, Atmos. Chem. Phys., 11, 6639-6662, 10.5194/acp-11-6639-2011, 2011.
- 734 Shrivastava, M., Berg, L. K., Fast, J. D., Easter, R. C., Laskin, A., Chapman, E. G., Gustafson Jr., W. I., Liu, Y., and
- Berkowitz, C. M.: Modeling aerosols and their interactions with shallow cumuli during the 2007 CHAPS field
- study, Journal of Geophysical Research: Atmospheres, 118, 1343-1360, <u>https://doi.org/10.1029/2012JD018218</u>,
 2013.
- 738 Shrivastava, M., Andreae, M. O., Artaxo, P., Barbosa, H. M. J., Berg, L. K., Brito, J., Ching, J., Easter, R. C., Fan, J.,
- 739 Fast, J. D., Feng, Z., Fuentes, J. D., Glasius, M., Goldstein, A. H., Alves, E. G., Gomes, H., Gu, D., Guenther, A.,
- 740 Jathar, S. H., Kim, S., Liu, Y., Lou, S., Martin, S. T., McNeill, V. F., Medeiros, A., de Sá, S. S., Shilling, J. E.,
- 741 Springston, S. R., Souza, R. A. F., Thornton, J. A., Isaacman-VanWertz, G., Yee, L. D., Ynoue, R., Zaveri, R. A.,
- 742 Zelenyuk, A., and Zhao, C.: Urban pollution greatly enhances formation of natural aerosols over the Amazon
- rainforest, Nature Communications, 10, 1046, 10.1038/s41467-019-08909-4, 2019.
- Shrivastava, M., Rasool, Q. Z., Zhao, B., Octaviani, M., Zaveri, R. A., Zelenyuk, A., Gaudet, B., Liu, Y., Shilling, J. E.,
- Schneider, J., Schulz, C., Zöger, M., Martin, S. T., Ye, J., Guenther, A., Souza, R. F., Wendisch, M., and Pöschl, U.:
- 746Tight Coupling of Surface and In-Plant Biochemistry and Convection Governs Key Fine Particulate Components
- over the Amazon Rainforest, ACS Earth and Space Chemistry, 6, 380-390, 10.1021/acsearthspacechem.1c00356,
- 748 2022.

- Shrivastava, M. K., Lane, T. E., Donahue, N. M., Pandis, S. N., and Robinson, A. L.: Effects of gas particle
 partitioning and aging of primary emissions on urban and regional organic aerosol concentrations, Journal of
 Geophysical Research: Atmospheres, 113, https://doi.org/10.1029/2007JD009735, 2008.
- 752 Stewart, G. J., Nelson, B. S., Acton, W. J. F., Vaughan, A. R., Farren, N. J., Hopkins, J. R., Ward, M. W., Swift, S. J.,
- 753 Arya, R., Mondal, A., Jangirh, R., Ahlawat, S., Yadav, L., Sharma, S. K., Yunus, S. S. M., Hewitt, C. N., Nemitz, E.,
- 754 Mullinger, N., Gadi, R., Sahu, L. K., Tripathi, N., Rickard, A. R., Lee, J. D., Mandal, T. K., and Hamilton, J. F.:
- Emissions of intermediate-volatility and semi-volatile organic compounds from domestic fuels used in Delhi,
 India, Atmos. Chem. Phys., 21, 2407-2426, 10.5194/acp-21-2407-2021, 2021a.
- 757 Stewart, G. J., Nelson, B. S., Acton, W. J. F., Vaughan, A. R., Hopkins, J. R., Yunus, S. S. M., Hewitt, C. N., Nemitz,
- E., Mandal, T. K., Gadi, R., Sahu, L. K., Rickard, A. R., Lee, J. D., and Hamilton, J. F.: Comprehensive organic
 emission profiles, secondary organic aerosol production potential, and OH reactivity of domestic fuel
- 759 combustion in Delhi, India, Environmental Science: Atmospheres, 1, 104-117, 10.1039/D0EA00009D, 2021b.
- 761 Tilmes, S., Hodzic, A., Emmons, L. K., Mills, M. J., Gettelman, A., Kinnison, D. E., Park, M., Lamarque, J.-F., Vitt, F.,
- 762 Shrivastava, M., Campuzano-Jost, P., Jimenez, J. L., and Liu, X.: Climate Forcing and Trends of Organic Aerosols
- in the Community Earth System Model (CESM2), Journal of Advances in Modeling Earth Systems, 11, 4323-4351,
 https://doi.org/10.1029/2019MS001827, 2019.
- Topping, D., Connolly, P., and McFiggans, G.: Cloud droplet number enhanced by co-condensation of organic vapours, Nat Geosci, 6, 443-446, 10.1038/ngeo1809, 2013.
- 767 Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E.,
- 768 Bellouin, N., Benedetti, A., Bergman, T., Berntsen, T. K., Beukes, J. P., Bian, H., Carslaw, K. S., Chin, M., Curci, G.,
- 769 Diehl, T., Easter, R. C., Ghan, S. J., Gong, S. L., Hodzic, A., Hoyle, C. R., Iversen, T., Jathar, S., Jimenez, J. L., Kaiser,
- J. W., Kirkevåg, A., Koch, D., Kokkola, H., Lee, Y. H., Lin, G., Liu, X., Luo, G., Ma, X., Mann, G. W., Mihalopoulos,
- N., Morcrette, J. J., Müller, J. F., Myhre, G., Myriokefalitakis, S., Ng, N. L., O'Donnell, D., Penner, J. E., Pozzoli, L.,
- Pringle, K. J., Russell, L. M., Schulz, M., Sciare, J., Seland, Ø., Shindell, D. T., Sillman, S., Skeie, R. B., Spracklen, D.,
- 773 Stavrakou, T., Steenrod, S. D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., van Noije, T., van Zyl, P. G., von Salzen,
- K., Yu, F., Wang, Z., Wang, Z., Zaveri, R. A., Zhang, H., Zhang, K., Zhang, Q., and Zhang, X.: The AeroCom
- evaluation and intercomparison of organic aerosol in global models, Atmos. Chem. Phys., 14, 10845-10895,
 10.5194/acp-14-10845-2014, 2014.
- Tsimpidi, A. P., Karydis, V. A., Zavala, M., Lei, W., Molina, L., Ulbrich, I. M., Jimenez, J. L., and Pandis, S. N.:
- Evaluation of the volatility basis-set approach for the simulation of organic aerosol formation in the Mexico City
 metropolitan area, Atmos. Chem. Phys., 10, 525-546, 10.5194/acp-10-525-2010, 2010.
- 780 Tsimpidi, A. P., Karydis, V. A., Pandis, S. N., and Lelieveld, J.: Global combustion sources of organic aerosols:
- 781 model comparison with 84 AMS factor-analysis data sets, Atmos. Chem. Phys., 16, 8939-8962, 10.5194/acp-16 782 8939-2016, 2016.
- von Schneidemesser, E., Monks, P. S., Allan, J. D., Bruhwiler, L., Forster, P., Fowler, D., Lauer, A., Morgan, W. T.,
 Paasonen, P., Righi, M., Sindelarova, K., and Sutton, M. A.: Chemistry and the Linkages between Air Quality and
- 785 Climate Change, Chemical Reviews, 115, 3856-3897, 10.1021/acs.chemrev.5b00089, 2015.
- 786 Watson, L. A., Shallcross, D. E., Utembe, S. R., and Jenkin, M. E.: A Common Representative Intermediates (CRI)
- 787 mechanism for VOC degradation. Part 2: Gas phase mechanism reduction, Atmos Environ, 42, 7196-7204,
 788 https://doi.org/10.1016/j.atmosenv.2008.07.034, 2008.
- 789 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire
- 790 INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning,
- 791 Geosci. Model Dev., 4, 625-641, 10.5194/gmd-4-625-2011, 2011.

- 792 Wild, O., Voulgarakis, A., O'Connor, F., Lamarque, J. F., Ryan, E. M., and Lee, L.: Global sensitivity analysis of
- chemistry–climate model budgets of tropospheric ozone and OH: exploring model diversity, Atmos. Chem.
- 794 Phys., 20, 4047-4058, 10.5194/acp-20-4047-2020, 2020.
- Willmott, C. J., Robeson, S. M., and Matsuura, K.: A refined index of model performance, International Journal
 of Climatology, 32, 2088-2094, <u>https://doi.org/10.1002/joc.2419</u>, 2012.
- 797 Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry
- 798 (MOSAIC), Journal of Geophysical Research: Atmospheres, 113, <u>https://doi.org/10.1029/2007JD008782</u>, 2008.
- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A.,
- 800 Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne,
- J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann,
- 802 S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J.
- 803 Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in
- anthropogenically-influenced Northern Hemisphere midlatitudes, Geophys Res Lett, 34, L13801, Artn L13801
 Doi 10.1029/2007gl029979, 2007.
- 206 Zhao, B., Shrivastava, M., Donahue, N. M., Gordon, H., Schervish, M., Shilling, J. E., Zaveri, R. A., Wang, J., Andreae, M.
- 807 O., Zhao, C., Gaudet, B., Liu, Y., Fan, J., and Fast, J. D.: High concentration of ultrafine particles in the Amazon free
- troposphere produced by organic new particle formation, Proc Natl Acad Sci U S A, 117, 25344-25351,
- 809 10.1073/pnas.2006716117, 2020.
- 810 Zhao, B., Fast, J. D., Donahue, N. M., Shrivastava, M., Schervish, M., Shilling, J. E., Gordon, H., Wang, J., Gao, Y., Zaveri,
- 811 R. A., Liu, Y., and Gaudet, B.: Impact of Urban Pollution on Organic-Mediated New-Particle Formation and Particle
- 812 Number Concentration in the Amazon Rainforest, Environ Sci Technol, 55, 4357-4367, 10.1021/acs.est.0c07465, 2021.