

Monitoring sudden stratospheric warmings under climate change (1980-2021) based on reanalysis data verified by radio occultation

Ying Li¹, Gottfried Kirchengast², Marc Schwaez², Yunbin Yuan¹

¹ State Key Laboratory of Geodesy and Earth's Dynamics, Innovation Academy for Precision Measurement Science and Technology (APM), Chinese Academy of Sciences, Wuhan, 430071, China

² Wegener Center for Climate and Global Change (WEGC) and Institute of Physics, University of Graz, 8010 Graz, Austria

Correspondence to: Ying Li (liying@asch.whigg.ac.cn), Gottfried Kirchengast (gottfried.kirchengast@uni-graz.at)

Abstract. We develop a new approach to monitor Sudden Stratospheric Warming (SSW) events under climate change since 1980 based on reanalysis data, verified by radio occultation data. We construct gridded daily-mean temperature anomalies

10 from the input fields at different vertical resolution (basic case full resolution; cross-check with reanalysis at 10 stratospheric standard pressure levels or 10hPa and 50hPa level only) and employed the concept of Threshold Exceedance Areas (TEAs), the geographic areas wherein the anomalies exceed predefined thresholds (such as 30 K) to monitor the phenomena. We derived main-phase TEAs, representing combined middle and lower stratospheric warming, to monitor SSW warming on a daily basis. Based on the main-phase TEAs, three key metrics, including main-phase duration, area, and strength are estimated and used

15 for the detection and classification of SSW events. An SSW is defined to be detected if the main-phase warming lasts at least 6 days. According to the strength, SSW events are classified into minor, major and extreme. An informative 42 winters' SSW climatology 1980-2021 was developed, including the three key metrics as well as onsets date, maximum-warming-anomaly location and other valuable SSW characterization information. Detection and validation against previous studies underpins that the new method is robust for SSW detection and monitoring and that it can be applied to any quality-assured reanalysis

20 and observational temperature data that cover the polar region and winter timeframes of interest, either using high vertical resolution input data (preferable basic case), coarser standard-pressure-levels resolution or (at least) 10hPa & 50hPa pressure level data. Within the 42 winters, 43 SSW events were detected for the basic case, yielding a frequency of about one event per year. In the 1990s, where recent studies showed gaps, we detected several events. Over 95% of event onset dates occurred in deep winter (Dec-Jan-Feb timeframe; about 50% in January) and three quarters have their onset location over Northern Eurasia

25 and the adjacent polar ocean. Regarding long-term change, we found a statistically significant increase in the duration of SSW main-phase warmings, by about 5(±2) days over the climate change period from the 1980s to the 2010s, raising the average duration by near 50% from about 10 to 15 days and inducing an SSW strength increase by about 40(±25) million km² days from about 100 to 140 million km² days. The results are robust (consistent within uncertainties) across using different input data resolution. They can hence be used as a reference for further climate change-related studies and be a valuable basis for

30 studying SSW impacts and links to other weather and climate phenomena, such as changes in polar vortex dynamics and in mid-latitude extreme weather.

删除的内容: and also a trailing-phase TEA to monitor potential upper stratospheric cooling in the wake of the warming phase.

删除的内容: Main-Phase Strength (MPS), Duration (MPD)

删除的内容: Area (MPA),

删除的内容: , enabling

删除的内容: ,

删除的内容: event categories.

删除的内容: 2020

删除的内容: strength and

删除的内容: .

删除的内容: 40

删除的内容: 0.95/

删除的内容: About

删除的内容: also

删除的内容: 4 days

删除的内容: 40

删除的内容: 14

删除的内容: near 30

删除的内容: (about 30 %)

删除的内容: 105

删除的内容: 135

删除的内容: long-term

55 1 Introduction

Sudden Stratospheric Warming (SSW) describes an atmospheric variability phenomenon at daily-to-monthly scale where temperature in middle stratosphere (about 30 km or 10 hPa) increase rapidly (>30 to 40 K) within a couple of days in sub-polar and polar regions (McInturff et al., 1978; Butler et al., 2015; Baldwin et al., 2020). In extreme cases, SSW temperature anomalies can reach more than 70 K relative to the long-term mean.

60 During a strong event, the westerly zonal mean zonal winds of the polar vortex can reverse, and the three-dimensional polar vortex can undergo a displacement or split (Charlton and Polvani, 2007; Hu et al., 2015; Butler and Gerber, 2018). SSWs are generally understood to be caused by tropospheric planetary waves, which penetrate into the stratosphere, and then influence the stratospheric polar vortex (McInturff et al., 1978; Thompson et al., 2002; Labizke and Kunze, 2009). Solar radiation are also believed to be one of the causes of stratospheric warming. SSWs usually occur in the polar regions of the northern hemisphere (beyond 60°N), while they rarely occur in the

65 southern polar region due to less tropospheric planetary wave activity (Van Loon et al., 1973). We therefore focus on SSWs of northern hemisphere in this SSW-ensemble-based analysis over multiple decades since 1980.

SSWs are an important indicator of polar winter variability. They strongly interact with the troposphere (Hitchcock and Simpson, 2014; Lehtonen and Karpechko, 2016), mesosphere (Vignon and Mitchell, 2015; Singh and Pallamraju, 2015) as well as the upper atmosphere and ionosphere (Jonah et al., 2013; Kakoti et al., 2020) through atmospheric circulations and

70 thermodynamics that mediate stratosphere-mesosphere-thermosphere couplings. The warming in the middle stratosphere will, on the one hand, propagate downwards to lower altitude levels and cause longer lasting warming in lower stratosphere (Hitchcock and Simpson, 2014). Some extreme events have impacts into deep troposphere and cause large area of blocking high pressure, and subsequently cause cold weather in northern Europe, eastern Asia and northern America regions (Cattiaux et al., 2010; Yu et al., 2015; Tyrlis et al., 2019; Hall et al., 2021). Some SSW events also cause the cooling of mesosphere and

75 elevated stratopause (Holt et al., 2013; Vignon and Mitchell, 2015; Singh and Pallamraju, 2015). In the ionosphere, the distribution of electron density are found to be changed as a response to SSW (Nayak and Yigit, 2019; Kakoti et al., 2020). Due to atmospheric meridional circulation, the tropical stratosphere is found to be cooling at the same time as there is polar stratospheric warming (Yoshida and Yamazaki, 2011; Dhaka et al., 2015). Regarding atmospheric composition and chemistry, such as the distribution of ozone, water vapour and energetic particle precipitation, these are found to be changed as well

80 (Kuttipurath and Nikulin, 2012; Ayarzaguna et al., 2013; Holt et al., 2013).

Given this variety of strong interactions of SSWs, it is important to accurately observe, detect, and monitor such events, including their possible transient changes under climate change. Accurate SSW observations require high quality data to be sufficiently dense in polar stratosphere. However, observations in these regions are notoriously sparse. Early studies used radiosonde or rocketsonde to observe SSWs. However, both datasets are generally land-limited and cannot provide high

85 vertical resolution and high quality data throughout the lower, middle, and upper stratosphere. With the advent of satellite era

刪除的內容: Many major SSWs occur accompanied by regular

刪除的內容: reversed and/or

刪除的內容: displaced or even

刪除的內容: by the Quasi-Biennial Oscillation (QBO)

刪除的內容: the Southern Oscillation (SO) in the tropics, which

刪除的內容: , and also by solar radiation

in the 1970s, it is possible to put instruments, such as microwave limb sounders, infrared spectrometers and radiometers, on satellites in order to observe the atmosphere globally (e.g., Hitchcock et al., 2013; Noguchi et al., 2020). However, satellite passive sounding data come in the form of radiances, which only allow coarse vertical resolution limiting the accurate 95 conversion to altitude-resolved temperature or wind profiles, which are key variables for reliable SSW monitoring. With the advances of atmospheric data assimilation systems, reanalysis data have become quite a reliable data source for long-term atmospheric analyses, due to their advantages of regular sampled in space and time and capability to provide reasonably reliable data up to the stratopause (e.g., Charlton and Polvani, 2007; Yoshida and Yamazaki, 2011; Butler et al., 2017, 2018; Hersbach et al., 2020). However, reanalysis data may have some inhomogeneities and irregularities in the long-term, due to episodic 100 observation system updates and adding in a diversity of new streams of observation datasets over multi-decadal time ranges; they are not a direct long-term consistent observation of the atmosphere. It is hence important and to verify results based on them by complementary use of observational data records with better long-term stability.

In addition to the sparsity of robust observation techniques, SSWs also have no community-agreed standard definition for reliable detection and monitoring. Butler et al. (2015) provided a detail overview on the history of various SSW definitions 105 and calculated SSW frequency to cross-evaluate nine different definitions based on reanalyses data. Their results suggest that frequencies obtained using different definitions vary a lot, from about 0.46 to 0.81 events per year, and the onset (or maximum anomaly) dates of major SSWs for each definition may differ substantially as well. Reasons of these discrepancies are mainly related to method design. Each definition has its own unique characteristic. For example, definitions based on wind reversal may more focus on polar vortex, and definitions based on stratosphere-troposphere impacts may more focus on the impacts of SSW on troposphere. Definitions based on one latitude or region are more sensitive to such variations than definitions based on larger domains (Butler et al., 2015). Also the details of implementation in selecting detection variable, latitude, altitude, thresholds and background climatology information can make the results different. These discrepancies of SSW definitions make consistent statistical assessments of SSWs more difficult. Furthermore, at the side of atmospheric physics and dynamics, the analysis of other weather and climate phenomena that relate to SSWs is more limited in scope if accurate SSW diagnostics 110 and monitoring cannot be given.

To help mitigate these current limitations, we here propose and apply a new method to monitor SSW events over the 1980 to 2021 northern hemisphere winter half-years, using Global Navigation Satellite System (GNSS) Radio Occultation (RO) data (Angerer et al., 2017) and ECMWF Reanalysis 5th generation (ERA5) data (Hersbach et al., 2020), developing a 42 years' SSW events tracking, and evaluating their characteristics.

120 GNSS RO is an atmospheric remote sensing technique to provide vertical atmospheric profiles, such as of temperature, density, and pressure (Kursinski et al., 1997; Kirchengast, 2004). RO data have distinctive advantages of high vertical resolution, high accuracy, long-term consistency, and global coverage (Anthes et al., 2011; Steiner et al., 2011). The vertical resolution of RO in the stratosphere is about 1 km, which is very high compared with other global observation techniques. Validation results against radiosonde and verification with (re)analysis data (that generally assimilate RO data) suggest that the data are of small 125 discrepancy (< 2 K) in the upper troposphere and lower stratosphere (Scherlin-Pirscher et al., 2011a, b; Ladst äler et al., 2015;

删除的内容: Reasons of these discrepancies include both data and method reasons. Regarding data aspects, reanalyses data, as stated above, may have inhomogeneities over time.

删除的内容: Regarding method aspects,

删除的内容: detail

删除的内容: quite

删除的内容: rather

删除的内容: also severely hindered

删除的内容: Motivated by the aim to

135 Steiner et al., 2020a, 2020b). RO data from different satellites can be combined without inter-calibration, which make them very suitable for climate-related studies (Foelsche et al., 2011; Angerer et al., 2017; Steiner et al., 2020a). Finally, since a multi-RO satellite observation record started in 2006 (Angerer et al., 2017), the geographic data coverage is sufficiently dense for monitoring and analysing regional-scale phenomena such as SSWs from that time onwards.

140 Complementary to reanalysis datasets, which also offer dense coverage, RO reprocessing datasets hence feature accurate and long-term stable observational records of climate benchmark quality (Steiner et al., 2020a), allowing for stable conditions for SSW monitoring over decades. Therefore, given the complementarity of these single-source long-term consistent benchmark observations to reanalyses (Bosilovich et al., 2013; Parker, 2016; Simmons et al., 2020; Hersbach et al., 2020), RO data are ideal for SSW studies.

145 An initial use of RO data for SSW study was by Klingler (2014) who was the first to use the data to examine the temperature changes during the 2009 SSW event. A couple of studies have also used RO data to analyse their impacts on gravity wave activity, the ionosphere, and also the tropical atmosphere (Yue et al. 2010, Lin et al. 2012, Dhaka et al., 2015). However, use for longer-term SSW detection and monitoring is a next step to be made. We have carried out an initial study (Li et al., 2021), where we used RO data and ERA5 data to develop a new threshold-exceedance-area-based approach to monitor and characterize the strong and well-known 2009 SSW event. We revealed, in principle, high potential for the new method to be 150 used for detection and monitoring of SSWs over multi-decadal timeframes as well.

155 In this study, building upon the initial Li et al. (2021) work, we apply the approach over 14 winters of RO data (2007 to 2020) and 42 ones (1980 to 2021) of ERA5 data, using the former reprocessing record to cross-verify the latter reanalysis dataset for the purpose. We derive robust SSW characterization metrics, a new definition based on temperature field data, and apply the new method for SSW detection, classification, and monitoring, and to explore long-term changes in their characteristics under the recent climate change.

The paper is structured as follows. Section 2 briefly assesses current SSW detection methods and definitions and then summarizes the features of our new method. Section 3 introduces the data and methodology of our method and Section 4 presents and discusses the results. Finally, conclusions are given in Section 5.

2 Assessment of current definitions and the new method

160 2.1 Current definitions

A Sudden Stratospheric Warming (SSW) was first observed by Richard Scherhag using radiosonde measurements in Berlin, Germany, in January/February 1952 (Scherhag, 1952). when he found an abrupt temperature increase in stratosphere. After about a decade, the World Meteorological Organization (WMO) Commission for Atmospheric Sciences (CAS) developed an international SSW monitoring program called STRATALERT based on available radiosonde and rocketsonde observations 165 (WMO/IQSY 1964). The WMO CAS suggested to provide an SSW warning, when a sudden and unusual increase in temperature at 30 km or above is detected.

删除的内容: basically very suitable

删除的内容:). He

With time ongoing and after more events were observed, it was well recognized that many SSWs occurred along with wind
170 reversals and/or polar vortex displacement or split (Johnson et al., 1969; Charlton and Polvani, 2007). Since the 1970s, many
studies combined temperature increase and wind reversals to detect SSWs, though detailed implementation and thresholds
used are different (e.g., Schoeberl, 1978; Labitzke, 1981). An often-used definition at this stage was suggested by McInturff
et al. (1978). He defined that a SSW event occurred if temperature increase more than 25 K and the event was defined as a
major one if a stronger temperature increase jointly observed with wind reversal.

175 Since wind reversal is one of the most important features of SSWs, many studies suggest using wind reversal for detecting
major SSW events (e.g., Charlton and Polvani, 2007; Hu et al., 2015; Butler et al., 2015, 2017; Butler and Gerber, 2018). One
of the most often used definitions is the one from Charlton and Polvani (2007) (denoted as CP07 hereafter): a major SSW
occurs, when the zonal mean zonal winds at 60 °N at the 10hPa level become easterly during wintertime. Wind reversal is a
simple and robust definition and is useful in studying many influences triggered by SSW. In addition to wind reversal
180 definitions, there are also studies that used vortex moment, which is another important characteristic of the warmings, to detect
SSW events (Seviour et al., 2013; Mitchell et al., 2013). Furthermore, polar-cap or zonal-mean stratospheric at 10hPa
geopotential height anomalies were used to detect SSW events (Baldwin and Thompson 2009; Gerber et al., 2010).

185 Butler et al. (2015) tested the sensitivity of SSW detection results to nine definitions and found SSW frequencies obtained
under these different definitions to vary from 0.46 to 0.81 events per year, and the onset dates also to vary substantially. There

are several reasons of these discrepancies. First of all, while SSWs usually occur along with wind reversal or polar vortex
change, this is not always the case. Several studies detected significant stratospheric warming but did not simultaneously detect
190 wind reversals at the commonly used 60 °N or 65 °N latitude. For example, Mitchell et al. (2013), who used vortex geometry
for diagnosing SSW, found half of their events inconsistent with those obtained by CP07. Therefore, definitions based on
different parameters can make detection results different. Secondly, single-latitude and/or single-altitude definitions such as
for wind reversal may miss some important SSWs that occurred primarily in other latitude-altitude domains (e.g., Manney et
al., 2015; Singh et al., 2015). Thirdly, definitions can be sensitive to background climatology and specific thresholds used,
especially if based on broad polar cap-mean anomalies. Finally, current definitions either use zonal-mean or polar cap-mean
195 results, which do not enable dynamical 3D tracking of such events, and therefore can only provide information of onset date.
However, the dynamic location and strength information is rather important for studies on SSW's interactions with other
phenomena, both regarding causes and impacts. Furthermore, SSW is also not a one level phenomena, its occurrence and
impacts are related to almost the complete stratosphere.

200 In view of the studies we surveyed and from our own initial study (Li et al., 2021), we suggest that a new SSW monitoring
method should build upon the temperature field that directly expresses the sudden stratospheric warming, for quantifying this
anomalous thermal behavior as the primary SSW fingerprint. It should robustly detect and characterize SSW events, from
minor to extreme ones, as a whole phenomenon, without being unduly sensitive to details. The method should also be readily
applicable to both observational and model data (as long as they are sampled sufficiently densely, preferably grid-based), and
not need adjustment to any specific suitable dataset (e.g., a particular reanalysis, atmospheric forecast, climate model

删除的内容: quite

删除的内容: who proposed

删除的内容: moderate

删除的内容: anomaly of

删除的内容:

删除的内容: is regarded a minor

删除的内容: , while

删除的内容: is regarded a major event.

删除的内容: to use

删除的内容: quite

删除的内容:

删除的内容: literature

删除的内容: standard

删除的内容: best directly

删除的内容: name-coining

删除的内容: as such (i.e., be temperature-field-based),

删除的内容: directly

删除的内容: the polar winter

删除的内容: variability

删除的内容: its

删除的内容: Secondly, it

删除的内容: Thirdly, the

225 simulation, or observational data record). Finally, upon detection, the SSW monitoring metrics should be informative on the duration, strength, and dynamic location of each SSW, in order to facilitate long-term change monitoring and effective use in cause and impact studies. Implementing these suggestions, we propose our new method and definitions in Section 2.2.

2.2 New method and its features

230 SSWs, as reflected by their name, were originally determined by their strong and [rapid](#) temperature increase. Therefore, in the method proposed here, we use temperature as the key variable for the diagnostics. Compared to wind- and polar vortex-based definitions, temperature is a more accessible parameter that can be obtained from various observations. In addition, temperature is a well-related parameter, when analyzing SSW relations to other phenomena in the troposphere as well as the mesosphere and thermosphere/ionosphere. Many studies chose to use temperature solely, or combined it with wind-field changes in studying impacts of SSWs (e.g., Zhou et al., 2002; Siskind et al., 2010 Manney et al., 2015; Jonah et al., 2013; Kakoti et al., 235 2020; Singh and Pallamraju, 2015; Vignon and Mitchell, 2015). Also, further thermodynamic variables, in particular air density and pressure, are often readily available for auxiliary co-information (Li et al., 2021).

240 Based on temperature changes, we designed the method to be fairly insensitive to temperature field details. Firstly, we use robust stratospheric temperature anomaly profiles [or pressure level data](#) at any data location (such as a grid point) as the basis for expressing the local warming; using an anomaly technique that has been proved useful and robust in diagnosing many other climate and atmospheric change phenomena such as related to tropical cyclones (Biondi et al., 2015), atmospheric blocking (Brunner et al., 2016) or thermal imprints of wildfires (Stocker et al., 2021). As described in detail by Li et al. (2021) and summarized in Section 3 below, we then calculate vertical mean anomalies in selected stratospheric layers, [or at selected levels](#), and categorize them into large-scale grid cells covering the polar region, based on which we compute, on a daily basis throughout wintertime, temperature threshold exceedance areas and related metrics, which serve as the basis for SSW detection 245 and monitoring.

250 In establishing SSW climatologies (i.e., cataloguing SSW events over a multi-decadal period), previous studies generally do only provide information about onset dates and vortex-split or displacement. In the climatology we build based on the new method, we can provide SSW event onset date (of maximum middle stratosphere warming), duration, exceedance area, and strength, as well as complementary day-by-day dynamic tracking of the center location, associated maximum warming, and areal extent of the exceedance area. Furthermore, [most previously](#) published climatologies [do not yet](#) reach [beyond](#) 2013 and [miss some events](#) over the 1990s decade, while we provide a climatology continuously extending from 1980 to 2021 and hence filling these gaps.

255 Compared to our initial method introduction and its careful evaluation in Li et al. (2021), which was based on the strong 2009 SSW event only, we focused and refined the diagnostics towards the metrics so that it now fully deploys for multi-decadal detection, classification, and monitoring. The details on data and methodology are described next in Section 3.

删除的内容: quick

删除的内容: previous

删除的内容: to

删除的内容: only

删除的内容: lack quality

3 Data and methodology

3.1 RO data

Since 2001, a continuous record of RO data is provided by GNSS RO missions, including the Challenging Mini-satellite Payload mission (CHAMP; Wickert et al., 2001), followed by the Gravity Recovery and Climate Experiment (GRACE,

265 Wickert et al., 2005), and the Constellation Observing System for Meteorology, Ionosphere, and Climate (COSMIC; Schreiner et al., 2007), the European Meteorological Operational satellites (MetOp; Luntama et al., 2008), the Chinese FengYun-3C operational satellite (Sun et al., 2018), and others. Since the launch of COSMIC near mid-2006, which was a constellation of six satellites, there was sufficient coverage with RO event observations for regional-scale studies such as on SSWs. Therefore, in this study, we use the RO data record from the wintertime 2006/07 onwards.

270 We use the atmospheric profiles from the Wegener Center for Climate and Global Change (WEGC), processed by its Occultation Processing System version 5.6 (denoted as OPSv5.6 hereafter). Several studies that introduced, validated and evaluated the OPSv5.6 record, showed that these data are of high quality (e.g., Ladst  ter et al., 2015; Angerer et al., 2017; Scherlin-Pirscher et al., 2017; Steiner et al., 2020a; Schw  rz et al., 2021). A detailed discussion of quality aspects is provided by Angerer et al. (2017). Based on the record available to end 2020, we use RO data from the winter seasons W06-07 until

275 W19-20, which are 14 winters ("W") in total, which we define to comprise the extended-winter season from November to March (hence, for example, "W06-07" contains the November 2006 to March 2007 timeframe). We use the OPSv5.6 multi-satellite data from COSMIC, CHAMP, GRACE, MetOp, SAC-C, in the form as available from the WEGC dataset (Schw  rz et al., 2021).

3.2 ECMWF Reanalysis 5 (ERA5) data

280 ERA5 is the fifth-generation ECMWF atmospheric reanalysis of the global weather and climate (Hersbach et al., 2019, 2020; Simmons et al., 2020). It was produced for the European Copernicus Climate Change Service (C3S) by ECMWF and replaces the ERA-Interim reanalysis (Dee et al., 2011), which stopped being produced by August 2019. ERA5 combines vast amounts of historical observations into global atmospheric gridded field estimates, using ECMWF's modeling and data assimilation system. The basic resolution of ERA5 used is about 30 km horizontal resolution and 137 vertical levels from the surface up to

285 an altitude of about 80 km (basic case with full resolution, also termed "full-res ERA5"). We use the ERA5 datasets from 1979 onwards, over the 42 winters from 1979/80 (W79-80) to 2020/21 (W20-21), fully encompassing also the RO data period.

For crosscheck of the dependence of detection and monitoring results on vertical resolution of the input temperature fields, we alternatively also used the data at coarser standard pressure levels (using the C3S 37 pressure levels dataset, also termed "37-

290 plevels ERA5") or, as a minimum-input case, at the 10 hPa and 50 hPa pressure levels only (termed "10 hPa & 50 hPa level ERA5"). These crosschecks are reported as part of the long-term monitoring results in Section 5 below, otherwise the basic-case resolution was used throughout for introducing, illustrating and describing the method. For the stratospheric focus of this study we note that the basic-case full resolution data cover the stratosphere with roughly 1 km vertical resolution (i.e., they are

删除的内容: the Constellation Observing System for Meteorology, Ionosphere, and Climate Experiment (GRACE; Wickert et al., 2005).

删除的内容: is

删除的内容: is

删除的内容: . Here we

删除的内容: quality-assured

删除的内容: that are provided

300 well altitude-resolved) while the 37 pressure levels include no more than 10 levels from 70 hPa (~19 km) to 1 hPa (~48 km) (70, 50, 30, 20, 10, 7, 5, 3, 2, 1 hPa). The latter hence correspond to a quite coarser resolution but still easily allow to compute reasonable finer-sampled vertical temperature profiles for our method's input, which we constructed by simple and robust vertical linear interpolation between the pressure level temperatures.

305 ERA5 data are used jointly with RO data for two purposes. On the one hand they are used as part of cross-checking the new method, to make sure that the method can be applied to both RO and reanalysis data. On the other hand (and even more relevant), since dense RO observations are only available from the year of 2006 onwards, we need to have reanalysis data, which provide much longer data records, to fully explore our method and to develop a long-term SSW climatological record.

310 In terms of horizontal resolution, we use them on a 2.5 °latitude \times 2.5 °longitude grid, to provide an adequate resolution that also roughly matches the RO horizontal resolution of about 300 km (e.g., Kursinski et al. 1997; Anthes et al. 2008). Temporal resolution used is 6 hours (four time layers per day, at 00, 06, 12 and 18 UTC), following the experience of many previous studies that intercompared and/or jointly used atmospheric (re)analysis and RO data (e.g., Gobiet et al., 2007; Scherlin-Pirscher et al., 2011; Angerer et al., 2017; Steiner et al., 2020a; Li et al., 2021).

315 Figure 1 illustrates characteristics of the RO and ERA5 profile datasets as relevant for the present SSW study, including on the daily number of events available (Fig. 1a) and for exemplary days during SSW events (Fig. 1b-d, see caption for details).

320 Evidently RO observational atmospheric profile data are comparatively sparse over the (northern high-latitude) region of interest, while ERA5 as a gridded dataset regularly provides its profiles at each and every of its grid cells without sparsity. Hence, while the number of RO profiles is of the order of several hundred per day, the ones of ERA5 amount to near 10000 per day. Furthermore, as Figure 1b-d shows, the spatial distribution of ERA5 data is regular-on-grid, while RO events occur with reasonable overall coverage but irregular sampling in detail. A few exemplary events ("Event1" to "Event3") are highlighted, against the back-plot of illustrated SSW temperature anomalies over the polar region, in order to use them next to explain the methodology.

3.3 Methodology

325 Since we recently provided a detailed basic introduction of the new SSW monitoring approach in Li et al. (2021) and discussed main overall features in Sect. 2.2 above already, we restrict to a brief summary here, supported by a schematic overview, concise tabular information, and focusing on updates and refinements since that introduction, which provides further technical details.

330 As a general overview, Figure 2 provides a schematic summary of the method's workflow. It highlights the computation sequence from starting with the temperature field input data via anomalies construction and gridded-maps generation in stratospheric layers or at pressure levels to extract the threshold exceedance areas (TEAs) of different SSW phases and finally deriving the SSW metrics then used for detection, classification, and characterization. Table 1, which is a condensed and refined update of Table 1 in Li et al. (2021), complements this schematic overview by summarizing the basic parameters and features of the method in more detail along with respective explanations (rightmost column). This ranges from definition of

删除的内容: Firstly,

删除的内容: Secondly,

删除的内容: We

删除的内容: then

删除的内容: ×

删除的内容: eg.

删除的内容: In the vertical, we use the model-level resolution (137 levels), which is sufficiently dense for our stratospheric altitude range over 20 to 50 km.

删除的内容: summarizes

删除的内容: ,

345 the temperature anomalies and the related daily TEAs in three characteristic stratospheric layers (lines (1)-(4)) via the anomaly-
maximum value and location (lines (5)-(6)) to the four derived TEA key variables during SSW events (lines (7)-(10)) that
monitor and characterize the different SSW phases. We note that the daily TEAs for the three layers (lines (2)-(4)) are computed
for a range of threshold values, for convenient closer insight to the depth of the anomalies (as illustrated in Section 4), while
only one basic threshold value (bold-faced in lines (2)-(4), middle column) is subsequently used for defining the TEA key
variables (lines (7)-(10)). All selections and settings summarized and explained in Table 1, including those in the footnotes a-
350 c related to the vertical resolution options, are based on very extensive sensitivity tests of all the choices, using the ERA5 and
RO data as the testing input datasets.

355 For aiding the understanding on how the profile and layer-main anomalies typically look like, Figure 3 graphically illustrates
the construction of the anomalies and variables by way of the three example RO events indicated in Figure 1 as well as
correspondingly for three ERA5 profiles from adjacent grid points. It can be seen that RO and ERA5 profiles are overall
consistent, with the latter profiles somewhat smoother in their resolution of vertical variability. Deviations of temperature and
corresponding climatological profiles are smallest for Event1 that is located in a non-warming area (cf. Fig. 1b), Event2, and
Event3 that are most affected, show larger deviations and anomalies than Event1 since these are located in the warming area
of the SSW (cf. Figs 1b, 1d). Largest anomalies for the latter two events are found in mid-stratosphere layer (30-35 km), with
values is about 45 K and 60 K, respectively. Maximum values in the other altitude layers are smaller. In this way, these few
360 examples are consistent with the broader and long-term picture over many SSW events (see Sect. 4 below), which show the
SSW warming to be strongest in the middle stratosphere (about 30 to 40 km).

365 While Li et al. (2021) introduced and initially tested the approach based on the single 2009 SSW event, we here made sure for
the long-term application that the four TEA key variables are captured and exploited in a way so that they reliably detect and
quantify actual SSW warmings, and cooling in the trailing phase if it occurs, among the on-going weaker and more “random”
370 polar variability due to other driving factors. The rightmost column of Table 1 (lines (7)-(10) therein) summarizes the criteria
that we chose for them, after comprehensive testing and sensitivity studies both with the RO and ERA5 data, including on the
different options of vertical resolution of the ERA5 temperature input fields. Based on these well-selected TEA key variables
and the auxiliary variables on maximum-values and locations, all prepared at daily sampling, we could finally define the
fundamental metrics and criteria that we use for the detection, classification and further characterization of SSWs. These
definitions were accompanied by another comprehensive portfolio of sensitivity tests and the choices distilled are summarized
in Table 2, including brief explanations (rightmost column therein and footnote a).

375 Overall, the extensive robustness and sensitivity testing provided us with due evidence and confidence that the new method,
effectively constructed from the dynamic temperature anomaly field as perturbed by the SSW, should enable a new level of
quality and quantitative insight into SSWs also in the long-term, maturing the methodology from our initial Li et al. (2021)
single-event study. We hence as a next step applied the method with the parameters and definitions summarized in Table 2 to
the complete RO and ERA5 datasets and discuss the results below.

删除的内容: threshold exceedance areas (

删除的内容:)

删除的内容: 2

删除的内容: this

删除的内容: similar

删除的内容: SSW

删除的内容: careful

删除的内容: qualification

删除的内容: These

删除的内容: Extensive

删除的内容: these characteristics

删除的内容: completing

删除的内容: new

390 4. SSW detection and monitoring results

4.1 Polar cap mean anomalies overview

Polar cap daily mean temperature anomalies over the 15 winters period from W06-07 to W20-21 are presented in Figure 4, where both the RO and ERA5 datasets overlap for 14 winters. The RO and ERA5 polar cap results are closely consistent, with anomalies from ERA5 typically about 5 K (occasionally up to 10 K) larger than RO data above 35 km. These differences can

395 mainly be attributed to the denser sampling of ERA5, leading to less spatial smoothing. In general, the overall close results of these independently produced and quite differently-sampled datasets (for detailed discussion see Li et al., 2021) lend confidence that we may use ERA5 data for the inspection of the multi-decadal time period from 1980 onwards.

Complementary back-extended polar cap temperature anomalies of ERA5 for the 27 winter periods W79-80 to W05-06 are shown in Figure 5. The results provide a neat first overview on which winter seasons hosted potentially strong SSWs (e.g.,

400 W84-85, W03-04, W18-19) and which comparatively quiet, including with little evidence of SSWs (e.g., W81-82, W93-94, W10-12). It also reveals to be a very salient feature already in this polar cap-mean inspection that strong SSW events often are entailed by a distinct upper stratospheric cooling, which is hence reflected in our metric definitions (see Table 2, lines (6)-(8)). Another interesting feature hinted here already is that in the 1990s, where existing SSW climatologies detect very few events (e.g., Charlton and Polvani., 2007; Hu et al., 2015; Butler and Gerber., 2018), we do detect several reasonably strong SSWs

405 that classify as major events (more details in Sect. 4.4).

Some distinct temperature anomalies are found in almost every winter in one or another form, which underlines the fact that the polar stratosphere is quite variable in winter. Regarding SSWs, some winters signal one strong warming, while some others show multiple moderately strong or minor warmings. Strong warming propagate to lower altitude levels and also cause longer-lasting warmings, and as noted above may be accompanied by distinct cooling (such as in the most recent decade W12-13 and

410 W18-19). A final observation from this basic synoptic view is that the altitude range of maximum warming varies somewhat from event to event. For example, the W11-12 warming is largest at about 35 to 40 km, while the W17-18 warming exhibits its largest anomalies at about 25 to 30 km. Our definition of main phase, combining middle and lower stratospheric TEA diagnostics (see Table 1, line (9)), robustly captures such different specific event dynamics.

4.2 SSW threshold-exceedance-area representative results

415 Figure 6 illustrates TEA spatial contour map results for typical daily temperature anomalies across the temporal evolution (top to bottom) of three representative SSW events of increasing strength (left to right; those already used for back-plot in Fig. 1). Evidently, both the temperature anomalies' magnitude and warming area increase from minor to extreme event, as expected based on our classification, which is in particular visible in the mid-stratosphere temperature anomaly on the event onset date (second row). Complementary to this, the snapshot day shown from the trailing phase (four weeks after onset date, bottom row) highlights that the extreme event (right) exhibit a very distinct upper stratosphere cooling anomaly, exceeding -40 K in a TEA of about 10 million km^2 size.

删除的内容: .

删除的内容: Figure 3 illustrates RO (Fig. 3a) and ERA5 (Fig. 3b) polar cap (60°N – 90°N)

删除的内容: and where RO data can hence be used to cross-verify ERA5 data.

删除的内容: appear rather similar

删除的内容: data

删除的内容: somewhat

删除的内容: closely similar

删除的内容: Figure 4 shows the complementary

删除的内容: The results

删除的内容: overall similar with

删除的内容: 3 and both together

删除的内容: appear

删除的内容: not any relevant sign

删除的内容: appears

删除的内容: metrics

删除的内容: Figures 3 and 4 also show that some

删除的内容: entailed

删除的内容: found

带格式的: 字体: 八号

删除的内容: 5 shows

删除的内容: maps of

Following this representative spatial view on individual daily TEAs, Figure 7 depicts, for the same events as shown in Figure 6, how our method leads from TEA time series in the three stratospheric layers (first three rows of Fig. 7) to the TEA key variables (fourth row) from which finally the SSW metrics for the event characterization according to Table 2 are derived.

Daily TEA values over positive thresholds quantify the size of exceedance areas of warming while those over negative thresholds diagnose the exceedance areas of cooling. It can be seen, for example, that MSTA-TEAs over positive thresholds of all events increase rapidly to maximum and then quickly decrease, indicating the typical sudden warming of the primary phase. LSTA-TEAs over positive thresholds are overall of smaller magnitude but longer duration, and with maxima delayed against MSTA-TEAs, from the SSW downward propagation. Regarding event strength, while MSTA-TEAs >30 K of major and extreme events are of small discrepancies within $2 \times 10^6 \text{ km}^2$, the duration of the extreme event is longer than for the major event. The USTA-TEA timeseries of the extreme event shows the distinct several-weeks-long cooling behavior in the trailing phase.

The TEA key variables (Fig. 7j to 7l) capture the essential daily TEA information per SSW event, in the form summarized in lines (7)-(10) of Table 1, as the basis for the three key metrics per event according to lines (1)-(3) of Table 2, the quantified values of which are shown in the panel legends. As already indicated by the polar-cap-mean view in Figs 4 and 5, it is well seen here that the ERA5 TEAs (heavy lines) are generally higher than from the RO data (thin lines), which in particular applies for the trailing-phase cooling of the extreme event (Fig. 7l) in the upper stratosphere, where the sparser sampling by RO events

leads to the relatively largest difference. In terms of magnitudes, it is clearly visible from the values that the MPS, MPD and MPA metrics reach that the event strength substantially grows from minor to extreme events (see also the strength class definitions based on the MPS in line (5) of Table 2).

The polar-map plots of Figure 7 (bottom row) finally illustrate the dynamic event tracking information of the SSW-PP-TEA, SSW-SP-TEA and SSW-TP-TEA timeseries for the three representative events. This view enables to see the geographic trajectory of the daily anomaly center location (maximum-value location, cf. line (6) of Table 1) together with an indication of the anomaly magnitude (color of corresponding TEA thresholds). This type of plots helps the detailed diagnostics and characterization of any specific event as introduced in Sect. 2 above (for details see also Li et al., 2021).

4.3 SSW detection and metrics-tracking results

Figures 8 and 9 employ the view introduced in Fig. 7j-7l to display the TEA diagnostics and MPS/MPD/MPA metrics results for all SSWs detected over the full multi-decadal period from 1980 to 2021 (Fig. 8 for the recent winters since 2001/02 and Fig. 9 for those before, up to 2000/01). In the recent two decades, typically one or two SSWs have occurred during almost every winter (except in W04-05 and W10-11), while in the two decades before there have been somewhat more SSW-quiet winters.

The strongest event during the entire period is the one in W08-09, with a main-phase strength (MPS) of over 360 million km^2 days for the ERA5 data (330 million based on the cross-verifying RO data). The second strongest event is the one of W18-19, where the MPS from ERA5 exceeded 290 million km^2 days. Additionally, the winters W01-02, W12-13 and W17-18, as well

删除的内容: from both warming and cooling anomalies, Fig.6 illustrates

删除的内容: three

删除的内容: (left to right)

删除的内容: timeseries

删除的内容: above polar-map plots

删除的内容: overall

删除的内容: relatively similar

删除的内容: 6j

删除的内容: 6l

删除的内容: 3

删除的内容: 4

删除的内容: 6l

删除的内容: 6

删除的内容: 7

删除的内容: 8

删除的内容: 6j-6l

删除的内容: 7

删除的内容: 8

删除的内容: Figure 7 shows that

删除的内容: Fig. 8 indicates that

删除的内容: in the two decades before

as the winters W84-85, W87-88, and W88-89 in the 1980s, hosted extreme events exceeding our classification threshold of
500 $180 \times 10^6 \text{ km}^2 \text{ days}$. Half of these eight extreme events also are seen to have caused a strong upper stratospheric cooling during
the trailing phase that lasted for more than a month.

In addition, several major events occurred (e.g., W02-03, W03-04, W05-06), with the MPS of such events varying from 90 to
180 $\times 10^6 \text{ km}^2 \text{ days}$ as defined in Table 2. Also two of the major events caused a long-lasting upper stratospheric cooling in the
trailing phase (in W03-04 and W05-06), which is an exception for these events, however. Together with these major events
505 also a range of minor events were detected and diagnosed, exhibiting an MPS smaller than $90 \times 10^6 \text{ km}^2 \text{ days}$. Based on these
long-term SSW tracking results shown in [Figures 8](#) and [9](#), we are now prepared to collect and analyze the long-term results in
a more climatological-statistical manner, including inspection for possible long-term transient changes in the record [under](#)
[climate change](#).

删除的内容: Figs 7

删除的内容: 8

5 SSW climatology and [trends](#) under climate change

5.1 SSW climatology

Based on the detection results in [Section 4](#), a climatological summary of the ERA5-based results for all SSW events detected
during the entire 42-years period [is provided in Table 3](#). It lists, for each event, the main characteristics as defined by Table 2
and [intercompares](#) the onset date to the onset dates found by the Butler and Gerber (2018) [work](#) (BG18) that extended to the
year 2013. The [intercomparison](#) reveals that a range of minor and major events was not part of the BG18 list, while several are
515 part of that list and not detected here. In general, the onset dates detected by the new method introduced here are consistent
with those in the BG18 climatology, with coincidence of the dates commonly within ± 1 day.

As a complement, the corresponding onset dates and key metrics as obtained from the RO data, [are shown in Table 4](#). Four
minor events (in W07-08 [W11-12](#) and [W16-17](#)) were not making the detection [criteria](#) for this dataset though they were
detected based on the ERA5 data. One of the two of W07-08, which overlaps with the BG18 data record, also was not detected
520 by [RO data](#).

In terms of count statistics, [43](#) events in the 42 winters [are detected](#), corresponding to a frequency of [1.02/year](#). [This is close](#)
to the frequency estimate of 0.9/year by McInturff (1978). [BG18 climatology, which only list major SSW events, yields a](#)
[frequency estimate of 0.6/year, which is close to a 0.50/year frequency we find for our major and extreme events \(i.e.,](#)
[discounting the minor events\)](#). In general, the high consistency of onset dates with the BG18 definition provides evidence that
525 our new temperature-anomalies-based method is robust in detecting SSW events and that the warming anomalies are strongly
related to wind reversals during SSWs. For those events detected by the BG18 definition but not by our method (e.g., the W06-
07 event in February 2007), we do find minor warmings [signalled](#) also in our TEA diagnostics, but they do not exceed
thresholds long enough (at least [six days](#)) for detection according to Table 2.

For the higher number of events that we detected, but which are not detected by the BG18 definition, [there are several reasons](#).
530 The main reason is that [a detection based on single altitudes and latitudes misses events occurring in other domains](#). Related

删除的内容: 4.4

带格式的: 标题 1

删除的内容: monitoring

删除的内容: Table 3 provides

删除的内容: , listing

删除的内容: intercomparing

删除的内容: work

删除的内容: latter

删除的内容: Table 4 shows

删除的内容: , which indicates that three

删除的内容: W19-20

删除的内容: threshold

删除的内容: the BG18 work

删除的内容: we detected 40

删除的内容: 0.95

删除的内容: Hence, compared to the frequency estimate of 0.6/year provided by the BG18 definition, the new approach detects about 35% more events,

删除的内容: signaled

删除的内容: one week

删除的内容: the

删除的内容: altitude

删除的内容: latitude miss

删除的内容: occurred

删除的内容: domain

to altitude, we found that a number of events occurred at levels higher than 10 hPa and therefore such events were not detected by the BG18 definition (e.g., in W92-93 and W00-01). In W07-08, we basically found four warmings (as can be seen in Fig. 3), of which three satisfied our detection criterion. This is in line with Singh et al. (2015), who also found four warmings based on a temperature increase definition. Related to latitude, Hu et al. (2015) used a wind reversal definition at 65 °N, detecting an event on 26 January 2010, which is close to our onset date of 29 January 2010 rather than the BG18 date of 9 February 2010. This indicates that selection of a specific latitude for detection also can hinder reliable detection. Finally, some temperature increase events may not be associated with wind reversals at 10hPa.

Furthermore, we detected a number of events in the 1990s, while BG18 detect only one in W98-99. This can be attributed to three reasons. Firstly, largest warmings occur at higher altitudes than the 10hPa level, and therefore may not be detected by data using such a single level. Secondly, a significant amount of radiosonde stations ceased operations in the 1990s and the availability of wind measurements to the BG18 study for SSW detection was degraded. Thirdly, several warming events are not highly related with wind reversal. Our method, based on temperature data records (here primarily from ERA5, verified by RO data) is robust against such observation network changes and, in addition, temperature is an easily available variable, including in form of multi-decadal records with reasonable long-term stability.

Figure 10 finally depicts summary statistics on the long-term results along various perspectives of interest, both for the full vertical resolution (basic case) and coarser resolution data (see the caption and the respective panel titles for explanation of the various panels). Regarding the upper half (Figs 10a-d), it is seen that the results from using the standard pressure level input data (“37-plevels ERA5”) are very similar to those from the basic case (“full-res ERA5”), with the coarser resolution case detecting only one event less and two events different. In addition, from simply using linear interpolation between the pressure level temperatures before computing layer-mean temperatures (cf. Fig. 3) from the interpolated coarser resolution profiles, the SSW strength (MPS) is found overall slightly smaller than for the basic case (see difference of adjacent color and gray symbols in Fig. 10a). The size of these differences relative to the basic case is typically smaller than 3%, however.

Furthermore, while the MPS appears widely variable as a time series (Fig. 10a), the inspection of its component metrics MPD and MPA reveals that the former (the duration) most clearly drives the strength (Fig. 10b, upper left subpanel) and that a salient trail-cooling phase, indicated by the TPD and TPA auxiliary metrics, is dominated by some of the strongest events (Fig. 10b, lower subpanels). Inspecting the SSW onsets, almost all of them (over 95%) are found within the deep winter (i.e., the Dec-Jan-Feb timeframe, with about half in January; Fig. 10c) and more than three-quarters are found to have their onset location over northern Eurasia and the adjacent polar ocean (Fig. 10d).

The crosscheck of using simple two-pressure-levels input data only (“10hPa & 50hPa level ERA5”), depicted as bottom half of Figure 10 (panels e-h), shows that the results are overall as well similar to the basic case but notably diluted in number of detected events (32 instead of 43) and providing systematically smaller MPS estimates, by about 20-30% (cf. panels a and e). This main reason is that the downward propagating warmings are captured at less strength at the 10hPa (~32 km) and 50hPa (~21 km) levels than by the layer-means taken over 30-35 km and 20-25 km, respectively. The respective minimum-threshold and MPS classification parameter adjustments for this simple two-levels approach (see Table 1 and 2 footnotes; TEA_{Min}

删除的内容: two

删除的内容: , which

删除的内容: our detection of

删除的内容: , is mainly because

删除的内容: also

删除的内容: 9

删除的内容:), in most cases (

删除的内容: 9a-9d) visualizing also the

删除的内容: class

删除的内容: the events according to the overarching MPS metric (by orange, red, dark-red

删除的内容:). While as a time series

删除的内容: 9a

删除的内容: 9b

删除的内容: 9b

删除的内容: Furthermore

删除的内容: SSW onsets (

删除的内容: 9c

删除的内容: 9d

610 reduced from 3 to 2 10^6 km^2 and classification boundaries lowered from 90/180 to 70/140 $10^6 \text{ km}^2 \text{ days}$) help to compensate for this “low bias” relative to the basic case. The overall detection frequency is nevertheless reduced to 0.76/year (Fig. 10g), while we find the frequency of the major and extreme events with 0.48/year closely similar to the one for the basic case (0.50/year). The relative under-detection of events is hence mainly attributable to minor events.

615 In summary we thus recommend to use temperature field data with adequately high vertical resolution as input to the method. At the same time the crosschecks demonstrate that the method is robust and reliable also for coarser resolution data, and as a minimum for 10hPa and 50hPa level data only, though somewhat less apt in its detection and monitoring capacity.

5.2 SSW long-term trends

Visually spotting a long-term “tendency” of possibly some strength intensification, in Figs. 10a, b and the numbers of Table 3, also the inspection of long-term trends in the SSW metrics during the climate change period from the 1980s to the 2010s is of 620 particular interest. We hence quantitatively inspected for multi-decadal trends in MPS, MPD, and MPA, based on the 33 decadal-mean values of these three metrics over 1984 to 2016 center years. For robustness crosscheck we performed this not only for the basic case with full-vertical-resolution input but also for the coarser resolution cases. In computing the decadal-mean value of a metric from all individual event values of a decade, events are allocated to a year based on their onset month, and the means obtained as 10-yr-aggregated values over center-yr – 4 yrs to center-yr + 5 yrs divided by 10 (we also cross- 625 checked to a simple averaging over the events in any 10-yr-window, which led to similar results but with somewhat more variability, due to depending in this case directly on the sometimes small event count per decade). For the trend fits we used ordinary-least-squares linear fitting and included uncertainty estimation for the trend rate accounting for reduced degrees-of-freedom due to autocorrelation and assuming small-sample t-distribution statistics (Santer et al., 2020; Loeb et al., 2022). Figure 11 shows that we indeed find a statistically significant positive trend in the MPS, with a best-estimate strength 630 intensification for the basic case (top row: “full-res ERA5”) by about 12(± 8) million $\text{km}^2 \text{ days}$ per decade (95% significance level; left subpanel), which is driven by a highly significant increase in the MPD by about 1.8(± 0.7) days/decade (99% significance level; middle subpanel). This implies an increase in the duration of SSW main-phase warmings by about 5(± 2) days from the 1980s to the 2010s, raising the average duration by near 50% from about 10 to 15 days and inducing an SSW strength increase by about 40(± 25) million $\text{km}^2 \text{ days}$ from about 100 to 140 million $\text{km}^2 \text{ days}$.

635 No significant trend was found in the MPA (right subpanel) as well as the associated threshold exceedance magnitude (the average warming strength of the main-phase temperature anomaly above threshold, as indicated by the Max ΔT threshold exceedance), for which sensitivity testing confirmed that it is well correlated with the MPA. As part of extensive sensitivity testing, we also found indication of an increasing trend in the number of major and extreme SSW events, by about 0.4 events per decade (near 90% significance level); an analysis for which the time series is still quite short, however, and which depends 640 for the minor event counts somewhat on the threshold definition for their detection.

The coarser-resolution-based results (middle and bottom row of Fig. 11: “37-levels ERA5” and “10hPa & 50hPa ERA5”) confirm that the trends found are robust across using different input data resolutions (i.e., consistent within the co-estimated

删除的内容: for perhaps

删除的内容: . 9a

删除的内容: we also

删除的内容: long-term

删除的内容: (Fig. 9e; events

删除的内容: ; decadal

删除的内容: are

删除的内容: yrs;

删除的内容: check

删除的内容: Using linear fitting (

删除的内容: trend fit, including

删除的内容: for

删除的内容: ;

删除的内容:),

删除的内容: found

删除的内容: by 9.3

删除的内容: Fig. 9e,

删除的内容: 3

删除的内容: Fig. 9e,

删除的内容: This implies an increase in the duration of SSW main-phase warmings by about 4 days from the 1980s to the 2010s, raising the average duration by 40 % from about 10 to 14 days and inducing an SSW strength increase by near 30 million $\text{km}^2 \text{ days}$ (about 30 %) from about 105 to 135 $\text{km}^2 \text{ days}$. No significant trend was found in the MPA (Fig. 9e,

删除的内容: (not shown). We also find

670 [uncertainties](#)). Though the mean estimates and associated uncertainty bounds quite vary across the three cases, reflecting the small-sample statistics with fairly low effective degrees-of-freedom (typically 8 to 12 only for the 33 decadal-mean values) of the rather short and volatile time series, the overall result of a clearly significant MPD increase that drives a corresponding MPS increase is robustly evident.

While a detailed interpretation and further study of the possible causes of this increase in warming duration, including cross-comparison with other temperature field datasets beyond ERA5, is beyond the scope of this study that focuses on introducing 675 the new monitoring method and related SSW climate data record, we speculate that it may be related to changes in the polar vortex dynamics over the recent decades that have led to transient change in prevalent vortex patterns, partly induced by anthropogenic climate change effects in the polar region (e.g., Kretschmer et al., 2018a;b). Since the MPS metric can be interpreted as an anomalous heat energy content contained in the exceedance warming of an SSW event (similar to the threshold exceedance metric of cooling degree days in the analysis of energy demand during hot days or even heat extremes; 680 Forster et al., 2021), the estimated increase by about [40%](#) since the 1980s corresponds to substantially more energy stored in and released by recent SSW events.

删除的内容: 30

6 Conclusions

删除的内容: 5

带格式的: 制表位: 18.59 字符, 左对齐

685 In this study we introduced and applied a new method for long-term monitoring of Sudden Stratospheric Warming (SSW) events based on metrics derived from daily stratospheric temperature anomaly threshold exceedance area data, refining upon the approach introduced in Li et al. (2021), which was based on the well-known 2009 SSW event only. We applied the new method over 1980 to 2021, including 14 winters using radio occultation (RO) data for verification (2006-2020), and all 42 winters within 1980-2021 from using ERA5 reanalysis data. Robust SSW characterization metrics including Main-Phase Duration (MPD), Area (MPA), and Strength (MPS) are derived, together with further auxiliary diagnostics.

Using these metrics, we proposed a new definition for SSW event detection and classification as well as explored multi-decadal 690 changes in their characteristics under the recent climate change. [An SSW is defined to be detected if main-phase warming duration lasts at least 6 days \(i.e., SSW-MPD > 6\).](#) According to MPS, SSWs are classified as minor, major, and extreme events. We also provide an informative SSW climatology over the four decades, recording valuable SSW event characterization information, including onset date, strength, duration, exceedance area, and type of event (minor, major, or extreme), complemented by the maximum-warming-anomaly geographic location and its associated maximum warming. In 695 addition, event trailing-phase metrics as well as day-by-day dynamic tracking of the warming-anomaly center location and associated maximum warming and of the areal extent of the exceedance area are available.

Detection results using RO and ERA5 are overall similar, suggesting that the new method can be applied using both RO and ERA5 data as well as any other quality-assured observational or reanalysis temperature (field) data covering the polar region and winter timeframes of interest. Comparison between our climatology with that from the recent BG18 climatology (Butler 700 and Gerber, 2018) reveals that a number of minor and major events was not part of the BG18 study, while several are part of

that one and not detected here. The coincidence of the onset dates of jointly detected events is commonly within ± 1 day, suggesting high detection consistency of the different methods and cross-verifying that our new method is robust.

删除的内容: 1day

705 In terms of event count statistics, we detected 43 events in the 42 winters, corresponding to an estimated event frequency of 1.02/year, close to the event frequency estimate of 0.9/year by McInturff (1978). Compared to the frequency estimate of 0.6/year provided by the BG18 study on major SSW events only, the new approach detects about 40% more events. Within the 1990s, where the BG18 study detected only two in W98-99, we detected seven events, (4 major and 3 minor ones). We also found that a salient upper stratospheric trailing-phase cooling occurs in the wake of the main warming phase for most, though 710 not all, of the strongest events. Regarding temporal and spatial occurrence, we found over 95% of the SSW onset dates in deep winter (i.e., Dec-Jan-Feb timeframe; about 50% in January) and more than three quarters of the associated onset locations over Northern Eurasia and the adjacent polar ocean.

删除的内容: 40

删除的内容: 0.95

删除的内容: . Hence, compared

删除的内容: 35

删除的内容: close to the event frequency estimate of 0.9/year by McInturff (1978).

删除的内容: a number of

删除的内容: .

删除的内容: about

删除的内容: using linear-trend fitting and statistical evaluation,

删除的内容: 9.3

删除的内容: 3

删除的内容: 4

删除的内容: 40

删除的内容: 14

删除的内容: near 30

带格式的: 字体: 八号

删除的内容: (about 30 %)

删除的内容: 105

删除的内容: 135

带格式的: 字体: 八号

删除的内容: 30

删除的内容: long-term

Regarding long-term changes, we found a statistically significant positive trend in the MPS metric, with a best-estimate strength intensification by about 12(±8) million km² days per decade, which is driven by a highly significant increase in the 715 MPD by about 1.8(±0.7) days/decade. This implies an increase in the duration of SSW main-phase warmings by about 5(±2) days from the 1980s to the 2010s, raising the average duration by near 50% from about 10 to 15 days and inducing an SSW strength increase by about 40(±25) million km² days from about 100 to 140 million km² days. These results are found robust (i.e., consistent within co-estimated uncertainties) across using different vertical resolutions of ERA5 temperature input data.

720 Since the MPS metric can be interpreted as an anomalous heat energy content contained in the exceedance warming of an event, such an increase by about 40% corresponds to substantially more energy stored in and released by recent SSW events. No significant trend was found in the MPA as well as the associated threshold exceedance warming magnitude.

It is hoped that the results of this study can be used as a reference for further complementary and climate change-related studies and, in particular, also be a basis for SSW impact studies related to other weather and climate phenomena linked to SSWs, such as changes in polar vortex dynamics and implications to mid-latitude extreme weather, among others. Follow-on work 725 will further investigate the SSW's long-term evolution over the recent decades and the causes of the evidenced trends. We also intend to investigate whether and how SSW events occurring in different regions have impacts on near-surface weather over middle latitudinal regions of the northern hemisphere.

730 **Code availability.** The code used to produce the results of this study is available from the first author (Y. L.) upon qualified request.

Data availability. The (numeric) data underlying the results of this study are available from the first author (Y. L.) upon qualified request. The ERA5 reanalysis data are available at full resolution via ECWMF's Mars archive (registered member state users) and publicly through the Copernicus Climate Change Services (C3S) via <https://cds.climate.copernicus.eu/>. The 735 OPSv5.6 RO data of WEGC are publicly available via <https://doi.org/10.25364/WEGC/OPSV5.6:2020.1>.

760 **Author contributions.** Ying Li implemented and refined the new method, performed the analysis, produced the figures, and wrote the initial draft of the manuscript. Gottfried Kirchengast conceived the method and served as primary coauthor, providing advice and guidance on all aspects of the design, analysis, and figure production, and significantly contributed to writing the manuscript. Marc Schwärz supported the setup and advancements of the OPS analysis system and provided input data as well as algorithms. Yunbin Yuan advised on analysis and algorithm comparisons. All authors commented on and agreed to the final manuscript.

765

刪除的內容: submitted

Competing interests. The authors declare that they have no conflict of interest.

770 **Acknowledgements.** We acknowledge ECMWF (Reading, UK) for providing access to their analysis and forecast data and the RO team at WEGC (Graz, Austria) for the preparation of the OPSv5.6 RO data; we specifically acknowledge Florian Ladstätter for supporting RO climatology provision and advising on its characteristics as well as on results interpretation. We furthermore thank the reviewers for their valuable comments that clearly helped to improve the final paper. The research at the APM (Wuhan, China) was funded by the Strategic Priority Research Program of Chinese Academy of Sciences (Grant No. XDA17010304) and the Chinese Natural Sciences Foundation (grant no. 41874040). At WEGC, the work was supported by the Aeronautics and Space Agency of the Austrian Research Promotion Agency (FFG-ALR) under the Austrian Space 775 Applications Programme (ASAP) projects ATROMSAF1 and ATROMSAF2 (proj.no. 859771 and 873696) funded by the Ministry for Climate, Environment, Energy, Mobility, Innovation, and Technology (BMK).

References

Angerer, B., Ladstätter, F., Scherllin-Pirscher, B., Schwärz, M., and Kirchengast, G.: Quality aspects of the Wegener Center multi-satellite GPS radio occultation record OPSv5.6, *Atmos. Meas. Tech.*, 10(12), 4845–4863, 780 <https://doi.org/10.5194/amt-10-4845-2017>, 2017.

Anthes, R. A.: Exploring Earth's atmosphere with radio occultation: contributions to weather, climate and space weather, *Atmos. Meas. Tech.*, 4, 1077–1103, <https://doi.org/10.5194/amt-4-1077-2011>, 2011.

Ayarzagüena, B., Langematz, U., Meul, S., Oberländer, S., Abalichin, J., and Kubin, A.: The role of climate change and ozone recovery for the future timing of major stratospheric warmings, *Geophys. Res. Lett.*, 40, 2460–2465, 785 <https://doi.org/10.1002/grl.50477>, 2013.

Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., Domeisen, D.I.V., Garfinkel, C. I., Garny, H., Gerber, E. P., Hegglin, M. I., Langematz, U., and Pedatella, M.: Sudden stratospheric warmings. *Reviews of Geophysics*, 59, <https://doi.org/10.1029/2020RG000708>, 2020.

Baldwin, M. P. and Thompson, D. W. J.: A critical comparison of stratosphere–troposphere coupling indices, *Quart. J. Roy. 790 Meteor. Soc.*, 135, 1661–1672, <https://doi.org/10.1002/qj.479>, 2009.

Biondi, R., Steiner, A. K., Kirchengast, G., and Rieckh, T.: Characterization of thermal structure and conditions for overshooting of tropical and extratropical cyclones with GPS radio occultation, *Atmos. Chem. Phys.*, 15, 5181–5193, <https://doi.org/10.5194/acp-15-5181-2015>, 2015.

795 Bosilovich M.G., Kennedy J., Dee D., Allan R., and O'Neill A.: On the Reprocessing and Reanalysis of Observations for Climate in Climate Science for Serving Society, edited by: Asrar, G. and Hurrell, J., Springer, Dordrecht, https://doi.org/10.1007/978-94-007-6692-1_3, 2013.

Brunner, L., Steiner, A.K., Scherlin-Pirscher, B., and Jury, M.W.: Exploring atmospheric blocking with GPS radio occultation observations, *Atmos. Chem. Phys.*, 16, 4593–4604, <https://doi.org/10.5194/acp-16-4593-2016>, 2016.

800 Butler, A. H., Sjoberg, J. P., Seidel, D. J., and Rosenlof, K. H.: A sudden stratospheric warming compendium, *Earth Syst. Sci. Data*, 9, 63–76, <https://doi.org/10.5194/essd-9-63-2017>, 2017.

Butler, A. H. and Gerber, E. P.: Optimizing the definition of a sudden stratospheric warming, *J. Climate.*, 31, 2337–2344, <https://doi.org/10.1175/JCLI-D-17-0648.1>, 2018.

805 Butler, A. H., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining sudden stratospheric warmings, *B. Am. Meteor. Soc.*, 96, 1913–1928, <https://doi.org/10.1175/BAMS-D-13-00173.1>, 2015.

Cattiaux, J. R., Vautard, R., Cassou, C., Yiou, P., and Codron, F.: Winter 2010 in Europe: a cold extreme in a warming climate, *Geophys. Res. Lett.*, 37, 114-122, <https://doi.org/10.1029/2010GL044613>, 2010.

Charlton, A. J. and Polvani, L. M.: A new look at stratospheric sudden warmings: Part I: Climatology and modeling benchmarks, *J. Climate.*, 20, 449–469, <https://doi.org/10.1175/JCLI3996.1>, 2007.

810 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksena, L., Källberg, P., Kállner, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut J.-N., and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q.J.R. Meteorol. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.

Dhaka, S. K., Kumar, V., Choudhary, R. K., Ho, S.P., Takahashi, M., and Yoden, S.: Indications of a strong dynamical coupling between the polar and tropical regions during the sudden stratospheric warming event January 2009, based on COSMIC/FORMASAT-3 satellite temperature data, *Atmos. Res.*, 166, 60–69, <https://doi.org/10.1016/j.atmosres.2015.06.008>, 2015.

820 Foelsche, U., Scherlin-Pirscher, B., Ladstädter, F., Steiner, A.K., and G. Kirchengast.: Refractivity and temperature climate records from multiple radio occultation satellites consistent within 0.05%, *Atmos. Meas. Tech.*, 4, 2007–2018, <https://doi.org/10.5194/amt-4-2007-2011>, 2011.

Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D.J., Mauritsen, T., Palmer, M.D., Watanabe, M., Wild, M., and Zhang, H.: The Earth's energy budget, climate feedbacks, and climate sensitivity (Chapter 7). In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment*

Report of the Intergovernmental Panel on Climate Change, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Pean, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, C., Waterfield, T., Yelekci, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 923–1054, <https://doi.org/10.1017/9781009157896.009>, 2021.

830 Gerber, E. P., Baldwin, M. P., Akiyoshi, H., Austin, J., and Dan, S.: Stratosphere-troposphere coupling and annular mode variability in chemistry-climate models, *J. Geophys. Res.*, 115, D00M06, <https://doi.org/10.1029/2009JD013770>, 2010.

Gobiet, A., Kirchengast, G., Manney, G. L., Borsche, M., Retscher, C., and Stiller, G.: Retrieval of temperature profiles from CHAMP for climate monitoring: intercomparison with Envisat MIPAS and GOMOS and different atmospheric analyses, *Atmos. Chem. Phys.*, 7, 3519–3536, <https://doi.org/10.5194/acp-7-3519-2007>, 2007.

835 Hall, R. J., Mitchell, D. M., Seviour, W. J. M., and Wright, C. J.: Tracking the stratosphere-to-surface impact of Sudden Stratospheric Warmings, *J. Geophys. Res. Atmos.*, 126, <https://doi.org/10.1029/2020JD033881>, 2021.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., 840 Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

Hersbach, H., Bell, W., Berrisford, P., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Radu, R., Schepers, D., Simmons, A., Soci, C., and Dee, D.: Global reanalysis: goodbye ERA-Interim, hello ERA5, *ECMWF Newslett.*, 159, 17–24, 845 <https://doi.org/10.21957/vf291hehd7>, 2019.

Hitchcock, P. and Simpson, I. R.: The downward influence of stratospheric sudden warmings, *J. Atmos. Sci.*, 71, 3586–3876, <https://doi.org/10.1175/JAS-D-14-0012.1>, 2014.

Hitchcock, P. and Shepherd, T. G.: Zonal-mean dynamics of extended recoveries from stratospheric sudden warmings, *J. Atmos. Sci.*, 70, 688–707, <https://doi.org/10.1175/JAS-D-12-0111.1>, 2013.

850 Holt, L. A., Randall, C. E., Peck, E. D., Marsh, D. R., Smith, A. K., and Harvey, V. L.: The influence of major sudden stratospheric warming and elevated stratopause events on the effects of energetic particle precipitation in WACCM, *J. Geophys. Res. Atmos.*, 118, 11,636–11,646, <https://doi.org/10.1002/2013JD020294>, 2013.

Hu, J., Ren, R., and Xu, H.: Occurrence of winter stratospheric sudden warming events and the seasonal timing of spring stratospheric final warming, *J. Atmos. Sci.*, 71(7), <https://doi.org/10.1175/JAS-D-13-0349.1> 2319–2334, 2015.

855 Johnson, K. W., Miller, A. J., and M. Gelman.: Proposed indices characterizing stratospheric circulation and temperature fields, *Mon. Wea. Rev.*, 97, 565–570, [https://doi.org/10.1175/1520-0493\(1969\)097<0565:PICSCA>2.3.CO;2](https://doi.org/10.1175/1520-0493(1969)097<0565:PICSCA>2.3.CO;2), 1969.

Jonah, O. F., de Paula, E. R., Kherani, E. A., Dutra, S. L. G., and Paes, R. R.: Atmospheric and ionospheric response to sudden stratospheric warming of January 2013, *J. Geophys. Res. Space Physics*, 119, 4973–4980, <https://doi.org/10.1002/2013JA019491>, 2014.

860 Kakoti, G., Kalita, B. R., Bhuyan, P. K., Baruah, S., and Wang, K.: Longitudinal and interhemispheric ionospheric response to 2009 and 2013 SSW events in the African-European and Indian-East Asian sectors, *J. Geophys. Res. Space Physics.*, 125, <https://doi.org/10.1029/2020JA028570>, 2020

865 Kirchengast, G.: Occultations for probing atmosphere and climate: Setting the scene, in: *Occultations for Probing Atmosphere and Climate*, edited by: Kirchengast, G., Foelsche, U., and Steiner, A.K., 1-8, Springer, Berlin-Heidelberg, https://doi.org/10.1007/978-3-662-09041-1_1, 2004.

870 Klingler, R.: Observing Sudden Stratospheric Warmings with Radio Occultation Data, with Focus on the Event 2009, MSc Thesis, 85 pp., University of Graz, Graz, Austria, URN <https://resolver.obvsg.at/urn:nbn:at:at-ubg:1-68069>, 2014.

Kretschmer, M., Coumou, D., Agel, L., Barlow, M., Tziperman, E., and Cohen, J.: More-persistent weak stratospheric polar vortex states linked to cold extremes, *Bull. Amer. Meteorol. Soc.*, 99, 49–60, <https://doi.org/10.1175/BAMS-D-16-0259.1>, 2018a.

875 Kretschmer, M., Cohen, J., Matthias, V., Runge, J., and Coumou, D.: The different stratospheric influence on cold-extremes in Eurasia and North America, *Clim. Atmos. Sci.*, 1, 44, <https://doi.org/10.1038/s41612-018-0054-4>, 2018b.

Kursinski, E. R., Hajj, G. A., Schofield, J. T., Linfield, R. P., and Hardy, K. R.: Observing Earth's atmosphere with radio occultation measurements using the Global Positioning System, *J. Geophys. Res.*, 102, 23429–23465, <https://doi.org/10.1029/97JD01569>, 1997.

Kuttippurath, J. and Nikulin, G.: A comparative study of the major sudden stratospheric warmings in the Arctic winters 2003/2004–2009/2010, *Atmos. Chem. Phys.*, 12, 8115–8129, <https://doi.org/10.5194/acp-12-8115-2012>, 2012.

880 Labitzke, K.: Stratospheric-mesospheric midwinter disturbances: A summary of observed characteristics, *J. Geophys. Res.*, 86, 9665–9678, <https://doi.org/10.1029/JC086iC10p09665>, 1981.

Labitzke, K. and Kunze, M.: On the remarkable Arctic winter in 2008/2009, *J. Geophys. Res. Atmos.*, 114, D00I02, <https://doi.org/10.1029/2009JD012273>, 2009.

885 Ladst  ter, F., Steiner, A. K., Schw  z, M., and Kirchengast, G.: Climate intercomparison of GPS radio occultation, RS90/92 radiosondes and GRUAN from 2002 to 2013, *Atmos. Meas. Tech.*, 8, 4, 1819–1834, <https://doi.org/10.5194/amt-8-1819-2015>, 2015.

Lehtonen, I. and Karpechko, A. Yu.: Observed and modeled tropospheric cold anomalies associated with sudden stratospheric warmings, *J. Geophys. Res. Atmos.*, 121, 1591–1610, <https://doi.org/10.1002/2015JD023860>, 2016.

Lin, J. T., Lin, C.H., Chang, L.C., Huang, H.H., Liu, J.Y., Chen, A. B., Chen, C.H., and Liu, C.H.: Observational evidence of ionospheric migrating tide modification during the 2009 stratospheric sudden warming, *Geophys. Res. Lett.*, 39, L02101, <https://doi.org/10.1029/2011GL050248>, 2012.

890 Li, Y., Kirchengast, G., Schwaerz, M., Ladst  ter, F., and Yuan, Y.-B.: Monitoring Sudden Stratospheric Warmings using radio occultation: a new approach demonstrated based on the 2009 event, *Atmos. Meas. Tech.*, 14, 2327–2343, <https://doi.org/10.5194/amt-14-2327-2021>, 2021.

Loeb, N.G., Mayer, M., Kato, S., Fasullo, J.T., Zuo, H., Senan, R., Lyman, J.M., Johnson, G.C., and Balmaseda, M.: Evaluating Twenty-Year Trends in Earth's Energy Flows From observations, *Earth. Space. Sci.*,
895 https://doi.org/10.1002/essoar.10510650.1, 2022.

Luntama, J.-P., Kirchengast, G., Borsche, M., Foelsche, U., Steiner, A., Healy, S., von Engeln, A., O'Clerigh, E., and Marquardt, C.: Prospects of the EPS GRAS mission for operational atmospheric applications, *B. Am. Meteorol. Soc.*, 89, 1863–1875, https://doi.org/10.1175/2008BAMS2399.1, 2008.

Manney, G. L., Lawrence, Z. D., Santee, M. L., Read, W. G., Livesey, N. J., Lambert, A., Froidevaux, L., Pumphrey, H. C.,
900 and Schwartz M. J.: A minor sudden stratospheric warming with a major impact: Transport and polar processing in the 2014/2015 Arctic winter, *Geophys. Res. Lett.*, 42, 7808–7816, https://doi.org/10.1002/2015GL065864, 2015.

McInturff, R. M.: Stratospheric warmings: Synoptic, dynamic and general-circulation aspects, *NASA Reference Publ.*, USA, NASA-RP-1017, 174, 1978.

Mitchell, D. M., Gray, L. J., Anstey, J., Baldwin, M. P., and Charlton-Perez A. J.: The influence of stratospheric vortex
905 displacements and splits on surface climate, *J. Climate*, 26, 2668–2682, https://doi.10.1175/JCLI-D-12-00030.1, 2013.

Nayak, C. and Yigit, E.: Variation of small-scale gravity wave activity in the ionosphere during the major sudden stratospheric warming event of 2009, *J. Geophys. Res. Space Physics.*, 124, 470–488, https://doi.org/10.1029/2018JA026048, 2019.

Noguchi, S., Kuroda, Y., Mukougawa, H., Mizuta, R., and Kobayashi, C.: Impact of satellite observations on forecasting sudden stratospheric warmings, *Geophys. Res. Lett.*, 47, https://doi.org/10.1029/2019GL086233, 2020.

910 Parker, W. S.: Reanalyses and observations: What's the difference, *Bull. Amer. Meteor. Soc.*, 97, 1565–1572, https://doi.org/10.1175/BAMS-D-14-00226.1, 2016.

Santer, B. D., Wigley, T., Boyle, J. S., Gaffen, D. J., Hnilo, J. J., and Nychka, D.: Statistical significance of trends and trend differences in layer-average atmospheric temperature time series, *J. Geophys. Res.*, 105, 7337–7356, https://doi.org/10.1029/1999jd901105, 2000.

915 Scherhag, R.: Die explosionsartige Stratosphärenerwärmung des Spätwinters 1951/52, *Ber. Deut. Wetterdienst*, 38, 51–63, 1952.

Scherllin-Pirscher, B., Kirchengast, G., Steiner, A. K., Kuo, Y.-H., and Foelsche, U.: Quantifying uncertainty in climatological fields from GPS radio occultation: an empirical-analytical error model, *Atmos. Meas. Tech.*, 4, 2019–2034, https://doi.org/10.5194/amt-4-2019-2011, 2011a.

920 Scherllin-Pirscher, B., Steiner, A. K., Kirchengast, G., Kuo, Y.-H., and Foelsche, U.: Empirical analysis and modelling of errors of atmospheric profiles from GPS radio occultation, *Atmos. Meas. Tech.*, 4, 1875–1890, doi:10.5194/amt-4-1875-2011, 2011b.

Scherllin-Pirscher, B., Steiner, A. K., Kirchengast, G., Schwaerz, M., and Leroy, S. S.: The power of vertical geolocation of atmospheric profiles from GNSS radio occultation, *J. Geophys. Res.-Atmos.*, 122, 1595–1616, https://doi.org/10.1002/2016JD025902, 2017.

Schoeberl, M. R.: Stratospheric warmings: Observations and theory, *Rev. Geophys.*, 16, 521–538, <https://doi.org/10.1029/RG016i004p00521>, 1978.

Schreiner, W., Rocken, C., Sokolovskiy, S., Syndergaard, S., and Hunt, D.: Estimates of the precision of GPS radio occultations from the COSMIC/FORMOSAT-3 mission, *Geophys. Res. Lett.*, 34, L04808, <https://doi.org/10.1029/2006GL027557>, 930 2007.

Schwarz, M., Kirchengast, G., Scherlin-Pirscher, B., Schwarz, J., Ladstädter, F., and Angerer, B.: Multi-mission validation by satellite radio occultation extension project—Final report, Tech. Rep. for ESA/ESRIN No. 01/2016, Wegener Center, University of Graz, Graz, Austria, available at: https://wegcwww.uni-graz.at/publ/wegcpubl/arsclisys/2016/Schwarz-etal_MMValRO-FinRep_Dec2016.pdf (last access: 12 Dec 2021), 2016

935 Seviour, W. J. M., Mitchell, D. M., and Gray, L.J.: A practical method to identify displaced and split stratospheric polar vortex events, *Geophys. Res. Lett.*, 40, 5268–5273, <https://doi.org/10.1002/grl.50927>, 2013.

Simmons, A., Soci, C., Nicolas, J., Bell, B., Berrisford, P., Dragani, R., Flemming, J., Haimberger1, L., Healy, S., Hersbach, H., Horányi, A., Inness, A., Muñoz-Sabater, J., Radu, R., and Schepers, D.: Global stratospheric temperature bias and other stratospheric aspects of ERA5 and ERA5.1, ECMWF Tech., Memo., No. 859, <https://doi.org/10.21957/rxqfmg0>, 940 2020.

Singh, R. P. and Pallamraju, D.: On the latitudinal distribution of mesospheric temperatures during sudden stratospheric warming events, *J. Geophys. Res. Space Physics.*, 120, 2926–2939, <https://doi.org/10.1002/2014JA020355>, 2015.

Siskind, D. E., Eckermann, S. D., McCormack, J. P., Coy, L., Hoppel, K. W., and Baker, N. L.: Case studies of the mesospheric response to recent minor, major, and extended stratospheric warmings, *J. Geophys. Res.*, 115, D00N03, 945 <https://doi.org/10.1029/2010JD014114>, 2010.

Steiner, A.K., Lackner, B.C., Ladstädter, F., Scherlin-Pirscher, B., Foelsche, U., and Kirchengast, G.: GPS radio occultation for climate monitoring and change detection, *Radio Sci.*, 46, RS0D24, <https://doi.org/10.1029/2010RS004614>, 2011.

950 Steiner, A. K., Ladstädter, F., Ao, C. O., Gleisner, H., Ho, S.-P., Hunt, D., Schmidt, T., Foelsche, U., Kirchengast, G., Kuo, Y.-H., Lauritsen, K. B., Mannucci, A. J., Nielsen, J. K., Schreiner, W., Schwarz, M., Sokolovskiy, S., Syndergaard, S., and Wickert, J.: Consistency and structural uncertainty of multi-mission GPS radio occultation records, *Atmos. Meas. Tech.*, 13, <https://doi.org/10.5194/amt-13-2547-2020>, 2020a.

Steiner, A. K., Ladstädter, F., Randel, W. J., Maycock, A. C., Fu, Q., Claud, C., Gleisner, H., Haimberger, L., Ho, S.-P., Keckhut, P., Leblanc, T., Mears, C., Polvani, L. M., Santer, B. D., Schmidt, T., Sofieva, V., Wing, R., and Zou, C.-Z.: Observed temperature changes in the troposphere and stratosphere from 1979 to 2018, *J. Climate.*, 33, 8165–8194, 955 <https://doi.org/202010.1175/JCLI-D-19-0998.1>, 2020b.

Stocker, M., Ladstädter, F., and Steiner, A. K.: Observing the climate impact of large wildfires on stratospheric temperature *Sci. Rep.*, 11, 22994, <https://doi.org/10.1038/s41598-021-02335-7>, 2021.

Sun, Y., Bai, W., Liu, C., Liu, Y., Du, Q., Wang, X., Yang, G., Liao, M., Yang, Z., Zhang, X., Meng, X., Zhao, D., Xia, J.,
Cai, Y., and Kirchengast, G.: The FengYun-3C radio occultation sounder GNOS: a review of the mission and its early
960 results and science applications, *Atmos. Meas. Tech.*, 11, 5797–5811, <https://doi.org/10.5194/amt-11-5797-2018>, 2018.

Thompson, D. W. J., Baldwin, M. P., and Wallace, J. M.: Stratospheric connection to northern hemisphere wintertime weather:
implications for prediction, *J. Climate*, 15(12), 1421–1428, [https://doi.org/10.1175/1520-0442\(2002\)015<1421:SCTNHW>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1421:SCTNHW>2.0.CO;2), 2002.

Tyrlis, E., Manzini, E., Bader, J., Ukita, J., Nakamura, H., and Matei, D.: Ural blocking driving extreme Arctic sea ice loss,
965 cold Eurasia, and stratospheric vortex weakening in autumn and early winter 2016–2017, *J. Geophys. Res.-Atmos.*, 124,
<https://doi.org/10.1029/2019JD031085>, 2019.

Vignon E. and Mitchell D.M.: The stratopause evolution during different types of sudden stratospheric warming event, *Clim
Dyn*, 44, 3323–3337, <https://doi.org/10.1007/s00382-014-2292-4>, 2015.

Van Loon, H., Jenne, R. L., and Labitzke, K.: Zonal harmonic standing waves. *J. Geophys. Res.*, 78, 4463–4471,
970 <https://doi.org/10.1029/JC078i021p04463>, 1973.

Wickert, J., Reigber, C., Beyerle, G., König, R., Marquardt, C., Schmidt, T., Grundwaldt, L., Galas, R., Meehan, T. K.,
Melbourne, W. G., and Hocke, K.: Atmosphere sounding by GPS radio occultation: First results from CHAMP, *Geophys.
Res. Lett.*, 28, 32633266, <https://doi.org/10.1029/2001GL013117>, 2001.

Wickert, J., Beyerle, G., König, R., Heise, S., Grunwaldt, L., Michalak, G., Reigber, Ch., and Schmidt, T.: GPS radio
975 occultation with CHAMP and GRACE: A first look at a new and promising satellite configuration for global atmospheric
sounding, *Ann. Geophys.*, 23, 653–658, <https://doi.org/10.5194/angeo-23-653-2005>, 2005.

WMO/IQSY.: International Years of the Quiet Sun (IQSY), 1964–1965: Alert messages with special references to stratwarms,
Secretariat of the WMO WMO/IQSY Rep. 6, 19, 1964.

Yoshida, K. and Yamazaki, K.: Tropical cooling in the case of stratospheric sudden warming in January 2009: focus on the
980 tropical tropopause layer, *Atmos. Chem. Phys.*, 11, 6325–6336, <https://doi.org/10.5194/acp-11-6325-2011>, 2011.

Yu, Y., Ren, R., and Cai, M.: Dynamic linkage between cold air outbreaks and intensity variations of the meridional mass
circulation, *J. Atmos. Sci.*, 72(8), 3214–3232, <https://doi.org/10.1175/JAS-D-14-0390.1>, 2015.

Yue, X., Schreiner, W.S., Lei, J., Rocken, C., Hunt, D.C., Kuo, Y.-H., and Wan, W.: Global ionospheric response observed by
COSMIC satellites during the January 2009 stratospheric sudden warming event, *J. Geophys. Res.*, 115, A00G09,
985 <https://doi.org/10.1029/2010JA015466>, 2010.

Zhou, S., Miller, A.J., Wang, J., and Angell, J.K.: Downward-propagating temperature anomalies in the preconditioned polar
stratosphere, *J. Climate*, 15(7), 781–792, [https://doi.org/10.1175/1520-0442\(2002\)015<0781:DPTAIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0781:DPTAIT>2.0.CO;2), 2002.

Table 1. Basic parameters and methodology of the new SSW monitoring approach (all parameters are updated daily; the boldfaced font in (2)-(4) and (7)-(10) marks key parameters for the monitoring as also shown in Figs. 6 to 8).

Parameter	Equation/Definition	Explanation/Description
(1) Temperature anomaly profile $\Delta T_{\text{Anomaly}}$	$\Delta T_{\text{Anomaly}} = T - T_{\text{Cli}} [\text{K}]$	T represents an individual RO or ERA5 profile, T_{Cli} is the collocated climatological profile. ^{a)}
(2) Middle Stratosphere Temperature Anomaly Threshold Exceedance Area: MSTA-TEA [10^6 km^2]	Altitude range: 30–35 km Thresholds <u>computed</u> : +50 K, +40 K, +30 K ; -30 K, -40 K, -50 K	Based on individual $\Delta T_{\text{Anomaly}}$ profiles in selected stratospheric altitude layers (e.g., 30–35 km for MSTA-TEA) first estimate layer-mean anomaly values from these profiles. The individual layer-mean values are then averaged into a suitable space-time-binned grid over 50–90°N (5°latitude \times 20°longitude grid). ^{b)}
(3) Lower Stratosphere Temperature Anomaly Threshold Exceedance Area: LSTA-TEA [10^6 km^2]	Altitude range: 20–25 km Thresholds <u>computed</u> : +30 K, +25 K, +20 K ; -20 K, -25 K, -30 K	The geographic areas wherein grid-cell anomalies exceed predefined thresholds (<u>see definition; the ones further used for (7)–(10) are bold-faced</u>) are finally calculated and stored as the Threshold Exceedance Area (TEAs).
(4) Upper Stratosphere Temperature Anomaly Threshold Exceedance Area: USTA-TEA [10^6 km^2]	Altitude range: 40–45 km Thresholds <u>computed</u> : +50 K, +40 K, +30 K; -30 K , -40 K, -50 K	Maximum (positive/negative) anomaly value of all grid cells within a TEA obtained by (2)–(4)
(5) Anomaly Maximum value	$\Delta T_{\text{AMax}} [\text{K}]$	Generate a contour that is 2 K smaller/larger than the positive/negative ΔT_{AMax} value; the center of the contour is then used as geographic location of the ΔT_{AMax} value.
(6) Geographic location (Lat, Lon) of Anomaly Maximum value	$\varphi^{\text{AMax}} [\text{N}], \lambda^{\text{AMax}} [\text{E}]$	Expresses the main and primary stratospheric warming anomaly strength; recorded if $\text{SSW-PP-TEA} > \text{TEA}_{\text{Min}} (3 \times 10^6 \text{ km}^2)$ for ≥ 3 days. ^{b)}
(7) SSW Primary-Phase Threshold Exceedance Area: SSW-PP-TEA [10^6 km^2]	$\text{SSW-PP-TEA} = (\text{MSTA-TEA} > +30 \text{ K})$	Expresses the secondary downward propagated warming anomaly strength; recorded if a SSW-PP-TEA is recorded (see (7)) and if during its presence the SSW-SP-TEA emerges and then exceeds $\text{TEA}_{\text{Min}} (3 \times 10^6 \text{ km}^2)$ for ≥ 5 days. ^{b)}
(8) SSW Secondary-Phase Threshold Exceedance Area: SSW-SP-TEA [10^6 km^2]	$\text{SSW-SP-TEA} = (\text{LSTA-TEA} > +20 \text{ K})$	Expresses the combined warming of primary- and secondary phase; it takes the higher value of SSW-PP-TEA and SSW-SP-TEA at any day.
(9) SSW Main-Phase Threshold Exceedance Area: SSW-MP-TEA [10^6 km^2]	$\text{SSW-MP-TEA} = \text{Max}(\text{SSW-PP-TEA}, \text{SSW-SP-TEA})$	Expresses the trailing upper stratospheric cooling anomaly strength; recorded if $\text{SSW-TP-TEA} > \text{TEA}_{\text{Min}} (3 \times 10^6 \text{ km}^2)$ for ≥ 21 days. ^{c)}
(10) SSW Trailing-Phase Threshold Exceedance Area: SSW-TP-TEA [10^6 km^2]	$\text{SSW-TP-TEA} = \text{Abs}(\text{USTA-TEA} < -30 \text{ K})$	^{a)} regarding vertical resolution of the (re)analysis, observational, or model simulation input datasets from which the anomaly profiles and layer-means are extracted, at least a resolution comparable to the 37 standard pressure level grid of the ERA5 (and other) data is recommended (with higher vertical resolution preferable, if available). This includes ten stratospheric level temperatures from 70 hPa (near 19 km) to 1 hPa (near 48 km), which through simple linear vertical interpolation of temperatures provide adequate profiles for extracting the desired SSW information according to the subsequent steps (2)–(10). Alternatively, as a minimum-input-data approach, temperature map data solely at 10 hPa (~32 km) and 50 hPa (~21 km) pressure levels (or similar altitudes) may be used as input, which substitute for the layer-mean values of steps (2) and (3). Step (4) is dropped in this simple two-levels approach, which restricts to middle and lower stratospheric TEA data only.

^{b)} in the simple two-levels approach (see note a), for which the 10 hPa and 50 hPa pressure levels (or corresponding altitudes) are located in the low part of the 5-km-layers of the profiles-based approach, TEA_{Min} is set to a reduced value of $2 \times 10^6 \text{ km}^2$ for primary and secondary phase TEA.

^{c)} in addition, in the simple two-levels approach, the MSTA-TEA replaces the USTA-TEA in step (10) and the cooling-anomaly threshold for computing the SSW-TP-TEA is accordingly set to a reduced value of $\text{Abs}(\text{MSTA-TEA} < -20 \text{ K})$.

删除的内容: selected

带格式的: 字体: 加粗

删除的内容: selected

带格式的: 字体: 加粗

删除的内容:

删除的内容: e.g., +30 K

删除的内容: selected

带格式的: 字体: 加粗

删除的内容:

Table 2. Metrics of the new SSW monitoring approach for detection, classification, and further qualification (the boldfaced font in (1)-(5) marks key metrics and criteria for the detection and classification also shown as main ones in Fig. 9).

Parameter	Equation/Definition	Explanation/Description
(1) Main-phase duration metric:	SSW-MPD [days]	Expresses SSW warming duration: number of days with SSW-MP-TEA available (at $> \text{TEA}_{\text{Min}}$; cf. Table 1)
(2) Main-phase area metric:	SSW-MPA [10^6 km^2]	Expresses SSW mean warming area: average daily area of SSW-MP-TEA during all SSW-MPD days
(3) Main-phase strength metric:	SSW-MPS [$10^6 \text{ km}^2 \text{ days}$] = (SSW-MPA \times SSW-MPD)	Express SSW warming strength: the larger this area-duration product, the stronger the event
(4) SSW detection criterion:	SSW-MPD \geq <u>6</u> days	SSW event adopted as detected and logged to the event count
	<u>SSW-MPS < 90</u> <u>from</u>	Minor SSW event
(5) SSW classification criteria: ^{a)}	<u>SSW-MPS ≥ 90</u> $10^6 \text{ km}^2 \text{ days}$ and ≤ 180 $10^6 \text{ km}^2 \text{ days}$	Major SSW event
	<u>SSW-MPS > 180</u> $10^6 \text{ km}^2 \text{ days}$	Extreme SSW event
(6) Trailing-phase duration metric:	SSW-TPD [days]	Expresses the trail-cooling duration: number of days with SSW-TP-TEA available (at $> \text{TEA}_{\text{Min}}$; cf. Table 1)
(7) Trailing-phase area metric:	SSW-TPA [10^6 km^2]	Express the mean trail-cooling area: Average daily area of SSW-TP-TEA during all SSW-TPD days
(8) Further SSW qualification:	<u>SSW-TPD < 21</u> days <u>SSW-TPD ≥ 21</u> days	Non-trail-cooling event Trail-cooling event (TC)
(9) SSW onset calendar date	Onset date [yyyy-mm-dd]	The day when the primary-phase exceedance area SSW-PP-TEA is largest (based on (7) in Table 1)
(10) SSW onset geographic location (latitude / longitude)	Onset location [°N / °E]	Location where ΔT_{Amax} occurs at the onset date (based on (6) in Table 1)
(11) SSW onset maximum warming anomaly	Max ΔT [K]	Maximum warming anomaly ΔT_{Amax} associated with the SSW-PP-TEA at onset date (based on (5) in Table 1)

^{a)} in the simple two-levels approach (see legend notes a-c of Table 1), where temperature input data solely at 10 hPa and 50 hPa standard pressure levels rather than from vertical profiles are used, the estimated threshold exceedance areas (TEAs) generally are smaller and hence the SSW-MPS classification boundary values are reduced in this case from 90 and 180 $10^6 \text{ km}^2 \text{ days}$ to 70 and 140 $10^6 \text{ km}^2 \text{ days}$, respectively.

删除的内容: of $3 \times 10^6 \text{ km}^2$

删除的内容: 7

带格式的: 字体: 9.5 磅, 字体颜色: 自定义颜色
(*Font/Color: 9.5 pt, Custom Color*)

删除的内容: $10^6 \text{ km}^2 \text{ days}$

删除的内容:

删除的内容: of $3 \times 10^6 \text{ km}^2$

删除的内容: .

Table 3. SSW climatology summary for 1979-2021 based on ERA5 (for definition and units of all parameters see Table 2: winters where no events are detected both by the new method and the BG18 study are skipped in this list).

Winters	Onset date	MPS	MPD	MPA	Type	Max ΔT	Onset location	Onset date BG18
W79-80	—	—	—	—	—	—	—	1980-02-29
W80-81	1981-02-04	88.2	9	9.8	Minor	45.2	83.6°N/46.4°E	1981-02-06
W81-82	—	—	—	—	—	—	—	1981-12-04
W82-83	1983-01-27	73.3	8	9.2	Minor	51.9	78.9°N/87.4°E	—
W83-84	1984-02-23	66.7	8	8.3	Minor	44.6	72.1°N/61.6°E	1984-02-24
W84-85	1985-01-01	192.3	16	12.0	Extreme(TC)	54.5	66.1°N/101.4°W	1985-01-01
W86-87	1987-01-24	51.8	7	7.4	Minor	46.8	65.6°N/39.1°E	1987-01-23
W87-88	1987-12-07	211.9	18	11.8	Extreme	68.4	68.9°N/49.0°E	1987-12-08
W88-89	1989-01-12	275.2	21	13.1	Extreme	49.6	84.9°N/24.2°W	—
	1989-02-20	49.8	7	7.1	Minor	57.6	61.2°N/60.7°W	1989-02-21
W89-90	1990-02-09	97.3	11	8.8	Major	61.6	70.6°N/80.3°E	—
W90-91	1991-01-09	121.8	10	12.2	Major	59.4	72.7°N/51.4°W	—
W91-92	1992-01-11	129.9	13	10.0	Major	64.7	77.6°N/71.9°E	—
W92-93	1993-02-20	33.2	7	4.7	Minor	45.2	73.3°N/72.2°E	—
W94-95	1994-12-30	57.7	7	8.2	Minor	56.9	70.3°N/140.6°E	—
	1995-01-26	144.9	15	9.7	Major	57.3	71.4°N/70.9°E	—
W98-99	1998-12-14	175.8	15	11.7	Major	57.9	72.7°N/100.3°E	1998-12-15
	1999-02-23	85.1	11	7.7	Minor	53.3	77.2°N/89.2°E	1999-02-26
W99-00	—	—	—	—	—	—	—	2000-03-20
W00-01	2000-12-07	83.1	10	8.3	Minor	56.6	68.7°N/68.7°E	—
	2000-12-18	43.9	7	6.3	Minor	39.8	69.1°N/53.9°W	—
	2001-01-28	51.1	7	7.3	Minor	49.6	70.3°N/77.9°E	2001-02-11
W01-02	2001-12-22	224.9	21	10.7	Extreme	64.9	73.0°N/48.9°E	2001-12-31
W02-03	2002-12-28	161.6	14	11.5	Major	74.8	70.8°N/38.3°E	2003-01-18
W03-04	2003-12-24	107.6	14	7.7	Major	48.0	80.7°N/82.1°E	—
	2004-01-04	120.6	15	8.0	Major (TC)	50.7	66.2°N/40.7°E	2004-01-05
W05-06	2006-01-11	43.6	7	6.2	Minor	51.0	72.5°N/59.4°E	—
	2006-01-21	114.6	12	9.5	Major (TC)	48.1	68.4°N/20.1°E	2006-01-21
W06-07	2007-01-01	72.4	10	7.2	Minor	43.1	84.6°N/97.5°E	2007-02-24
	2008-01-23	73.9	8	9.2	Minor	55.8	80.7°N/87.5°E	—
W07-08	2008-02-05	46.7	6	7.8	Minor	57.4	79.0°N/65.7°E	—
	2008-02-23	68.3	8	8.5	Minor	56.9	65.7°N/18.7°E	2008-02-22
W08-09	2009-01-23	366.8	24	15.3	Extreme(TC)	65.9	76.8°N/48.6°W	2009-01-24
W09-10	2010-01-29	68.5	8	8.6	Minor	58.8	68.3°N/59.7°E	2010-02-09
W11-12	2012-01-01	24.7	6	4.1	Minor	39.9	79.4°N/166.2°W	—
W12-13	2012-01-17	112.1	12	9.3	Major	55.2	72.5°N/42.6°E	—
	2013-01-05	191.2	22	8.7	Extreme(TC)	48.5	67.9°N/62.5°E	2013-01-07
W13-14	2014-02-08	100.4	11	9.1	Major	69.1	70.1°N/39.4°W	—
W14-15	2015-01-06	146.5	16	9.2	Major	54.9	70.2°N/30.9°W	—
W15-16	2016-03-05	74.9	9	8.3	Minor	51.1	77.8°N/59.2°E	—
W16-17	2017-01-28	59.4	10	5.9	Minor	56.8	75.8°N/88.9°E	—
W17-18	2017-02-24	27.8	6	4.6	Minor	46.3	73.1°N/71.7°E	—
	2018-02-16	207.5	18	11.5	Extreme	60.9	61.8°N/102.0°W	—
W18-19	2018-12-25	290.8	31	9.4	Extreme(TC)	60.2	77.9°N/72.5°E	—
W19-20	2020-03-21	57.1	7	8.2	Minor	41.7	86.5°N/22.8°E	—
W20-21	2021-01-03	110.2	11	10.0	Major	55.8	75.7°N/11.2°E	—

Table 4. SSW climatology summary 2006-2020 based on RO, for main metrics, for enabling quantitative intercomparison to the results based on ERA5 as summarized in Table 3 (for definition and units of the parameters see Table 2).

Winters	Onset date	MPS	MPD	MPA	Winters	Onset date	MPS	MPD	MPA
W06-07	2007-01-01	49.7	7	7.1	W13-14	2014-02-08	84.0	10	8.4
W07-08	2008-01-23	56.4	6	9.4	W14-15	2015-01-06	128.8	16	8.0
W08-09	2009-01-22	329.7	22	15.0	W15-16	2016-03-05	70.1	9	7.8
W09-10	2010-01-29	56.3	7	8.0	W16-17	2017-01-28	52.2	9	5.8
W10-11	—	—	—	—	W17-18	2018-02-15	186.0	17	10.9
W11-12	2012-01-17	98.2	11	8.9	W18-19	2018-12-25	275.3	30	9.2
W12-13	2013-01-05	164.5	21	7.8	W19-20	2020-03-21	43.0	6	7.2

带格式的：段落间距段后：3 磅

删除的内容: —

删除的内容: —

删除的内容: —


删除的内容: —

刪除的內容

删除的内容:

删除的内容: —

删除的内容: . .






Figure 1. Number and distribution of RO and ERA5 profile data over the Northern polar region. (a): Daily number of RO events over 50-90 °N (blue dots) and 60-90 °N (green dots) for 14 winters from W06-07 to W19-20, with the blue and green horizontal lines showing the related long-term average number, as well as the (constant) daily number of ERA5 grid-point profiles (four analysis times per grid point per day; red and magenta lines); (b): illustrative distribution of RO event locations on 01 Jan 2007 (blue dots), and on the previous and next days (light blue dots), over-plotted on the middle-stratosphere temperature anomaly (MSTA) of the day (color bar), on which a minor SSW prevailed; (c): distribution of the regular ERA5 grid-point profile locations (2.5 °latitude \times 2.5 °longitude grid), over-plotted on the MSTA of 09 Jan 1991, where a major SSW prevailed; (d): illustrative distribution of RO event locations on 25 Dec 2018 in the same style as in (b), over-plotted on the MSTA of the day, on which an extreme SSW prevailed. The green diamonds / yellow circles in (b) and (d) show the location of three exemplary RO events / ERA5 profiles (Event1 to Event3) that are located in different SSW anomaly strength conditions and used in Fig. 2 to illustrate the anomalies construction concept.





Figure 2.



1080 [Figure 2. Schematic overview of the new SSW detection and monitoring approach and its main algorithmic steps from the temperature input data \(top\) to the final SSW metrics \(in particular SSW-MPD, SSW-MPA, SSW-MPS\) and their use for detection, classification and further event characterization \(bottom\). For implementation details see Tables 1 and 2 and the description in Section 3.3.](#)

1085




Figure 3. Illustration of anomaly construction, layer selection, and computation of layer-mean anomaly values based on the three example RO events / ERA5 profiles indicated in Fig. 1b, d. **(a):** Event 1 to 3 temperature profiles from RO and collocated climatological profiles RO_{clim}; **(b):** RO temperature anomaly profiles from the difference of RO to RO_{clim} profiles as well as indication of the lower-stratosphere (20–25 km), middle stratosphere (30–35 km), and upper stratosphere (40–45 km) layers and associated layer-mean anomaly values; **(c):** Profile 1 to 3 temperature and corresponding climatological profiles from ERA5 in same style as (a); **(d):** ERA5 temperature anomaly profiles and layer-mean values in the same style as (b). The RO satellites and event times, and ERA5 analysis time layers, of these examples are (for the locations see the panel headers): Event1: COSMIC-FM1 event 01 Jan 2007 04:28 UTC; Event2: COSMIC-FM1 event 01 Jan 2007 22:44 UTC; Event3: MetOp-A event 25 Dec 2018 17:04 UTC; Prof1: ERA5 profile 01 Jan 2007 06:00 UTC; Prof2: ERA5 profile 02 Jan 2007 00:00 UTC; Prof3: ERA5 profile 25 Dec 2018 18:00 UTC. The climatological profiles are extracted (and interpolated to the needed locations and times) from long term-averaged (2006–2020 for RO and 1979–2020 for ERA5) monthly mean $2.5^{\circ} \times 2.5^{\circ}$ temperature fields.




Figure 4. Temporal evolution of polar-cap mean (60° - 90° N) daily mean temperature anomaly profiles from RO (a) and ERA5 (b), over four winter months each (December, January, February, March), for the winters from 2006-07 (W06-07) to 2019-20 for RO (W19-20) and 2020-21 for ERA5 (W20-21). The RO dataset did not yet cover the W20-21 time period.

删除的内容: 3

带格式的: 字体: 非加粗

删除的内容: .





Figure 5. Temporal evolution of polar-cap mean (60° - 90° N) daily mean temperature anomaly profiles from ERA5, over four winter months each (December, January, February, March), for the winters from W78-80 to W91-92 (a) and from W92-93 to W05-06 (b), respectively. The W78-79 time period was not covered by the ERA5 dataset used.

删除的内容: 4



删除的内容: 5

Figure 6. Polar-view (50° - 90° N) contour maps of W06-07 Jan Minor (left), W90-91 Jan Major (middle) and W18-19 Dec

Extreme (right) SSW example events (cf. Figs. 1 and 6), illustrating ERA5-based Middle Stratosphere Temperature Anomalies (MSTA), Lower Stratosphere Temperature Anomalies (LSTA), and Upper Stratosphere Temperature Anomalies (USTA).

1115 MSTA (top and second row) is shown two days before and on the defined onset date (see Table 2 for definition), LSTA (third row) 4 days after the onset date, and USTA (bottom) 4 weeks after the onset date, respectively.







Figure 7. Time evolution of the daily ERA5 MSTA (a, b, c), LSTA (d, e, f) and USTA (g, h, i) Threshold Exceedance Areas (TEAs) for the same Minor (left), Major (middle) and Extreme (right) SSW events as illustrated in Fig. 5. Panels (j), (k), and (l) depict the four derived SSW TEAs (SSW-PP-TEA, SSW-SP-TEA, SSW-MP-TEA, and SSW-TP-TEA in case it occurs) according to Table 1, (7)-(10); the SSW metrics MPS, MPD, and MPA (see Table 2) are also noted in each panel, and heavy and light lines denote ERA5 and RO results, respectively (differences especially visible for the extreme event). Panels (m), (n), and (o) illustrate geographical tracks (numbered by day-of-winter as of 1st Nov) of maximum positive/negative anomaly values of SSW-PP-TEA (red), SSW-SP-TEA (orange) and SSW-TP-TEA (blue).



刪除的內容:

刪除的內容: 7

Figure 8. Time evolution of the SSW TEAs (SSW-PP-TEA, SSW-SP-TEA, SSW-MP-TEA, SSW-TP-TEA as applicable; definitions see Table 1) for all recorded SSW events of the winters W01-02 to W20-21; the SSW metrics MPS, MPD, and MPA (definitions and units see Table 2) are noted in the panels (E1, E2 are SSW event numbers, ordered according to the occurrence time in a winter). ERA5 results (heavy lines) are complemented by RO results (light lines, especially visible for stronger events) for the winters W06-07 to W19-20.






Figure 9. Time evolution of the SSW TEAs (SSW-PP-TEA, SSW-SP-TEA, SSW-MP-TEA, SSW-TP-TEA) and related SSW metrics (MPS, MPD, MPA) for all recorded SSW events of the winters W80-81 to W00-01; same plotting style as Fig. 8.



删除的内容:

删除的内容: 8... Time evolution of the SSW TEAs (SSW-PP-TEA, SSW-SP-TEA, SSW-MP-TEA, SSW-TP-TEA) and related SSW metrics (MPS, MPD, MPA) for all recorded SSW events of the winters W80-81 to W00-01; same plotting style as Fig. 7






Figure 10. Overview of main characteristics of the SSW events recorded over the 42 winters from 1980 to 2021 using the new monitoring approach based on ERA5 temperature data input at full 137-model-levels resolution ("full-res ERA5") (upper-half part, a-d), coarser standard 37-pressure-levels resolution ("37-plevels ERA5") (co-illustrated in a-b with non-filled symbols),


- 删除的内容: 9
- 删除的内容: 40
- 删除的内容: based on ERA5 data
- 删除的内容: (see Table 3 for
- 带格式的: 字体: 加粗
- 删除的内容: tabular summary). (

1160 [or simple two-levels data extraction \(“10hPa & 50hPa level ERA5”\)](#) (bottom-half part, [e-h](#), respectively, [\(a\)](#) and [\(e\)](#)): time evolution of main-phase strengths (MPS) (year tick marks denote January-of-year); [\(b\)](#) and [\(f\)](#): relation of main-phase and trailing-phase duration and area (MPD, MPA; TPD, TPA) to main-phase strength (MPS); [\(c\)](#) and [\(g\)](#): distribution of the number of events over the winter months (showing the strength types by the same color as in [a\(e\)](#), [b\(f\)](#)); [\(d\)](#) and [\(h\)](#): spatial distribution of onset locations, indicating the clustering of more than 75% of the events over the Northern Eurasia/Polar ocean region by a circle (using [again](#) the same strength-type colors as in [a\(e\)](#), [b\(f\)](#)).

删除的内容: three

删除的内容: (e):

1165



1170 **Figure 11.** Monitoring and assessment of long-term trends in the three core SSW metrics (main-phase strength, duration, and area) along the recent climate change period from the 1980s decade to the 2010s decade, noting also main statistical numbers within the respective panel legends. The results derive from the metrics as illustrated in Figure 10 and are intercompared for the ERA5 temperature data input either taken in at full 137-model-levels resolution (“full-res ERA5 data”) (top row), coarser standard 37-pressure-levels resolution (“37-plevels ERA5 data”) (middle row), or simple two-levels data extraction (“10hPa & 50hPa ERA5 data”) (bottom row), respectively. For description see Section 5.2.

带格式的：正文

删除的内容：results

删除的内容：panels (

删除的内容：text).