

An attribution of the low single-scattering albedo of biomass-burning aerosol over the southeast Atlantic

Amie Dobracki¹, Paquita Zuidema¹, Steve Howell², Pablo Saide³, Steffen Freitag², Allison C. Aiken⁴, Sharon P. Burton⁵, Arthur J. Sedlacek III⁶, Jens Redemann⁷, and Robert Wood⁸

¹Department of Atmospheric Sciences, Rosenstiel School, University of Miami, Miami, Florida, USA

²University of Hawai'i at Mānoa, Honolulu, Hawaii, USA

³University of California Los Angeles, Los Angeles, California, USA

⁴Earth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, New Mexico, USA

⁵NASA Langley Research Center, Hampton, VA, USA

⁶Brookhaven National Laboratory, Upton, New York, USA

⁷University of Oklahoma, Norman, Oklahoma, USA

⁸University of Washington, Seattle, WA, USA

Correspondence: Paquita Zuidema (pzuidema@miami.edu) and Amie Dobracki (amie.dobracki@rsmas.miami.edu)

Abstract. Aerosol over the remote southeast Atlantic is some of the most sunlight-absorbing aerosol on the planet: the *in-situ* free-tropospheric single-scattering albedo at the 530 nm wavelength (SSA_{530nm}) ranges from 0.83 to 0.89 within ORACLES (ObseRVations of Aerosols above CLouds and their intERactions) aircraft flights from late August-September. Here we seek to explain the low SSA. The SSA depends strongly on the black carbon (BC) number fraction, which ranges from 0.15 to 0.4. Low organic aerosol (OA) to BC mass ratios of 8-14 and modified combustion efficiency values > 0.975 point indirectly to the dry, flame-efficient combustion of primarily grass fuels, with back trajectories ending in the miombo woodlands of Angola. The youngest aerosol, aged 4-5 days since emission, occupied the top half of a 5 km thick plume sampled directly west of Angola with a vertically-consistent BC: Δ CO (carbon monoxide) ratio, indicating a homogenization of the source emissions. The younger aerosol, transported more quickly off of the continent by stronger winds, overlaid older, slower-moving aerosol with a larger mean particle size and fraction of BC-containing particles. This is consistent with ongoing gas condensation and the coagulation of smaller non-BC particles upon the BC-containing particles. The particle volumes and OA:BC mass ratios of the older aerosol were smaller, attributed primarily to evaporation following fragmentation, instead of dilution or thermodynamics. The CLARIFY (CLoud-Aerosol-Radiation Interaction and Forcing: Year-2017) aircraft campaign sampled aerosols that had traveled further to reach the more remote Ascension Island. CLARIFY reported higher BC number fractions, lower OA:BC mass ratios, lower SSA yet larger mass absorption coefficients compared to this study's. Values from one ORACLES-2017 flight, held midway to Ascension Island, are intermediate, confirming the long-range changes. Overall the data are most consistent with continuing oxidation through fragmentation releasing aerosols that subsequently enter the gas phase, reducing the OA mass, rather than evaporation through dilution or thermodynamics. The data support the following best-fit: $SSA_{530nm} = 0.801 + 0.0055 * (OA:BC)$ ($r = 0.84$). The fires of southern Africa emit approximately one-third of the world's carbon; the emitted aerosols are distinct from other regional smoke emissions and their composition needs to be represented appropriately to realistically depict regional aerosol radiative effects.

1 Introduction

Biomass burning, the largest source of carbon to the atmosphere globally, is fundamental to the Earth's global carbon cycle (Bowman et al., 2009; Bond et al., 2013). The emissions include carbon monoxide, carbon dioxide, methane and carbonaceous aerosols, significantly altering the atmospheric composition over large regions of the globe (Andreae, 2019). This in turn influences all of the gaseous, aerosol and aerosol-cloud interaction radiative forcing terms considered within the IPCC Assessments. Despite the importance of biomass burning events on climate, smoke properties after long-range transport are still poorly characterized. These include the effluents from northern European and Russian forest fires reaching the Arctic basin (Cubison et al., 2011), wildfire smoke from western continental north America observed over Europe (Zheng et al., 2020; Baars et al., 2021), and aerosols from fires in southern Africa reaching south America (Holanda et al., 2020). Without wet or dry scavenging, the aerosol's areal coverage is increased through transport, increasing the aerosol's radiative impact.

Southern Africa region produces approximately one-third of the world's fire-emitted carbon (van der Werf et al., 2010). The global maximum of absorbing aerosol above cloud occurs above the southeast Atlantic (Waquet et al., 2013), a combination that produces a direct radiative warming of the regional climate (Keil and Haywood, 2003; Graaf et al., 2014; Zuidema et al., 2016; Mallet et al., 2021; Doherty et al., 2022). Biomass-burning aerosol (BBA) from this region is unusual for being highly absorbing of sunlight, with SSA values of 0.85 or less at the green wavelength (Zuidema et al., 2018; Chylek et al., 2019; Pistone et al., 2019; Holanda et al., 2020; Taylor et al., 2020; Denjean et al., 2020b; Mallet et al., 2020; Shinozuka et al., 2020; Carter et al., 2021; Brown et al., 2021). More absorbing aerosol will reduce the need for latent heat release as a balance to longwave radiative cooling within the world's energy distribution (Pendergrass and Hartmann, 2012) and alters regional circulation and precipitation patterns (Mallet et al., 2020; Solmon et al., 2021; Chaboureau et al., 2022). While climate models discern an ensemble-mean direct radiative warming, individual models disagree strongly on magnitude and even sign (Myhre et al., 2013; Zuidema et al., 2016; Haywood et al., 2021; Mallet et al., 2021). In addition, the direct aerosol radiative effect estimated from satellites typically exceeds model estimates (de Graaf et al., 2020). That the measured SSAs are lower than what is currently implemented in many models (Shinozuka et al., 2020; Mallet et al., 2021; Doherty et al., 2022), suggests one cause for an underestimated modeled direct radiative warming is a model SSA that is too high.

This study's goal is to examine the optical properties and composition of *in-situ* smoke sampled in the free troposphere during six flights of the NASA Earth Venture Suborbital-2 ORACLES (ObseRvations of Aerosols above CLouds and their intERactionS; Redemann et al., 2021) deployment, primarily from September, 2016 (Fig. 1). Complementary observations were taken in this region by the UK CLARIFY (CLoud-Aerosol-Radiation Interaction and Forcing: Year-2017) aircraft campaign (Haywood et al., 2021) on Ascension Island (8°S, 14.5°W), from August 17 - September 7, 2017, and the DACCIIWA (Dynamics-Aerosol-Chemistry-Clouds Interactions in West Africa) airborne campaign over southern west Africa during June-July 2016 (Knippertz et al., 2015). Both campaigns have already revealed that southern African biomass-burning aerosol (BBA) is highly absorbing of sunlight because the fractional black carbon content is high in both number and mass of total particles (Taylor et al., 2020; Denjean et al., 2020b), with loss of particle coating also contributing (Sedlacek et al., 2022). We further strengthen the attribution to aerosol composition, fire source characteristics and indicators of chemical aging and seek to

place the ORACLES data within the context of these other measurements. The aerosol sampled during the Southern African Regional Science Initiative (SAFARI) campaign (Haywood et al., 2003) held in 2002 near Namibia was less than 3 days of age, while the CLARIFY-2017 campaign over Ascension sampled aerosol approximately a week old (Wu et al., 2020; Taylor et al., 2020). The ORACLES-2016 model-derived age estimates place the flights uniquely within the time record available on
60 chemical aging over the southeast Atlantic.

The paper is organized as follows: Section 2 describes the Methodology, including description of the flights and relevant details about the datasets, with the more technical details relegated to the Supplement. Section 3 presents the chemical characterization, including the aerosol age estimates. Section 4 discusses the chemical optical and physical properties of the smoke plumes, including the likelihood of brown carbon. Section 5 investigates how the organic aerosol component varies between the
65 flights and if OA differences can be explained as a aging process, with Section 6 incorporating a comparison to measurements made at Ascension. Section 7 provides a summary and discussion.

2 Methodology

Six flights were selected, based on the availability of at least 20 minutes of organic aerosol (OA) masses exceeding $>20 \mu\text{g m}^{-3}$ (a threshold justified in section 2.4.2), at altitudes above 1.5 km and relative humidities (RH) $< 80\%$. The latter is applied
70 to reduce the likelihood of aqueous-phase reactions. The flights occur within 30 days of each other in the seasonal cycle, spanning August 31 through all of September, to preferentially select for similar composition of the fire source emissions. Five flights come from 2016 and one from 2017, with their tracks shown relative to the satellite-derived above-cloud aerosol optical depths for September 2016 in Fig. 1. Spatial maps of the aerosol forecasts used to guide the flight planning are shown overlaid
75 (explained further in Section 2.2) on individual altitude-latitude flight track projections in the Supplementary Figs. S1-S2.

More description of the flights is followed by descriptions of the datasets that are more central to the results (listed in Table 1), with the more technical details provided in the Supplement.

2.1 Flight Description

The aircraft flew along a routine southeast to northwest track on three flights (31 August, 4 and 25 September of 2016), and
80 performed three target-of-opportunity flights sampling more aerosol-rich locations (6 and 24 September of 2016, 31 August 2017), all shown in Figs. S1-S2. The flight tracks make clear that the aircraft sampled widely, but never near the fire emission sources, with the 9/24/2016 flight coming closest (Fig. 2). The aerosol spatial distribution is strongly influenced by the strength of the free-tropospheric easterly winds, with the aerosol either constrained to be near the coast when the winds are weak, or elongated zonally along 10°S when the winds are strong. On the 24-25 September 2016 and 31 August 2017 flights, the zonal
85 easterly winds exceeded 6 m s^{-1} along $\sim 10^\circ\text{S}$ at altitudes between 3-5 km, forming a wind isotach known as the African Easterly Jet-South (Nicholson and Grist, 2003; Adebisi and Zuidema, 2016). Overall, September 2016 was climatologically representative (Ryoo et al., 2021), with more synoptic detail on individual flight days available in Ryoo et al. (2022).

An aerosol forecast model was used to seek out smoke layers to sample during the target-of-opportunity flights. The six flights intersect aerosol of different ages, but all model-estimated ages exceed 4 days. Table S1 lists all of the ORACLES-2016 flights and includes comments on their flight pattern, the number of seconds with $OA > 20 \mu\text{g m}^{-3}$, and other selection considerations. We highlight two flights further: one from 9/24/2016, because it sampled a thick, younger smoke plume as close as possible to a fire emission source (Fig. 2), and the 9/31/2017 flight, which sampled an aerosol layer of significant mass and extremely stable OA:BC halfway between Ascension and the African continent, and helps connect interpretations of ORACLES to CLARIFY aerosol characteristics. The focus on these two flights has been added since (Dobracki et al., 2022). Optical properties were primarily examined for data from level legs, for which further time averaging could reduce the measurement uncertainty. Table S2 provides flight dates, location, time span and altitude of the level [legs leg data we analyze further, ranging four to ten minutes in length.](#)

2.2 Determination of physical aerosol age

Model-released tracers tagged to CO at the fire source for each day of the campaign's operational two-week aerosol forecast, made using the Weather Research Forecasting - Aerosol Aware Microphysics (WRF-AAM) Model (Thompson and Eidhammer, 2014), were used to estimate the physical age of the aerosol. The regional model has a 12-km spatial resolution and encompasses a domain (41°S - 14°N , 34°W - 51°E) sufficiently large to capture almost all contributing fires (Saide et al., 2016). The model is driven by the National Center for Environmental Prediction Global Forecasting System (NCEP GFS) meteorology, using daily smoke emissions from the Quick Fire Emissions Dataset (Darmenov and da Silva, 2013) released into the model surface layer. These are advected thereafter according to the model physics, with their spatial distribution constrained near real-time with satellite-derived optical depths. Most fires in southern Africa occur during the day, and the satellite constraint captures this diurnal cycle. The model fire emissions rely on a burned-area product of 500 m spatial resolution from the Moderate Resolution Imaging Spectrometer (Giglio et al., 2006). This may miss up to $\sim 40\%$ of the total burned area coming from smaller fires (Ramo et al., 2021). Larger fires, with more protected cores, contribute more to the emissions reaching higher altitudes for the boreal northern hemisphere (Martin et al., 2010). We are unsure how well this same vertical selection applies for the smaller agricultural fires of southern Africa, although stronger zonal winds aloft will aid lofting of the smoke underneath (Adebisi and Zuidema, 2016). The tracer-derived estimates tend to keep the smoke emissions near the surface until the aerosol is eventually carried aloft, shown for 24 September 2016 in Fig. S3. The time lag allows the emissions from nearby fire sources to mix, homogenizing local differences in, e.g., grass versus leaf-litter, moisture content, and surface burn history. Conditions can nevertheless still change from day to day.

Backtrajectories based on the HYSPLIT model (Stein et al., 2015), [also driven by the same National Centers for Environmental Prediction Global Forecasting System \(NCEP GFS\) meteorology used to drive WRF-AAM](#) [NCEP GFS meteorology](#), further illuminate the pathway [taken by the BBA. The backtrajectories end when they the BBA traveled prior to its sampling on 24 September 2016 \(shown later in Fig. 12d\). These backtrajectories reach the location of a the fire emission source, and typically generate younger aerosol age estimates than the aerosol forecasts; after approximately three days, at higher altitudes than where the aerosol was sampled. The aerosol age estimate is younger than the WRF-AAM aerosol forecasts would suggest. This is](#)

because the time needed for the aerosol ~~vertical transport is unaccounted for~~, to travel from the surface to the higher altitude is not accounted for in the HYSPLIT age estimate. The vertical advection timescale is highly model-dependent on boundary layer parameterizations and we merely note the difference in the two age estimates here.

125 2.3 Modified combustion efficiency

CO and carbon dioxide (CO₂) are used to infer fire emission conditions through the modified combustion efficiency (MCE) metric (Collier et al., 2016; Yokelson et al., 1997):

$$MCE = \frac{\Delta CO_2}{\Delta CO + \Delta CO_2} = \frac{1}{1 + \Delta CO / \Delta CO_2} \quad (1)$$

An MCE of 0.9 marks the 50% threshold between flaming and smoldering combustion (Akagi et al., 2011), a threshold that is largely insensitive to fuel type (May et al., 2014). Higher values of MCE (>0.9), more associated with flaming combustion, preferentially produce more BC, whereas an MCE < 0.9 is more typical of a smoldering fire that emits more organic aerosol for the same amount of fuel (Yokelson et al., 2009; Vakkari et al., 2018). A regression is used to estimate the $\Delta CO / \Delta CO_2$ with ΔCO and ΔCO_2 calculated from the measured CO and CO₂ amounts, in moles, relative to background values. Adopted background values were 65 (77) ppbv for CO, and 397 (404) ppmv for CO₂, in September 2016 (August 2017), based on measurements in the free troposphere taken above the smoke plumes (~7000 m).

2.4 Aerosol Composition

2.4.1 Black carbon

~~BC mass and number concentrations~~ The rBC size distributions are derived from a 4-channel single particle soot photometer (SP2, Droplet Measurement Technology) deployed by the Hawaii Group for Environmental Aerosol Research HiGEAR in 2016, and an 8-channel SP2, of which only the incandescent channels were functional for the August 31, 2017 flight, deployed by Art Sedlacek of Brookhaven National Laboratory. No scattering data are available, precluding information on coating thicknesses. The intensity of laser-induced incandescent emission at 1064 nm can be quantitatively related to the mass of the refractory black carbon (rBC) particles for mass-equivalent diameters between approximately ~~80-500 nm. This size range can successfully capture 99%~~ 0.08-0.5 μ m. The SP2 was calibrated using fullerene soot using effective density estimates from Gysel et al. (2011). Calibration uncertainty dominates the nominal mass uncertainty of $\pm 17%$ (Laborde et al., 2012).

Size distributions by number and mass (assuming a BC density of 1.8 g cm^{-3} (Bond and Bergstrom, 2006)) for the level legs detailed in Table S2 indicate particle diameters remain well below the upper SP2 detection limit (Fig. 3). Lognormal fits help visualize a drop-off in detection efficiency for diameters $< 0.08 \mu\text{m}$ for the samples weighted towards the smaller sizes (Fig. S4, for 9/24/2016 and 9/25/2016). The SP2 size range nevertheless captures almost all of the black carbon mass (Taylor et al., 2020) ~~, close to the 99% value reported for CLARIFY (Taylor et al., 2020). rBC total number concentrations vary between 530 to 1370 cm^{-3} (Fig. 3a), and undercounting of the mass and number through coincidence is estimated to be less than 3% based on Taylor et al. (2020).~~ Throughout, we use BC to refer to the SP2-derived refractory black carbon, following other literature,

although the two are not entirely the same (Petzold et al., 2013). ~~BC number concentrations were almost always below 1000~~
~~em⁻³, and undercounting of the mass and number through coincidence is estimated to be less than 2% (Taylor et al., 2020).~~
155 ~~The SP2 was calibrated using fullerene soot using effective density estimates from Gysel et al. (2011). Calibration uncertainty~~
~~dominates the nominal mass uncertainty of ±17% (Laborde et al., 2012).~~

Ratios of $\frac{\Delta BC}{\Delta CO}$, reducing to $\frac{BC}{\Delta CO}$ because the background BC is zero in clean conditions, serve to assess homogeneity of the aerosol composition at the emission source. The ratios are non-dimensionalized by using the ideal gas law at standard temperature (273K) and pressure (1000 hPa) to convert the CO concentrations from ppb to ng m⁻³.

160 2.4.2 Aerosol Mass Spectrometer measurements

HiGEAR operated an Aerodyne High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS, referred to as AMS), building on previous experience in the southeast Pacific (Yang et al., 2011; Shank et al., 2012) and the Arctic (Howell et al., 2014). This measured masses of organic aerosol (OA), nitrate (NO₃), sulfate (SO₄) and ammonium (NH₃). Chloride, a small component of the total aerosol mass in the free troposphere, was not considered because of its inconsistent ionization
165 signature. The native time resolution is approximately five seconds, with the data interpolated onto a one-second temporal grid to facilitate integration with other datasets. The overall uncertainty in the reported aerosol mass concentrations is estimated at 33% to 37% at a one-minute time resolution, based on Bahreini et al. (2009), generating a combined uncertainty in the OA to BC mass ratio of close to 40% (since the background OA in clean conditions is also zero, $\frac{\Delta OA}{\Delta BC}$ reduces to OA:BC).

Means over ~~level legs, ranging from 4 to 10 minutes in length, the level legs~~ listed in Table S2, further reduce OA mass
170 uncertainties to 19%-10 %, and to 25%-19% for the OA:BC mass ratio. The instrument inter-comparison flight with CLARIFY sampled a clean troposphere but a polluted boundary layer (BC of ~ 300 ng m⁻³), during which the ORACLES OA and nitrate mass concentrations were 80% of those measured by the UK plane, within each other's standard deviations (Barrett et al., 2022).

A threshold of 20 μg m⁻³ is applied to the OA mass to select for the heart of the smoke plumes. This is one approach to minimizing dilution effects, by which OA evaporates through mixing with cleaner environmental air (e.g., Hodshire et al.,
175 2021). This threshold was selected based on when a stabilization of the OA:BC mass ratio occurs as a function of the OA mass concentrations (Fig. S4 4). The OA:BC mass ratio is significantly less for air with OA > 3 μg m⁻³ than for air with OA > 20 μg m⁻³, particularly for younger aerosol (Fig. S4a S5), consistent with evaporation through dilution. The choice of threshold is inherently arbitrary, and some analysis is repeated using an OA > 10 μg m⁻³ threshold to make sure our findings are not sensitive to the choice of OA mass threshold (Figs. S11, S13). The BC:ΔCO and OA:BC ratios are shown for individual flights
180 as a function of f44 for both thresholds in Figs. S11-S14 in the interest of full documentation. We also account for dilution effects by normalizing OA with respect to BC or ΔCO, two quantities that do not change with dilution. We stress that the aerosol plumes over the southeast Atlantic, termed 'rivers of smoke' within Swap et al. (2003), are typically much larger and homogeneous than the fire plumes sampled in the western northern hemisphere, which are often linked to named, individual fires and sampled close to the source. The plume sampled on 8/31/2017, halfway to Ascension Island, is an example of a wide,
185 homogeneous smoke plume (Fig. 3). OA mass concentrations often remained highly stable over level legs (see Fig. S2, bottom row, for an example) ~~and aerosol plume dilution should be much less of a concern than for the overall less smoky northern~~

hemisphere. Further justification for a threshold is that the OA mass uncertainty is smaller at higher signal-to-noise ratios. Additionally, model-observational disparities in the smoke plume locations have less impact on further aging-related analyses if based on the plume centers. For the same reason we exclude aerosol with physical ages > 10 days as the model skill in
190 predicting smoke age is likely poorer by then.

Other AMS measurements include the fraction of the OA mass spectrum signal at m/z 44, 43 and 60 relative to the total OA mass concentration, termed f_{44} , f_{43} , f_{60} , and hydrogen (H), oxygen (O), and organic carbon (OC). f_{44} indicates the presence of the CO_2^+ ion, a form of oxidation resulting from chemical aging (Canagaratna et al., 2015). f_{43} indicates the presence of $\text{C}_2\text{H}_3\text{O}^+$ and C_3H_7^+ , also representative of oxygenated OA. f_{60} indicates $\text{C}_2\text{H}_4\text{O}_2$, a fragment of levoglucosan and a known
195 tracer for biomass burning aerosol (Cubison et al., 2011). Elemental analysis, yielding H, O and OC rely on algorithms from Aiken et al. (2007). The calibration constants differ between the two years but this change does not quantitatively impact any differences shown here.

2.5 Determination of organic/inorganic nitrate contribution

Farmer et al. (2011) provide an approach for estimating the contribution to the total nitrate signal from organic nitrate (ON) using the $\text{NO}^+:\text{NO}_2^+$ ratio, building on the observation that organic nitrates typically fragment into larger proportions of NO^+
200 than do inorganic nitrates. In their study, organic NO^+ ratios vary between 1.8 to 4.6 for different organonitrates, compared to 1.5 for NH_4NO_3 . Their Equation 1, reproduced below, provides an estimate of the ON fraction that can be readily applied to the ORACLES AMS data, assuming enough ON is present that it can be resolved. The success of this approach also assumes that the inorganic nitrates capable of providing a large NO^+ ratio, such as mineral nitrates, are not present. Both assumptions
205 are justified for the [SEA-southeast Atlantic \(SEA\)](#) free troposphere.

$$X(\text{ON}\%) = \frac{(R_{obs} - R_{\text{NH}_4\text{NO}_3})(1 + R_{ON})}{(R_{ON} - R_{\text{NH}_4\text{NO}_3})(1 + R_{obs})} \quad (2)$$

R_{obs} is the ORACLES m/z ratio of ion fragments 30 to 46, $R_{\text{NH}_4\text{NO}_3}$ is the ionization efficiency (IE) calibration-derived ratio (1.26 for 2016 and 1.545 for 2017) and an R_{ON} value of 3.41 is a reference ratio based on the average fragmentation pattern into the $\text{NO}^+:\text{NO}_2^+$ ratios for the OIA-HN, OIA-CN and OIA-olig standards evaluated within Table S1 of Farmer et al. (2011).
210 The inorganic nitrate (IN) fraction is 1-ON. We use this approach to estimate the inorganic nitrate (IN, primarily NH_4NO_3) fraction, keeping in mind that it is an indirect inference. The CLARIFY campaign relied on assessing m/z 30 and m/z 46 (primarily NO^+ and NO_2^+) to assess the organic to inorganic nitrate contribution.

2.6 Aerosol sizing

Total aerosol number concentrations from a Condensation Particle Counter (CPC; TSI 3010, marked 'ACN' in Fig. [S7S8](#))
215 establish the fraction of BC-containing particles. The CPC counter applies a size threshold of 10 nm with no upper bound. Aerosol size distribution measurements rely on a long differential mobility analyzer (LDMA; heavily modified from a TSI 3071A). The aerosol sizing from an LDMA and a thermal DMA (TDMA) are preferred to those from a Ultra-High Sensitivity Aerosol Spectrometer (UHSAS; DMT) because of UHSAS sizing uncertainties (Howell et al., 2021). A correction for a known

undersizing by the UHSAS, put forward in Howell et al. (2021), is evaluated in Fig. S5S7, in which the LDMA and UHSAS median diameters are compared for the level-leg plumes (Table S2) with $OA > 20 > 20 \mu\text{g m}^{-3}$. Two particle populations emerge: one with diameters between 0.15-0.2 μm that is more likely to contain black carbon (see Fig. 3), and another with median diameters $< 0.1 \mu\text{m}$, speculated to consist mostly of OA. For the larger particles containing BC, the UHSAS correction reduces the undersizing bias to 15% compared to LDMA median diameters $> 150 \text{ nm}$, but for the smaller particles that are less likely to contain BC, the UHSAS particle sizes are now overcorrected. For this reason, and because a particle cavity aerosol spectrometer probe (PCASP) underperformed, we only show aerosol sizes based on the LDMA data. Analysis duplicated using UHSAS data did not contradict our findings.

The LDMA measures mostly singly-charged particles between 10-550 nm in mobility diameter, with multiply-charged particles occurring at diameters $> 200 \text{ nm}$ (Howell et al., 2021). The inversions include a size-dependent charging efficiency that accounts for the multiple charges and for size-dependent losses (Zhou, 2001). The HiGEAR LDMA operated in a scanning mode at ambient temperature and pressure, drawing in desiccated air ($RH < 30\%$) from an aluminum lagged-aerosol grab chamber for 60 seconds. The total number uncertainty is estimated to be $\pm 30 \%$ due to errors in sizing of non-spherical particles along with uncertainties of flow rate. All size distributions and concentrations are corrected to STP ($T = 25^\circ\text{C}$, $p = 1000 \text{ hPa}$).

The black carbon core mass-median diameter is ~~also estimated using the SP2-provided mass and number concentrations, towards inferring used to infer~~ fire conditions at the source. Larger BC sizes ~~can~~ correspond to more woody fuels than grasses (Holder et al., 2016) and ~~in Holder et al. (2016), while~~ larger BC cores are associated with more flaming conditions in Pan et al. (2017), attributed to less oxygen reaching the interior flame zone. ~~The mass-median diameter calculation assumes a BC density of 1.8 g cm^{-3} (Bond and Bergstrom, 2006).~~ An estimate of the fraction of total particles containing black carbon (FrBC) is also constructed from the total number of SP2-derived BC particles divided by the total CPC particles.

2.7 Optical Measurements

Scattering from all particles is measured continuously by a nephelometer (TSI 3563) at the (450, 550, 700) nm wavelengths (λ), from which scattering coefficients (σ_s) are retrieved. The spectral light absorption coefficients (σ_a) of the total aerosol are estimated from Particle Soot Absorption Photometer (PSAP; Radiance Research) measurements at the (470, 530, and 660) nm wavelengths. The nephelometer scattering measurements are interpolated to the PSAP wavelengths. The extinction (scattering+absorption) and absorption measurements compare well at the blue and green wavelengths to the more sophisticated measurements made by the CLARIFY EXtinction SCattering and Absorption of Light for AirBorne Aerosol Research (EXS-CALABAR) instrument (Davies et al., 2019; Barrett et al., 2022). More detail on the algorithmic treatment of the filter-based measurements is provided in Section 7 of the Supplement.

The absorption Ångström exponents (AAE) are calculated from the linear fit of $\log(\sigma_a)$ to $\log(\lambda)$. The mass absorption cross-section at 660 nm ($MAC_{BC,660}$) is based on $\sigma_{a,660}$ divided by the BC mass concentration. Following Carter et al. (2021), we also evaluate the MAC relative to the BC+OA mass concentration at $\lambda = 470 \text{ nm}$ ($MAC_{BC+OA,470}$), to assess absorption contributions from both OA-induced brown carbon as well as other wavelength-dependent absorbers (Zhang et al., 2022). The single-scattering albedo is examined at 530 nm ($SSA_{530}; = \frac{\sigma_{a,530}}{\sigma_{s,530} + \sigma_{a,530}}$) to support comparisons to other published values.

3 Chemical composition and age distribution within the six flights

The mean submicron mass fractions of the six flights combined are 66% OA, 10% nitrate (NO₃), 11% sulfate (SO₄), 5% ammonium (NH₄), and 8% BC, with the masses for each species and flight in Fig. 5, thresholded for OA > 20 g m⁻³. Flight-mean submicron mass totals typically exceed 35 μg m⁻³. This is much more than measured in the free troposphere above Ascension during CLARIFY (Wu et al., 2020), although the OA mass fraction during CLARIFY still remained > 50% of the total aerosol mass.

Fig. 6 provides an overview of the *f*₄₄, OA to BC mass ratio, model-derived time since emission (age), MCE, non-dimensionalized $\frac{BC}{\Delta CO}$ ratios and ozone values for each flight. *f*₄₄ flight-mean values range from 0.18 to 0.22, on par with *f*₄₄ values of Asian/Siberian smoke after a two-week transport to Alaska (Cubison et al., 2011). They are also similar to CLARIFY values (Wu et al., 2020), suggesting a maximum *f*₄₄ value of ~ 0.22 for this aerosol regime. The *f*₄₄ values indicate highly-oxidized aerosol but their range may still contain information on the relative aerosol age: the lowest flight-mean *f*₄₄ value from the 9/24/2016 flight, corresponds to the youngest aerosol (Fig. 6c), although the other flight-mean aerosol ages since emission do not correlate well to *f*₄₄.

Flight-mean OA:BC mass ratios range from 7 to 13. MCE values are above 0.97 for each flight. These clearly indicate flame-efficient fires (Collier et al., 2016; Zhou et al., 2017), whose emissions can also more easily reach higher altitudes than can emissions from smoldering fires (Kondo et al., 2011). This may explain why the ORACLES-2016 MCE values exceed the September-mean estimate of ~ 0.89 from a source emission-based model (Zheng et al., 2018). Mean non-dimensionalized $\frac{BC}{\Delta CO}$ ratios vary between 0.007 to 0.011, with a minimum on 24 September. These ratios are among the highest surveyed in the literature (Table 2). Overall, $\frac{BC}{\Delta CO}$ ratios do not increase with increasing MCE as expected based on Kondo et al. (2011), but this likely reflects our study's small range of MCE values, for which Vakkari et al. (2018) also do not find a correlation. The mean $\frac{BC}{\Delta CO}$ values hint at a decrease throughout September, with the flight-mean OA:BC mass ratios also increasing to 13 later in September. The flight-mean ozone levels range from 80-105 ppbv, possibly decreasing as September progresses. Flights with more ozone correspond to flights with lower MCEs: less flaming fires will emit more ozone along with more OA. The changes over the course of the month are consistent with more combustible fuel being ignited earlier (Eck et al., 2013), but none of the trends are statistically-insignificant.

We interpret the high MCE values to reflect a large contribution from dry and dead grasses, rather than green grass or more woody materials, for the following reasons. MCE varies inversely with the moisture content for grasses (Korontzi et al., 2003), with leaf litter and woody fuels tending to dry more slowly than do grasses. For this reason woody fuels are more prone to smoldering than flaming combustion. The burning of dry grass produces relatively low emissions of carbon monoxide (Scholes et al., 1996) and higher emissions of black carbon than do agricultural or woodland fires (Andreae, 2019), elevating the $\frac{BC}{\Delta CO}$ ratios, as seen here (Table 2). That the $\frac{BC}{\Delta CO}$ values measured at offshore locations exceeds those measured previously over land (Table 2) could be because emissions from more intense, larger, flaming fires can more easily reach higher altitudes (Martin et al., 2010; Holder et al., 2016), where they can be dispersed further afield through the stronger winds aloft.

Daily maps of fire locations for the flight days (not shown, see Redemann et al. (2021) for the monthly-mean distributions) indicate the BBA sources are primarily fire emissions from miombo woodlands, which contain a significant fraction of savanna grasses and some agricultural fields (Shea et al., 1996; Christian et al., 2003; Korontzi et al., 2003; Vakkari et al., 2018; Huntley, 2019), distributed over a broad geographic region encompassing Angola, Zambia and the Congo. The miombo shrubbery is fire-adapted and less likely to burn than the grass. A survey of the published emission factors for the vegetation types typical of southern Africa - savannahs, grasslands, agricultural fields, and at times tropical forest indicates that the high $\frac{BC}{\Delta CO}$ ratios reported in Table 1 and Fig. 6 are primarily representative of grass fires. Overall, these metrics indicate aged, oxidized aerosol emanating from flame-efficient fires, without any strong outliers amongst the flights (flight-mean $\frac{BC}{\Delta CO}$ ratios vary from $7-11 \times 10^{-3}$), typical values for grasslands and savannahs (Janhäll et al., 2010; Vakkari et al., 2018).

Southern African fires can still produce significant near-source secondary organic aerosol (SOA), depending on the burning conditions (Vakkari et al., 2018; Pokhrel et al., 2021). The comparison of f_{44} to f_{43} for all the flights (Fig. 7a) indicates a mixture of semi-volatile and low-volatile oxygenated organic aerosol (Ng et al., 2011). A PIKA analysis reveals the dominant peak at f_{43} is from $C_2H_3O^+$, representative of oxygenated organic aerosol (Ng et al., 2011). f_{60} values are relatively constant and below 0.005 (Fig. 7b), and f_{44} values lie between 0.2 and 0.22. Chamber studies report lifetimes for f_{44} and f_{60} of approximately 20 days and 10 hours, respectively (George and Abbatt, 2010; Hodshire et al., 2019), but little change is evident in f_{44} after 6 days since emission (Fig. 7c), with f_{44} values of 0.2-0.22 also reported at Ascension (Wu et al., 2020), suggesting a steady state has been reached.

The H:C versus O:C mass ratios occur close to the -1 slope line (Fig. 8; based on vanKrevelen (1950)), also inferred at Ascension (Wu et al., 2020). This slope relationship is common to many laboratory and field studies (Heald et al., 2010), with the narrow distribution, particularly within individual flights, suggesting either a limit to the number of oxidation pathways and molecular structures, or, a dominant few. Most of the oxidation states (OS), defined as $2 \times O:C-H:C$ (Kroll et al., 2011), lie between -0.2 and 0.5, which Kroll et al. (2011) categorize as “aged” and semi-volatile oxygenated OA (OS between -0.5 to 0). Only the 31 August, 2017 flight has some aerosol that is oxidized enough to be considered low-volatile (OS > 0.5). We are only able to report the end product of the aerosol chemical properties, and different SOA precursors may also contribute to the range of the observed H:C and O:C ratios (Jimenez et al., 2009; Ng et al., 2011). Nevertheless, Kroll et al. (2009) show aerosol with O:C > 0.4 are dominated by fragmentation pathways, in which further oxidation occurs through the loss of a carbon atom (as opposed to functionalization, which adds an oxygen atom). Fragmentation generates relatively small changes in H:C. The fragmentation process releases small amounts of volatile aerosol and we speculate this pathway is suggested by Fig. 8 for the continuing oxidation of ORACLES-2016 BBA.

Flight-mean O:C mass ratios range between 0.61 to 0.69 for the 2016 flights, with small within-flight standard deviations (0.03-0.06, not shown). Overall, the average (\pm standard deviation) plume values of H:C, O:C, and the organic-aerosol-to-organic-carbon mass ratio (OA:OC) are 1.2 ± 0.1 , 0.7 ± 0.1 , and 2.2 ± 0.1 , respectively, over all six flights. The OA:OC mass ratio, a measure of the oxygen content that is useful for model evaluation (Hodzic et al., 2020; Lou et al., 2020), are on par with measurements from the Atmospheric Tomography (ATom) campaign made in the same region (Hodzic et al., 2020) and during CLARIFY (Wu et al., 2020). The mean value of 2.2 is substantially higher than common model-applied values of 1.4-

1.8 (Aiken et al., 2008; Tsigaridis et al., 2014; Hodzic et al., 2020) and of primary near-source OA:OC ratios of 1.6 (Andreae, 2019).

4 Optical and Physical Properties

Here we discuss relationships between the aerosol optical properties to their chemical and physical composition, and examine their spatial distribution, using the more statistically-robust level-leg mean (\pm standard deviations) values (Figs. 9-11).

4.1 Mean Relationships

~~Absorption of sunlight primarily depends on BC, and as expected, the~~

~~The bulk mass absorption coefficients ($MAC_{BC,660}$) and SSA values depend strongly on decrease with~~ the estimate of the fraction of particles containing black carbon (~~FrBC for the ORACLES-2016 flights (Fig. 9a and), with the SSA values~~
330 ~~decreasing more systematically with FrBC for all four flights (Fig. 9b)~~). The BC-containing particle fraction varies from 0.2 to 0.4, more than the 0.1-0.2 range shown for July south of remote western Africa (Denjean et al., 2020b), and less than the 0.3-0.45 range at Ascension (Taylor et al., 2020). The total particle number was drawn from the full aerosol size distribution within Denjean et al. (2020b), and by a PCASP (0.1-3 μm) at Ascension. These size ranges are comparable enough to support the comparison across the three campaigns.

335 Independent electron microscopy on 2017 filter samples found that almost all BC is at least partially coated, meaning the BC particles are dominated by internal mixing (Dang et al., 2022). Nevertheless, the majority of particles cannot include BC, since $FrBC < 0.5$. As the fraction of BC-containing particles increase, the bulk OA:BC mass ratio tends to decrease. $MAC_{BC,660}$ ranges from 9-12 $\text{m}^2 \text{g}^{-1}$, and $MAC_{BC+OA,470}$ from 13-18 $\text{m}^2 \text{g}^{-1}$, corresponding to absorption enhancement factors of 1.2-1.6 (1.7-2.4) at the 660 nm (470 nm) wavelengths, assuming an MAC of 7.5 $\text{m}^2 \text{g}^{-1}$ for uncoated black carbon (Bond and Bergstrom, 2006) (and greatly exceeding the MAC value of 6.25 $\text{m}^2 \text{g}^{-1}$ for strongly light-absorbing carbon (Bond and Bergstrom, 2006)). The mean $MAC_{BC,660}$ of 10.8 $\text{m}^2 \text{g}^{-1}$ is slightly higher than the median value of 9.3 $\text{m}^2 \text{g}^{-1}$ reported in
340 Carter et al. (2021), likely because the BC-enriched 31 August 2017 flight contributes strongly to the mean value reported here. Median LDMA-inferred particle diameters range from 120-210 nm, with no clear relationship to $MAC_{BC+OA,470}$. This indicates the absorption enhancements are governed more by composition than particle size, similar to Denjean et al. (2020a) for
345 June-July BBA close to the near-equatorial African coast. The 8/31/2017 flight, for an FrBC of 0.3, has a higher $MAC_{BC,660nm}$ (by $\sim 2 \text{m}^2 \text{g}^{-1}$), lower OA:BC mass ratio, larger ~~particle/BC core sizes, and more coating (crudely estimated as the LDMA-BC core diameter difference divided by two, primarily intended as a relative measure),~~ LDMA-inferred particle and mass-median
BC core size, compared to values from the 9/6/2016 flight of comparable FrBC. The larger BC core size for 8/31/2017 may come from a woodier fuel, supported by backtrajectories emanating from further north (not shown). Woodier material ~~can has~~
350 been shown to generate larger BC sizes irrespective of MCE (Holder et al., 2016). ~~We do not know how to reconcile a lower OA:BC mass ratio with a thicker coating for the same FrBC, however. within Holder et al. (2016).~~

The single scattering albedos (SSA) at $\lambda=530$ nm range from 0.83 to 0.89, consistent with the ORACLES-2016 mean SSA of 0.86 (inter-quartile range of ~ 0.028) based on all the flight data (Pistone et al., 2019). These SSA values are lower than previously documented *in situ* values over land or coastal (Haywood et al., 2003; Formenti et al., 2003; Dubovik et al., 2002), on par with AERONET September-mean values at Mognu (Eck et al., 2013), and higher than those reported at Ascension Island (Zuidema et al., 2018; Wu et al., 2020). An SSA best-fit regression on OA:BC provides a straightforward connection between the BBA chemical and optical properties: $SSA_{530}=0.801+0.0055*(OA:BC)$ (Fig. 10a, correlation coefficient r of 0.84). The dependence on BC:TC (total carbon) following Brown et al. (2021) is: $SSA_{530}=0.929-0.389*(BC:TC)$ (Fig. 10b; $r=-0.79$) ~~using the same calculation for OC (Aiken et al., 2007)~~; where $TC=BC+\text{organic carbon (OC)}$ and OC is estimated from $OA:OC=1.26*O:C+1.18$ (Aiken et al., 2008). The dependence on BC:TC is not as pronounced as in Brown et al. (2021) primarily because our dataset has a smaller SSA range, with no SSAs > 0.9 . The variance in SSA is explained slightly better by OA:BC than BC:TC.

4.2 Is there evidence of brown carbon?

Taylor et al. (2020) place an upper estimate of 11% on shortwave absorption by brown carbon (BrC) at 405 nm wavelength by the time the BBA plume reaches Ascension Island. Zhang et al. (2022) indicate that other non-BrC materials such as iron oxides absorb sunlight over the southeast Atlantic, so that BrC may contribute even less than $< 10\%$ of the total absorption at sub-500 nm wavelengths. By four days since emission, the primary organic aerosol has mostly converted to secondary OA (SOA), which typically absorbs little light (Bond and Bergstrom, 2006; Laskin et al., 2015). Nevertheless, if oxidation can continue to produce new chromophores (O'Brien and Kroll, 2019) that absorb differently based on wavelength, that could be interpreted as SOA-induced BrC. Ozone Monitoring Instrument UltraViolet Aerosol Index values do suggest OA-produced brown carbon should be present east of the prime meridian (Carter et al., 2021), however. Laboratory studies find more BrC absorption for lower OA:BC mass ratios (Saleh et al., 2014; Holder et al., 2016; McClure et al., 2020), because more intense fires also produce more primary OA and BrC. One important difference is that the reported primary OA fraction and AAEs are much higher within Saleh et al. (2014) than we would expect over the southeast Atlantic.

Motivated by Carter et al. (2021) we examine if distance from the continent has a detectable influence on the absorption Ångström exponents (AAE) calculated over the 470-660 nm wavelength range for the level leg data (Fig. 11a), recognizing that the 470 nm wavelength may already be too long to be responsive to additional absorption by OA-produced BrC (Zhang et al., 2022). The AAEs span 1.1-1.3 south of 8°S irrespective of distance from the coast, and are close to one further north for the more remote 8/31/2017 flight (Fig. 11a). Such AAE values typically indicate a lack of BrC (e.g., Forrister et al., 2015). AAE is weakly positively correlated with OA:BC ($r = 0.27$; not shown), but the relationship is statistically insignificant. Brown carbon absorption is also assessed using $MAC_{BC+OA,470}$ following Carter et al. (2021) (Fig. 11b). These range from 0.94-1.2 $\text{m}^2 \text{g}^{-1}$ south of 8°S (with one exception) to 1.4-1.7 $\text{m}^2 \text{g}^{-1}$ further north. $MAC_{BC+OA,470}$ is anticorrelated with OA:BC ($r = -0.86$). Although consistent with Saleh et al. (2014), the small sample size, dominated by one flight north of 8°S with less OA:BC, precludes much interpretation. We primarily conclude a lack of a longitudinal dependence, although the sample size is too small to say this with confidence. Other work has found that co-emitted sulfate can contribute to increasing overall

absorption (Christian et al., 2003), but we do not find a correlation between either MAC estimate and the sulfate fraction (not shown).

5 Is there evidence for ongoing loss of organic aerosol? 24 September 2016 case study

The data for the youngest aerosol, aged 4-5 days since emission, stems from the 9/24/2016 flight (Fig. 2). Since this aerosol
390 may be more susceptible to aging we examine its features more closely. Backtrajectories from the profile at 12.3°S, 11°E show
the aerosols are coming from similar source regions (Fig. 12d), and become distributed vertically primarily by variations in
the advection speed. One main aerosol layer is centered on 5 km, aged ~4 days since emission, and a slightly older smoke
layer of ~5 days in age is centered on 3 km (Fig. 12b,c). The younger aerosol aloft is connected to stronger upper-altitude
winds also transporting moisture (Fig. 12a), consistent with climatological expectations (Adebisi et al., 2015; Adebisi and
395 Zuidema, 2016; Pistone et al., 2021). These generate relative humidities exceeding 80% above 4 km when combined with the
cooler high-altitude temperatures (Fig. 12a). Although there are two main aerosol plumes, the potential temperature profile is
of a thermally stratified atmosphere containing many thinner seemingly well-mixed layer separated by discrete stability jumps
(Fig. 12c). The water vapor mixing ratio profile (Fig. 12c) indicates there is only one truly well-mixed layer, capping the upper
aerosol plume between 5.3-5.8 km with slackening winds. The lack of vertical mixing indicates the smoke plume heights are
400 likely set above land. The upper-level aerosol plume registered both the highest OA:BC mass ratio and the highest SSA of the
ORACLES-2016 campaign. More intense fires, with lower OA:BC are typically able to reach higher altitudes (Martin et al.,
2010), but the higher OA:BC and NO₃:BC mass ratios aloft (Fig. 12b) may instead indicate more condensation of the emitted
vapors aloft, aided by cooler temperatures and higher relative humidities (Li et al., 2018).

Secondary aerosol formation is expected to proceed more quickly when $\frac{BC}{\Delta CO}$ ratios are lower (Vakkari et al., 2018), because
405 the precursor gases needed for nucleation may be more available (Yokelson et al., 2009). We first confirm that the flight's $\frac{BC}{\Delta CO}$
values remain statistically-similar as a function of f_{44} : these remain within $7.5-7.9 \times 10^{-3}$ independent of f_{44} (Fig. 13a; see
Figs. [S9-S10](#) [S10-S11](#) for the same analysis for the other flights). We interpret this to mean that the aerosol are emitted from
similar sources over a two-day time span, with no wet deposition throughout. The corresponding OA:BC mass ratio decreases
from 14.2 to 9.8 (Fig. 13b) - an approximate 35% decline in OA:BC over a span of 1-2 days. The corresponding SSA reduces
410 from 0.89 to 0.865. The mean $AAE_{470-660}$ decreases from 1.25 to 1.21 ($\pm 0.07-0.08$) as a function of the three f_{44} bins, a
statistically insignificant decline.

An evaluation of the changes to the particle size distribution with f_{44} , normalized with respect to BC as a control for dilution,
indicates the processes of condensation and coagulation - and volume loss consistent with the mass loss. As the aerosol ages
chemically, the LDMA median diameter increases from ~ 170 nm to ~ 205 nm (Fig. 14a), mostly because the number of
415 particles with diameters < 100 nm declines. BC particles are typically larger than OA particles (e.g., Fig. [S5S6](#)), and the
reduction in the number of small particles indicates coagulation of the OA particles upon the larger BC particles. It also
indicates that most of the vapors are condensing on the larger BC particles, as opposed to forming new particles by nucleation.
The total LDMA and CPC particle number concentrations reduce from approximately 1200 cm^{-3} to 500 cm^{-3} with respect

to BC, and 2400 cm^{-3} to 1500 cm^{-3} , respectively (Fig. 14c). The large difference in the two number concentrations likely
420 reflects an instrument difference; both instruments agree there is no net production of the smaller particles. The combined effect
of condensation and coagulation results in an increase in the fraction of BC-containing particles from 0.18, to 0.23 and then
0.27 as f_{44} increases. The evolution towards larger sizes would increase the SSA, all else equal. Instead, the SSA decreases in
response to the decrease in OA, indicating again that changes in particle size do not dominate the SSA changes.

At the same time, the LDMA-determined particle volume decreases (Fig. 14b), indicating genuine particle mass loss that is
425 consistent with the decrease in OA:BC. One mechanism for the mass loss could be evaporation through dilution. The selection
for data samples with $\text{OA} > 20 \mu\text{g m}^{-3}$ focuses the analysis on the aerosol plume center, and a normalization by BC provides
an additional control, leading us to discount this mechanism. Aqueous phase reactions and mid-level cloud processing could
potentially also contribute to the oxidation increase and loss of free-tropospheric OA mass concentration. This is partially
controlled for by only selecting free-tropospheric data samples with $\text{RH} < 80\%$. Mid-level clouds, produced by dry convection
430 saturating the top of the land boundary layer, can occur (Adebiyi et al., 2020), but are not a dominant presence on this day or
other ORACLES flight days. This suggests to us that the reduction of free-tropospheric OA through aqueous phase reactions
is of secondary importance (becoming even more so with distance from the continent).

Instead, we speculate the dominant loss mechanism can be increasing oxidation through fragmentation, which can release
higher-volatility particles that can then be subsequently removed (e.g., Jimenez et al., 2009). Figs. 14c-d support that interpre-
435 tation: both the LDMA and CPC total particle number concentrations decrease with f_{44} , consistent with processes occurring
at the surface of the larger particles - either coagulation or surface reactions. The oxidative environment, inferred from $\frac{O_3}{\Delta CO}$,
remains constant with f_{44} , but these measures for oxidation may have reached their upper limit. The reduction in the total
non-BC aerosol mass, which reflects a reduction in the combined $\text{OA} + \text{NH}_4 + \text{NO}_3$ mass but not SO_4 is nevertheless in concert
with the OA:BC decrease. The constancy of SO_4 :BC with f_{44} (not shown) confirms the aerosol is aged, as the lifetime of SO_2
440 is 1-2 days, after which its conversion to SO_4 will have ended. In summary we interpret Fig. 14 to reflect changes in the particle
size distribution induced by condensation, coagulation and mass loss through fragmentation .

The backtrajectories do not show lower-level westerlies, in contrast to the *in-situ* profiles. We conjecture that the daytime
aircraft sampling sampled a land breeze below 4 km that is converging above a warming continental surface and is not rep-
resented in the GFS meteorology. The ERA5 dataset, which has an hourly temporal resolution, might be able to address this
445 hypothesis. Not shown, the marine boundary layer top is at one km, and the boundary layer did not include any BC, consistent
with a slow entrainment time scale for aerosols (Diamond et al., 2018).

~~A third~~

6 Does comparing to aerosol measured at Ascension Island indicate ongoing compositional changes?

A third possible mechanism for the loss of the overall particle mass ~~can~~could be through thermodynamics, consistent with the
450 increase in NO_3 :BC with altitude (Fig. 12b). ~~This mechanism is assessed in more detail within the next Section 6.-~~

7 Does comparing to aerosol measured at Ascension Island indicate ongoing compositional changes?

A comparison to the aerosol properties measured at the more remote location of Ascension island by CLARIFY (Table 3) supports the speculation that fragmentation of oxidized aerosol may be contributing to mass loss, by ultimately releasing some aerosol that can evaporate through photochemistry, similar to the younger aerosol sampled on 9/24/2016. Table 3 compares values for the aerosol parameters derived from the six ORACLES flights to the free-tropospheric values reported within Wu et al. (2020) and Taylor et al. (2020). The ORACLES 8/31/2017 flight coincided with CLARIFY and occurred halfway to the island (Fig. 1). Important to this comparison, the ORACLES and CLARIFY $\frac{OA}{\Delta CO}$, BC and SSA values compared well on the intercomparison flight held on 18 August 2017 (Barrett et al., 2022). Their similar $\frac{BC}{\Delta CO}$ ratios (Table 2) also equal the maximum values inferred from the surface-based measurements at Ascension between June to October (Che et al., 2022).

Overall, CLARIFY sampled more BC-enriched particles in the free troposphere above Ascension in both number and mass, with slightly lower SSA, higher MAC_{660s} , and larger particle sizes. The OA:BC mass ratios are lower, mostly because the OA masses themselves are much lower, typically $< 4 \mu\text{g m}^{-3}$ (Wu et al., 2020). Interestingly, September African BBA sampled near the Amazon Tall Tower Observatory indicated BC mass and number fractions on par with CLARIFY, with OA:BC mass ratios between 2.5-5.0, although the sampling is only from one aircraft flight (Holanda et al., 2020). The H:C, O:C and OA:OC CLARIFY values (1.2-1.4, 0.7-0.8, 2.3) indicate slightly more oxidized aerosol at Ascension (Wu et al., 2020) than for ORACLES, with the exception of 31 August 2017 (Fig. 8). ORACLES 8/31/2017 flight values tend to be intermediate to the CLARIFY and ORACLES-2016 numbers.

Other prior field campaigns have highlighted a small net OA loss as BBA ages beyond a day (e.g., Capes et al., 2008; Jolleys et al., 2012, 2015; Hodzic et al., 2015; Konovalov et al., 2019; Farley et al., 2022), attributed to evaporation through dilution in Jolleys et al. (2012). The extent of the OA reduction reported here - a factor of at least two between ORACLES and CLARIFY OA masses - suggests a different process must be dominant. Nevertheless, we examine if dilution could be factor, through comparing the number and volume size distributions measured by the TDMA during the 8/31/2017 flight at STP and at 150°C (Fig. S6S7). The heating is used as a proxy for dilution because processes respond to aerosol volatility. Although the TDMA size distribution does not extend beyond 0.2 μm at most, missing the bulk of the aerosol mass, the size distributions change little between the temperatures, supporting the inference that evaporation through dilution does not explain differences in OA:BC between the two campaigns.

Another significant difference between ORACLES-2016 and CLARIFY is the inorganic nitrate (IN) fraction. Nitrate only contributes 10% to the total aerosol mass analyzed here, and only 8% of the total free-tropospheric aerosol mass during CLARIFY (Wu et al., 2020). The fraction that is inorganic nitrate is even smaller. Interestingly, the 8/31/2017 inorganic nitrate fraction was 51% - intermediate to the ORACLES-2016 and CLARIFY values (Table 3). This suggests that organic nitrate may become converted to inorganic nitrate as the aerosol ages. The air sampled during ORACLES was mildly acidic (Fig. S9S10), based on a simplification of the $NH_{4,measured}/NH_{4,predicted}$ relationship put forth in Zhang et al. (2007). A mild acidity will slow the rate of inorganic acid formation, and may help explain the lower IN fraction for ORACLES (20%-25%). Inorganic ammonium nitrate is held responsible for an increase in SSA with height at Ascension (Wu et al., 2020), as thermodynamic

485 partitioning favors the particle phase at higher altitudes. The nitrate fraction is never large, and the coating on the BC will be dominated by OA by mass, so that the IN fraction may be more valuable as an indicator of ongoing oxidation that is also capable of increasing the MAC (Shrivastava et al., 2017). The sulfate fraction is similar between the two campaigns, including the 8/31/2017 flight, and seems unlikely to explain the differences in the MAC between the campaigns. Increasing coagulation of smaller particles upon the BC particles could help explain why the particle diameters are larger at Ascension compared to
490 ORACLES-2016, at the same time that evaporation through photochemistry increases the fraction of BC-containing particles while decreasing the overall OA:BC mass ratio. It is also possible that the non-Lagrangian sampling is introducing a bias. A Lagrangian analysis of filter samples did find increased aerosol volatility and continued OA loss in ORACLES-CLARIFY resampled aerosol (Dang et al., 2022).

We also examine if a portion of the OA can thermodynamically repartition. We composite OA:BC, NO₃:BC and aerosol
495 age by the free-tropospheric RH for all six flights to illuminate how much thermodynamical partitioning by altitude may be occurring. The higher altitudes in the free troposphere are also often more humid (e.g., Fig 12, shown more statistically in Adebisi et al. (2015) and Pistone et al. (2021)). Physically-younger aerosol is more likely to occupy a more (relatively) humid, colder free troposphere at higher altitudes than is older aerosol (e.g., Fig. 12). The mean NO₃:BC ratio decreases by almost 50% as the free troposphere RH decreases from 70% to 30% (Fig. 15a), consistent with a thermodynamic repartitioning to
500 the gas phase. The mean OA:BC mass ratio only reduces from 10.5 ± 2.6 for RH = 60-80% to 9.9 ± 2.1 for 20% < RH < 40%, a statistically insignificant decrease. A thermodynamical repartition can only explain a relative change in OA:BC of less than 10%, if that. The small change in OA:BC, if real, may also reflect moisture contributing to the OA mass loss through increasing OH uptake and/or fragmentation (Li et al., 2018), or because at higher altitudes, less-oxidized material is continuing to condense upon the pre-existing organics, ultimately favoring the evaporation of OA into the gas phase.

505 7 Discussion

This study extends and strengthens an earlier analysis begun within Dobracki et al. (2022). One flight, on 9/24/2016, has consistent BC:CO ratios as a function of the *f44* chemical aging marker, interpreted to mean emissions from similar sources over a few days with no wet deposition. At the lower altitudes with weaker offshore winds, condensation and coagulation explain an increased particle size for the slightly older aerosol. Increasing oxidation through fragmentation, which can then
510 release higher-volatility particles through photochemistry, can explain the reduced overall mass. Dilution is not considered as influential as for northern hemisphere boreal fires, because the smoke distribution is so broad and loadings so large, composed of many small continental small fires that become homogenized before the smoke is advected offshore. An increase in SSA with height is primarily explained by more OA aloft. This contrasts with the increase in SSA with height at Ascension attributed to an increase in ammonium nitrate (Wu et al., 2020), which may reflect changes in the aerosol composition occurring with
515 further transport offshore. Level-leg measurements from six flights demonstrate how optical properties relate to chemical and physical composition and can be compared to values from Wu et al. (2020) and Taylor et al. (2020) made at Ascension

Island. Further focus on data from the 8/31/2017 flight helps connect interpretations of ORACLES versus CLARIFY aerosol characteristics.

520 The total aerosol concentration exceeds the total SP2-derived BC number by a factor of 2.5 to 7, from which we infer that at least one-half of the non-BC aerosol remains externally mixed with the BC. The BC itself, because of its transport within multiple days within broad, dense smoke plumes, is most likely internally mixed with other aerosols, confirmed by independent electron microscopy measurements (Dang et al., 2022). Because the BBA is at least 4 days old, and as already shown within Taylor et al. (2020) and Denjean et al. (2020b), the BC can be treated as compacted. Taylor et al. (2020) find a better fit to the CLARIFY MAC measurements using the semi-empirical models of Liu et al. (2017) and Chakrabarty and Heinson (2018) 525 than to a core-shell Mie model, but Lee et al. (2022) conclude a core-shell mode can be successfully applied once particle-by-particle differences are accounted for. The 2016 data from ORACLES lacks SP2 mixing state data with which to better evaluate optical fits, but an independent assessment could be pursued using the SP2 coating-resolved ORACLES data from 2017 and 2018 (Sedlacek et al., 2022).

A highlight of this study is its SSA_{530} best-fit regression on OA:BC: $SSA_{530}=0.801+0.0055*(OA:BC)$. The range of OA:BC 530 of 7 through 14 equates to an SSA variability of 0.83 to 0.89. This provides a straightforward connection between the BBA chemical and optical properties that is useful for the modeling of the direct aerosol radiative effect. Of course, use of such a best-fit is only effective if the model OA:BC mass ratios are realistic. Given that OA:BC mass ratios are often too low in models, their absorption of sunlight will also be overestimated (Brown et al., 2021). This study adds to literature indicating that OA model estimates made by multiplying the organic carbon by a factor of 1.4 will underestimate OA in this (and other) 535 regions (Aiken et al., 2008; Tsigaridis et al., 2014; Shinzuka et al., 2020; Doherty et al., 2022). This study's OA:OC mass ratios of 2.2 ± 0.1 is also shown for the Atomic Tomography mission (Hodzic et al., 2020).

More sophisticated aerosol schemes can, in contrast, overestimate OA:BC mass ratios over the southeast Atlantic (Chylek et al., 2019), suggesting the loss of OA with aging or slower SOA production processes (Kroll and Seinfeld, 2008; McFiggans et al., 2019) may also be under-accounted for. For the southeast Atlantic region, far removed from urban and industrial sources 540 of pollution, continued production of SOA after 1-2 days is expected to remain minor (e.g., O'Brien and Kroll, 2019). This contrasts with northern hemisphere fire emission sources. Brown carbon production has been linked to low OA:BC ratios (Saleh et al., 2014; McClure et al., 2020), but this does not seem supported by the admittedly-limited ORACLES measurements of AAE and MAC, perhaps because brown carbon is typically more closely linked to primary than to secondary organic aerosol. Many prior studies find continuing oxidation of OA (see review by Shrivastava et al., 2017). This will be more important for 545 remote environments containing thick smoke layers lacking additional pollution sources producing the precursor gases for additional SOA. Further work is needed to better support the process conclusion of this study, named that evaporation through fragmentation/photochemistry is the dominant chemical aging process over the southeast Atlantic, but nevertheless this study indicates the importance of developing sophisticated SOA schemes (e.g., Lou et al., 2020) for this regional climate.

September is unique in that meteorology and fire processes conspire to accentuate the direct radiative warming of the south- 550 east Atlantic. August is likely the month with the most fires in southern Africa (Scholes et al., 1996), but the upper-level winds that transport the aerosol depend on a strong heat low over southern Africa, and don't become well-established until September

(Adebisi and Zuidema, 2016; Kuete et al., 2020). The winds occur to the north of the heat low, and only dry convection lofts the aerosols to their altitude. The winds distribute aerosol as far away as south America (Holanda et al., 2020), so that the entire south Atlantic is covered by a blanket of highly-absorptive aerosol, with submicron BC mass fractions far exceeding the 2-10% estimated for western north America (Garofalo et al., 2019). The strong September upper-level winds also discourage subsidence (Chaboureau et al., 2022), and the cloud cover and height are also affected directly by meteorology during this month (Adebisi and Zuidema, 2018; Zhang and Zuidema, 2021). Less of the aerosol reaches the cloud top, reducing aerosol entrainment into the cloud layer (Zuidema et al., 2018; Shinozuka et al., 2020; Doherty et al., 2022) and ability to influence the cloud top inversion strength (Herbert et al., 2020). The net radiative impact will primarily be the direct aerosol radiative effect of the aerosol aloft then, lending further weight on model representation of SSA (Mallet et al., 2021). The remaining ORACLES data from 2016 and 2017, for which SP2 mixing state data are available, will be used to support further investigation of SSA-aerosol composition relationships in a follow-up study.

Data availability. The data are available through doi=10.5067/Suborbital/ORACLES/P3/2016_V2 and doi=10.5067/Suborbital/ORACLES/P3/2017_V2.

Author contributions. The present work was conceived by PZ, AD, PS and SH. SF contributed to the HiGEAR data analysis, AS provided the BC datasets and PS the WRF-AAM model age estimates. Portions of this work first appeared in the M.S. thesis of A.D at U. of Hawaii. PZ led the writing with AD providing most of the figures. All authors contributed to the final writing.

Competing interests. PZ is a guest editor for the ACP Special Issue: “ACP special issue: New observations and related modelling studies of the aerosol–cloud–climate system in the Southeast Atlantic and southern Africa regions”. The other authors declare no competing interests.

Acknowledgements. ORACLES is a NASA Earth Venture Suborbital-2 investigation, funded by the US National Aeronautics and Space Administration (NASA)’s Earth Sciences Division and managed through the Earth System Science Pathfinder Program Office (grant no. NNH13ZDA001N-EVS2). This work was further supported by the US Department of Energy grant DE-SC0018272 to PZ and PS and DE-SC0021250 to PZ. We thank Hugh Coe, Huihui Wu and Jonathan Taylor for interesting initial conversations that in particular encouraged us to examine the inorganic nitrate fraction.

References

- 575 Adebisi, A. A. and Zuidema, P.: The role of the southern African easterly jet in modifying the southeast Atlantic aerosol and cloud environments, *Q. J. Roy. Meteor. Soc.*, 142, 1574–1589, <https://doi.org/10.1002/qj.2765>, 2016.
- Adebisi, A. A. and Zuidema, P.: Low cloud sensitivity to biomass-burning aerosols and meteorology over the southeast Atlantic, *J. Climate*, 31, 4329–4346, <https://doi.org/10.1175/JCLI-D-17-0406.1>, 2018.
- Adebisi, A. A., Zuidema, P., and Abel, S. J.: The Convolution of Dynamics and Moisture with the Presence of Shortwave Absorbing Aerosols over the Southeast Atlantic, *J. Climate*, 28, 1997–2024, <https://doi.org/10.1175/JCLI-D-14-00352.1>, 2015.
- Adebisi, A. A., Zuidema, P., Chang, I., Burton, S. P., and Cairns, B.: Mid-level clouds are frequent above the southeast Atlantic stratocumulus clouds, *Atmos. Chem. Phys.*, pp. 11 025–11 043, <https://doi.org/10.5194/acp-20-11025-2020>, 2020.
- Aiken, A. C., DeCarlo, P. F., and Jimenez, J. L.: Elemental analysis of organic species with electron ionization high-resolution mass spectrometry, *Anal. Chem.*, 79, 8350–8358, 2007.
- 585 Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D., Huffman, J. A., Docherty, K. S., Ulbrich, I., and et al., C. M.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ Sci Technol.*, 42, 4478–4485, 2008.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-11-4039-2011)
- 590 11-4039-2011, 2011.
- Anderson, T. L. and Ogren, J. A.: Determining Aerosol Radiative Properties Using the TSI 3563 Integrating Nephelometer, *Aer. Sci. Tech.*, 29, 57–69, <https://doi.org/10.1080/02786829808965551>, 1998.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- 595 Baars, H., Radenz, M., Floutsi, A., Engelmann, R., Althausen, D., and et al., B. H.: Californian wildfire smoke over Europe: A first example of the aerosol observing capabilities of Aeolus compared to ground-based lidar, *Geophys. Res. Lett.*, 48, <https://doi.org/10.1029/2020GL092194>, e2020GL092194, 2021.
- Bahreini, R., Ervens, B., Middlebrook, A. M., Warneke, C., de Gouw, J. A., DeCarlo, P. F., and et al., J. L. J.: Organic aerosol formation in urban and industrial plumes near Houston and Dallas, Texas, *J. Geophys. Res.*, 114, <https://doi.org/10.1029/2008jd011493>, 2009.
- 600 Barrett, P. A., Abel, S. J., Coe, H., Crawford, I., Dobracki, A., Haywood, J. M., Howell, S., Jones, A., Langridge, J., McFarquhar, G., Nott, G., Price, H., Redemann, J., Shinzuka, Y., Szpek, K., Taylor, J., Wood, R., Wu, H., Zuidema, P., Bauguitte, S., Bennett, R., Bower, K., Chen, H., Cochrane, S. P., Cotterell, M., Davies, N., Delene, D., Flynn, C., Freedman, A., Freitag, S., Gupta, S., Noone, D., Onasch, T. B., Podolske, J., Poellot, M. R., Schmidt, S. K., Springston, S., III, A. J. S., Trembath, J., Vance, A., Zawadowicz, M., and Zhang, J.: Intercomparison of airborne and surface-based measurements during the CLARIFY, ORACLES and LASIC field experiments, *Atmos.*
- 605 *Meas. Tech.*, 15, 6329–6371, <https://doi.org/10.5194/amt-15-6329-2022>, 2022.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, *Aer. Sci. Tech.*, 40, 27–67, <https://doi.org/10.1080/02786820500421521>, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., and et al., Y. K.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.*, 118,
- 610 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.

- Bowman, D. M., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M., and et al., R. S. D.: Fire in the Earth System, *Science*, 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.
- Brown, H., Liu, X., Pokhrel, R., Murphy, S., Lu, Z., Saleh, R., Mielonen, T., Kokkola, H., Bergman, T., Myhre, G., Skeie, R. B., Watson-Paris, D., Stier, P., Johnson, B., Bellouin, N., Schulz, M., Vakkari, V., Beukes, J. P., vanZyl, P. G., Liu, S., and Chand, D.: Biomass burning aerosols in most climate models are too absorbing, *Nat. Comm.*, 12, 277, <https://doi.org/10.1038/s41467-020-20482-9>, 2021.
- 615 Canagaratna, M. R., Jimenez, J. L., and et al., J. H. K.: Elemental ratio measurements of organic compounds using aerosol mass spectrometry: characterization, improved calibration, and implications, *Atmos. Chem. Phys.*, 15, 253–272, <https://doi.org/10.5194/acp-15-253-2015>, 2015.
- Capes, G., B.Johnson, McFiggans, G., Williams, P. I., Haywood, J., and Coe, H.: Aging of biomass burning aerosols over West Africa: Aircraft measurements of chemical composition, microphysical properties, and emission ratios, *J. Geophys. Res.*, 113, <https://doi.org/10.1029/2008jd009845>, 2008.
- 620 Carter, T. S., Heald, C. L., Cappa, C. D., Kroll, J. H., Campos, T. L., Coe, H., Cotterel, M. I., Davies, N. W., Farmer, D. K., Fox, C., Garofalo, L. A., Hu, L., Langridge, J. M., Levin, E. J. T., Murphy, S. M., Pokhrel, R. P., Shen, Y., Szpek, K., Taylor, J. W., and Wu, H.: Investigating Carbonaceous Aerosol and its Absorption Properties from Fires in the western US (WE-CAN) and southern Africa (ORACLES and CLARIFY), *J. Geophys. Res.*, 126, e2021JD034984, 2021.
- 625 Chaboureaud, J.-P., Labbouz, L., Flamant, C., and Hodzic, A.: Acceleration of the southern African easterly jet driven by radiative effect of biomass burning aerosols and its impact on transport during AEROCLO-sA, *Atmos. Chem. Phys.*, 22, 8639–8658, <https://doi.org/10.5194/acp-22-8639-2022>, 2022.
- Chakrabarty, R. K. and Heinson, W. R.: Scaling Laws for Light Absorption Enhancement Due to Nonrefractory Coating of Atmospheric Black Carbon Aerosol, *Phys. Rev. Lett.*, 121, <https://doi.org/10.1103/PhysRevLett.121.218701>, 2018.
- 630 Che, H., Segal-Rozenhaimer, M., Zhang, L., Dang, C., Zuidema, P., III, A. J. S., Zhang, X., and Flynn, C.: Seasonal variations in fire conditions are important drivers to the trend of aerosol optical properties over the south-eastern Atlantic, *Atmos. Chem. Phys.*, 22, 8767–8785, <https://doi.org/10.5194/acp-22-8767-2022>, 2022.
- Christian, T. J., Kleiss, B., Yokelson, R. J., Holzinger, R., Crutzen, P. J., Hao, W. M., Saharjo, B. H., and Ward, D. E.: Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from Indonesian, African, and other fuels, *J. Geophys. Res.*, 108, 4719, <https://doi.org/10.1029/2003JD003704>, 2003.
- 635 Chylek, P., Lee, J. E., Romonosky, D. E., Gallo, F., Lou, S., and et al., M. S.: Mie Scattering Captures Observed Optical Properties of Ambient Biomass Burning Plumes Assuming Uniform Black, Brown, and Organic Carbon Mixtures, *J. Geophys. Res.*, 124, 11406–11427, <https://doi.org/10.1029/2019jd031224>, 2019.
- 640 Collier, S., Zhou, S., Onasch, T. B., Jaffe, D. A., Kleinman, L., A. J. Sedlacek, I., Briggs, N. L., Hee, J., Fortner, E., and et al., J. E. S.: Regional Influence of Aerosol Emissions from Wildfires Driven by Combustion Efficiency: Insights from the BBOP Campaign, *Environ Sci Technol.*, 50, 8613–8622, <https://doi.org/10.1021/acs.est.6b01617>, 2016.
- Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., and et al., D.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049–12064, <https://doi.org/10.5194/acp-11-12049-2011>, 2011.
- 645 Dang, C., Segal-Rozenhaimer, M., Che, H., Zhang, L., Formenti, P., Taylor, J., Dobracki, A., Purdue, S., Wong, P.-S., Nenes, A., Sedlacek, A., Coe, H., Redemann, J., Zuidema, P., and Haywood, J.: Biomass burning and marine aerosol processing over the southeast Atlantic Ocean: A TEM single particle analysis, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-2021-724>, 2022.

- Darmenov, A. and da Silva, A.: The Quick Fire Emissions Dataset (QFED) - Documentation of Versions 2.1, 2.2, and 2.4, 32, 2013.
- 650 Davies, N. W., Fox, C., Szpek, K., Cotterell, M. I., Taylor, J. W., Allan, J. D., Williams, P. I., Trembath, J., Haywood, J. M., and Langridge,
J. M.: Evaluating biases in filter-based aerosol absorption measurements using photoacoustic spectroscopy, *Atmos. Meas. Tech.*, 12,
3417–3434, <https://doi.org/10.5194/amt-12-3417-2019>, 2019.
- de Graaf, M., Schulte, R., Peers, F., Waquet, F., Tilstra, L. G., and Stammes, P.: Comparison of south-east Atlantic aerosol direct radiative
effect over clouds from SCIAMACHY, POLDER and OMI-MODIS, *Atmos. Chem. Phys.*, 20, 6707–6723, [https://doi.org/10.5194/acp-
20-6707-2020](https://doi.org/10.5194/acp-
655 20-6707-2020), 2020.
- Denjean, C., Bourriane, T., Burnet, F., Mallet, M., Maury, N., Colomb, A., Dominutti, P., Brito, J., Dupuy, R., Sellegri, K., Schwarzen-
boeck, A., Flamant, C., and Knippertz, P.: Overview of aerosol optical properties over southern West Africa from DACCIWA aircraft
measurements, *Atmos. Chem. Phys.*, 20, 4735–4756, <https://doi.org/10.5194/acp-20-4735-2020>, 2020a.
- Denjean, C., Brito, J., Libois, Q., Mallet, M., Bourriane, T., and et al., F. B.: Unexpected Biomass Burning Aerosol Absorption Enhancement
660 Explained by Black Carbon Mixing State, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2020gl089055>, 2020b.
- Diamond, M. S., Dobracki, A., Freitag, S., Small Griswold, J. D., Heikkila, A., Howell, S. G., Kacarab, M. E., Podolske, J. R., Saide, P. E.,
and Wood, R.: Time-dependent entrainment of smoke presents an observational challenge for assessing aerosol–cloud interactions over
the southeast Atlantic Ocean, *Atmos. Chem. Phys.*, 18, 14623–14636, <https://doi.org/10.5194/acp-18-14623-2018>, 2018.
- Dobracki, A., Zuidema, P., Howell, S., Saide, P., Freitag, S., Aiken, A. C., Burton, S. P., III, A. J. S., Redemann, J., and Wood, R.: Non-
665 reversible aging can increase solar absorption in African biomass burning aerosol plumes of intermediate age, *Atmos. Chem. Phys.*
Discuss., <https://doi.org/10.5194/acp-2021-1081>, preprint, 2022.
- Doherty, S. J., Saide, P., Zuidema, P., Shinozuka, Y., Ferrada, G., and et al., H. G.: Modeled and observed properties related to the direct
aerosol radiative effect of biomass burning aerosol over the Southeast Atlantic, *Atmos. Chem. Phys.*, 22, 1,46, [https://doi.org/10.5194/acp-
22-1-2022](https://doi.org/10.5194/acp-
22-1-2022), 2022.
- 670 Dubovik, O., Holben, B. N., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanre, D., and Slutsker, I.: Variability of absorption and
optical properties of key aerosol types observed in worldwide locations, *J. Atmos. Sci.*, 59, 590–608, 2002.
- Eck, T. F., Holben, B. N., Reid, J. S., Mukelabai, M. M., Piketh, S. J., and et al., O. T.: A seasonal trend of single scattering albedo in southern
African biomass-burning particles: Implications for satellite products and estimates of emissions for the world’s largest biomass-burning
source, *J. Geophys. Res.*, 118, <https://doi.org/10.1002/jgrd.50500>, 2013.
- 675 Farley, R., Bernays, N., Jaffe, D. A., Ketcherside, D., Hu, L., Zhou, S., Collier, S., and Zhang, Q.: Persistent influence of wildfire emissions
in the western United States and characteristics of aged biomass burning organic aerosols under clean air conditios, *Env. Sci. Tech.*, 56,
3645–3657, <https://doi.org/10.1021/acs.est.1c07301>, 2022.
- Farmer, D., Matsunaga, A., Docherty, K. S., Surratt, J. D., Seinfeld, J. H., Ziemann, P. J., and Jimenez, J. L.: Response of an aerosol mass
spectrometer to organonitrates and organosulfates and implications for atmospheric chemistry, *Proc Natl Acad Sci USA*, 107, 6670–6675,
680 <https://doi.org/10.1073/pnas.0912340107>, 2011.
- Formenti, P., Elbert, W., Maenhaut, W., Haywood, J., Osborne, S., and Andreae, M. O.: Inorganic and carbonaceous aerosols during the
Southern African Regional Science Initiative (SAFARI 2000) experiment: Chemical characteristics, physical properties, and emission
data for smoke from African biomass burning, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JD002408>, 2003.
- Forrister, H., Liu, J., Scheuer, E., Dibb, J., and et al., L. Z.: Evolution of brown carbon in wildfire plumes, *Geophys. Res. Lett.*, 42, 4623–4630,
685 <https://doi.org/10.1002/2015GL063897>, 2015.

- Garofalo, L. A., Pothier, M. A., Levin, E. J., Campos, T., Kreidenweis, S. M., and Farmer, D. K.: Emission and Evolution of Submicron Organic Aerosol in Smoke from Wildfires in the Western United States, *ACS Earth Space Chem.*, 3, 1237–1247, <https://doi.org/10.1021/acsearthspacechem.9b00125>, 2019.
- George, I. J. and Abbatt, J. P. D.: Chemical evolution of secondary organic aerosol from OH-initiated heterogeneous oxidation, *Atmos. Chem. Phys.*, 10, 5551–5563, <https://doi.org/10.5194/acp-10-5551-2010>, 2010.
- 690 Giglio, L., van der Werf, G. R., Randerson, J. T., Collatz, G., and Kasibhatla, P.: Global estimation of burned area using MODIS active fire observations, *Atmos. Chem. Phys.*, 6, 957–974, <https://doi.org/10.5194/acp-6-957-2006>, 2006.
- Graaf, M. D., Bellouin, N., Tilstra, L. G., Haywood, J., and Stammes, P.: Aerosol direct radiative effect of smoke over clouds over the southeast Atlantic Ocean from 2006 to 2009, *Geophys. Res. Lett.*, 41, 7723–7730, <https://doi.org/10.1002/2014GL061103>, 2014.
- 695 Gysel, M., Laborde, M., Olfert, J. S., Subramanian, R., and Gröhn, A.: Effective density of Aquadag and fullerene soot black carbon reference materials used for SP2 calibration, *Atmos. Meas. Tech. Disc.*, 4, 4937–4955, <https://doi.org/10.5194/amtd-4-4937-2011>, 2011.
- Haywood, J. M., Osborne, S. R., Francis, P. N., Keil, A., Andreae, P. F. M. O., and Kaye, P. H.: The mean physical and optical properties of regional haze dominated by biomass burning aerosol measured from the C-130 aircraft during SAFARI 2000, *J. Geophys. Res.*, 108, 8473–8481, <https://doi.org/10.1029/2002JD002226>, 2003.
- 700 Haywood, J. M., Abel, S. J., Barrett, P. A., Bellouin, N., Blyth, A., Bower, K. N., Brooks, M., Carslaw, K., Che, H., Coe, H., Cotterell, M. I., Crawford, I., Cui, Z., Davies, N., Dingley, B., Field, P., Formenti, P., Gordon, H., de Graaf, M., Herbert, R., Johnson, B., Jones, A. C., Langridge, J. M., Malavelle, F., Partridge, D. G., Peers, F., Redemann, J., Stier, P., Szpek, K., Taylor, J. W., Watson-Parris, D., Wood, R., Wu, H., and Zuidema, P.: Overview: The CLOUD-Aerosol-Radiation Interaction and Forcing: Year-2017 (CLARIFY-2017) measurement campaign, *Atmos. Chem. Phys.*, 21, 1049–1084, <https://doi.org/10.5194/acp-21-1049-2021>, 2021.
- 705 Heald, C. L., Kroll, J. H., Jimenez, J. L., Docherty, K. S., DeCarlo, P. F., Aiken, A. C., Chen, Q., Martin, S. T., Farmer, D. K., and Artaxo, P.: A simplified description of the evolution of organic aerosol composition in the atmosphere, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010gl042737>, 2010.
- Herbert, R. J., Bellouin, N., Highwood, E. J., and Hill, A. A.: Diurnal cycle of the semi-direct effect from a persistent 1300 absorbing aerosol layer over marine stratocumulus in large-eddy simulations, *Atmos. Chem. Phys.*, 20, 1317–1340, [https://doi.org/10.5194/acp-20-1317-](https://doi.org/10.5194/acp-20-1317-2020)
- 710 2020, 2020.
- Hodshire, A. L., Akherati, A., Alvarado, M., Brown-Steiner, B., Jathar, S., Jimenez, J. L., Kreidenweis, S., Lonsdale, C., Onasch, T., Ortega, A., and Pierce, J.: Aging Effects on Biomass Burning Aerosol Mass and Composition: A Critical Review of Field and Laboratory Studies, *Environ Sci Technol*, 53, 10 007–10 022, <https://doi.org/10.1021/acs.est.9b02588>, 2019.
- Hodshire, A. L., Ramnarine, E., Akherati, A., Alvarado, M. L., Farmer, D. K., Jathar, S. H., and et al., S. M. K.: Dilution impacts on smoke
- 715 aging: evidence in Biomass Burning Observation Project (BBOP) data, *Atmos. Chem. Phys.*, 21, 6839–6855, <https://doi.org/10.5194/acp-21-6839-2021>, 2021.
- Hodzic, A., Madronich, S., Kasibhatla, P. S., Tyndall, G., Aumont, B., Jimenez, J. L., Lee-Taylor, J., and Orlando, J.: Organic photolysis reactions in tropospheric aerosols: effect on secondary organic aerosol formation and lifetime, *Atmos. Chem. Phys.*, 15, 9253–9269, <https://doi.org/10.5194/acp-15-9253-2015>, 2015.
- 720 Hodzic, A., Campuzano-Jost, P., Bian, H., Chin, M., Colarco, P. R., Day, D. A., Froyd, K. D., Heinold, B., and et al., D. S. J.: Characterization of organic aerosol across the global remote troposphere: a comparison of ATOM measurements and global chemistry models, *Atmos. Chem. Phys.*, 20, 4607–4635, <https://doi.org/10.5194/acp-20-4607-2020>, 2020.

- Holanda, B. A., Pöhlker, M. L., Walter, D., Saturno, J., Sörgel, M., Ditas, J., Ditas, F., Schulz, C., Franco, M. A., Wang, Q., Donth, T., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Braga, R., Brito, J., Cheng, Y., Dollner, M., Kaiser, J. W., Klimach, T., Knote, C., Krüger, O. O., Fütterer, D., Lavrič, J. V., Ma, N., Machado, L. A. T., Ming, J., Morais, F. G., Paulsen, H., Sauer, D., Schlager, H., Schneider, J., Su, H., Weinzierl, B., Walser, A., Wendisch, M., Ziereis, H., Zöger, M., Pöschl, U., Andreae, M. O., and Pöhlker, C.: Influx of African biomass burning aerosol during the Amazonian dry season through layered transatlantic transport of black carbon-rich smoke, *Atmos. Chem. Phys.*, 20, 4757–4785, <https://doi.org/10.5194/acp-20-4757-2020>, 2020.
- Holder, A. L., Hagler, G. S., Aurell, J., Hays, M. D., and Gullett, B. K.: Particulate matter and black carbon optical properties and emission factors from prescribed fires in the southeastern United States, *J. Geophys. Res.*, 121, 3465–3483, <https://doi.org/10.1002/2015JD024321>, 2016.
- Howell, S. G., Clarke, A. D., Freitag, S., McNaughton, C. S., Kapustin, V., Brekovskikh, V., Jimenez, J.-L., and Cubison, M. J.: An airborne assessment of atmospheric particulate emissions from the processing of Athabasca oil sands, *Atmos. Chem. Phys.*, 14, 5073–5087, <https://doi.org/10.5194/acp-14-5073-2014>, 2014.
- Howell, S. G., Freitag, S., Dobracki, A., Smirnow, N., and III, A. J. S.: Undersizing of aged African biomass burning aerosol by an ultra-high-sensitivity aerosol spectrometer, *Atmos. Meas. Tech.*, 14, 7381–7404, <https://doi.org/10.5194/amt-14-7381-2021>, 2021.
- Huntley, B. J.: Angola in outline: physiography, climate and patterns of biodiversity, chap. 2, pp. 15–42, Springer, <https://doi.org/10.1007/978-3-030-03083-4>, 2019.
- Janhäll, S., Andreae, M. O., and Pöschl, U.: Biomass burning aerosol emissions from vegetation fires: particle number and mass emission factors and size distributions, *Atmos. Chem. Phys.*, 10, 1427–1439, 2010.
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., and al., J. D. A.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, <https://doi.org/10.1126/science.1180353>, 2009.
- Jolleys, M. D., Coe, H., McFiggans, G., Capes, G., Allan, J. D., Crosier, J., Williams, P. I., Allen, G., Bower, K. N., Jimenez, J. L., Russell, L. M., Grutter, M., and Baumgardner, D.: Characterizing the aging of biomass burning organic aerosol by use of mixing ratios: a meta-analysis of four regions, *Environ. Sci. Technol.*, 46, 13 093–13 102, 2012.
- Jolleys, M. D., Coe, H., McFiggans, G., Taylor, J. W., O’Shea, S. J., Breton, M. L., Bauguitte, S. J.-B., Moller, S., and et al., P. D. C.: Properties and evolution of biomass burning organic aerosol from Canadian boreal forest fires, *Atmos. Chem. Phys.*, 15, 3077–3095, <https://doi.org/10.5194/acp-15-3077-2015>, 2015.
- Keil, A. and Haywood, J. M.: Solar radiative forcing by biomass burning aerosol particles during SAFARI 2000: A case study based on measured aerosol and cloud properties, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JD002315>, 2003.
- Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Liousse, C., Mari, C., Allan, R. P., Brooks, B., Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCIWA project: Dynamics-Aerosol-Chemistry-Cloud Interactions in West Africa, *B. Am. Meteorol. Soc.*, 96, 1451–1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, 2015.
- Kondo, Y., Matsui, H., Moteki, N., Sahu, L., Takegawa, N., Kajino, M., Zhao, Y., Cubison, M. J., Jimenez, J. L., Vay, S., Diskin, G. S., Anderson, B., Wisthaler, A., Mikoviny, T., Fuelberg, H. E., Blake, D. R., Huey, G., Weinheimer, A. J., Knapp, D. J., and Brune, W. H.: Emissions of black carbon, organic, and inorganic aerosols from biomass burning in North America and Asia in 2008, *J. Geophys. Res.*, 116, 2011.
- Konovalov, I. B., Beekmann, M., Golovushkin, N. A., and Andreae, M. O.: Nonlinear behavior of organic aerosol in biomass burning plumes: a microphysical model analysis, *Atmos. Chem. Phys.*, 19, 12 091–12 119, <https://doi.org/10.5194/acp-19-12091-2019>, 2019.

- 760 Korontzi, S., Ward, D. E., Susott, R. A., Yokelson, R. J., Justice, C. O., Hobbs, P. V., Smithwick, E. A. H., and Hao, W. M.: Seasonal variation and ecosystem dependence of emission factors for selected trace gases and PM_{2.5} for southern African savanna fires, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2003JD003730>, 2003.
- Kroll, J., Smith, J. D., Che, D. L., Kessler, S. H., Worsnop, D. R., and Wilson, K. R.: Measurement of fragmentation and functionalization pathways in the heterogeneous oxidation of oxidized organic aerosol, *Phys. Chem. Chem. Phys.*, 11, 8005–8014, <https://doi.org/10.1039/B905289E>, 2009.
- 765 Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Env.*, 42, 3593–3624, 2008.
- Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R., Wilson, K. R., and et al., K. E. A.: Carbon oxidation state as a metric for describing the chemistry of atmospheric organic aerosol, *Nat Chem.*, 3, 133–139, <https://doi.org/10.1038/nchem.948>, 2011.
- 770 Kuete, G., Mba, W. P., and Washington, R.: African Easterly Jet South: control, maintenance mechanisms and link with Southern subtropical waves, *Clim Dyn.*, 54, 1539–1552, <https://doi.org/10.1007/s00382-019-05072-w>, 2020.
- Laborde, M., Schnaiter, M., Linke, C., Saathoff, H., Naumann, K., Mohler, O., Berlenz, S., Wagner, U., Taylor, J., Liu, D., Flynn, M., Allan, J., Coe, H., Heimerl, K., Dahlkotter, F., Weinzierl, B., Wollny, A., Zanatta, M., Cozic, J., Laj, P., Hitznerberger, R., Schwarz, J., and Gysel, M.: Single Particle Soot Photometer intercomparison at the AIDA chamber, *Atmos. Meas. Tech.*, 5, 3077–3097, <https://doi.org/10.5194/amt-5-3077-2012>, 2012.
- 775 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of atmospheric brown carbon, *Chem. Rev.*, 155, 4335–4382, <https://doi.org/10.1021/cr5006167>, 2015.
- Lee, J. E., Gorkowski, K., Meyer, A., Benedict, K., Aiken, A. C., and Dubey, M. K.: Wildfire smoke demonstrates significant and predictable black carbon light absorption enhancements, *Geophys. Res. Lett.*, 49, <https://doi.org/10.1029/2022GL099334>, e2022GL099334, 2022.
- 780 Li, Z., Smith, K. A., and Cappa, C. D.: Influence of relative humidity on the heterogeneous oxidation of secondary organic aerosol, *Atmos. Chem. Phys.*, 18, 14 585–14 608, <https://doi.org/10.5194/acp-18-14585-2018>, 2018.
- Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S., Williams, P. I., , Ting, Y.-C., Haslett, S., Taylor, J., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe, H., and Allan, J. D.: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, *Nat. Geosci.*, 10, 184–188, <https://doi.org/10.1038/ngeo2901>, 2017.
- 785 Lou, S., Shrivastava, M., Easter, R. C., Yang, Y., Ma, P.-L., Wang, H., Cubison, M., Campuzano-Jost, P., and et al., J. L. J.: New SOA Treatments Within the Energy Exascale Earth System Model (E3SM): Strong Production and Sinks Govern Atmospheric SOA Distributions and Radiative Forcing, *J. Adv. Mod. Earth Sys.*, 12, <https://doi.org/10.1029/2020ms002266>, 2020.
- Mallet, M., Solmon, F., Nabat, P., Elguindi, N., Waquet, F., and et al., D. B.: Direct and semi-direct radiative forcing of biomass-burning aerosols over the southeast Atlantic (SEA) and its sensitivity to absorbing properties: a regional climate modeling study, *Atmos. Chem. Phys.*, 20, 13 191–13 216, <https://doi.org/10.5194/acp-20-13191-2020>, 2020.
- 790 Mallet, M., Nabat, P., Johnson, B., Michou, M., Haywood, J. M., Chen, C., and Dubovik, O.: Climate models generally underrepresent the warming by Central Africa biomass-burning aerosols over the Southeast Atlantic, *Sci. Adv.*, 7, eabg9998, <https://doi.org/10.1126/sciadv.abg9998>, 2021.
- Martin, M. V., Logan, J. A., Kahn, R. A., Leung, F.-Y., Nelson, D. L., and Diner, D. J.: Smoke injection heights from fires in North America: Analysis of 5 years of satellite observations, *Atmos. Chem. Phys.*, 10, 1491–1510, <https://doi.org/10.5194/acp-10-1491-2010>, 2010.
- 795

- May, A. A., McMeeking, G. R., Lee, T., Taylor, J. W., Craven, J. S., Burling, I., Sullivan, A. P., Akagi, S., Collett, J. L., and Flynn, M.: Aerosol emissions from prescribed fires in the United States: A synthesis of laboratory and aircraft measurements, *J. Geophys. Res.*, 119, 11 826–11 849, <https://doi.org/10.1002/2014JD021848>, 2014.
- McClure, C. D., Lim, C. Y., Hagan, D. H., Kroll, J., and Cappa, C.: Biomass-burning-derived particles from a wide variety of fuels – Part 1: Properties of primary particles, *Atmos. Chem. Phys.*, 20, 1531–1547, <https://doi.org/10.5194/acp-20-1531-2020>, 2020.
- McFiggans, G., Mentel, T. F., and Wildt, J.: Secondary organic aerosol reduced by mixture of atmospheric vapours, *Nature*, 565, 587–593, <https://doi.org/10.1038/s41586-018-0871-y>, 2019.
- Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., and et al., T. K. B.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmospheric Chemistry and Physics*, 13, 1853–1877, <https://doi.org/10.5194/acp-13-1853-2013>, 2013.
- 805 Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., and et al., U.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, <https://doi.org/10.5194/acp-10-4625-2010>, 2010.
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Chhabra, P. S., Seinfeld, J. H., and Worsnop, D. R.: Changes in organic aerosol composition with aging inferred from aerosol mass spectra, *Atmos. Chem. Phys.*, 11, 6465–6474, <https://doi.org/10.5194/acp-11-6465-2011>, 2011.
- Nicholson, S. E. and Grist, J. P.: The seasonal evolution of the atmospheric circulation over West Africa and equatorial Africa, *J. Clim.*, 16, 810 1013–1030, 2003.
- O'Brien, R. E. and Kroll, J. H.: Photolytic Aging of Secondary Organic Aerosol: Evidence for a Substantial Photo-Recalcitrant Fraction, *J. Phys. Chem. Lett.*, 10, 4003–4009, <https://doi.org/10.1021/acs.jpcclett.9b01417>, 2019.
- Pan, X., Kanaya, Y., Taketani, F., Miyakawa, T., Inomata, S., Komazaki, Y., Tanimoto, H., Wang, Z., Uno, I., and Wang, Z.: Emission characteristics of refractory black carbon aerosols from fresh biomass burning: a perspective from laboratory experiments, *Atmos. Chem. Phys.*, 17, 13 001–13 016, <https://doi.org/10.5194/acp-17-13001-2017>, 2017.
- 815 Pendergrass, A. G. and Hartmann, D. L.: Global-mean precipitation and black carbon in AR4 simulations, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2011GL050067>, 101703, 2012.
- Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S.-M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, X.-Y.: Recommendations for reporting black carbon measurements, *Atmos. Chem. Phys.*, 13, 820 8365–8379, <https://doi.org/10.5194/acp-13-8365-2013>, 2013.
- Pistone, K., Redemann, J., Doherty, S., Zuidema, P., Burton, S., Cairns, B., Cochrane, S., Ferrare, R., Flynn, C., Freitag, S., Howell, S., Kacenenbogen, M., LeBlanc, S., Liu, X., Schmidt, K. S., Sedlacek III, A. J., Segal-Rosenhaimer, M., Shinozuka, Y., Stamnes, S., van Diedenhoven, B., Van Harten, G., and Xu, F.: Intercomparison of biomass burning aerosol optical properties from in-situ and remote-sensing instruments in ORACLES-2016, *Atmos. Chem. Phys.*, 19, 9181–9208, <https://doi.org/10.5194/acp-19-9181-2019>, 2019.
- 825 Pistone, K., Zuidema, P., Wood, R., Diamond, M., da Silva, A. M., Ferrada, G., Saide, P., Ueyama, R., Pfister, J.-M. R. L., Podolske, J., Noone, D., Bennett, R., Stith, E., Carmichael, G., Redemann, J., Flynn, C., LeBlanc, S., Segal-Rosenhaimer, M., and Shinozuka, Y.: Exploring the elevated water vapor signal associated with the free-tropospheric biomass burning plume over the southeast Atlantic Ocean, *Atmos. Chem. Phys.*, 21, 9643–9668, <https://doi.org/10.5194/acp-21-9643-2021>, 2021.
- Pokhrel, R. P., Gordon, J., Fiddler, M., and Bililign, S.: Determination of emission factors of pollutants from biomass burning of African 830 fuels in laboratory measurements, *J. Geophys. Res.*, 126, <https://doi.org/10.1029/2021JD034731>, e2021JD034731, 2021.
- Ramo, R., Roteta, E., Bistinas, I., van Wees, D., Bastarrika, A., Chuvieco, E., and der Werf, G. R.: African burned area and fire carbon emissions are strongly impacted by small fires undetected by coarse resolution satellite data, *Proc. Nat. Ac. Sci.*, 118, e2011160118, <https://doi.org/10.1073/pnas.2011160118>, 2021.

- Redemann, J., Wood, R., Zuidema, P., Doherty, S. J., Luna, B., LeBlanc, S. E., Diamond, M. S., Shinozuka, Y., Chang, I. Y., Ueyama, R.,
835 Pfister, L., Ryoo, J.-M., Dobracki, A. N., da Silva, A. M., Longo, K. M., Kacenenbogen, M. S., Flynn, C. J., Pistone, K., Knox, N. M.,
Piketh, S. J., Haywood, J. M., Formenti, P., Mallet, M., Stier, P., Ackerman, A. S., Bauer, S. E., Fridlind, A. M., Carmichael, G. R., Saide,
P. E., Ferrada, G. A., Howell, S. G., Freitag, S., Cairns, B., Holben, B. N., Knobelspiesse, K. D., Tanelli, S., L'Ecuyer, T. S., Dzambo,
A. M., Sy, O. O., McFarquhar, G. M., Poellot, M. R., Gupta, S., O'Brien, J. R., Nenes, A., Kacarab, M., Wong, J. P. S., Small-Griswold,
J. D., Thornhill, K. L., Noone, D., Podolske, J. R., Schmidt, K. S., Pilewskie, P., Chen, H., Cochrane, S. P., Sedlacek, A. J., Lang, T. J.,
840 Stith, E., Segal-Rozenhaimer, M., Ferrare, R. A., Burton, S. P., Hostetler, C. A., Diner, D. J., Seidel, F. C., Platnick, S. E., Myers, J. S.,
Meyer, K. G., Spangenberg, D. A., Maring, H., and Gao, L.: An overview of the ORACLES (ObseRvations of Aerosols above CLouds
and their intEractionS) project: aerosol–cloud–radiation interactions in the southeast Atlantic basin, *Atmos. Chem. Phys.*, 21, 1507–1563,
<https://doi.org/10.5194/acp-21-1507-2021>, 2021.
- Ryoo, J.-M., Pfister, L., Ueyama, R., Zuidema, P., Wood, R., Chang, I., and Redemann, J.: A meteorological overview of the ORACLES
845 (ObseRvations of Aerosols above CLouds and their intEractionS) campaign over the southeast Atlantic during 2016–2018: Part 1 -
Climatology, *Atmos. Chem. Phys.*, 21, 16 689–16 707, <https://doi.org/10.5194/acp-21-16689-2021>, 2021.
- Ryoo, J.-M., Pfister, L., Ueyama, R., Zuidema, P., Wood, R., Chang, I., and Redemann, J.: A meteorological overview of the ORACLES
(ObseRvations of Aerosols above CLouds and their intEractionS) campaign over the southeast Atlantic during 2016–2018: Part 1 - Daily
and Synoptic Characteristics, *Atmos. Chem. Phys. Disc.*, 22, <https://doi.org/10.5194/acp-2022-256>, 2022.
- 850 Saide, P., Thompson, G., Eidhammer, T., da Silva, A. M., and Carmichael, R. B. P. G. R.: Assessment of biomass burning smoke influence
on environmental conditions for multi-year tornado outbreaks by combining aerosol-aware microphysics and fire emission constraints, *J
Geophys. Res.*, 121, 10 294–10 311, <https://doi.org/10.1002/2016JD025056>, 2016.
- Saleh, R., Robinson, E., and et al., D. T.: Brownness of organics in aerosols from biomass burning linked to their black carbon content,
Nature Geosci., 7, 647–650, <https://doi.org/10.1038/ngeo2220>, 2014.
- 855 Scholes, R. J., Ward, D. E., and Justice, C. O.: Emissions of trace gases and aerosol particles due to vegetation burning in southern hemisphere
Africa, *J. Geophys. Res.*, 101, 23 677–23 682, <https://doi.org/10.1029/95JD02049>, 1996.
- Sedlacek, A., Lewis, E., Onasch, T., Zuidema, P., Redemann, J., Jaffee, D., and Kleinman, L.: Using the black carbon particle mixing state
to characterize the lifecycle of biomass burn aerosols, *Env. Sci. Tech.*, 56, 14 315–14 325, <https://doi.org/10.1021/acs.est.2c30851>, 2022.
- Shank, L. M., Howell, S., Clarke, A. D., Freitag, S., Brekhovskikh, V., Kapustin, V., McNaughton, C., Campos, T., and Wood, R.: Organic
860 matter and non-refractory aerosol over the remote Southeast Pacific: oceanic and combustion sources, *Atmos. Chem. Phys.*, 12, 557–576,
<https://doi.org/10.5194/acp-12-557-2012>, 2012.
- Shea, R. W., Shea, B. W., Kauffman, J. B., Ward, D. E., Haskins, C. I., and Scholes, M. C.: Fuel biomass and combustion factors associated
with fires in savanna ecosystems of South Africa and Zambia, *J. Geophys. Res.*, 101, 23,551– 23,568, 1996.
- Shinozuka, Y., Saide, P. E., Ferrada, G. A., Burton, S. P., Ferrare, R., Doherty, S. J., Gordon, H., Longo, K., Mallet, M., Feng, Y., Wang, Q.,
865 Cheng, Y., Dobracki, A., Freitag, S., Howell, S. G., LeBlanc, S., Flynn, C., Segal-Rosenhaimer, M., Pistone, K., Podolske, J. R., Stith,
E. J., Bennett, J. R., Carmichael, G. R., da Silva, A., Govindaraju, R., Leung, R., Zhang, Y., Pfister, L., Ryoo, J.-M., Redemann, J., Wood,
R., and Zuidema, P.: Modeling the smoky troposphere of the southeast Atlantic: a comparison to ORACLES airborne observations from
September of 2016, *Atmos. Chem. Phys.*, 20, 11,491–11,526, <https://doi.org/10.5194/acp-20-11491-2020>, 2020.
- Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A., Jimenez, J. L., and et al., C. K.: Recent advances in understanding
870 secondary organic aerosol: Implications for global climate forcing, *Rev. Geophys.*, 55, 509–559, <https://doi.org/10.1002/2016RG000540>,
2017.

- Solmon, F., Elguindi, N., and et al., M. M.: West African monsoon precipitation impacted by the South Eastern Atlantic biomass burning aerosol outflow, *npj Clim Atmos Sci.*, 4, <https://doi.org/10.1038/s41612-021-00210-w>, 2021.
- 875 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bull. Am. Meteor. Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Swap, R. J., Annegarn, H. J., Suttles, J. T., King, M. D., Platnick, S., Privette, J. L., and Scholes, R. J.: Africa burning: A thematic analysis of the Southern African Regional Science Initiative (SAFARI 2000), *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2003JD003747>, 8465, 2003.
- 880 Taylor, J. W., Wu, H., Szpek, K., Bower, K., Crawford, I., Flynn, M. J., Williams, P. I., Dorsey, J., Langridge, J. M., and et al.: Absorption closure in highly aged biomass burning smoke, *Atmos. Chem. Phys.*, 20, 11 201–11 221, <https://doi.org/10.5194/acp-20-11201-2020>, 2020.
- Thompson, G. and Eidhammer, T.: A study of aerosol impacts on clouds and precipitation development in a large winter cyclone, *J. Atmos. Sci.*, 71, 3636–3658, <https://doi.org/10.1175/JAS-D-13-0305.1>, 2014.
- 885 Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E., Bellouin, N., and Benedetti, A. e. a.: The AeroCom evaluation and intercomparison of organic aerosol in global models, *Atmos. Chem. Phys.*, 14, 10 845–10 895, <https://doi.org/10.5194/acp-14-10845-2014>, 2014.
- Vakkari, V., Beukes, J. P., Maso, M. D., Aurela, M., Josipovic, M., and van Zyl, P. G.: Major secondary aerosol formation in southern African open biomass burning plumes, *Nature Geo.*, 11, 580–583, <https://doi.org/10.1038/s41561-018-0170-0>, 2018.
- 890 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., and Morton, D. C. e. a.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11 707–11 735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- vanKrevelen, D. W.: Graphical-statistical method for the study of structure and reaction processes of coal, *Fuel*, 24, 269–284, 1950.
- Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), *Aerosol Sci. Tech.*, 44, 706–712, 2010.
- 895 Waquet, F., Peers, F., Ducos, F., Goloub, P., Platnick, S., Riedi, J., Tanré, D., and Thieuleux, F.: Global analysis of aerosol properties above clouds, *Geophys. Res. Lett.*, 40, 5809–5814, <https://doi.org/10.1002/2013GL057482>, 2013.
- Wu, H., Taylor, J. W., Szpek, K., Langridge, J. M., Williams, P. I., Flynn, M., Allan, J. D., Abel, S. J., Pitt, J., Cotterell, M. I., Fox, C., Davies, N. W., Haywood, J., and Coe, H.: Vertical variability of the properties of highly aged biomass burning aerosol transported over the southeast Atlantic during CLARIFY-2017, *Atmos. Chem. Phys.*, 20, 12 697–12 719, <https://doi.org/10.5194/acp-20-12697-2020>, 2020.
- 900 Yang, M., Huebert, B. J., Blomquist, B. W., Howell, S. G., Shank, L. M., McNaughton, C. S., Clarke, A. D., Hawkins, L. N., and et al., L. M. R.: Atmospheric sulfur cycling in the southeastern Pacific – longitudinal distribution, vertical profile, and diel variability observed during VOCALS-REx, *Atmos. Chem. Phys.*, 11, 5079–5097, <https://doi.org/10.5194/acp-11-5079-2011>, 2011.
- Yokelson, R. J., Susot, R. A., Ward, D. E., Reardon, J., and Griffith, D. W. T.: Emissions from Smoldering Combustion of Biomass Measured by Open-Path Fourier Transform Infrared Spectroscopy, *J. Geophys. Res.*, 102, 865–877, 1997.
- 905 Yokelson, R. J., Crounse, J. D., DeCarlo, P. F., Karl, T., Urbanski, S., and et al., E. A.: Emissions from biomass burning in the Yucatan, *Atmos. Chem. Phys.*, 9, 5785–5812, 2009.
- Zhang, J. and Zuidema, P.: Sunlight-absorbing aerosol amplifies the seasonal cycle in low-cloud fraction over the southeast Atlantic, *Atmos. Chem. Phys.*, 21, 11 179–11 199, <https://doi.org/10.5194/acp-21-11179-2021>, 2021.

- Zhang, L., Segal-Rozenhaimer, M., Che, H., Dang, C., III, A. J. S., Lewis, E. R., Dobracki, A., Wong, J. P. S., Formenti, P., Howell, S. G., and Nenes, A.: Light Absorption by Brown Carbon over the South-East Atlantic Ocean, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2021-1000>, 2022.
- Zhang, Q., Jimenez, J. L., Worsnop, D. R., and Canagaranta, M.: A Case Study of Urban Particle Acidity and Its Influence on Secondary Organic Aerosol, *Environ. Sci. Technol.*, 41, 3213–3219, <https://doi.org/10.1021/es061812j>, 2007.
- Zheng, B., Chevallier, F., Ciaia, P., Yin, Y., and Wang, Y.: On the role of the flaming to smoldering transition in the seasonal cycle of African fire emissions, *Geophys. Res. Lett.*, 45, 11,998–12,007, <https://doi.org/10.1029/2018GL079092>, 2018.
- Zheng, G., Sedlacek, A. J., Aiken, A. C., Feng, Y., B. Watson, T., Raveh-Rubine, S., Uin, J., R. Lewis, E., and Wang, J.: Long-range transported north American wildfire aerosols observed in marine boundary layer of eastern North Atlantic, *Env. Inter.*, 139, <https://doi.org/10.1016/j.envint.2020.105680>, 2020.
- Zhou, J.: Hygroscopic properties of atmospheric aerosol particles in various environments, Ph.D. thesis, [https://doi.org/ISBN 91-7874-120-3](https://doi.org/ISBN%2091-7874-120-3), LUTFD2/(TFKF-1025)/1-166/, 166 pp., Lund University, Lund, Sweden, 2001.
- Zhou, S., Collier, S., Jaffe, D. A., Briggs, N. L., Hee, J., III, A. J. S., Kleinman, L., Onasch, T. B., and Zhang, Q.: Regional influence of wildfires on aerosol chemistry in the western US and insights into atmospheric aging of biomass burning organic aerosol, *Atmos. Chem. Phys.*, pp. 2477–2493, <https://doi.org/10.5194/acp-17-2477-2017>, 2017.
- Zuidema, P., Redemann, J., Haywood, J., Wood, R., Piketh, S., Hipondoka, M., and Formenti, P.: Smoke and Clouds above the South-east Atlantic: Upcoming Field Campaigns Probe Absorbing Aerosol's Impact on Climate, *Bull. Am. Meteor. Soc.*, 97, 1131–1135, <https://doi.org/10.1175/bams-d-15-00082.1>, 2016.
- Zuidema, P., Sedlacek III, A. J., Flynn, C., Springston, S., Delgadoillo, R., Zhang, J., Aiken, A. C., Koontz, A., and Muradyan, P.: The Ascension Island Boundary Layer in the Remote Southeast Atlantic is Often Smoky, *Geophys. Res. Lett.*, 45, 4456–4465, <https://doi.org/10.1002/2017gl076926>, 2018.

Table 1. Instrument Table

Measurement	Instrument (Manufacturer)	Notes
OA, NO ₃ , SO ₄ , NH ₄ masses, <i>f44</i> , <i>f60</i> , O,C,H,OC	HR-ToF-AMS (Aerodyne Inc.)	chloride excluded
BC mass, number	SP2 (DMT)	operated by HiGEAR in 2016, A. Sedlacek in 2017
particle size distribution	LDMA (TSI 3071A)	10-550 nm mobility diameter
particle number concentration	CPC (TSI 3010)	diameter > 10nm
aerosol absorption (470, 530, 660 nm)	PSAP (Radiance Research)	Virkkula (2010) wavelength-averaged correction
aerosol scattering (450, 550, 700 nm)	nephelometer (TSI 3563)	Anderson and Ogren (1998) correction
CO, CO ₂ , O ₃	Los Gatos Research	

abbreviations provided in text

Table 2. Comparison to other published $\frac{BC}{\Delta CO}$ ratios

Fuel/Geographic Source	$\frac{BC}{\Delta CO} * 10^{-3}$	reference
savannah	2-15	Vakkari et al. (2018)
grass	10-17	Vakkari et al. (2018)
savannah	7.9	Andreae (2019)
agriculture	5.6	Andreae (2019)
savannah	5.9	Akagi et al. (2011)
agriculture (crop residue)	7.4	Akagi et al. (2011)
NW African agriculture, smouldering	7.2	Capes et al. (2008)
southern Africa (SAFARI)	7.0	Formenti et al. (2003)
Ascension Island, August	8.7-13.4	Wu et al. (2020)
this study	9.6	

all $\frac{BC}{\Delta CO}$ values are dimensionless. Methods for deriving the BC mass concentration may vary between the studies. Most CMIP6 models rely on the Akagi et al. (2011) emission factors.

Table 3. Comparison of level-leg mean values to CLARIFY

	CLARIFY	September 2016	31 August 2017
BC mass frac. (%)	13-15	5.4-9.2	7
BC num. frac. (%)	39 ± 7	15-40	30-35
SSA_{530}	~ 0.84	0.85-0.88	0.83-0.86
$MAC_{BC,660}$ ($m^2 g^{-1}$)	11-12	9.5-11.5	10-11.5
OA:BC mass	4-5	$10-14 \pm 2$	8-10
LDMA median diam. (nm)	232	140-180	180-200
BC core diam. (nm)		130-150	150-160
IN frac. (%)	100	~ 25	~ 50
f_{44}	0.19-0.22	0.18-0.22	0.215
$\frac{OA}{\Delta CO} * 10^{-2}$	4.2-6.4	6-11	6.5-11.5

CLARIFY free-tropospheric values taken from Wu et al. (2020) and Taylor et al. (2020), based on CLARIFY 033-039 and 045-051 flights. CLARIFY BC number fraction calculated relative to PCASP-derived total number concentrations.




Figure 1. Terra MODIS Above Cloud Aerosol Optical Depth (Meyer, 2015) for September 2016 overlaid with the tracks of the 6 flights selected for this study. The location of the profile shown in Fig. 12 is indicated with red diamonds. ST=Sao Tome; WB=Walvis Bay; ASI=Ascension Island.




Figure 2. 24 September 2016 flight track with colorized OA:BC mass ratios superimposed on High Spectral Resolution Lidar-2 523 nm aerosol backscatter imagery collected along 11°E by the overflying ER-2 plane, near in time to the P-3 plane location's at 10°S .




Figure 3. Leg-averaged SP2 size by number (top) and mass (bottom) distributions for the level legs detailed in Table S2. Total leg-averaged SP2 number concentrations included in top plot. The kink at 2.5 μm mass-mean diameter for the 2016 data is an instrument artifact.




Figure 4. OA:BC mass ratio, shown using 10, 25, 50, 75 and 90th percentiles with means in red, composited by OA mass.




Figure 5. Distribution of the bulk chemical species masses for each flight, for five-second samples with $\text{OA} > 20 \mu\text{g m}^{-3}$.




Figure 6. a) f_{44} , b) OA:BC mass ratio, c) model-derived time since emission (age), d) MCE, e) $\frac{BC}{\Delta CO}$ as a non-dimensionalized ratio, and f) ozone, all for each indicated flight. Whiskers represent the 10th and 90th percentiles, boxes illustrate the 75th and 25th percentiles with a line indicating the median and yellow (2017) and blue (2016) filled circles representing the mean. OA > 20 $\mu\text{g m}^{-3}$ only.




Figure 7. a) $f44$ versus $f43$ for the six flights where $OA > 20 \mu\text{g m}^{-3}$. Averages (\pm standard deviation) are colored by flight date, grey boxes indicate individual data points. b) similar to a), for $f44$ vs $f60$. Blue and red dashed lines define the parameters for ambient oxygenated OA, following Ng et al. (2010). c) $f44$ versus the model-derived physical age for the six flights combined.




Figure 8. Hydrogen to carbon (H:C) molar ratio versus oxygen to carbon (O:C) molar ratio, colored by flight date, shown at the native 5-second time resolution. Superimposed are lines of constant oxidation state (OS, defined as $2 \cdot O:C - H:C$; Kroll et al., 2011), used to define semi-volatile (SV), aged and low-volatile (LV) oxidized organic aerosol (OOA) regimes.




Figure 9. a) Mass absorption coefficient ($MAC_{BC,660}$; units of $m^2 g^{-1}$) at 660 nm wavelength versus the fraction of black-carbon-containing particles (FrBC), colored by flight day. b) same as a) but for SSA_{530} versus FrBC. c) same as a) but for OA:BC mass versus FrBC. d) $MAC_{BC+OA,470}$ ($m^2 g^{-1}$) versus LDMA median and mass-median BC core mass-median-diameter. All for the level legs listed in Table S2.




Figure 10. a) Level-leg-mean \pm standard deviation values for SSA_{530nm} versus the OA:BC mass ratio, colored by flight. The best-fit line is represented by $SSA=0.801+0.0055*(OA:BC)$ ($r=0.84$). b) same as a) but for SSA_{530nm} versus the BC:TC mass ratio, where TC=BC+organic carbon. The best-fit line is $SSA=0.93-0.39*(BC:TC)$, ($r=-0.79$). Times and spatial ranges of the level-legs provided in Table S1. Also shown is the SSA parameterization put forth within Brown et al. (2021), namely $SSA_{530nm}=0.969-0.779*(BC:TC)$, where TC=BC+organic carbon (OC) and OC is estimated from $OA:OC=1.26*O:C+1.18$ (Aiken et al., 2008).




Figure 11. a) Absorption Ångström exponent (470-660 nm), b) $MAC_{OA+BC,470}$ ($m^2/(-^1)$), for the same level legs shown in Figs. 9-10, similarly colored by flight date.




Figure 12. 24 September, 2016 (12.34°S, 11°E) vertical profiles of a) relative humidity (%; blue), zonal winds (m s^{-1} ; grey) and temperature ($^{\circ}\text{C}$), and b) organic aerosol to black carbon mass ratio (OA:BC; green), total nitrate to black carbon ratio (NO_3 :BC; blue) averaged every 500 m (approximately 2 minutes of data), and c) total mass concentration (OA + BC + SO_4 + NO_3 + NH_4 in $\mu\text{g m}^{-3}$; 1Hz resolution) colored by aerosol age. d) HYSPLIT trajectories, [initialized on 12 UTC of 24 September, 2016 with markers placed at 0 UTC of preceding days](#), superimposed on map of fires detected between 9/17/2016-9/24/2016.




Figure 13. a) $\frac{BC}{\Delta CO}$ ratios (dimensionless) as a function of f_{44} for the 9/24/2016 flight. The number of minutes contributing to each f_{44} bin is stated at bottom of panel. b) same for OA:BC. Whiskers represent the 10th and 90th percentiles, boxes illustrate the 75th and 25th percentiles with a line indicating the median and a red filled circle the mean. $OA > 20 \mu\text{g m}^{-3}$ only.




Figure 14. a) 24 September, 2016 LDMA-derived number particle size distribution for three *f44* bins (0.15-0.18 - red; 0.18-0.21-blue; 0.21-0.24 - purple), divided by the BC mass. b) same as a) but for the LDMA-derived **volume**-particle size distribution. c) LDMA and CPC particle number concentration as a function of *f44* and d) ratio of non-BC total mass ($\text{OA} + \text{SO}_4 + \text{NO}_3 + \text{NH}_4$) to BC, as a function of the three *f44* bins. All data are selected from $\text{OA} > 20 \mu\text{g m}^{-3}$.




Figure 15. a) $\text{NO}_3:\text{BC}$ and c) $\text{OA}:\text{BC}$ mass ratios for the 6 selected flights as a function of relative humidity, for $\text{OA} > 20 \mu\text{g m}^{-3}$ at STP. The 10th, 25th, median, 75th and 90th percentiles are indicated using box-whiskers, the mean with solid red circle and marker. b) corresponding distribution of aerosol ages within each relative humidity range, with the y-axis indicating the number of 1-sec samples.