



Strong particle production and condensational growth in the upper troposphere sustained by biogenic VOCs from the canopy of the Amazon Basin

5 Yunfan Liu¹, Hang Su², Siwen Wang¹, Chao Wei², Wei Tao², Mira L. Pöhlker^{2,6,7},
Christopher Pöhlker², Bruna A. Holanda², Ovid O. Krüger², Thorsten Hoffmann⁵, Manfred
Wendisch⁷, Paulo Artaxo⁸, Ulrich Pöschl², Meinrat O. Andreae^{3,4}, and Yafang Cheng¹

¹Minerva Research Group, Max Planck Institute for Chemistry, 55128 Mainz, Germany

10 ²Multiphase Chemistry Department, Max Planck Institute for Chemistry, 55128 Mainz, Germany

³Max Planck Institute for Chemistry, 55128 Mainz, Germany

⁴Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, USA

⁵Institute of Inorganic and Analytical Chemistry, Johannes Gutenberg University Mainz, Mainz, Germany

15 ⁶Experimental Aerosol and Cloud Microphysics Department, Leibniz Institute for Tropospheric Research,
Leipzig, Germany

⁷Faculty of Physics and Earth Sciences, Leipzig Institute for Meteorology, Leipzig University, Leipzig,
Germany

⁸Institute of Physics, University of São Paulo, São Paulo, Brazil

20 * Correspondence to: yafang.cheng@mpic.de

Abstract. Nucleation and condensation associated with biogenic volatile organic compounds (BVOCs) are important aerosol formation pathways, yet their contribution to the upper tropospheric aerosols remains inconclusive, hindering the understanding of aerosol climate effects. Here, we develop new schemes describing
25 these organic aerosol formation processes in the WRF-Chem model and investigate their impact on the abundance of cloud condensation nuclei (CCN) in the upper troposphere (UT) over the Amazon Basin. We find that the new schemes significantly increase the simulated CCN number concentrations in the UT (e.g., up to ~400 cm⁻³ at 0.52% supersaturation) and greatly improve the agreement with the aircraft observations. Organic condensation enhances the simulated CCN concentration by 90% through promoting particle growth, while
30 organic nucleation, by replenishing new particles, contributes an additional 14%. Deep convection determines the rate of these organic aerosol formation processes in the UT through controlling the upward transport of biogenic precursors (i.e., BVOCs). This finding emphasizes the importance of the biosphere-atmosphere coupling in regulating upper tropospheric aerosol concentrations over the tropical forest and calls for attention to its potential role in anthropogenic climate change.



35 1 Introduction

Atmospheric aerosol particles can influence the Earth's climate by acting as cloud condensation nuclei (CCN), among other pathways. The CCN residing in the upper troposphere (UT), which have been repeatedly observed in a substantial amount over the globe (e.g., Minikin et al., 2003; Andreae et al., 2018, and references therein; Williamson et al., 2019), not only constitute an important aerosol source for the lower troposphere
40 (Wang et al., 2016a; Williamson et al., 2019), but also can directly be activated into cloud droplets through in-cloud activation and thus alter the cloud properties (Paluch and Knight, 1984; Fan et al., 2018). However, the formation mechanisms for the upper tropospheric CCN are poorly understood, which impedes their representation in models and the assessment of their climate effects (Heald et al., 2011; Watson-Parris et al., 2019).

Large concentrations of cloud active aerosol particles were detected in the UT over the Amazon by aircraft observations during the ACRIDICON-CHUVA campaign (Wendisch et al., 2016; Andreae et al., 2018). Chemical analysis demonstrated that their composition is dominated by organic compounds with signatures of secondary organic aerosol (SOA) related to the oxidation of biogenic volatile organic compounds (BVOCs; Schulz et al., 2018), yet, detailed processes driving the biogenic SOA formation remain inconclusive. Generally,
50 two mechanisms may promote the CCN production from biogenic SOA in the UT. The first relates to organic new particle formation (NPF), where aerosol particles can form out of nucleation of highly oxygenated molecules (HOMs) oxidized from biogenic organic vapors such as α -pinene and β -pinene (Burkholder et al., 2007; Kirkby et al., 2016) and subsequently grow to larger sizes. The pure organic NPF can notably affect the atmospheric CCN budget in the planetary boundary layer (PBL; Gordon et al., 2016). Alternatively, if there are
55 enough preexisting fine particles in the UT from transport or inorganic nucleation, the condensation of low volatile organic compounds (LVOCs) produced by BVOCs oxidation onto the preexisting particles can also increase the CCN number (D'Andrea et al., 2013). However, a quantitative assessment of the BVOC-driven nucleation and condensation processes is lacking (Tröstl et al., 2016; Williamson et al., 2019). To what extent these two processes account for the CCN production in the Amazon UT, and whether these processes and CCN
60 formation proceed in the UT or if CCN form in the lower troposphere and then are transported upwards, is not known.

Motivated by these questions, this study implements the laboratory-based organic nucleation (HOMs-induced nucleation) and condensation processes into the WRF-Chem model and conducts simulations to quantify the CCN production from these BVOCs-driven SOA formation pathways in the Amazon UT. We
65 explore the upper tropospheric biogenic SOA formation mechanisms in terms of the involved atmospheric physical and chemical processes and on a diurnal variation scale.

2 Results

2.1 Simulation of number concentration of CCN and total aerosol particles in the upper troposphere

The organic nucleation mechanism in this study focuses on pure organic nucleation, including neutral and
70 ion-induced processes, triggered by HOMs from α -pinene and β -pinene oxidation (Kirkby et al., 2016), as it was found dominant among organic nucleation pathways (Zhu et al., 2019). The organic condensation mechanism addresses the dynamic condensation of LVOCs oxidized from α -pinene, β -pinene, and isoprene (Mann et al.




2010) as well as HOMs. For an accurate representation of HOMs concentrations, we adopted the Common
Reactive Intermediates gas-phase Mechanism (CRIMEch) scheme (Jenkin et al., 2008) with an explicit
75 description of α -pinene and β -pinene oxidation and calculated the HOMs concentration dynamically, which
circumvented the uncertainties related to species approximation in other chemical schemes (Riccobono et al.,
2014; Zhu et al., 2019) and the bulk assumption of an equilibrium state of HOMs (Gordon et al., 2016; Tröstl et
al., 2016), respectively. The temperature effects on nucleation rate and LVOC yields were included in the model
according to a combination of nucleation theory (Yu et al., 2017) and experimental results (Sahhaf et al., 2008).
80 Details of the model description can be found in Appendix Sections A1.1 and A1.2.

To disentangle the organic nucleation effect from the organic condensation influence, we performed the
following sensitivity simulations:

- BASE, the default WRF-Chem simulation with H₂SO₄-H₂O binary nucleation and without biogenic
nucleation or condensation;
- 85 - CTRL, in which both the newly developed nucleation and condensation modules were added;
- OCD, which only added the organic condensation;
- BNUnoT and OCDnoT, which excluded the temperature effect on the nucleation rate and the LVOC
yields, respectively;
- NoOH which was based on CTRL but without the HOMs formation from OH oxidation.

90 Settings for all scenarios are summarized in Table A3. The simulations were conducted for two nested domains
covering the Amazon with a horizontal resolution of 75 km and 15 km, respectively (Fig. A1) from 1 September
to 1 October 2014. Aircraft measurements of aerosol number concentration profiles reaching up to 15 km
altitude, close to the tropopause (18 km, Wendisch et al., 2016), sampled during the ACIRIDICON-CHUVA
campaign (Wendisch et al., 2016; Andreae et al., 2018) were used to evaluate the model results. Details of the
95 model configuration and observation dataset are documented in Appendix Section A1.3–A2.




100 **Figure 1.** Comparison of observed and simulated number concentrations of (a) condensation nuclei (CN, total
aerosol population) with diameters above 20 nm, and (b) cloud condensation nuclei (CCN) at 0.52%
supersaturation. The aerosol concentrations are at standard temperature and pressure (STP; 273.15 K and 1000
hPa). Planetary boundary layer (PBL), middle troposphere (MT), and upper troposphere (UT) are defined as the
altitude range of 0–4 km, 5–8 km, 9–15 km, respectively. The standard deviations of the observations are
105 provided in Table A4.

Figure 1 shows the average profiles of the number concentration of aerosol particles (also called
condensation nuclei, CN) with a diameter above 20 nm and of the CCN at a supersaturation of 0.52% from
aircraft observation (Fig. A1) and model simulations. The size of the CCN is approximately 90 nm in diameter,
calculated according to the algorithm of Su et al. (2010; Section A3). Compared with the observations, the
110 BASE case appears to reproduce the vertical distribution of CN in general (Section A3), but the CCN
concentration in this simulation shows noticeable biases, especially in the UT (9–15 km) where the model
underestimates the observed CCN number by up to 500 cm^{-3} (58%). Considering the different size ranges in
which CN and CCN reside, the large model underestimation in CCN may suggest insufficient growth of the
smaller particles in the UT.
115

When adding the particle growth from the LVOCs condensation into the model (i.e., the OCD case), the
simulated CCN number in the UT rises notably, with an increase of 310 cm^{-3} (about 90% relative to the BASE
case; Table A4). However, the larger particles from the condensation growth meanwhile deplete the nano-sized
particles, causing a dramatic drop in CN number concentrations from BASE to OCD. Though the non-cloud-
120 resolving resolution of the simulations may cause an excessive mixing of ultrafine particle-laden fresh cloud
outflows and their surrounding airmasses (Andreae et al., 2018) and thus aggravate the particle scavenging, the
considerable underestimation of the averaged CN number under a reasonable condensation growth in OCD is
strongly indicative of some missing NPF mechanisms (Zhu et al., 2019; Zhao et al., 2020). As expected, by
further taking into account the organic nucleation (i.e., the CTRL case), the simulated CN number




125 concentrations are enhanced substantially (2100 cm^{-3} , over 50%; Table A4) relative to the OCD case and in
markedly better agreement with the observation, while the CCN number concentrations in the model show a
relatively weaker increase (90 cm^{-3} , about 14%; Table A4). Thus, in total, both the BVOCs-driven organic
nucleation and condensation play important roles in maintaining the particle population and size distribution in
the UT (Fig. A7). The HOMs nucleation effectively increases the CN number by replenishing new nano-sized
130 particles, yet its contribution to the CCN, which are mainly in accumulation mode, is relatively limited. In
contrast, the organic condensation causes efficient particle growth and, therefore, greatly enhances the CCN
population.



135 **Figure 2.** Profiles of (a) contribution to organic aerosol (OA) mass from secondary organic aerosol production
processes and (b) observed and simulated OA mass averaged along all the flight tracks. The aerosol
concentrations are at STP.

140 Consistent with the CCN behavior, organic aerosol (OA), the dominant aerosol component over the
simulated region and period (Andreae et al., 2018), is also underestimated in the UT in default WRF-Chem
(BASE), but improved close to observation when the biogenic SOA formation is included (Fig. 2b). The
condensation of LVOCs plays a predominant role in the OA mass production among all processes, while the
other two formation pathways, especially the HOMs nucleation, contribute little (Fig. 2a). This also explains
why the organic condensation can cause profound particle growth while the HOMs nucleation works mainly to
145 increase the number of small particles. The OA production from the SOA processes in the UT shows a similar
vertical pattern to that of the OA mass (Fig. 2b), implying local origins of the upper tropospheric CCN.

2.2 Factors influencing organic-driven particle formation and growth in the upper troposphere



150 **Figure 3.** Spatial distribution of (a) α -pinene, (b) HOMs, (c) organic nucleation rate, (d) isoprene, (e) SOA production rate by LVOCs, (f) CCN at 0.52% supersaturation, and (g) condensation sink of HOMs in the upper troposphere (UT) averaged over 1 Sep–1 Oct 2014. Also shown are (h) upward α -pinene flux to the UT and (i) precipitation rate averaged over the same period. The black rectangle in (a) denotes the region of the Central Amazon for further analysis in this study.


155

Figure 3 shows the horizontal distributions of the monthly mean biogenic organic precursors (α -pinene and isoprene; α -pinene is used here as a surrogate for the organic nucleation precursors) and the HOMs nucleation (J_{org}) and LVOCs condensation rates (SOA_LVOCs) in the UT. The organic nucleation and condensation distributions closely follow that of the biogenic precursors (Fig. 3a, c, d, e) but not the oxidants (O_3 and OH; Fig. A8), suggesting the upper tropospheric BVOCs concentration as the limiting factor for the organic nucleation/condensation in the Amazon UT. The region with high BVOC concentrations in the UT is different from the α -pinene distribution in the PBL (Fig. A9), but identical to the precipitation pattern as well as the large upward α -pinene flux (Fig. 3i, h), showing a necessary role of deep convection transport in the BVOCs availability in the UT.

160



165 The J_{org} can reach over $0.1 \text{ cm}^{-3} \text{ s}^{-1}$ in the UT (Fig. 3) while the PBL (0–4 km) and middle troposphere (MT;
5–8 km) show low values (Fig. A9–A10), even though the α -pinene concentration in PBL is a magnitude larger
than in the UT. Such high upper tropospheric J_{org} is favored by not only the low sink of HOMs (CS) but also the
low temperature in the UT (Fig. 3g). When the temperature dependence of J_{org} (Section A1.2) is not considered,
the J_{org} in the UT is much lower than in the model run with the temperature effect (Fig. A11). The overall
170 magnitude of J_{org} is lower than simulated previously in the Amazon (Zhu et al., 2019), possibly due to the
consideration of the ion sink in this study. For the SOA production from LVOCs condensation, the bulk
assumption of the LVOC yields used previously in the boreal forest (Scott et al. 2014) fails to reproduce the
observed OA mass due to different conditions in the tropics (Section A4). A temperature-dependent correction
of LVOC yields based on laboratory experiments (Saathoff et al., 2009) is necessary for correcting the OA
175 simulation bias associated with the bulk LVOC yields assumption (Fig. 2b). The low temperature in the UT also
serves as a favorable condition for the SOA production from LVOCs condensation.



180 **Figure 4.** Modeled temporal variation of the rate and precursors of organic nucleation and condensation as well
 as meteorological variables. (a) Daily variation of regionally averaged nucleation and condensation rates,
 number concentration of CN and CCN at 0.52% supersaturation (CCN(0.52%)), vertical wind at 8 km (w), and
 precipitation. The dots above the figure mark the upper troposphere (UT) biogenic SOA episodes. The diurnal
 patterns of nucleation and condensation rate, number concentration of CN and CCN(0.52%), condensation sink
 185 of HOMs, and mixing ratio of α -pinene, isoprene, O_3 , and OH average in the UT as well as the vertical wind at 8
 km and precipitation for the (b) UT biogenic SOA episode days and (c) non-episode days. The analyzed region
 is indicated by the black rectangle in Fig. 3a. The bars denote the standard error.

As the biogenic SOA formation in the UT exhibits prominent daily variation, we defined days with J_{org}
 190 greater than $1\text{e}^{-3} \text{cm}^{-3} \text{s}^{-1}$ and simultaneous condensational SOA production rate larger than $0.05 \mu\text{g m}^{-3} \text{h}^{-1}$ as UT
 biogenic SOA episodes (Fig. 4a) to identify their characteristics. The biogenic SOA episodes in the UT emerge



with stronger convection where strong vertical wind effectively transports biogenic precursors to the UT and fosters SOA formation (Fig. 4b). In contrast, the biogenic SOA formation in the PBL tends to be suppressed by deep convection (Fig. A14) due to the restrained emission and oxidation of BVOCs by reduced incident solar radiation in cloudy weather (Fig. A13) as well as the BVOCs dilution near the surface by low-BVOCs air from the MT.

The organic nucleation and condensation rate demonstrate a clear diurnal cycle, with significant occurrence in the daytime. As the oxidant in the HOMs and LVOCs production, OH has reaction rates several orders faster than O_3 (Section A1.2; Atkinson et al., 2006), making it the dominant oxidant, which is also evidenced as the sensitivity study switching off the OH oxidation (NoOH) shows significantly reduced HOMs concentrations (Fig. A15). Thus, the photolysis origin of OH (Fig. 4b–4c) explains the daytime burst of the J_{org} and condensation rate. A detailed mechanism tracking the diurnal variation of the upper tropospheric CCN production can be drawn. The nighttime convective activity as an extension of the former daytime convection contributes to the upper tropospheric accumulation of biogenic precursors. With the onset of solar radiation, the photochemical reactions start to produce OH, which efficiently oxidizes BVOCs to form HOMs and LVOCs and then triggers the organic nucleation and condensation. The CN and CCN concentrations increase accordingly and reach high levels in the afternoon which is also the typical time for the vigorous development of local convective clouds and thereby favors potential interactions between upper tropospheric CCN and clouds.

3 Conclusions

In this work, we developed a new organic nucleation and condensation scheme for the WRF-Chem model to investigate the CN and CCN production in the UT (upper troposphere) by BVOCs-driven SOA formation over a forest canopy region, the Amazon Basin.

The model evaluation against aircraft measurements shows that including the BVOCs-driven SOA formation significantly improves the model agreement with the measured upper tropospheric CCN (at 0.52% supersaturation) number concentrations by elevating the simulated values up to $\sim 400 \text{ cm}^{-3}$. Individually, the organic condensation drives efficient particle growth and promotes the CCN concentration in the UT by about 90%. With the nano-sized particles from H_2SO_4 - H_2O binary nucleation scavenged under sufficient particle growth, the organic nucleation serves to replenish nano-particles and enhances the upper tropospheric CN and CCN number concentration by over 50% and 14%, respectively. The rates of SOA processes in the UT depend on deep convection for its vertical transport, and are favored by low condensation sink and temperature at high altitudes.

The considerable CCN production in the UT by BVOCs-driven organic processes underlines the important regulation of biospheric BVOCs on the high-altitude aerosol concentrations. Considering the climate significance of these upper tropospheric aerosols, the biosphere-atmosphere coupling should be emphasized in the context of climate change, not only for its possible impact on the preindustrial reference state (Gorden et al., 2016), but also for its feedback to climate under the future anthropogenic influence (e.g., deforestation) and climate change.



Author contributions

YC designed and led the study. YL and YC conducted the model development. YL performed the model simulation and analyzed the data. YL, YC, and HS interpreted the results. MOA and UP contributed to discussing the results. SW, CW, and WT supported model simulation and data visualization. MOA, UP, PA, and MW coordinated the ACRIDICON-CHUVA observation field campaign. MLP, CP, OOK, and BAH provided the observation data from HALO for model comparison. YL wrote the paper with input from all coauthors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This study is supported by the Max Planck Society (MPG). YC and YL acknowledge the Minerva Program of MPG. We also gratefully acknowledge the observational data and data providers from the ACRIDICON-CHUVA campaign, which was supported by the Max Planck Society (MPG), the German Science Foundation (DFG Priority Program SPP 1294), the German Aerospace Center (DLR), the FAPESP (Sao Paulo Research Foundation) grants 2009/15235-8, 2013/05014-0, and 2017/17047-0, and a wide range of other institutional partners.

Reference

- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523–8546, doi:10.5194/acp-19-8523-2019, 2019.
- Andreae, M. O., Acevedo, O. C., Araújo, A., Artaxo, P., Barbosa, C. G. G., Barbosa, H. M. J., Brito, J., Carbone, S., Chi, X., Cintra, B. B. L., da Silva, N. F., Dias, N. L., Dias-Júnior, C. Q., Ditas, F., Ditz, R., Godoi, A. F. L., Godoi, R. H. M., Heimann, M., Hoffmann, T., Kesselmeier, J., Könemann, T., Krüger, M. L., Lavric, J. V., Manzi, A. O., Lopes, A. P., Martins, D. L., Mikhailov, E. F., Moran-Zuloaga, D., Nelson, B. W., Nölscher, A. C., Santos Nogueira, D., Piedade, M. T. F., Pöhlker, C., Pöschl, U., Quesada, C. A., Rizzo, L. V., Ro, C. U., Ruckteschler, N., Sá, L. D. A., de Oliveira Sá, M., Sales, C. B., dos Santos, R. M. N., Saturno, J., Schöngart, J., Sörgel, M., de Souza, C. M., de Souza, R. A. F., Su, H., Targhetta, N., Tóta, J., Trebs, I., Trumbore, S., van Eijck, A., Walter, D., Wang, Z., Weber, B., Williams, J., Winderlich, J., Wittmann, F., Wolff, S., and Yáñez-Serrano, A. M.: The Amazon Tall Tower Observatory (ATTO): overview of pilot measurements on ecosystem ecology, meteorology, trace gases, and aerosols, *Atmos. Chem. Phys.*, 15, 10723–10776, doi:10.5194/acp-15-10723-2015, 2015.
- Andreae, M. O., Afchine, A., Albrecht, R., Holanda, B. A., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Cecchini, M. A., Costa, A., Dollner, M., Fütterer, D., Järvinen, E., Jurkat, T., Klimach, T., Konemann, T., Knote, C., Krämer, M., Krisna, T., Machado, L. A. T., Mertes, S., Minikin, A., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Sauer, D., Schlager, H., Schnaiter, M., Schneider, J., Schulz, C., Spanu, A., Sperling, V. B., Voigt, C., Walser, A., Wang, J., Weinzierl, B., Wendisch, M., and Ziereis, H.: Aerosol



- characteristics and particle production in the upper troposphere over the Amazon Basin, *Atmos. Chem. Phys.*, 18, 921–961, doi:10.5194/acp-18-921-2018, 2018.
- 265 Andreae, M. O. and Andreae, T. W.: The cycle of biogenic sulfur compounds over the Amazon Basin: 1. Dry season, *J. Geophys. Res.-Atmos.*, 93, 1487–1497, <https://doi.org/10.1029/JD093iD02p01487>, 1988.
- Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., Hodnebrog, Ø., Denier van der Gon, H., and McFiggans, G.: Gaseous chemistry and aerosol mechanism developments for version 3.5.1 of the online regional model, WRF-Chem, *Geoscientific Model Development*, 7, 2557–2579, doi:10.5194/gmd-7-2557-2014, 2014.
- 270 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, doi:10.5194/acp-6-3625-2006, 2006.
- Burkholder, J. B., Baynard, T., Ravishankara, A. R., and Lovejoy, E. R.: Particle nucleation following the O₃ and OH initiated oxidation of α -pinene and β -pinene between 278 and 320 K, *Journal of Geophysical Research: Atmospheres*, 112, doi:10.1029/2006jd007783, 2007.
- 275 Chapman, E. G., Gustafson, W. I., Easter, R. C., Barnard, J. C., Ghan, S. J., Pekour, M. S., and Fast, J. D.: Coupling aerosol-cloud-radiative processes in the WRF-Chem model: Investigating the radiative impact of elevated point sources, *Atmos. Chem. Phys.*, 9, 945–964, doi:10.5194/acp-9-945-2009, 2009.
- 280 Cui, Y. Y., Hodzic, A., Smith, J. N., Ortega, J., Brioude, J., Matsui, H., Levin, E. J. T., Turnipseed, A., Winkler, P., and de Foy, B.: Modeling ultrafine particle growth at a pine forest site influenced by anthropogenic pollution during BEACHON-RoMBAS 2011, *Atmos. Chem. Phys.*, 14, 11011–11029, doi:10.5194/acp-14-11011-2014, 2014.
- D’Andrea, S. D., Häkkinen, S. A. K., Westervelt, D. M., Kuang, C., Levin, E. J. T., Kanawade, V. P., Leaitch, 285 W. R., Spracklen, D. V., Riipinen, I., and Pierce, J. R.: Understanding global secondary organic aerosol amount and size-resolved condensational behavior, *Atmos. Chem. Phys.*, 13, 11519 – 11534, doi:10.5194/acp-13-11519-2013, 2013.
- Dunne, E. M., Gordon, H., Kurten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle, K. J., Adamov, A., Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., 290 Curtius, J., Dommen, J., Donahue, N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J., Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C. L., Riccobono, F., Richards, N. A., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M., Sipila, M., Smith, J. N., Stozkhov, Y., Tome, A., Trostl, J., 295 Wagner, P. E., Wimmer, D., Winkler, P. M., Worsnop, D. R., and Carslaw, K. S.: Global atmospheric particle formation from CERN CLOUD measurements, *Science*, 354, 1119–1124, doi:10.1126/science.aaf2649, 2016.
- Dunne, E. M., Gordon, H., Kurten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle, K. J., Adamov, A., Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., 300 Curtius, J., Dommen, J., Donahue, N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J.,



- Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C. L., Riccobono, F., Richards, N. A., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M., Sipila, M., Smith, J. N., Stozhkov, Y., Tome, A., Trostl, J., Wagner, P. E., Wimmer, D., Winkler, P. M., Worsnop, D. R., and Carslaw, K. S.: Global atmospheric particle formation from CERN CLOUD measurements, *Science*, 354, 1119–1124, doi:10.1126/science.aaf2649, 2016.
- Easter, R. C., Ghan, S. J., Zhang, Y., Saylor, R. D., Chapman, E. G., Laulainen, N. S., Abdul-Razzak, H., Leung, L. R., Bian, X. D., and Zaveri, R. A.: MIRAGE: Model description and evaluation of aerosols and trace gases, *J. Geophys. Res.-Atmos.*, 109, doi:10.1029/2004jd004571, 2004.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J. F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010.
- Fan, J., Rosenfeld, D., Zhang, Y., Giangrande, S. E., Li, Z., Machado, L. A. T., Martin, S. T., Yang, Y., Wang, J., Artaxo, P., Barbosa, H. M. J., Braga, R. C., Comstock, J. M., Feng, Z., Gao, W., Gomes, H. B., Mei, F., Pöhlker, C., Pöhlker, M. L., Pöschl, U., and de Souza, R. A. F.: Substantial convection and precipitation enhancements by ultrafine aerosol particles, *Science*, 359, 411–418, doi:10.1126/science.aan8461, 2018.
- Fast, J. D., Gustafson, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *J. Geophys. Res.*, 111, doi:10.1029/2005jd006721, 2006.
- Freitas, S. R., Longo, K. M., Chatfield, R., Latham, D., Dias, M. A. F. S., Andreae, M. O., Prins, E., Santos, J. C., Gielow, R., and Carvalho, J. A.: Including the sub-grid scale plume rise of vegetation fires in low resolution atmospheric transport models, *Atmos. Chem. Phys.*, 7, 3385–3398, doi:10.5194/acp-7-3385-2007, 2007.
- Gordon, H., Sengupta, K., Rap, A., Duplissy, J., Frege, C., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Trostl, J., Nieminen, T., Ortega, I. K., Wagner, R., Dunne, E. M., Adamov, A., Amorim, A., Bernhammer, A. K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J. S., Dias, A., Ehrhart, S., Fischer, L., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Kirkby, J., Krapf, M., Kurten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Monks, S. A., Onnela, A., Perakyla, O., Piel, F., Petaja, T., Praplan, A. P., Pringle, K. J., Richards, N. A., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sharma, S., Sipila, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tome, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Curtius, J., and Carslaw, K. S.: Reduced anthropogenic aerosol radiative forcing caused by biogenic new particle formation, *Proc. Natl. Acad. Sci. U S A*, 113, 12053–12058, doi:10.1073/pnas.1602360113, 2016.



- 340 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmospheric Environment*, 39, 6957–6975, 10.1016/j.atmosenv.2005.04.027, 2005.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and
345 updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.
- Hardacre, C., Mulcahy, J. P., Pope, R. J., Jones, C. G., Rumbold, S. T., Li, C., Johnson, C. and Turnock, S. T.: Evaluation of SO₂, SO₄²⁻ and an updated SO₂ dry deposition parameterization in UKESM1. *Atmos. Chem. Phys.*, 21 (24), 18465–18497, doi:10.5194/acp-21-18465-2021, 2021.
- 350 Heald, C. L., Coe, H., Jimenez, J. L., Weber, R. J., Bahreini, R., Middlebrook, A. M., Russell, L. M., Jolleys, M., Fu, T.-M., Allan, J. D., Bower, K. N., Capes, G., Crosier, J., Morgan, W. T., Robinson, N. H., Williams, P. I., Cubison, M. J., DeCarlo, P. F., and Dunlea, E. J.: Exploring the vertical profile of atmospheric organic aerosol: comparing 17 aircraft field campaigns with a global model, *Atmos. Chem. Phys. Discuss.*, 11, 25371 – 25425, doi:10.5194/acpd-11-25371-2011, 2011.
- 355 Jenkin, M. E., Watson, L. A., Utembe, S. R., and Shallcross, D. E.: A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 1: Gas phase mechanism development, *Atmospheric Environment*, 42, 7185–7195, doi:10.1016/j.atmosenv.2008.07.028, 2008.
- Kerminen, V.-M., Anttila, T., Lehtinen, K., and Kulmala, M.: Parameterization for Atmospheric New-Particle Formation: Application to a System Involving Sulfuric Acid and Condensable Water-Soluble Organic
360 Vapors, *Aerosol Science and Technology*, 38, 1001–1008, doi:10.1080/027868290519085, 2004.
- Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Trostl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A. K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J.,
365 Kim, J., Krapf, M., Kurten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Perakyla, O., Piel, F., Petaja, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipila, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tome, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N.
- 370 M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521–526, doi:10.1038/nature17953, 2016.
- Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from isoprene photooxidation under high-NO_x conditions, *Geophys. Res. Lett.*, 32, L18808, doi:10.1029/2005gl023637, 2005.
- 375 Kulmala, M.: How particles nucleate and grow, *Science*, 302, 1000–1001, doi:10.1126/science.1090848, 2003.
- Minikin, A., Petzold, A., Ström, J., Krejci, R., Seifert, M., Velthoven, P. v., Schlager, H., and Schumann, U.: Aircraft observations of the upper tropospheric fine particle aerosol in the Northern and Southern Hemispheres at midlatitudes, *Geophys. Res. Lett.*, 30, 1503, <https://doi.org/10.1029/2002gl016458>, 2003.



- 380 Paluch, I. R. and Knight, C. A.: Mixing and the evolution of cloud droplet size spectra in a vigorous continental cumulus, *J. Atmos. Sci.*, 41, 1801–1815, 1984.
- Paralovo, S. L., Barbosa, C. G. G., Carneiro, I. P. S., Kurzlop, P., Borillo, G. C., Schiochet, M. F. C., Godoi, A. F. L., Yamamoto, C. I., de Souza, R. A. F., Andreoli, R. V., Ribeiro, I. O., Manzi, A. O., Kourtchev, I., Bustillos, J. O. V., Martin, S. T., and Godoi, R. H. M.: Observations of particulate matter, NO₂, SO₂, O₃, H₂S and selected VOCs at a semi-urban environment in the Amazon region, *Sci. Total Environ.*, 650, 996–1006, <https://doi.org/10.1016/j.scitotenv.2018.09.073>, 2019.
- 385 Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, 2007, <http://www.atmos-chem-phys.net/7/1961/2007/>.
- Ramsay, R., Di Marco, C. F., Sörgel, M., Heal, M. R., Carbone, S., Artaxo, P., de Araújo, A. C., Sá, M., Pöhlker, C., Lavric, J., Andreae, M. O., and Nemitz, E.: Concentrations and biosphere–atmosphere fluxes of inorganic trace gases and associated ionic aerosol counterparts over the Amazon rainforest, *Atmos. Chem. Phys.*, 20, 15551–15584, <https://doi.org/10.5194/acp-20-15551-2020>, 2020.
- 390 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kurten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petaja, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipila, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tome, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R., and Baltensperger, U.: Oxidation products of biogenic emissions contribute to nucleation of atmospheric particles, *Science*, 344, 717–721, doi:10.1126/science.1243527, 2014.
- 400 Scott, C. E., Rap, A., Spracklen, D. V., Forster, P. M., Carslaw, K. S., Mann, G. W., Pringle, K. J., Kivekas, N., Kulmala, M., Lihavainen, H., and Tunved, P.: The direct and indirect radiative effects of biogenic secondary organic aerosol, *Atmospheric Chemistry and Physics*, 14, 447–470, [10.5194/acp-14-447-2014](https://doi.org/10.5194/acp-14-447-2014), 2014.
- 405 Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer, A. K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kurten, A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Mohler, O., Nieminen, T., Onnela, A., Petaja, T., Piel, F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipila, M., Smith, J. N., Steiner, G., Tome, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop, D. R., Donahue, N. M., and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533, 527–531, doi:10.1038/nature18271, 2016.
- 415 Wang, J., Krejci, R., Giangrande, S., Kuang, C., Barbosa, H. M., Brito, J., Carbone, S., Chi, X., Comstock, J., Ditas, F., Lavric, J., Manninen, H. E., Mei, F., Moran-Zuloaga, D., Pöhlker, C., Pöhlker, M. L., Saturno, J.,



- 420 Schmid, B., Souza, R. A., Springston, S. R., Tomlinson, J. M., Toto, T., Walter, D., Wimmer, D., Smith, J. N., Kulmala, M., Machado, L. A., Artaxo, P., Andreae, M. O., Petaja, T., and Martin, S. T.: Amazon boundary layer aerosol concentration sustained by vertical transport during rainfall, *Nature*, 539, 416–419, doi:10.1038/nature19819, 2016a.
- 425 Wang, Q. Q., Saturno, J., Chi, X., Walter, D., Lavric, J. V., Moran-Zuloaga, D., Ditas, F., Pohlker, C., Brito, J., Carbone, S., Artaxo, P., and Andreae, M. O.: Modeling investigation of light-absorbing aerosols in the Amazon Basin during the wet season, *Atmospheric Chemistry and Physics*, 16, 14775–14794, 10.5194/acp-16-14775-2016, 2016b.
- Watson, L. A., Shallcross, D. E., Utembe, S. R., and Jenkin, M. E.: A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 2: Gas phase mechanism reduction, *Atmospheric Environment*, 42, 7196–7204, doi:10.1016/j.atmosenv.2008.07.034, 2008.
- 430 Watson-Parris, D., Schutgens, N., Reddington, C., Pringle, K. J., Liu, D. T., Allan, J. D., Coe, H., Carslaw, K. S., & Stier, P. (2019). In situ constraints on the vertical distribution of global aerosol. *Atmospheric Chemistry and Physics*, 19(18), 11765–11790. <https://doi.org/10.5194/acp-19-11765-2019>
- 435 Wendisch, M., Pöschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araújo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jäkel, E., Järvinen, E., Jurkat, T., Kanter, S., Kästner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kölling, T., Krämer, M., Krüger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Münch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: ACRIDICON-CHUVA Campaign: Studying Tropical Deep Convective Clouds and Precipitation over Amazonia Using the New German Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 97, 1885–1908, doi:10.1175/bams-d-14-00255.1, 2016.
- 445 Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmospheric Environment* (1967), 23, 1293–1304, doi:10.1016/0004-6981(89)90153-4, 1989.
- Wexler, A. S., Lurmann, F. W., and Seinfeld, J. H.: Modelling urban and regional aerosols—I. model development, *Atmospheric Environment*, 28, 531–546, doi:10.1016/1352-2310(94)90129-5, 1994.
- 450 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, doi:10.5194/gmd-4-625-2011, 2011.
- 455 Williamson, C. J., Kupc, A., Axisa, D., Bilsback, K. R., Bui, T., Campuzano-Jost, P., Dollner, M., Froyd, K. D., Hodshire, A. L., Jimenez, J. L., Kodros, J. K., Luo, G., Murphy, D. M., Nault, B. A., Ray, E. A., Weinzierl, B., Wilson, J. C., Yu, F., Yu, P., Pierce, J. R., and Brock, C. A.: A large source of cloud condensation nuclei from new particle formation in the tropics, *Nature*, 574, 399–403, doi:10.1038/s41586-019-1638-9, 2019.



- 460 Yu, F. Q., Luo, G., Nadykto, A. B., and Herb, J.: Impact of temperature dependence on the possible contribution
of organics to new particle formation in the atmosphere, *Atmos. Chem. Phys.*, 17, 4997–5005,
doi:10.5194/acp-17-4997-2017, 2017.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and
Chemistry (MOSAIC), *J. Geophys. Res.-Atmos.*, 113, D13204, doi:10.1029/2007jd008782, 2008.
- 465 Zhao, B., Shrivastava, M., Donahue, N. M., Gordon, H., Schervish, M., Shilling, J. E., Zaveri, R. A., Wang, J.,
Andreae, M. O., Zhao, C., Gaudet, B., Liu, Y., Fan, J., and Fast, J. D.: High concentration of ultrafine
particles in the Amazon free troposphere produced by organic new particle formation, *P. Natl. Acad. Sci.*,
117, 25344, <https://doi.org/10.1073/pnas.2006716117>, 2020.
- Zhu, J. L., and Penner, J. E.: Global Modeling of Secondary Organic Aerosol With Organic Nucleation, *J.*
Geophys. Res.-Atmos., 124, 8260–8286, doi:10.1029/2019jd030414, 2019.
- 470 Zhu, J. L., Penner, J. E., Yu, F. Q., Sillman, S., Andreae, M. O., and Coe, H.: Decrease in radiative forcing by
organic aerosol nucleation, climate, and land use change, *Nat. Commun.*, 10, 423, doi:10.1038/s41467-019-
08407-7, 2019.



Appendix:

A1 Model development

475 A1.1 Model description

The Weather Research and Forecasting model coupled with chemistry (WRF-Chem), version 3.9.1, was utilized in this study to investigate the biogenic organic nucleation and condensation over the Amazon. WRF-Chem is a meteorology-chemistry online coupled regional model, which integrates meteorological, gas-phase, and aerosol fields by calculating transport of chemical species under the same dynamical system as
480 meteorological elements at each time step and considering complicated feedbacks between meteorological field and trace gases and aerosols (Grell et al., 2005).

Table A1. WRF-Chem configuration.

Atmospheric Process	WRF-Chem Option
Longwave radiation	RRTMG
Shortwave radiation	RRTMG
Surface layer	Monin-Obukhov
Land surface	RUC
Boundary layer	YSU
Microphysics	Lin et al.
Cumulus	Grell-Devenyi ensemble
Gas-phase chemistry	CRIMech
Aerosol module	MOSAIC
Aqueous-phase chemistry	Fahey and Pandis
Photolysis	Fast-J
Anthropogenic emissions	EDGAR-HTAP V2
Biogenic emissions	MEGAN
Biomass burning emissions	FINNv1.5

485 **Table A2.** Description of aerosol size bins in MOSAIC.

Bin name	Default bins			Modified bins			
	Low ^a	High ^b	Center ^c	Bin name	Low	High	Center
				01	0.6	2.4	1.2
				02	2.4	10	5
				03	10	39	20
01	39	156	78	04	39	156	78
02	156	625	312	05	156	625	312
03	625	2500	1250	06	625	2500	1250
04	2500	10000	5000	07	2500	10000	5000

a, b, c: low boundary, high boundary, and geometric mean diameter of the bin in nm, respectively.

The WRF-Chem configurations used are listed in Table A1. We chose the Common Reactive Intermediates gas-phase Mechanism (CRIMech) scheme (Jenkin et al., 2008; Watson et al., 2008) with the KPP interface to
490 simulate the gas-phase chemistry. CRIMech contains up to 112 non-methane volatile organic compounds (VOCs), including α -pinene, β -pinene, and isoprene (Archer-Nicholls et al., 2014; Watson et al., 2008), and adopts 652 chemical reactions involving inorganic species, organic vapors, and their oxidation intermediates, based on the Master Chemical Mechanism (MCM). These chemical settings enable it to directly provide the gas-




phase precursors, i.e. α -pinene, β -pinene and isoprene, and chemical reactions for the biogenic SOA formation.

495 The Model for Simulating Aerosol Interactions and Chemistry (MOSAIC; Zaveri et al., 2008) was utilized to simulate aerosols, which uses discrete size bins to represent the aerosol size distribution. This study employed the 4-bin version with the size bins distributed as listed in Table A2. Aerosol species in MOSAIC include 5 inorganic ions, i.e. sulfate, nitrate, ammonium, sodium, chloride, and 3 unreactive primary aerosol species, i.e., black carbon (BC), particulate organic matter (POM), and other inorganics (OIN; Fast et al., 2006; Zaveri et al., 500 2008). In the WRF-Chem version 3.9.1 MOSAIC aerosol module, the binary nucleation of $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ is included (Wexler et al., 1994), while the organic nucleation and condensation are not accounted for. The coagulation process of particles and gas-particle partitioning were parameterized as described in Zaveri et al. (2008). The dry deposition of aerosol is parameterized by the updated resistance-in-series approach of Wesely (1989). The in- and below-cloud wet deposition takes place by activating aerosol from an interstitial state into 505 cloud-borne particles and calculating the washout rate due to precipitation, respectively (Chapman et al., 2009; Easter et al., 2004). The WRF-Chem model configurations used in this study allow aerosol-cloud interactions, following the way described by Fast et al. (2006) and Chapman et al. (2009); while the aerosol-radiation interactions were not included in the model simulations.

A1.2 Model improvement

510 Based on the aforementioned CRIMEch gas-phase chemistry scheme and MOSAIC aerosol scheme, a new module resolving the purely organic nucleation and condensation associated with BVOCs has been added to WRF-Chem, which provides a modeling tool to investigate biogenic SOA formation and its contribution to the upper tropospheric CCN.

The implementation of the new organic nucleation includes integrating the production of HOMs in the 515 CRIMEch gas-phase chemistry scheme, resetting the sectional bins in the aerosol model, and adding parameterizations of pure organic nucleation mechanisms by HOMs. Specifically, 4 reactions regarding HOM production from the oxidation of α -pinene and β -pinene by O_3 and OH were added to the CRIMEch mechanism based on reaction coefficients and yields suggested by laboratory experiments (Atkinson et al., 2006). Then, the condensation sink of HOMs was represented according to the algorithm of Kerminen et al. (2004). Instead of 520 parameterizing the HOMs concentration as the ratio of its production and condensation sink with the assumption that the HOMs were in a thermal equilibrium state (Kirkby et al., 2016; Gordon et al., 2016), the kinetic calculation of the HOMs production and condensation sink in this study enables a more accurate representation of the HOMs concentration.



530 The 4-bin MOSAIC scheme in WRF-Chem addresses aerosols with diameters from 39 nm to 10 μm , which does not cover the size range of newly formed particles, whose diameters are in the nanometer range (Kulmala, 2003). To explicitly represent the nucleation of vapor into particles, we extended the lower end of the aerosol



size range in the MOSAIC scheme from 39 nm to 0.6 nm by introducing 3 additional size bins whose boundaries are set following the same lognormal size distribution law as the original 4 bins (Table A2). Thus, the newly developed 7-bin MOSAIC scheme can resolve the formation and initial growth of new particles and assures a high computation efficiency. The 4-bin MOSAIC scheme includes the H₂SO₄-H₂O binary nucleation using a thermodynamic equilibrium parameterization where a critical concentration of H₂SO₄ is calculated based on air temperature and relative humidity and then the extra H₂SO₄ beyond this threshold is parameterized into aerosols centered at 78 nm (i.e., the 39–156 nm bin). The equilibrium method for describing H₂SO₄-H₂O binary nucleation is validated for the aerosol size above 10 nm. With extended aerosol size bins in the 7-bin MOSAIC scheme, we now applied the H₂SO₄-H₂O binary nucleation parameterization to the third bin (i.e. 10–39 nm) to not only assure the practical application of this nucleation parameterization (Wexler et al., 1994) but also keep the aerosol size range in agreement with the observations (i.e., starting from 20 nm).

Then, in addition to the existing H₂SO₄-H₂O binary nucleation, pure biogenic nucleation mechanisms induced by HOMs were integrated into the MOSAIC module. The mechanisms of organic nucleation were investigated in CLOUD (Cosmics Leaving OUTdoors Droplets) Chamber experiments (Kirkby et al., 2016), which suggested that the HOMs-induced pure organic nucleation rate (J_{org} , unit: cm⁻³ s⁻¹) can be represented by the combination of the neutral (J_n) and the ion-induced (J_{iin}) nucleation rate. The detailed parameterization of the J_n and J_{iin} is as follows:

$$J_{org} = J_n + J_{iin} \quad (\text{E. A1})$$

$$J_n = a_1 [HOM]^{a_2 + \frac{a_5}{[HOM]}} \quad (\text{E. A2})$$

$$J_{iin} = 2[n_{\pm}]a_3[HOM]^{a_4 + \frac{a_5}{[HOM]}} \quad (\text{E. A3})$$

where HOMs concentrations are in units of 10⁷ molecules per cubic centimeter and obtained by chemical kinetic calculations as described above; the a_i represent free parameters whose values were suggested by Kirkby et al. (2016) where a_1 , a_2 , a_3 , a_4 and a_5 equaled to 0.04001, 1.848, 0.001366, 1.566 and 0.1863, respectively.

n_{\pm} is the ion concentration produced from radon and galactic cosmic rays and is parameterized as:

$$[n_{\pm}] = \frac{(k_i^2 + 4\alpha q)^{0.5} - k_i}{2\alpha} \quad (\text{E. A4})$$

where q (in cm⁻³ s⁻¹) represents the ion-pair production rate and adopts the value of 10 cm⁻³ s⁻¹ (Horrak et al., 2008). α is the ion-ion recombination coefficient (in cm³ s⁻¹) and was set to 1.6e⁻⁶ cm³ s⁻¹ here. The ion loss rate, k_i , is due to the ion condensation sink (CS) onto aerosols and the ion-induced nucleation:

$$k_i = CS + \frac{J_{iin}}{2[n_{\pm}]} \quad (\text{E. A5})$$


In this study, the condensation sink term was calculated according to the empirical parameterization proposed by Tamm et al. (1991). The HOMs nucleation rate, J_{org} , is then modulated by temperature. Unlike the approximated temperature correction suggested in Dunne et al. (2016), a temperature dependence associated with the Gibbs free energy for forming the critical cluster based on the classical homogeneous nucleation theory (Yu et al., 2017) is used here. We applied temperature corrections to J_n and J_{iin} by multiplying them by a correction factor, $\exp(\Delta G_n/k^*(1/T-1/278))$ and $\exp(\Delta G_{iin}/k^*(1/T-1/278))$, respectively, and the ΔG_n and ΔG_{iin} are



570 based on smog chamber results (Kirkby et al., 2016). After organic nucleation, the newly formed particles were added into the smallest bin and underwent subsequent processes such as coagulation, transport, and deposition.

Additionally, a new module addressing the condensation of LVOCs was integrated into WRF-Chem. The LVOCs were oxidation products of α -pinene, β -pinene, and isoprene by O_3 , OH, and NO_3 . A yield of 13% for monoterpene oxidation products and 3% for isoprene oxidation products were used in Scott et al. (2014). Laboratory chamber experiments found a temperature dependence of the SOA yield from α -pinene oxidation (Saathoff et al., 2009). Therefore, instead of constant LVOC yields, temperature-corrected yields based on these laboratory experiment results (Saathoff et al., 2009) were applied in the model here.

A1.3 Numerical experiment design



580 **Figure A1.** Model domain and tracks of flight AC07 to AC20 from the ACRIDICON-CHUVA campaign. The outer map represents the parent domain with 75 km horizontal grid spacing, and the embedded square shows the extent of the 15 km resolution (d02) domain. The black rectangle is the same as the one marked in Fig. 3a, denoting the region of the Central Amazon for further analysis in this study.

585 In this study, two nested domains with a horizontal spacing of 75 km and 15 km were set up over South America (Fig. A1), with Domain1 covering most of the South American continent, while Domain2 is over the Amazon Basin area. Vertical layers of 29 levels extending from the ground surface to a height of 50 hPa were applied for all domains. The initial and boundary meteorological and chemical conditions were from the 6-hour



National Centers for Environmental Prediction (NCEP) Final Analysis (FNL) data and Model for Ozone and
 Related Chemical Tracers, version 4 (MOZART-4) global chemical transport model output (Emmons et al.,
 590 2010), respectively. The anthropogenic emissions with a resolution of $0.1^\circ \times 0.1^\circ$ were obtained from the global
 air pollution emission dataset, EDGAR-HTAPv2 (http://edgar.jrc.ec.europa.eu/htap_v2; Janssens-Maenhout et
 al., 2015). The Fire Inventory from NCAR version 1.5 (FINNv1.5; Wiedinmyer et al., 2011) provided the
 biomass burning emission, which is updated daily at 1-km resolution. The rise of fire plumes after emission was
 represented by a plume-rise parameterization (Freitas et al., 2007). The primary organic matter (POM) emission
 595 rate was calculated based on the OC emission by applying a ratio of 1.6 between the mass of POM and OC
 (Andreae, 2019). The biogenic emissions of NO and volatile organic compounds (VOCs) were generated online
 by the Model of Emissions of Gases and Aerosols from Nature (MEGAN; Guenther et al., 2012). Among the
 biogenic VOCs are the precursors, i.e. α -pinene, β -pinene, and isoprene, for the organic nucleation and
 condensation which are investigated in this study. MEGAN calculated the emission of biogenic gases based on
 600 the United States Geological Survey (USGS) land use category, temperature, and radiation, which were
 subsequently put into the corresponding chemical species in the CRIMEch gas-phase scheme as a source term.
 The simulation was conducted from 24 August to 1 October 2014, and the first 8 days of the simulation were
 used as spin up. The comparisons between model outputs and aircraft measurements in Section 2.1 are made
 with the results from Domain2. A rectangular area focusing on the Central Amazon, as shown in Fig. A1, was
 605 used in the analysis in Section 2.2.

Table A3. Experiment design description.

Experiment identification	Aerosol size range	Inorganic nucleation	Biogenic HOMs-induced nucleation (BNU)	Temperature effect on BNU	LVOCs organic condensation	Temperature effect on LVOC yields
BASE	0.6 nm–10 μ m	Wexler et al. (1994)	No	No	No	No
BASEnoNUC	0.6 nm–10 μ m	No	No	No	No	No
CTRL	0.6 nm–10 μ m	Wexler et al. (1994)	Yes	Yes	Yes	Yes
OCD	0.6 nm–10 μ m	Wexler et al. (1994)	No	No	Yes	Yes
BNUnoT	0.6 nm–10 μ m	Wexler et al. (1994)	Yes	No	Yes	Yes
OCDnoT	0.6 nm–10 μ m	Wexler et al. (1994)	Yes	Yes	Yes	No
NoOH	0.6 nm–10 μ m	Wexler et al. (1994)	Only with HOMs from O ₃ oxidation but without that form OH oxidation	Yes	Yes	Yes




To characterize the pure organic nucleation and condensation and investigate its controlling factors, a series of sensitivity simulations were performed as listed in Table A3. A baseline simulation (BASE) was conducted based on the default WRF-Chem, except the binary nucleation-generated aerosols were put into the third bin as described above. Simulation using the improved version of WRF-Chem, CTRL, was conducted, where new particles can be formed by organic nucleation in addition to the default H₂SO₄-H₂O binary nucleation scheme in BASE, and where the organic condensation process was also taken into consideration. In order to examine the effect of atmospheric vertical temperature variation on the organic nucleation and condensation growth, sensitivity simulations were performed using the modified WRF-Chem model, but without temperature influence on the nucleation rate and the yields of LVOCs, namely BNU_{noT} and OCD_{noT}, respectively. For the purpose of distinguishing the influence from the organic nucleation and the condensation of organics, an additional sensitivity simulation was made where only the condensation of organics was included in the BASE case, which was termed OCD. To examine the relative importance of O₃ and OH in the HOMs-generating oxidation reactions, NoOH was conducted based on CTRL but with the HOMs formation from OH oxidation turned off.

A2. Data

The ACRIDICON-CHUVA (ACRIDICON stands for “Aerosol, Cloud, Precipitation, and Radiation Interactions and Dynamics of Convective Cloud Systems” and CHUVA is the acronym for “Cloud Processes of the Main Precipitation Systems in Brazil: A Contribution to Cloud Resolving Modeling and to the GPM (global precipitation measurement)”; Wendisch et al., 2016) campaign was conducted in the Amazon region in 2014. It aimed at in-depth investigations of the properties of the aerosols and clouds in this area and the explorations of interactions between aerosols, radiation, clouds, and precipitation. Fourteen flights were operated between 6 September 2014 and 1 October 2014, encompassing comprehensive measurements of meteorology, trace gases, and aerosols with ceiling heights up to 15 km, close to the top of the troposphere. The measurements of meteorological parameters (air temperature, relative humidity, and wind speed), O₃, total aerosol number concentration, CCN number concentration, and black carbon and organic aerosol mass conducted on the 14 flights were used in this study. The total aerosol particles, also called condensation nuclei (CN) focus on aerosols with a diameter above 20 nm. The observed CCN are the CCN at a supersaturation of 0.52% (CCN(0.52%); Andreae et al., 2018). The flight tracks are shown in Fig. A1. Overviews of the ACRIDICON-CHUVA campaign and observation are documented by Wendisch et al. (2016) and Andreae et al. (2018).

A3. model evaluation



640 **Figure A2.** Comparison of (a) air temperature (Temp), (b) relative humidity (RH), and (c) horizontal wind
645 speed (WS) averaged from all flight measurements (OBS) and WRF-Chem simulations (Model).




Figure A3. Comparison of (a) O₃ mixing ratio, and (b) black carbon (BC) mass concentration averaged from all
645 flight measurements (OBS) and WRF-Chem simulations (Model).




Figure A4. Simulated (a) vertical profiles and (b) time series of α -pinene, β -pinene, and isoprene mixing ratios at the location of ATTO.

650

To compare against the flight observation, the hourly model outputs at the corresponding location of the observed data within the hour were used. The model reasonably reproduced the meteorological conditions (Fig. A2), the O_3 vertical distributions, and the black carbon concentrations (Fig. A3), showing its ability to capture the meteorological processes, basic atmospheric chemical processes, and primary aerosol emission and transport processes. The simulated concentrations of the biogenic organic vapors, α -pinene, β -pinene, and isoprene (Fig. A4) are of comparable magnitude to previous observations (Kuhn et al., 2010), demonstrating a reasonable model simulation of the biogenic emissions.

655




Figure A5. Simulated vertical profile of the SO₂ mixing ratio at the location of ATTO.

660




Figure A6. Vertical profiles of the simulated number concentrations of CN and CCN at 0.52% supersaturation averaged along the observation trajectories within the planetary boundary layer (PBL).

665





Figure A7. Simulated size distributions of (a) particle number concentration, and (b) surface area in the upper troposphere (UT) along the flight tracks.



670

Figure A8. Horizontal distribution of (a) OH and (b) O₃ averaged over 1 Sep–1 Oct 2014 from the CTRL case.





Figure A9. Same as Figure 3 (a-g), but for the planetary boundary layer (PBL). Note that the scales in (a) and (d) are 3 times of those in Fig 3a and Fig. 3d, respectively.



675

Figure A10. Same as Figure 3 (a-g), but for the middle troposphere (MT).




Figure A11. Horizontal distribution of (a) HOMs and (b) organic nucleation rate averaged over 1 Sep–1 Oct 2014 from the BNUoT case.

680

The simulated SO_2 concentration at the location of the ATTO site (Fig. A5) showed a fair agreement with the observations in the free troposphere of the Central Amazon (Andreae & Andreae, 1988; Parolovo et al., 2019; Ramsay et al., 2020), which is an important prerequisite for an accurate simulation of the H_2SO_4 - H_2O binary nucleation in this model (Wexler et al., 1994). The modeled SO_2 concentration in the PBL, especially near the ground surface, was higher than the observations (Ramsay et al., 2020), which could be related to several factors, e.g. overestimated SO_2 emission (Andreae, 2019) and/or inadequate scavenging (Hardacre et al., 2021), and requires further investigations. The influence of the SO_2 overestimation in the PBL on the simulated aerosol concentration was examined by conducting a sensitivity study, namely, BASE_ SO_2 _constrain, where all the settings were the same as the BASE case except that the SO_2 concentration in the PBL was fixed to 30 ppt in accordance with the lower end of the range of published measurements during the dry season. The simulation results show that the difference in the aerosol concentration within the PBL between BASE and BASE_ SO_2 _constrain is minor (Fig. A6), which indicates an insignificant influence of the SO_2 overestimation in the PBL on the simulated aerosol concentration.


690



695 **Table A4.** Averaged values of modeled and observed aerosol particle number concentration.

	CN (cm ⁻³)			CCN(0.52%) (cm ⁻³)		
	PBL*	MT*	UT*	PBL*	MT*	UT*
Observation (Andreae et al., 2018)	1650±1030	2130±3070	7700±7970	880±630	410±150	840±440
BASE	2230	2490	6130	800	490	350
BASEnoNUC	2150	1530	390	800	440	160
CTRL	2700	2220	6010	1100	580	750
OCD	2390	2140	3900	1090	570	660
BNUnoT	3280	2180	3980	1170	580	660
OCDnoT	2660	2430	5950	1370	540	460
NoOH	2470	2170	4000	1090	580	670

*PBL, MT, and UT are defined as the altitude range of 0–4 km, 5–8 km, 9–15 km, respectively.



700 **Figure A12.** Horizontal distribution of (a) neutral organic nucleation rate, (b) ion-induced organic nucleation rate, and (c) H₂SO₄-H₂O binary homogeneous nucleation rate at the planetary boundary layer (PBL, left panel), middle troposphere (MT, middle panel), and upper troposphere (UT, right panel) averaged over 1 Sep–1 Oct 2014 from the CTRL case.



In this study, the H₂SO₄-H₂O binary nucleation mechanism was simulated as it has been widely used for describing inorganic nucleation in the free troposphere (Cui et al., 2014; Gordon et al., 2016; Zhu & Penner, 2019). As shown from the nucleation rate in Fig. A12, H₂SO₄-H₂O binary nucleation mainly occurs in the free troposphere, which is consistent with the vertical distribution of binary nucleation simulated for the Amazon region in Zhao et al. (2020). The H₂SO₄-H₂O binary nucleation causes a CN increase of over 3000 cm⁻³ in the UT under sufficient particle condensational growth as approximately estimated from the difference between OCD and BASEnoNUC (Table A4). It is of a comparable magnitude to the CN increase of 2100 cm⁻³ by organic nucleation. A higher rate of the H₂SO₄-H₂O binary nucleation over the organic nucleation was also found by Zhao et al. (2020) in the Amazon from 9 to 13 km altitude but the overall H₂SO₄-H₂O binary nucleation in the UT was insignificant in Zhao et al. (2020) which is different from the result in this study. This is expected as the result in Zhao et al. (2020) was for a low-SO₂ area and there was competition for H₂SO₄ by other H₂SO₄-involving inorganic nucleation processes in Zhao et al. (2020). In a global simulation where the inorganic nucleation was represented only by the H₂SO₄-H₂O binary nucleation, the column-integrated H₂SO₄-H₂O binary nucleation in the Amazon is of the same magnitude as but somewhat lower than the organic nucleation (Zhu & Penner, 2019). Considering the H₂SO₄-H₂O binary nucleation occurs mainly in the upper troposphere and the organic nucleation in Zhu & Penner (2019) includes the hetero-molecular organic nucleation, the relative importance of H₂SO₄-H₂O binary nucleation to pure organic nucleation in the UT should be greater than shown in the column-integrated results. Therefore the simulated H₂SO₄-H₂O binary nucleation in this study should be generally reasonable.

The model can generally reproduce the vertical distribution of CN (Fig. 1) as described in Section 2.1, yet a systematic overestimation of CN exists in model simulations below 5 km. This could be associated with uncertainties in the fire emission inventories (Andreae, 2019), as aerosols near the ground surface during the Amazon dry season are overwhelmingly influenced by persistent biomass burning (Andreae et al., 2015). In addition, the comparison of a grid-average value in the model with an observation on a spot may also contribute to the discrepancy. To make the CCN comparison, the modeled aerosols of a size consistent with that of the observed CCN(0.52%) were used. The cut-off size of the CCN(0.52%) was calculated to be approximately 90 nm in diameter (Su et al., 2010), based on an observed average hygroscopicity value of 0.12 derived from the aerosol component observations. This hygroscopicity level is close to those of organic aerosols (Petters and Kreidenweis, 2007), which is expected since organic aerosols dominate the aerosol components in this area (Andreae et al., 2018). As the main aerosol component (i.e., organic aerosols) can be well reproduced by the developed model version (Fig. 2b), it is justifiable to use this observed hygroscopicity to calculate the CCN size in models. The comparison shows that the CCN number was underestimated by the BASE case. The model underestimation of CCN number in the UT reaches -58% as shown in Section 2.1. Compared with the UT, the biases in CCN below 4 km are much smaller and lie around -16%. The negative biases in CCN number in both the UT and PBL are however corrected in the CTRL case, suggesting the lack of SOA production and inadequate particle growth in the BASE case as the reason for the CCN underestimation.



740




Figure A13. The diurnal variation of radiation flux (upward shortwave radiation at the top of the atmosphere (SWUPT) and downward shortwave radiation at the surface (SWDNB)) and α -pinene emission rate for the average of (a) upper troposphere (UT) biogenic SOA episode days and (b) non-episode days.

745




Figure A14. Same as Fig. 4b and Fig. 4c, but for the planetary boundary layer (PBL).




Figure A15. Horizontal distribution of (a) HOMs and (b) organic nucleation rate averaged over 1 Sep–1 Oct 2014 from the NoOH case.

750

A4. Sensitivity simulations of LVOCs condensation

Figure 2b compares the simulated OA mass from CTRL and BASE with the observed data. The BASE case using the MOSAIC aerosol module and FINN biomass burning emission inventory shows a reasonable performance of OA representation in the PBL, which was also confirmed by previous evaluations for this region (Archer-Nicholls et al., 2015; Wang et al., 2016b); however, the OA in the UT is significantly underestimated. This negative bias of OA mass in the UT in the BASE case is greatly improved in the CTRL case by considering the organic aerosol processes driven by the biogenic precursors, among which the LVOCs condensation plays a dominant role (Fig. 2a).

To further examine the uncertainty of the LVOCs condensation in terms of the LVOC yields, sensitivity simulations regarding the temperature dependence of the LVOC yields were performed. The OCDnoT case adopted a bulk assumption of a yield of 13% from monoterpene oxidation and 3% from isoprene oxidation (OCDnoT), as suggested by Scott et al. (2014); while in the CTRL case, the temperature dependence of LVOC yields based on an α -pinene oxidation experiment (Saathoff et al., 2009) was applied to the LVOC yields. The OCDnoT case produces a larger amount of boundary layer OA than the CTRL case, causing a higher bias in the model compared with the observations and suggesting an excessive SOA production. A significant difference between the environment where the LVOC yields were originally based (Kroll et al., 2005) and the region investigated here may be the reason for the poor performance of the bulk yields in the studied area, as the temperature in previous applications is much lower than the tropical forest boundary layer conditions. On the

760

765



770 other end, the temperature difference could also explain the underestimation of OA mass with bulk yields in the
OCDnoT for the UT (Fig. 2b), where the temperature is far below the freezing point (Fig. A2). With the
temperature dependence correction, i.e., LVOC yields increasing with colder temperature, the OA
underestimation in the UT in the OCDnoT case can be effectively corrected in the CTRL case.