
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

**A change in the relation between the Subtropical Indian Ocean Dipole and the
South Atlantic Ocean Dipole indices in the past four decades**

Lejiang Yu^{1*}, Shiyuan Zhong², Timo Vihma³, Cuijuan Sui⁴, and Bo Sun¹

1 MNR Key Laboratory for Polar Science, Polar Research Institute of China,
Shanghai, China,

2 Department of Geography, Environment and Spatial Sciences, Michigan State
University, East Lansing, MI, USA,

3 Finnish Meteorological Institute, Helsinki, Finland

4 National Marine Environmental Forecasting Center, Beijing, China

*Corresponding Author's address Dr. Lejiang Yu

MNR Key Laboratory for Polar Science, Polar Research Institute of China, Shanghai,
China

Jinqiao Road 451, 200136, Shanghai, China

Phone: 0086-020-58712034, email: yulejiang@sina.com.cn

23 **Abstract**

24 We utilized the global atmospheric reanalysis (ERA5) and reconstructed sea
25 surface temperature (SST) data from 1979 through 2020 to examine the stability of
26 the relationship between the SST oscillations in the southern Indian and the Atlantic
27 Oceans described by the Subtropical Indian Ocean Dipole (SIOD) and the South
28 Atlantic Ocean Dipole (SAOD) indices. We note a significant positive correlation
29 between the two indices prior to the year 2000 but practically no correlation
30 afterwards. We show that in the two decades prior to 2000, a positive phase of SAOD
31 is associated with more convective activities over the subtropical southern Atlantic
32 Ocean and eastern Brazil, which trigger a stronger upper-atmosphere wavetrain, and
33 further produces stronger southern subtropical highs and surface anti-cyclonic
34 circulations and therefore a stronger correlation between the two indices. The
35 situation is reversed after 2000. Our results are potentially applicable for predictions
36 of precipitation in southern Africa and South America.

37 **1 Introduction**

38 A southwest-northeast-oriented dipole mode characterizes the anomalous sea
39 surface temperature (SST) patterns over the subtropical South Indian and Atlantic
40 Oceans (Wang, 2010). The former is referred to as the Subtropical Indian Ocean
41 Dipole (SIOD) mode (Behera and Yamagata, 2001) and the latter is named as the
42 South Atlantic Ocean Dipole (SAOD) mode (Venegas et al., 1997). The two
43 subtropical modes display similar seasonal variability with their peaks in austral
44 summer (Morioka et al., 2012). Surface latent heat flux anomalies play a vital role in

45 their variability (Sterl and Hazeleger, 2003; Suzuki et al., 2004; Hermes and Reason,
46 2005). Moreover, the interannual variability of the two modes has been linked to the
47 El Niño–Southern Oscillation (ENSO) (Boschat et al., 2013). The two subtropical
48 modes exert a great influence on precipitation in Africa and South America (Reason,
49 2001; 2002; Vigaud et al., 2009; Nnamchi and Li, 2011; Morioka et al., 2012; Wainer
50 et al., 2020) and, therefore, understanding the relationship between the two modes has
51 practical implications for precipitation forecasts in Africa and South America on
52 seasonal scale and beyond.

53 Using observational data, Fauchereau et al. (2003) noted the co-variability of the
54 SIOD and SAOD indices in austral summer. Hermes and Reason (2005) confirmed
55 the co-variability of the two indices and attributed it to an anomalous subtropical high.
56 Both studies suggested a linkage between the two indices and an atmospheric zonal
57 wavenumber-4 pattern in the Southern Hemisphere. Lin (2019) also suggested the
58 atmospheric zonal wavenumber-4 pattern controlling the South Atlantic–South Indian
59 Ocean SST pattern. The atmospheric wavenumber-4 was also observed in other
60 studies (Chiswell, 2021; Senapati, et al., 2021). The global wavenumber-4 pattern in
61 SST includes southern subtropical Indian and Atlantic Ocean components that
62 resemble the two subtropical dipole modes (Senapati et al., 2021). The linkage
63 between the SST patterns in the Southern Hemisphere ocean basins and their relation
64 with atmospheric wavenumber-4 pattern is a challenging and active research topic
65 worthy of further investigation.

66 Although previous studies have suggested that a relationship exists between the

67 SIOD and the SAOD indices, few have focused on the stability of the relationship. In
68 this study, we examine the SIOD-SAOD relationship over the past four decades from
69 1979 through 2020. We underscore a change in the relationship that occurred around
70 2000 and provide a physical explanation for the change.

71 **2 Datasets and methods**

72 Monthly SST data from the United States National Oceanic and Atmospheric
73 Administration (NOAA) Extended Reconstructed SST V5 (Huang et al., 2017) is the
74 primary dataset utilized to calculate the SIOD and SAOD indices. A secondary SST
75 data, the Kaplan Extended SST V2 data set from the UK Met Office (Kaplan et al.,
76 1998), is also used to confirm the results. Following previous studies, we derive the
77 SIOD index as the difference in the SST anomalies between the western (55-65 °E,
78 37-27 °S) and eastern (90-100 °E, 28-18 °S) subtropical Indian Ocean (Behera and
79 Yamagata, 2001) and the SAOD index as the difference of SST anomalies between
80 the south-western (10-30 °W, 30-40 °S) and north-eastern (0-20 °W, 15-25 °S) South
81 Atlantic Ocean (Morioka et al., 2011). Atmospheric data from the European Centre
82 for Medium-Range Weather Forecasts (ECMWF) fifth-generation reanalysis (ERA5,
83 Hersbach et al., 2020) provide the upper-level (200-hPa) and surface atmospheric
84 variables used in our analyses except for the monthly top-of-atmosphere (TOA)
85 outgoing longwave radiation (OLR) that is from the NOAA Interpolated OLR dataset
86 (Liebmann and Smith, 1996). For SST or atmospheric variables, the anomalies refer
87 to the departure from their climatology computed as the 42-year averaged value.

88 Correlation and regression analyses are utilized to examine the relationship

89 between the SIOD and the SAOD indices. The confidence levels are determined by
90 the two-tailed Student's *t* test. Before the correlation or regression analyses are
91 applied to the data, the variables and indices are detrended. We also remove the
92 influence from the ENSO signal using the method proposed by An (2003), where the
93 ENSO signal is represented by the Niño 3.4 index. The generation and propagation of
94 planetary waves are identified on the basis of the Rossby wave source (RWS) and the
95 wave activity flux (WAF). The RWS is calculated following Sardeshmukh and
96 Hoskins (1988) and the WAF is derived using the method of Takaya and Nakamura
97 (2001). Notice that due to the peak of the SIOD and SAOD in February (Morioka et
98 al., 2012), the austral seasons in this study refer to summer (January-March), autumn
99 (April-June), winter (July-September), and spring (October-December).

100 **3 Results**

101 The regressed SIOD and SAOD indices on detrended SST anomalies display
102 southwest-northeast-oriented dipoles in the subtropical southern Indian Ocean (Figure
103 1a) and Atlantic Ocean (Figure 1b). The correlations between the SIOD and SAOD
104 indices over the 42-year period are 0.56 for austral summer ($p < 0.05$), becoming
105 insignificant in other seasons with correlation coefficients dropping by nearly half to
106 0.23 for austral autumn and winter and 0.25 for austral spring. Removing the ENSO
107 signal resulted in small changes in the correlations and their seasonal variations, with
108 summer being the only season when the two indices are significantly correlated (0.45,
109 $p < 0.01$) (Figure 1c). The Kaplan Extended SST V2 data yielded similar results, with
110 slightly lower summertime correlation coefficients of 0.49 (with ENSO signal) and

111 0.38 (without ENSO signal, $p < 0.05$). Henceforth, we focus on the summer time series
112 without the ENSO signal.

113 To assess the stability of the SIOD-SAOD correlation over the past four decades,
114 we calculate moving correlation of the two indices using 15-year and 20-year sliding
115 windows (Figure 1d). For the 15-year window, the correlation is above (below) the 95%
116 confidence level before (after) 1998 and for the 20-year sliding window the shifting
117 occurs in 2003. Similar results are obtained using the Kaplan Extended SST V2 data
118 (Figure 2). There is a remarkable difference in the correlation between the two indices
119 prior to and after 1999 (Figure 1d). For the 1979-1999 period, the correlation
120 coefficient is 0.64 ($p < 0.01$), dropping sharply to only 0.19 ($p > 0.05$) for the 2000-2020
121 period. Results derived using the Kaplan Extended SST V2 data are very similar, with
122 the correlation coefficients of 0.60 ($p < 0.01$) for the 1979-1999 period and 0.20
123 ($p > 0.05$) for the 2000-2020 period. This notable drop in the correlation between the
124 SIOD and SAOD indices from the first two decades to the next two warrants further
125 investigation. Below we explore the reasons behind the change.

126 We compare the regression maps of the Southern Hemisphere SST anomalies on
127 the summertime SAOD and SIOD indices for the 1979-1999 period with those for the
128 2000-2020 period (Figure 3). There are clear differences in the anomalous SST
129 patterns between the two periods. As a response to the positive phase of the SAOD
130 index, significant SST anomalies occur in the southern subtropical Indian Ocean
131 during the 1979-1999 period, with a spatial pattern (Figure 3a) closely resembling the
132 positive phase SIOD index (Figure 1a); however, the SST anomalies for the

133 2000-2020 period are not significant in the southern subtropical Indian Ocean (Figure
134 3b). Similarly, corresponding to the SIOD index, a dipole of significant SST
135 anomalies appears in the South Atlantic Ocean (Figure 3c) for the 1979-1999 period
136 that bear strong resemblance to the positive phase SAOD pattern (Figure 1b), whereas
137 for the 2000-2020 period, the SST anomalies are insignificant (Figure 3d). These
138 results confirm the strong correlation between the SAOD and SIOD indices during the
139 first two decades and the lack of correlation in the last two decades, separated by the
140 turn of the century. The SST anomalies in Figure 3 display the appearance of the SST
141 wavenumber-4 mode (Senapati et al., 2021), including the SIOD and SAOD pattern.
142 Senapati et al. (2022) suggested that the weakening of the SST wavenumber-4 pattern
143 after 2000 is related to South Pacific Meridonal Mode. In addition, the weaker
144 SIOD-SAOD relationship after 2000 may be related to the decadal variability of a
145 warm pool dipole, with opposite SST anomalies in the southeastern Indian Ocean and
146 the western-central tropical Pacific Ocean (Zhang et al., 2021).

147 Lin (2019) related a South Atlantic-South Indian Ocean pattern to a wavetrain
148 induced by the South Atlantic Convergence Zone anomaly. We hypothesize that the
149 stability of the SAOD-SIOD relation may also be related to the strength of the
150 wavetrain. To test this hypothesis, we examine the regression patterns of several
151 atmospheric variables related to convective and wave activities (OLR, RWS, WAF,
152 200-hPa divergent wind, and streamfunction) to the SAOD index in austral summer
153 separately for the 1979-1999 period and the 2000-2020 period (Figure 4). Over the
154 1979-1999 period, corresponding to the positive phase of the SAOD index,

155 convective activities are enhanced over the southern subtropical Atlantic Ocean and
156 eastern Brazil, which are flanked by suppressed convective activities over tropical and
157 mid-latitude South Atlantic Ocean (Figure 4a). The convective activities over western
158 subtropical southern Atlantic Ocean and eastern Brazil produce positive RWS and
159 200-hPa divergent wind (Figure 4c), which trigger a wavetrain propagating
160 southeastwards into the South Atlantic Ocean, and then eastwards into the South
161 Indian Ocean, Australia and the South Pacific Ocean (Figure 4e). The wavetrain
162 generates negative streamfunction anomalies over the South Indian and Atlantic
163 Oceans (Figure 4e). In contrast, over the 2000-2020 period, the magnitude of the
164 anomalous OLR is less significant than that over the 1979-1999 period (Figure 4b).
165 Weaker RWS and upper level divergent wind (Figure 4d) indicate a weaker wavetrain,
166 which results in weaker streamfunction anomalies over the South Atlantic and Indian
167 Oceans (Figure 4f). To show clearly the RWS, upper-level divergent wind and Rossby
168 wavetrain, we show the significant areas in Figures 4c-f in supplementary file
169 (Figures S1)

170 Although the magnitudes of the OLR anomalies related to the SAOD index are
171 comparable over the two periods (Figure 4a and 4b), the anomalous OLR and RWS
172 and the related wavetrain associated with the SAOD index are substantially different
173 between the two periods. The differences in the climatological conditions over the two
174 periods may provide a plausible explanation. For example, over the subtropical
175 southern Atlantic Ocean and most of Brazil, the climatological OLR anomalies are
176 generally negative during the 1979-1999 period, suggesting stronger convective

177 activity favorable for the generation of the wavetrain (Figure 5a), but in contrast, OLR
178 anomalies are mostly positive during 2000-2020, indicating suppressed convective
179 activities unfavorable for the formation of the wavetrain (Figure 5b). Thus, the
180 interdecadal variability of the OLR activities can modulate the effect of the SAOD
181 mode on atmospheric circulation patterns over other ocean basins.

182 The SAOD and SIOD modes are related to the subtropical highs in the South
183 Atlantic and Indian Oceans, with stronger high corresponding to the positive phase of
184 the two indices due to wind-induced evaporation (Wang, 2010; Behera and Yamagata,
185 2001; Venegas et al., 1997). We proceed to examine the climatological mean sea level
186 pressure and surface wind field related to the aforementioned wavetrain over the two
187 periods (Figure 6). The position and strength of the climatological subtropical highs
188 and the associated surface winds in the southern Indian and the Atlantic Oceans show
189 little difference over the two periods (Figure 6a and 6b). However, the regression of
190 the mean sea level pressure to the SAOD index for the two periods show considerably
191 stronger subtropical highs and anti-cyclonic circulations in the South Atlantic and the
192 Indian Oceans over the 1979-1999 period than the 2000-2020 period (Figure 6c and
193 6d). According to the study of Hermes and Reason (2005), a stronger subtropical high
194 favors larger magnitude of the SST anomalies represented by the SAOD and SIOD
195 indices. The large decrease in the strength of the summertime subtropical high
196 associated with SAOD from the first two decades to the next two (Figure 6c, 6d)
197 corroborates the sharp drop in the SAOD-SIOD correlation (Figure 1d). Prior to 2000,
198 the stronger wavetrain associated with SAOD induced stronger summertime

199 subtropical highs, which produced larger subtropical SST anomalies in both basins,
200 strengthening the SASD-SIOD relationship. After 2000, the weaker wavetrain
201 induced weaker subtropical highs. The SST anomalies in each basin are largely
202 determined by the physical processes in each basin, which is not directly related to the
203 wavetrain. Thus, the SAOD-SIOD relationship is weaker after 2000.

204 Similarly, we have also obtained the patterns of the aforementioned atmospheric
205 circulation variables associated with the SIOD index separately for the two periods
206 (Figure 7). During 1979-1999, negative OLR anomalies occur over the northern South
207 America, corresponding to upper-level divergent wind and positive RWS anomalies,
208 while positive OLR anomalies exist over the southern Atlantic Ocean, leading to
209 upper-level convergent wind and negative RWS anomalies (Figure 7a and 7c). Those
210 anomalous RWSs produce an anomalous Rossby wavetrain propagating from the
211 southern Atlantic Ocean to southern Indian Ocean (Figure 7e). During 2000-2020,
212 negative (positive) OLR anomalies over the tropical (subtropical) central Pacific
213 Ocean generate anomalous upper-level winds and RWSs, which excite a wavetrain
214 propagating from the Pacific to South America and the southwestern South Atlantic
215 Ocean (Figure 7b, 7d, and 7f). Meanwhile, stronger convective activities over the
216 southwestern Indian Ocean and weaker convective activities over central Indian
217 Ocean also produce anomalous RWSs, which trigger a local wavetrain propagating
218 eastwards into Australia. However, the two wavetrains are controlled by different
219 factors and are not connected to each other over the South Atlantic Ocean. We also
220 examine the MSLP and surface wind field related to the SIOD index in austral

221 summer for the 1979-1999 and 2000-2020 periods (Figure 8). Over the 1979-1999
222 period, stronger subtropical highs develop over the South Indian and Atlantic Oceans,
223 which induce the positive phase of the SIOD and SAOD modes, respectively (Figure
224 8a), suggesting that the SIOD and SAOD index is connected to each other through the
225 aforementioned wavetrain (Figure 7e). Over the 2000-2020 period, positive MSLP
226 anomalies and anomalous anticyclonic circulation dominate over the South Indian
227 Ocean, though negative MSLP anomalies and anomalous cyclonic circulation occur
228 over the southwestern South Indian Ocean (Figure 8b). The atmospheric circulation
229 anomalies over the South Indian Ocean are related to the OLR anomalies and induced
230 a local wavetrain (Figure 7b, 7d and 7f). Similarly, the significant areas in Figures
231 7c-f are shown in supplementary file (Figures S1). The positive MSLP anomalies and
232 anticyclonic circulation anomalies are absent over the South Atlantic Ocean (Figure
233 8b). These results indicate that the SIOD mode over the 2000-2020 period is related to
234 local convective activities, not to those over the South Atlantic Ocean.

235 **4 Conclusion and discussion**

236 In this study, we examined the relation between the oscillations of the SST in the
237 subtropical South Indian and the Atlantic Oceans described by the SIOD and SAOD
238 indices and the stability of the relation using the ERA5 global atmospheric reanalysis
239 and reconstructed SST data from 1979 through 2020. We found significant relation
240 between the two indices in austral summer. Through moving correlation analyses, we
241 discovered that the relation in austral summer was not stable for the past four decades.
242 Specifically, the correlation between the two indices was significant prior to 2000 but

243 insignificant afterwards. The change in the relation between the two indices is
244 attributed to a change in the strength of the atmospheric wavetrain induced by
245 anomalous convective activity over the subtropical southern Atlantic Ocean and
246 eastern Brazil. More frequent and stronger convective activities prior to 2000 excited
247 stronger wavetrain, which produced stronger subtropical highs during the positive
248 phase of SAOD, resulting in a stronger relation between the two indices. The opposite
249 occurred after 2000.

250 The interdecadal variability of OLR over the subtropical South America and
251 Atlantic Ocean is the key to the relation between the SAOD and SIOD indices. What
252 determined the OLR anomalies in the region prior to and after 2000 needs to be
253 further investigated. Hermes and Reason (2005) suggested that the southern
254 subtropical high is related to the Antarctic Oscillation (AAO) and the linkage
255 strengthened after mid-1970s. The influence of the change in the AAO index on the
256 relation between the SAOD and the SIOD indices needs to be assessed. Yu et al.
257 (2017) noted a phase change of the Atlantic Multidecadal Oscillation (AMO) and the
258 Pacific Decadal Oscillation (PDO) indices in the late 1990s, with PDO shifting from
259 positive to negative and AMO switching from negative to positive around 1999. Dong
260 and Dai (2015) noted the influence of IPO on precipitation in Brazil. However, the
261 influence from the same phase of the IPO has great uncertainty and depends on the
262 period and dataset (Dong and Dai, 2015). Jones and Carvalho (2018) suggested more
263 precipitation in Brazil during the negative phase of the AMO than during its positive
264 phase. Longer datasets are utilized to examine the effect of the IPO and AMO on

265 convective activity over the subtropical South America and Atlantic Ocean on the
266 interdecadal time scale. Although our results are only based on statistical analyses,
267 they have potential for improving the prediction of precipitation in southern Africa
268 and South America.

269 *Code/Data availability*

270 The monthly SST data from the U.S. NOAA Extended Reconstructed Sea Surface
271 Temperature (ERSST) version 5 (ERSST v5) are available online
272 (<https://www1.ncdc.noaa.gov/pub/data/cmb/ersst/v5/netcdf/>). Kaplan Extended SST
273 V2 data are derived from below website
274 ([https://psl.noaa.gov/cgi-bin/db_search/DBSearch.pl?Dataset=Kaplan+Extended+SST
275 +V2&Variable=Sea+Surface+Temperature](https://psl.noaa.gov/cgi-bin/db_search/DBSearch.pl?Dataset=Kaplan+Extended+SST+V2&Variable=Sea+Surface+Temperature)). The monthly ERA5 reanalysis data are
276 available from the Copernicus Climate Data Store
277 (<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5>). The monthly
278 OLR data are derived from the NOAA Interpolated OLR
279 ([https://psl.noaa.gov/cgi-bin/db_search/DBSearch.pl?Dataset=NOAA+Interpolated+O
280 LR&Variable=Outgoing+Longwave+Radiation](https://psl.noaa.gov/cgi-bin/db_search/DBSearch.pl?Dataset=NOAA+Interpolated+OLR&Variable=Outgoing+Longwave+Radiation)).

281 Codes are available from the corresponding author on reasonable request.

282 *Financial support*

283 This study is financially supported by the National Key R&D Program of China
284 (2022YFE0106300), the National Science Foundation of China (41941009), and the
285 European Commission H2020 project Polar Regions in the Earth System (PolarRES;
286 Grant101003590).

287 *Author contributions.* LY designed the research, analyzed the data, and wrote the first
288 draft of the paper. S Z and TV revised the first draft and provided useful insights
289 during various stages of the work. CS and BS provided some comments and helped
290 with editing the paper.

291 *Competing interests.* The authors declare that they have no conflict of interest.

292 **References**

- 293 An, S. I.: Conditional maximum covariance analysis and its application to the
294 tropical Indian Ocean SST and surface wind stress anomalies, *J. Climate*, 16,
295 2932–2938, 2003.
- 296 Behera, S. K., and Yamagata, T.: Subtropical SST dipole events in the southern
297 Indian Ocean, *Geophys. Res. Lett.*, 28, 327-331, 2001.
- 298 Boschat, G, Terray, P., and Masson, S.: Extratropical forcing of ENSO, *Geophys. Res.*
299 *Lett.*, 40, 1605-1611, 2013.
- 300 Chiswell, S. M.: Atmospheric wavenumber-4 driven South Pacific marine heat waves
301 and marine cool spells, *Nature Communication*, 12, 4779, 2021.
- 302 Dong, B., and Dai, A.: (2015).The influence of the Interdecadal Pacific Oscillation on
303 temperature and precipitation over the global, *Clim. Dyn.*, 45, 2667-2681.
- 304 Fauchereau, N., Trzaska, S., Richard, Y., Roucou, P., and Camberlin, P.: Sea
305 Surface temperature co-variability in the Southern Atlantic and Indian Oceans
306 And its connections with the atmospheric circulation in the Southern
307 Hemisphere, *Int. J. Climatol.*, 23, 663–677, 2003.
- 308 Hermes, J. C., and Reason, C. J. C.: Ocean model diagnosis of interannual coevolving
309 SST variability in the South Indian and South Atlantic Oceans, *J. Climate*, 18,
310 2864-2882, 2005.
- 311 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J.,
312 Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla,
313 S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M.,
314 De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming,

315 J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, J. R.,
316 Hřm, E., Janiskov ř M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti,
317 G., de Rosnay, P. Rozum, I., Vamborg, F., Villaume, S., and Th řpaut J.-N.: The
318 ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*,
319 146, 1999-2049, 2020.

320 Huang, B., Thorne, P. W., Banzon, V. F., and Zhang, H. M.: Extended Reconstructed
321 SeaSurface Temperature version 5 (ERSSTv5), Upgrades, validations, and
322 Intercomparisons, *J. Climate*, 30, 8179-8205, 2017.

323 Jones, C., and Carvalho, L. M. V.: The influence of the Atlantic multidecadal
324 oscillation on the eastern Andes low-level jet and precipitation in South
325 America, *npj Climate and Atmospheric Science*, 1, 40, 2018.

326 Kaplan, A., Cane, M., Kushnir, Y., Clement, A., Blumenthal, M., and Rajagopalan, B.:
327 Analyses of global sea surface temperature 1856-1991, *J. Geophys. Res.*, 103,
328 18567-18589, 1998.

329 Liebmann, B., and Simth, C. A.: Description of a complete (interpolated)
330 outgoing longwave radiation dataset, *Bulletin of the American Meteorological*
331 *Society*, 77, 1275–1277, 1996.

332 Morioka, Y., Tozuka, T., and Yamagata, T.: On the growth and decay of the
333 subtropical dipole mode in the south Atlantic. *J. Climate*, 24, 5538-5554, 2011.

334 Morioka, Y., Tozuka, T., Masson, S., Terray, P., Luo, J.-J., and Yamagata, T.:
335 Subtropical dipole modes simulated in a coupled general circulation mode, *J.*
336 *Climate*, 25, 4029-4047, 2012.

337 Nnamchi, H. C., and Li, J. P.: Influence of the South Atlantic Ocean Dipole on West
338 African summer precipitation, *J. Climate*, 24, 1184-1197, 2011.

339 Reason, C. J. C.: Subtropical Indian Ocean SST dipole events and southern
340 African rainfall, *Geophys. Res. Lett.*, 28, 2225–2227, 2001.

341 Reason, C. J. C.: Sensitivity of the southern African circulation to dipole sea-surface
342 temperature patterns in the South Indian Ocean, *Int. J. Climatol.*, 22, 377–393,
343 2002.

344 Sardeshmukh, P. D., and Hoskins, B. J.: The generation of global rotational flow

345 by steady idealized tropical divergence, *J. Atmos. Sci.*, 45, 1228–1251, 1988.

346 Senapati, B., Deb, P., Dash, M. K., and Behera, S. K.: Origin and dynamics of
347 global atmospheric wavenumber-4 in the Southern mid-latitude during austral
348 summer. *Climate Dynamics*, doi:10.1007/s00382-021-06040-z, 2021.

349 Senapati, B., Dash, M. K., and Behera, S. K.: Global wave number-4 pattern in
350 the southern subtropical sea surface temperature, *Scientific Reports*, 11, 142,
351 doi:10.1038/s41598-020-80492-x, 2021.

352 Senapati, B., Dash, M. K., and Behera, S. K.: Decadal variability of southern
353 subtropical SST wavenumber-4 pattern and its impact. *Geophysical Research*
354 *Letters*, e2022GL099046. doi:10.1029/2022GL099046, 2022.

355 Sterl, A., and Hazeleger, W.: Coupled variability and air-sea interaction in the
356 South Atlantic Ocean. *Climate Dyn.*, 21, 559–571, 2003.

357 Suzuki, R., Behera, S. K., Iizuka, S., and Yamagata, T.: Indian Ocean Subtropical dipoles
358 simulated using a coupled general circulation model. *J. Geophys. Res.*, 109,
359 C09001, doi:10.1029/2003JC001974, 2004.

360 Takaya, K., and Nakamura, H.: A formulation of a phase in dependent wave-activity
361 flux for stationary and migratory quasi geostrophic eddies on a zonally varying
362 basic flow. *J. Atmos. Sci.*, 58, 608–627, 2001.

363 Vigaud, N., Richard, Y., Rouault, M., and Fauchereau, N.: Moisture transport between
364 The South Atlantic Ocean and southern Africa: Relationships with summer
365 rainfall and associated dynamics. *Climate Dyn.*, 32, 113–123, 2009.

366 Venegas, S. A., Mysak, L. A., and Straub, D. N.: Atmosphere–ocean coupled
367 variability in the South Atlantic, *J. Climate*, 10, 2904–2920, 1997.

368 Wainer, I., Prado, L. F., Khodri, M., and Otto-Bliesner, B.: The South Atlantic
369 subtropical dipole mode since the last deglaciation and changes in rainfall,
370 *Climate Dynamics*, 56, 109–122, doi:10.1007/s00382-020-05468-z, 2020.

371 Wang, F.: Subtropical dipole mode in the Southern Hemisphere: A global view.
372 *Geophys. Res. Lett.*, 37, L10702, doi:10.1029/2010GL042750, 2010.

373 Yu, L., Zhong, S., Winkler, J. A., Zhou, M., Lenschow, D. H., Li, B., Wang, X., and

374 Yang, Q.: Possible connection of the opposite trends in Arctic and Antarctic

375 sea ice cover. *Scientific Reports*, 7, 45804, 2017.


376 Zhang, L., Han, W., Karnauskas, K. B., Li, Y., and Tozuka, T.: Eastward shift of

377 Interannual climate variability in the South Indian Ocean since 1950.

378 *Journal of Climate*, 35, 561-575.

379

380



381

382

383 Figure 1. Regression patterns of austral summer (JFM) SST anomalies ($^{\circ}\text{C}$) on the positive phase

384 of the summertime indices of (a) the Subtropical Indian Ocean Dipole (SIOD), (b) the South

385 Atlantic Ocean Dipole (SAOD), (c) their time coefficients, and (d) the moving correlations

386 between the detrended and ENSO-signal-removed SIOD and SAOD indices (time coefficients)

387 using a 20-year (black solid line) and a 15-year (blue solid line) sliding window. In (d), the dashed


388 lines denote the correlation coefficients with the 95% confidence level for 20 (black) and 15 (blue)

389 samples and the abscissa indicates the end year of the moving correlations. The above results are

390 derived using the NOAA Extended Reconstructed SST V5 data.

391

392



393

394 Figure 2 Moving correlations of the detrended and ENSO-signal-removed SIOD and SAOD
395 indices using a 20-year (black solid line) and a 15-year (blue solid line) sliding window. Dashed
396 lines denote the correlation coefficients with the 95% confidence level for 20 (black) and 15 (blue)
397 samples. Abscissa indicates the end year of the moving correlations. The above results are
398 obtained using the Kaplan Extended SST V2 data.

399

400

401

402

403

404

405

406

407

408

409

410

411

412


413

414

415

416

417



418

419

Figure 3. Regression maps of the SST anomalies ($^{\circ}\text{C}$) onto the summertime indices of (a), (b) SAOD and (c), (d) SIOD, over the periods of (a), (c) 1979-1999 and (b), (d) 2000-2020. Dots denote the regions of above 95% confidence level.

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449


450

451

452


453

454



455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485

Figure 4. Regression maps of (a), (b) the anomalous outgoing longwave radiation (OLR) (W m^{-2}) at the top of the atmosphere, (c), (d) Rossby wave source (RWS) (10^{-10}s^{-2}) and 200-hPa divergent wind (vector), and (e), (f) wave activity flux (WAF) (vector) and streamfunction ($\text{m}^2 \text{s}^{-1}$) onto the summertime SAOD index over the periods of (a), (c), (e) 1979-1999 and (b), (d), (f) 2000-2020.



486

487 Figure 5. Climatological OLR anomalies (W m^{-2}) during (a) 1979-1999 and (b) 2000-2020, with
488 respect to the 42-year climatology over the 1979-2020 period.

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509


510

511

512


513

514



515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537

Figure 6. Climatological mean sea level pressure (MSLP, in hPa) and 10-m wind field (vector) over the periods of (a) 1979-1999 and (b) 2000-2020, and regression maps of MSLP (in Pa) and 10-m wind field (vector) onto the summertime SAOD index over the periods of (c) 1979-1999 and (d) 2000-2020. Shaded regions and red vectors indicate above 95% confidence level.



538

539 Figure 7. Regression maps of (a), (b) OLR ($W m^{-2}$), (c), (d) RWS ($10^{-10} s^{-2}$) and 200-hPa
 540 divergent wind (vector), (e), (f) WAF (vector) and streamfunction ($m^2 s^{-1}$) onto the summertime
 541 SIOD index over the 1979-1999 period (a), (c), (e) and the 2000-2020 period (b), (d), (f).

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558


559

560

561

562

563



564

565 Figure 8. Regression maps of MSLP (in Pa) and 10-m wind field (vector) onto the summertime
 566 SIOD index over the periods of (a) 1979-1999 and (b) 2000-2020. Shaded regions and red vectors
 567 indicate above 95% confidence level.

568

569

570

571

572

573

574

575