

Momentum fluxes from airborne wind measurements in three cumulus cases over land

Ada Mariska Koning¹, Louise Nuijens¹, and Christian Mallaun²

¹Delft University of Technology

²Deutsches Zentrum für Luft- und Raumfahrt

Correspondence: Mariska Koning (A.M.Koning@tudelft.nl)

Abstract. Measurements of wind and momentum fluxes are not typically at the center of field studies on (shallow) cumulus convection, but the mesoscale organization of convection is likely closely tied to patterns in wind. This study combines in situ high-frequency turbulence measurements from a gust probe onboard a Cessna aircraft with downward profiling Doppler Wind Lidar (DWL) measurements onboard a Falcon aircraft to study variability in the wind profile and momentum fluxes in regions of convection. The dual-aircraft measurements were made during three prototype flights in shallow convective regimes over German agricultural areas (two of which had hilly topography, one flat) in late Spring 2019, including forced cumulus humilis under weak winds and "popcorn" cumuli during stronger wind and wind shear after front passages.

All flights show pronounced meso-gamma (2-20 km) scale variability in the wind, with largest wind variance (on the order of 2 - 4 m²s⁻²) towards cloud base and in the cloud layer on flights with large vertical wind shear. The wind and wind variance profiles measured in-situ and by lidar compare very well, despite the DWL's coarse (~ 8 km) horizontal footprint. This highlights the presence of wind fluctuations on scales larger than a few km and that wind lidars can be used more deliberately in field studies to map (mesoscale) flows.

Cloudy transects are associated with more than twice the momentum flux compared to cloud-free transects. The contribution of the updraft to the total momentum flux, typically a third to two-thirds, is far less than the typical contribution of the updraft to buoyancy flux. Even on the same flight day, momentum flux profiles can differ per track, with one case of counter-gradient momentum transport when the updraft does carry substantial momentum flux. Scales beyond 1 km contribute significantly to the momentum flux and there is clear evidence for compensating flux contributions across scales. The results demonstrate that momentum flux profiles and their variability require understanding of motions across a range of scales, with non-negligible contributions of the clear-sky fluxes and of mesoscales that are likely coupled to the convection.

20 1 Introduction

Observations of the vertical profile of wind are valuable for reducing forecast errors and for advancing the understanding of processes that influence wind variability, including large-scale and mesoscale dynamics and small-scale turbulent processes. In this paper we combine state-of-the-art airborne Doppler wind lidar (DWL) combined with traditional in situ turbulence measurements to measure the profile of wind and turbulent wind fluctuations within cloud-topped boundary layers, in which

25 thermally-driven (convective) plumes are thought to play an important role in transporting wind. By measuring wind profiles at levels beyond meteorological towers and ground-based operational DWLs, we aim to investigate wind and momentum flux variability beyond the surface layer.

On local and regional scales, the growing wind energy industry has boosted wind profiling observations in the lowest layers of the atmosphere through the deployment of Doppler wind lidars (DWLs). DWLs conventionally measure high-resolution
30 wind to the top of the wind turbine or to hub height (the centre of the wind turbine's rotor, up to 250 m). Such measurements are used to understand turbulent wind fluctuations in the surface layer that are influenced by weather, terrain, turbine wake effects or shear across the rotor-swept area (Bakhshi and Sandborn, 2020; Banta et al., 2013; Iungo and Porté-Agel, 2013; Krishnamurthy et al., 2013; Mann et al., 2010).

One source of wind in the surface layer is the (downward) mixing of momentum from higher levels. Not only dry convection,
35 but also moist convection plays an important role in this process, because clouds extend the boundary layer height, tapping in regions aloft with faster moving winds. This transport of momentum (momentum fluxes) by convective eddies (thermals) and through clouds is broadly called convective momentum transport (CMT). Like small-scale turbulence, CMT is an unresolved process in forecast models that contributes to uncertainties in local wind predictions. However, unlike the turbulent wind fluctuations measured in the surface layer through most commercial DWLs, few high-resolution wind profiles extend beyond
40 the surface layer (> 200 m) to target wind fluctuations and momentum transport at heights where moist convection develops.

Our understanding of turbulent wind fluctuations throughout the boundary layer largely stem from a handful of in situ turbulence measurements during research aircraft flights at selected height levels in subtropical settings. A seminal study is that by LeMone and Pennell (1976), where flight tracks below and through cumulus fields near Puerto Rico were used to derive wind and flux profiles. This work highlighted that the momentum flux profile can take a very different shape depending on
45 clouds overhead. In particular, they found that in fields of cumulus clouds organized in rolls, the rolls were responsible for a significant amount of the momentum transport even though clouds were extremely shallow. In fields of more significant and randomly distributed clouds, the linear flux dependence disappeared, becoming counter-gradient at various altitudes. However, some doubt remained as to whether the wind profile could have evolved during the flight, because the profile itself was only sampled by the turbulence measurements at selected legs.

Large-eddy simulations (LES) have revealed the very different nature that momentum flux profiles can take depending on
50 the scales and domains considered (Zhu, 2015; Schlemmer et al., 2017; Saggiorato et al., 2020). LES output suggests that turbulent fluctuations on scales larger than 400 m explain a considerable part of the momentum flux, in particular above the surface layer towards the mixed-layer top and within the cloud layer. Recently, Dixit et al. (2021) suggest that the absence of mesoscale circulations in idealised periodic-boundary LESs lead to an underestimation of momentum flux that tends to be
55 counter-gradient in the cumulus layer. So far, we lack understanding about the contribution of mesoscale horizontal flows to the momentum flux and its net effect on the wind. However, both observations and simulations clearly show the spatial variability in wind that is large in areas of convection, especially in the presence of precipitation whose evaporation can lead to cold pools and associated gust fronts (Zuidema et al., 2012; Li et al., 2014; Helfer and Nuijens, 2021).

Three prototype flights were carried out focusing on measuring the wind environment in convective situations to evaluate
60 turbulent to mesoscale (up to 7 km) wind fluctuations and implications for the momentum flux profile. The campaign in-
volved dual-aeroplane flights over Germany using the Falcon and Cessna research aircrafts from the German Aerospace Center
(Deutsches Luft- und Raumfahrt e.V., DLR) in Oberpfaffenhofen. The Falcon flew at high altitudes deploying a downward
looking 2 micron Doppler wind lidar (DWL), used here, as well as the sideward looking ADM Aeolus demonstrator (A2D)
lidar - the airborne prototype to the space-borne Aeolus measurements. A comparison of the A2D, DWL and Aeolus measure-
65 ments can be found in Witschas et al. (2020) and Lux et al. (2020). The smaller Cessna aeroplane flew below the Falcon at
lower altitudes in the boundary layer collecting in situ wind and turbulence measurements. The three collocated flights captured
conditions ranging from fair-weather shallow cumulus developing over hilly terrains to pre- and post-frontal convection with
popcorn convection over flat terrains. Land use below the flight track was often agricultural with grass or low crops with very
few bare lands. Few patches of trees were encountered, especially during the flight on 24 May 2019, as well as few villages.

70 The amount of statistics collected with just three flights is limited, but demonstrate the value of wind profiling by the DWL
by showing on the one hand how robust the wind profile is as derived by the in-situ measurements, and on the other hand by
revealing variability in winds carried at different scales. The questions we address in this manuscript are as follows:

- How do the DWL wind profiles match with in situ wind measurements?
- Are the measured momentum flux profiles in the sub-cloud layer and in the cloud layer in line with our expectation and
75 can they be explained by convective updrafts?
- Which scales contribute significantly to wind variance and momentum flux?

The manuscript includes a description of the flight strategy, measurement techniques in section 2, which also includes a
short explanation of down-gradient turbulent momentum transport and the updraft detection method. Section 3 describes the
meteorological conditions during the three flights. Section 4 explores the different momentum fluxes in relation to the wind
80 profiles and convection, including an analysis of the contributions of updrafts and scales that carry most of the momentum. Our
findings are summarized in section 5.

2 Flight measurements and data processing

2.1 Flight strategy and measurements

The CloudBrake measurement campaign took place in Germany at the end of May and beginning of June 2019, a period that
85 is known to often display (shallow) cumulus clouds. Starting from around noon until 13:30 or 14:30 local time (CEST), the
flights targeted a time of the day during which cumulus clouds are typically well developed. An impression of the cumulus and
weather conditions during each flights is given in Figure 1. The first flight on May 24 (2019) was a typical shallow cumulus
day: starting out with clear skies and weak winds, local shallow cumulus started forming over the hilly parts of the Swabian
Jura. Clouds remained shallow, reaching a thickness of approximately 500 m. The second flight was under the influence of an




Figure 1. Photographs (upper row) and Modis satellite images from NASA Worldview Snapshots (lower row) of the cumulus fields during the flights on 2019-05-24 (left), where white is the cloud free area and black the cloudy area, 2019-05-27 (middle), where the west track is indicated in black, the east track in red and the southern track in yellow, and 2019-06-04 (right), where the west track is indicated in black and east in red. In each satellite picture, the horizontal black and white bar indicates a total distance of 20 km and the mean wind direction during the flight is drawn in the windrose.

90 approaching cold front, providing an interesting and dynamic mixture of shallow cumulus- and stratocumulus-topped boundary layers. Above the shallow cumulus that were around 1 km thick, mid-level alto-cumulus and stratus layers were present. The third flight, on June 4th 2019, experienced post cold front conditions. There was a large cumulus field with very diverse cloud tops. Clouds were at most 800 m thick and were typically thicker at the northern part of the leg.

During the 2–2.5-hour-flights, the two aeroplanes flew back and forth across the same pre-defined tracks. Because the two
 95 planes have different cruising speed (the Cessna about 70 m s^{-1} , for the Falcon about 200 m s^{-1}), the pre-defined tracks ensure overlap in space and time to the degree possible. Flight legs were mostly flown cross wind and ranged from 50 to 100 km in length to ensure sufficient low-frequency wind variability. During some of the flights the tracks were moved or shortened with respect to the original plan to ensure cumulus clouds were captured. All flown tracks are shown in Figure 1. The terrain below was mostly used for agriculture with low crops, occasionally encountering patches of trees or villages. On the first two flights
 100 a hilly topography was present, whereas the last flight was above flat land.

Turbulence measurements using an in situ (3D) turbulence probe aboard the DLR Cessna Grand Caravan were taken at 100 Hz along the track at four different altitudes: within the mixed-layer, near cloud base, within the cloud layer and through the tops of only the thickest clouds. Employing the downward staring Doppler wind LiDARs at a measurement rate of 40 s, the DLR Falcon remained around 11 km altitude throughout the flight. The instruments are described next.

105 2.1.1 In situ turbulence probe

The DLR Cessna Grand Caravan was equipped with (i) a meteorological sensor package (METPOD) that measures temperature, humidity, pressure, and wind, and (ii) the IGI systems' AEROcontrol system, which combines measurements of a Differential Global Positioning System (DGPS) with a high-accuracy inertial reference system (IRS). Calibration of the devices before the flight and applying corrections afterwards result in a horizontal wind measurement uncertainty of 0.3 m s^{-1} and 0.2 m s^{-1} for the vertical wind component. Further details on the instrument specifics, calibration, correction procedure, and uncertainties can be found in Mallaun et al. (2015).

The high-frequency 100 Hz wind measurements, taken with a boom-mounted Rosemount model 858 AJ air velocity probe, are used for flux calculations. The aircraft movements are corrected using IGI. A linear fit is subtracted from the data before flux calculations. All scales from 10^{-2} Hz are included in this calculation, unless stated otherwise.

115 2.1.2 Energy spectra

To check the quality of the measurements, we calculated the power spectral density (based on the Fast Fourier Transform), aftersubtracting a linear trend from the data. Welch method was used with a Hann window with 1000 samples and 50% overlap to reduce noise in the spectrum. The spectra of the streamwise, cross and vertical wind components at four heights are displayed in Figure 2 for the western track on 4 June. Note that the streamwise and cross wind do not change much from zonal and meridional winds. The legs flown in the sub-cloud layer and in cloud layer contain more energy than the legs flown near cloud base and cloud top. Comparing the three wind components, turbulence appears to behave anisotropic: from 0.01-1 Hz, w contains similar (or in the sub-cloud layer more) energy than u and v . Between 1-10 Hz, the streamwise component has most energy, and w least. The characteristic $5/3$ slope of the inertial sub-range (dashed line) is seen from $\sim 0.2 - 15$ Hz (equivalent to a spatial resolution of 350 m down to 5 m, assuming a typical cruising speed of $65-75 \text{ m s}^{-1}$). From 15 Hz onward, the dampening of the fluctuations in the tube becomes visible and the signal falls of faster, except for one peak at 30 Hz, which is attributed to propeller effects (Mallaun et al., 2015). For calculations of the (eddy-covariance) fluxes and variances, we apply a high-pass filtering after linear detrending, removing the contributions by eddies with a horizontal length scale larger than 7 km (having a frequency lower than 0.01 Hz), a method that is also followed by Brilouet et al. (2021). Filtering out these frequencies will lead to an error because we lose information, however the random error that is generated by the finite sample size will be reduced (Lenschow et al., 1994).

2.1.3 Eddy-covariance fluxes

The time series are partitioned in leg-averaged values $\bar{\phi}$ and fluctuating parts ϕ' by subtracting a linear trend from the time series and applying a high-pass filter, a method that has also been applied by Brilouet et al. (2021). Doing so removes the influence of larger scales having frequencies lower than the cut-off frequency and will lead to a loss of information (Lenschow




Figure 2. Power spectrum of the streamwise, orthogonal and vertical wind components for the western legs flown on 2019-06-04. Each altitude is shown in an individual panel. The dashed line represents the $-5/3$ slope corresponding to the inertial sub-range. Welch method with a Hann window of 1000 samples and 50% overlap has been applied to reduce noise.

135 et al., 1994). Fluxes and variances are then calculated by multiplying and averaging the fluctuations of w and ϕ over a specific time window, known as the eddy-covariance method. For instance, the leg average flux of ϕ is given by:

$$\overline{w'\phi'} = 1/N \sum_{i=1}^N w'_i \phi'_i \quad (1)$$

The smallest resolved frequency depends on the length of the leg, *i.e.* on the number of samples N : $f_{min} = f_s/N$, in which f_s is the sampling rate in Hz. Flying at a cruising speed of $\sim 65\text{-}75 \text{ m s}^{-1}$ at a constant height, and with constant ground speed,
 140 it is reasonable to assume that a static turbulent field is sampled. However, the statistical representation of the low frequencies is poor and therefore needs cautious interpretation.

2.1.4 Airborne Doppler wind LiDAR

Doppler wind LiDARs (DWLs) are the international standard for wind measurements and have been used for among other things 1) data assimilation experiments (Horányi et al., 2015; Pu et al., 2017; George et al., 2021, e.g.), 2) to study for instance
 145 turbulence, gravity waves, orographic effects (Yuan et al., 2020; Gisinger et al., 2020; Baidar et al., 2020, e.g.), and 3) to monitor the flow in wind farms (Käsler et al., 2010; Wagner et al., 2017; Zhan et al., 2020; Schneemann et al., 2021, e.g.). The coherent detection DWL employed in this study has a wavelength of 2022.54 nm (approximately $2 \mu\text{m}$), being eye-safe and

operating in the Rayleigh scattering regime. The (vertical) resolution of the wind measurements depends on both the duration of the pulse, also called pulse width, and the distance that the signal can travel during the sampling time. The shorter the pulse, the better the spatial resolution, although a reasonable sampling duration is needed to ensure sufficient accuracy of the velocity estimation (Liu et al., 2019). With a pulse width of ~ 400 ns and an averaging time of 1 s, we have a vertical resolution of 100 m (Witschas et al., 2017) (i.e. along-beam resolution approx. 94 m). Furthermore, the aircraft speed influences the horizontal resolution. Flying with approximately 200 m s^{-1} and having a sampling frequency of ~ 40 s, the horizontal resolution (distance traveled between two measurements) is about 8 km. Pulsed LiDARs have a blind spot of tens to hundreds of meters near the beam source, depending on the pulse duration and range gate width (Liu et al., 2019). Therefore, although flying at 11 km, the first wind velocities are obtained from approximately 7 km altitude down to about 500 m. The DWL employed in this study has previously been compared to dropsonde measurements, in which the systematic error has been found to remain below 0.1 m s^{-1} and the random error to vary between 0.92 and 1.5 m s^{-1} (Weissmann et al., 2005; Chouza et al., 2016; Schaeffler et al., 2018; Witschas et al., 2020).

The Velocity-Azimuth Display technique (Browning and Wexler, 1968) with an off-nadir angle of 20 degrees, is used to retrieve all three wind components. The processing algorithm that is applied to retrieve the wind vectors from one revolution of line-of-sight measurements is described in Witschas et al. (2017).

Figure 3 shows an example of the wind anomalies (i.e. the wind measurements of which the average wind during the measurement flight is subtracted) on June 4 2019. The turning points indicating reverse heading on the same leg are indicated with white vertical lines, revealing similar but mirrored wind structures on subsequent legs. On this particular flight, the track was moved further to the east around 11:40 UTC, where different structures are visible. Data gaps, which can be associated with clouds, are indicated in black.

The top of the boundary layer that is around 2 km altitude is clearly visible in the w fluctuations, with larger fluctuations below, and smaller above. The top of the boundary layer is marked by predominantly blue colours, indicating negative velocities produced by overshooting thermals that become negatively buoyant. Within the boundary layer updrafts generate the largest fluctuations, while a few downdrafts extending to the surface are also evident. It appears that the DWL can at least to some extent observe the coherent convective features that are responsible for mass transport of scalars and momentum.

For one of the legs on 4 June, the histograms of the sub-cloud layer u , v , and w wind are compared in Figure 4. Mean horizontal wind measurements over this leg are comparable for the DWL and in-situ measurements. Despite its much coarser resolution and its missing v winds $< 2.5 \text{ m s}^{-1}$, the wind variance observed by the DWL is only slightly smaller (0.1 m s^{-1} less than in the in-situ measurements). This gives us confidence that the DWL can provide complementary information of the (horizontal) wind profile at heights where in situ measurements are absent. It also tells us that horizontal wind fluctuations are largely set by scales of 8 km or larger and that cloud convection scales of 1-2 km are less important. On the other hand, the vertical wind shows much less variation than the in situ measurements. This is explained by the much larger area that is measured by the DWL: it can only see the average vertical velocity in this area, which on average is much lower than the vertical velocity of vertical transient small eddies than can be better captured by the in situ measurements.




Figure 3. Anomalies of zonal (u), meridional (v), and vertical (w) wind measurements from the DWL, zoomed in on the lowest 5 km. Measurements taken on 2019-06-04. Missing values are indicated in black and often correspond to clouds (1-2 km altitude). White vertical lines indicate turning points at the ends of each the track.




Figure 4. Distribution of u , v , and w wind in the sub-cloud layer of the western track at 617 m altitude on 2019-06-04, as measured by the Doppler Wind LiDAR (top panels, blue) and the in situ turbulence probe (lower panels, black). The DWL range bin closest to the in situ flight height have been used.

2.2 Updraft detection algorithm

Using conditional sampling we identify updrafts, following the method described and tested by Lenschow and Stephens (1980). We conditionally sample on updrafts ($w' > 0$ & $w > 0$) that are wider than 100 m, and that have an excess in absolute humidity

Table 1. This table shows the number of updrafts, relative updraft area, average updraft size, and average updraft speed for the legs flown on 4 June 2019.

Updraft statistics	Number of updrafts	Updraft area [% of leg]	Chord length [m]	Updraft velocity [m s^{-1}]
West (thicker clouds)				
Cloud top	8	2.3	264	1.6
Cloud layer	12	5.4	412	2.2
Cloud base	20	8.1	328	1.4
Mixed layer	16	8.2	333	1.7
East (thinner clouds)				
Cloud top	1	0.6	372	2.1
Cloud base	3	1.8	412	1.5
Mixed layer	10	4.7	289	1.4

185 $\rho'_v > 0.5 \sigma_{\rho'_v}$. This method is more robust than using virtual temperature or buoyancy, and can be applied both in the sub-cloud and cloud layer.

Table 1 shows the updraft statistics of the legs flown on 4 June 2019. It lists the number of updrafts, the relative length of the leg that they occupy, the average horizontal size and the average updraft velocity. We find that the fraction of the leg that is covered by updrafts (updraft area) decreases with height, although the average updraft chord length (the length of the updraft slice that we passed through) peaks at cloud base for the thinner clouds on the eastern track on June 4th as well for the clouds on 2019-05-24 (not shown). On May 24th, we find more and stronger updrafts in the cloud-topped mixed layer than under clear-skies, whereas updraft chord length is comparable. This is in correspondance with findings of *e.g.* Nicholls and LeMone (1980), who found stronger sub-cloud and lower cloud-layer vertical velocity standard deviation in more cloudy conditions. The largest average updraft velocity is found at cloud base, suggesting that the stronger mixed layer updrafts reach the lifted condensation level and benefit from the energy released at condensation. On June 4th, fastest average updraft speeds are found in the cloud layer in the case of thicker clouds. With the thinner clouds fastest updraft speed is found at cloud top, although we must be careful as this includes only one sample.

3 Flight conditions: wind and thermodynamic profiles

Although the three flight days all captured a shallow cloud regime, they differed substantially in their characteristics of the wind (wind speed, wind shear and directional shear), providing a set of diverse case studies, whose wind and thermodynamic profiles are described next. For the wind, an entire profile of the mean and variance are shown from the DWL, with the in situ measurements denoted on top (Figure 5). Except for the wind direction on May 24th, which varied greatly in the sub-cloud layer, the mean DWL and in situ winds compare very well.




Figure 5. Average wind speed and wind direction profiles for each flight date. Average DWL profile indicated in blue, shading indicates the range between the first and third quartile. Average in situ measurement indicated with squares (measurements in/below cloud field areas) and circles (cloud free areas). Cloud base height (cbh) has been estimated during the flight and is indicated with a horizontal dashed line.

The first flight (May 24) took place after a number of overcast days and heavy rain. Southern Germany was under influence of a broad area of high pressure west of Europe and over the Northern Atlantic and the conditions were very stable with hardly any clouds in Southern Germany. The northern part of the leg was flown over the Swabian Alps, where numerous gliders were making use of the thermal structures that typically develop here and shallow cumulus with cloud bases near 2 km (dashed horizontal lines) and tops near 2.5 km developed. These were the focus of our measurements. Winds were weak and reasonably well mixed up to 1400 m, topped by a layer with strong wind turning near 1.5 km (some 500 m below cloud base), and wind speed increased up to 2500 m in a layer extending through cloud base (Figure 5). In contrast, temperature and humidity were very well-mixed vertically. The atmosphere was relatively dry, with a pronounced inversion in temperature and moisture starting near 2.2 km (Figure 6), as well as a typical region of negative buoyancy below cloud base (Figure 6) that has not been captured in the other two flights.




Figure 6. Virtual temperature and mixing ratio during the three flight days. On the first flight there were three legs that were partly below clear-sky and partly below cloudy sky. They have been separated and are represented by open circles and closed squares, respectively. The other two cases had multiple tracks that are indicated with different colours. The raw 100 Hz data is indicated in light blue. Cloud base is indicated by the dashed horizontal line.

Considerably stronger wind speeds, but far less wind shear were present during the second flight (May 27) when we sampled
 215 air masses ahead of a cold front located SW-NE across eastern Germany (Figures 1(b), 5 (b,e)). The air masses were somewhat warmer and moister, but with a thermodynamic structure and a cloud base similar to that of the first flight (Figure 6. Besides shallow convection, there was plenty of mid- and upper levels cloud, which we encountered at the end of the first flight leg towards the north. Later, the front seemed to break up and skies were clearer, especially towards the southeast. Eventually, also in the southeastern area of our operations, shallow cumulus made way for stratocumulus layers, with only rare sights of clear
 220 sky and sunshine.

During the third flight (June 4th), we measured an extended field of shallow cumulus clouds that developed behind a cold front over northwestern Germany in air masses that were considerably colder and moister (Figure 6 (c,f)), with much lower cloud bases near 1000 (western N-S leg) and 1200 m (eastern N-S leg) and very diverse cloud top heights (see Figure 1(c)), with maximum tops near 2 km. The cloud field was organised in patches of alternating cloudy and cloud-free air masses. As
 225 the clouds were getting deeper towards the northern parts of the leg, the relative sizes of the patches increased. Difference in the inversion strenght of the temperature profile is clearly visible between the western and easter leg (Figure 6(c), as well as a




Figure 7. Profiles of buoyancy and the moisture flux for the three flight days. On the first flight there were three legs that were partly below clear-sky and partly below cloudy sky. They have been separated and are represented by open circles and closed squares, respectively. The other two cases had multiple tracks that are indicated with different colours. The average flux below cloudy skies has been indicated by a solid line, of which two are present in the last flight day to show the difference between the western and eastern track. Cloud base is indicated by the dashed horizontal line.

large difference in moisture flux (Figure 7). Near- surface winds were weak and from the south, with strong shear and a turning from southeasterly to southwesterly winds right around cloud base (Figure 5(c,f)).

Based on the wind profiles, the three flights could be classified as having weak wind and strong shear either in the sub-cloud layer (Flight 1) or in the cloud layer (Flight 3), and having strong wind but little shear (Flight 2). In the next section, we will explore the associated turbulent statistics of these flights and evaluate whether the derived momentum flux profiles are in line with our expectations *e.g.*, that momentum fluxes throughout the mixed layer and cloud layer increase with wind shear as predicted by K-theory.

4 Momentum flux profiles

230 4.1 Sub-cloud and cloud layer profiles

In Figures 8 and 9 we consider the profiles of wind and momentum flux for the vector wind components u and v separately. As in Figure 5, the wind speed is shown for both the DWL (in blue) and the in situ turbulence probe at the flight levels




Figure 8. Average u (a-d) and $\overline{u'w'}$ (e-h) profiles for each flight date. The u -axis is positive eastward. To obtain these profiles, we applied the same averaging procedure as in Figure 5 and the eddy covariance method as described in Section 2.1.3. On 2019-05-24 (a,e), the clear-sky measurements are indicated with open circles, whereas measurements in cloud fields are indicated with filled squares. Pink triangles represent the wind speed within updrafts (upper panels) and updraft contribution to the total flux (lower panels).

(circles, squares). A guideline for the flux profiles in cloudy conditions are indicated with solid black lines, which are linearly interpolated between leg averaged values at the different flight levels (and are sometimes averaged over two legs the same
 240 level).

When fluxes are dominated by small-scale turbulent diffusion, it may be modeled (parametrized) by using so-called flux-gradient relationships, also known as K-diffusion. We find that most of the fluxes and their relationship with the wind gradient lead to a K-value that is in line with down-gradient diffusion, acting to reduce the wind gradient.

On May 24th, ignoring the strong gradients in u below ~ 700 m, $\partial_z u > 0$. This implies that air parcels that are displaced
 245 upward ($w' > 0$) generally have a negative u perturbation compared to their environment ($u' < 0$), acting to remove the wind gradient, in line with K-theory. This holds generally for all flight days. Negative u perturbations are in particular evident from the actual wind in air masses sampled within updrafts, which tend to be several m s^{-1} slower (pink triangles in Figure 8 and 9). Similarly, the meridional momentum fluxes are also down-gradient. For example, the gradient $\partial_z v < 0$ above 1 km on May 24th, corresponding to a positive meridional momentum flux ($\overline{v'w'} > 0$), and $\partial_z v > 0$ on June 4th, corresponding to $\overline{v'w'} < 0$.




Figure 9. Same as Figure 8, but then for v and $v'w'$. The v -axis is positive northward.

250 The profiles of $\overline{u'w'}$, $\overline{v'w'}$ reveal that larger fluxes are measured on May 24th than on May 27th, in line with the stronger shear present in u and (to a lesser extent) in v . Fluxes typically decrease towards the boundary layer height (cloud top or mixed-layer top in case of clear sky), as the variance of vertical velocity decreases towards the top of the boundary layer (Figure 10).

Evidently, on May 24th the momentum fluxes throughout the mixed-layer increased considerably from the first transect of the flight, which captured a dry convective boundary layer (open circles), to the second transect, which is when cumulus clouds
 255 developed on top of the mixed layer (filled squares), although both transects have comparable u, v profiles (Figure 8, 9). The larger fluxes reflect the presence of stronger turbulent eddies. Especially just below and at cloud base, much larger variances are present in u and v and throughout the mixed layer in w (Figure 10).

While on May 24th, 27th and the eastern leg on June 4th the fluxes decreased towards cloud base, with little flux remaining in the cloud layer, the western leg on June 4th shows an increase in momentum fluxes with height (in particular $\overline{u'w'}$ but also
 260 $\overline{v'w'}$). Whereas the flux in the mixed-layer below clouds is almost negligible, one of the largest fluxes was measured in the cloud layer ($\overline{u'w'} \sim 0.4 \text{ m}^2\text{s}^{-2}$). Clouds on this western leg had a lower base (just above 1 km) and higher cloud tops (up to 2 km) and thus were thicker than the clouds that were encountered on the eastern track. The thicker clouds do not only have larger momentum transport in the cloud layer, but also a much larger (percentage) contribution of the updraft to the total flux than any of the other measurements (Figure 6). The fraction of the leg that was occupied by updrafts was also significantly




Figure 10. Variance of u' , v' , w' for each flight.

265 larger than in all other cases. The deeper clouds may have been accompanied by wider updrafts with better protected cores that may be responsible for carrying larger fluxes.

Figure 11 shows a time series of turbulence measured at 600 m in the mixed layer on the western track on June 4th. In grey the unfiltered turbulence statistics are shown, while black shows the linearly detrended and high-pass filtered ($f > 0.01$ Hz) statistics. Cloudy updrafts can have vertical speeds up to 5 m s^{-1} , in both altitudes. Evidently, large buoyancy fluxes ($w'\theta'_v$, top
270 row) are associated with large momentum fluxes (bottom row), which reveals the importance of convection in generating a large




Figure 11. Raw (unfiltered) and fluctuating (filtered: linear detrended, high-pass filter cut-off 0.01 Hz) time series of (a) buoyancy flux, (b) vertical velocity, (c) zonal wind, (d) meridional wind, (e) convergence and divergence of the wind speed in the streamwise direction calculated using a low-pass filter that considers all scales larger than 700 m ($f_c < 0.1$ Hz), and (f) momentum fluxes measured on 2019-06-04 at 600 m, in the middle of the mixed layer of the western leg. Updrafts are indicated with light-blue shading.

momentum flux. Sometimes, for instance at 11:33 UTC, the convergence (panel (e)) is closely tied to convection. Typically, updrafts carry wind speeds that are much slower (up to 5 m s^{-1} for cloudy updrafts) than the environment. Looking carefully, one might see that u' and w' peak at different times and that u' has a different sign in various updrafts. This could explain a much lower momentum flux. We discuss this further in the next sections, where we explore the fluxes sampled on (cloudy) updrafts, as well as how eddies of different scales contribute to the fluxes.

4.2 Scale contributions to flux

Observations in the surface layer, for instance during the Kansas experiment, show that the cospectra of the $\overline{u'w'}$ flux follow a fixed slope, with the large scales more significant (?). However, throughout the boundary layer, Large Eddy Simulations of various cases indicate that the momentum flux carried by small-scale shear-driven turbulent eddies (with a size smaller than ~ 200 m) can contribute more than 50% of momentum fluxes. Small scale turbulence may also transport momentum in an opposite direction than larger more coherent eddy structures (Zhu, 2015). This is particularly true for the lower mixed-layer and near cloud tops. However, in shallow cumulus cases, especially from the middle of the mixed-layer (sub-cloud layer) to

the middle of the cloud layer, the net momentum fluxes are almost entirely carried by eddies with scales greater than 400 m (Zhu, 2015).

285 In Figure 12, the total (net) momentum flux is shown for the legs in the sub-cloud layer, near cloud base, within the cloud layer and near cloud top for June 4th. The momentum flux at different scales is calculated using a high-pass filter filters that removes larger scales with increasing cut-off frequency. The flux all the way to the right is for instance carried by eddies up to a scale of $\sim 7\text{km}$, corresponding to a cut-off frequency of 0.01 Hz. When the flux magnitude increases rapidly from left to right, it implies that larger-scale eddies contribute more to the momentum flux than smaller scales. If considerable flux is
290 already at smaller scales (higher cut-off frequencies), as is the case in the zonal flux at cloud top in the eastern leg, the smaller eddies play a more important role.

The results suggest that momentum fluxes carried by eddies of different scales can counteract to reduce the overall flux. The relatively small flux (in the profiles, Figures 8,9) for instance in $\overline{u'w'}$ in the mixed layer on the western leg with thicker clouds, is produced by a positive $\overline{u'w'}$ carried by scales larger than 2.8 km up to maximally 7 km ($f_c = 0.025 - 0.01\text{Hz}$), which almost
295 compensates for the negative $\overline{u'w'}$ flux carried by turbulence on scales less than 2.8 km ($f_c = 0.025\text{Hz}$).

The same is true for $\overline{u'w'}$ near cloud top in the eastern leg with thinner clouds, and to a lesser extent, in the flux of $\overline{v'w'}$ in that leg within the mixed layer and near cloud tops. In the leg with thick clouds, the change in sign of the $\overline{v'w'}$ flux takes place already between 0.7 - 2.8 km. In other words: the profiles deviate from a profile where fluxes linearly decrease with height when scales beyond 1-2 km (cloud scale) play an important role.

300 Having less shear, for instance on 27 May, the fluxes are much smaller in general (Figure 8 and 9). There, smaller scales do not matter much and the flux is solely generated by the largest scales (not shown).

5 Conclusions

In this paper we aimed to investigate the variability in wind profiles and momentum fluxes under convective conditions, motivated by a lack of knowledge on the nature of the momentum flux profile in cloud-topped boundary layers. As in seminal
305 marine cumulus field studies, we used traditional in-situ turbulence measurements on board a Cessna aircraft that flew 50 - 100 km tracks at different legs below and within the cloud layer. These measurements were complemented with downward profiling Doppler Wind Lidar (DWL) measurements onboard a Falcon aircraft that flew the same track at higher altitudes. DWL are not typically employed in field studies of convection and clouds, but can help elucidate flows on mesoscales that accompany (organized) convection.

310 We carried out a limited number of flights in different wind and convective conditions: from a day with calm winds, strong directional shear and clear-sky and convective cumulus humilis (of maximum 500 m extent) to two days with pre- and post-frontal convection that was deeper and embedded in flows with strong winds and weak shear respectively strong shear.

First, we aimed to answer how the wind profiles and variance in the in-situ and DWL measurements compare. Both the mean horizontal wind and the variance of horizontal wind at given heights compared well despite the much larger horizontal
315 sampling scale of the DWL (7 km opposed to 70 m). Wind fluctuations on scales larger than turbulence and (cloudy) updrafts




Figure 12. Scale contributions to the momentum flux $\overline{u'w'}$ (upper panels) and $\overline{v'w'}$ (lower panels) for different heights in the atmosphere. The bars in each panel represent the flux contribution, in which the left-most bar only includes small scales (frequencies exceeding 0.2 Hz) and the right-most bar includes all scales that are represented in the measurements limited to 7 km (with a cut-off frequency of 0.01 Hz, excluding lower frequencies). Filter scales are 0.20, 0.15, 0.10, 0.05, 0.025, and 0.001 Hz that, when assuming a cruising speed of 70 m s^{-1} , correspond to length scales of 350, 467, 700, 1400, 2800, and 7000 m. The black bars represent the western leg with thicker clouds, the red ones the eastern leg with thinner clouds.

(on meso-gamma scales from 2 - 20 km) were found to be up to $\pm 5 \text{ m s}^{-1}$. This emphasises that flight legs need to be sufficiently long to capture mesoscale fluctuations in the mean wind (and its variance).

The profiles of wind derived from the few sampling legs of the Cessna compared reasonably well with the DWL profiles, in other words, the wind profile did not evolve majorly during the course of the flight as the Cessna transitioned from one leg
 320 altitude to another. The DWL also revealed the location of up- and downdrafts, such as negative vertical velocities at the top of the boundary layer where thermal plumes encounter a warmer environment and become negatively buoyant.

Second, we asked whether the measured momentum flux profiles in the sub-cloud layer and in the cloud layer are in line with our expectation, as well as to what extent we can explain them from convective updrafts alone. On the same flight day and even
 325 on the same track, substantial differences in the momentum flux could be seen associated with regions of convection. Most of the momentum flux profiles revealed down-gradient momentum transport that was generally strongest within the mixed-layer and decreasing towards cloud tops, but counter-gradient transport was also observed on the third flight day with post-frontal

convection and strong shear, on which distinct alternating patches of clear-sky and sheared clouds of about 1 km deep were seen. Different flight segments, even on that same flight day, had a very different momentum flux profile that was not explained by turbulent transport across the local vertical gradient in the wind.

330 Although momentum fluxes generally increased in areas with (cloudy) updrafts, the contribution of the updraft to the total momentum flux was typically (only) a third to two-thirds, which is comparatively small considering the contribution of the updraft to the buoyancy flux (pending). Horizontal momentum perturbations carried by updrafts below and within clouds were clearly distinguishable on all flights, but especially large on the third flight (up to almost 5 m s^{-1} in the cloud layer).

Finally, we asked which scales contribute significantly to wind variance and the momentum flux. We found that scales
335 beyond 1 km contribute significantly to the momentum flux and there is clear evidence for compensating flux contributions from different scales. In the post-frontal convection day, cancellation of fluxes of different signs may have explained a differing momentum flux profile with relatively small fluxes in the well-mixed sub-cloud layer (compared to the fluxes in the cloud layer).

The limited number of flights do not allow us to draw a general conclusion across a wide range of convective states. Nev-
340 ertheless, they highlight that momentum flux profiles and their variability require understanding of motions across a range of scales, with non-negligible contributions of the clear-sky fluxes and of mesoscales that may be coupled to the convection. Wind lidars can help elucidate the flows on larger than cloud scales and should be used more deliberately in studies of clouds and their spatial organization.

Data availability. Data is made publically available on the 4TU Data Repository. DOI 10.4121/18614102

345 *Author contributions.* M.K. and L.N. conceptualized the study, M.K. was responsible for the data analysis, and writing of the manuscript. C.M. was responsible for the technical preparation of the flight data and his experience aided the decision for an approach for eddy covariance estimation significantly. L.N. was also involved in the supervision. All authors provided critical comments on the quality of the work.

Competing interests. No competing interests are present.

Acknowledgements. This project has received funding from the European Research Council (ERC) under the European Union's Horizon
350 2020 research and innovation program (Starting Grant Agreement 714918). We want to thank Margaret LeMone and Chris Fairall for taking the time to review this paper. Their constructive comments led to improvements in this paper.

References

- Baidar, S., Bonin, T., Choukulkar, A., Brewer, A., and Hardesty, M.: Observation of the urban wind island effect, EPJ Web Conferences, <https://doi.org/10.1051/epjconf/202023706009>, 2020.
- 355 Bakhshi, R. and Sandborn, P.: Maximizing the returns of LIDAR systems in wind farms for yaw error correction applications, *Wind Energy*, 23, 1408–1421, <https://doi.org/https://doi.org/10.1002/we.2493>, 2020.
- Banta, R., Pichugina, Y., Kelley, N., Hardesty, R., and Brewer, W.: Wind Energy Meteorology: Insight into Wind Properties in the Turbine-Rotor Layer of the Atmosphere from High-Resolution Doppler Lidar, *Bulletin of the American Meteorological Society*, 94, 883 – 902, <https://doi.org/10.1175/BAMS-D-11-00057.1>, 2013.
- 360 Brilouet, P.-E., Lothon, M., Etienne, J.-C., Richard, P., Bony, S., Lernoult, J., Bellec, H., Vergez, G., Perrin, T., Delanoë, J., Jiang, T., Pouvesle, F., Lainard, C., Cluzeau, M., Guiraud, L., Medina, P., and Charoy, T.: The EUREC⁴A turbulence dataset derived from the SAFIRE ATR 42 aircraft, *Earth System Science Data*, 13, 3379–3398, <https://doi.org/10.5194/essd-13-3379-2021>, 2021.
- Browning, K. A. and Wexler, R.: The Determination of Kinematic Properties of a Wind Field Using Doppler Radar, *Journal of Applied Meteorology and Climatology*, 1, 105–113, <https://doi.org/10.1175/1520-0450>, 1968.
- 365 Chouza, F., Reitebuch, O., Jähn, M., Rahm, S., and Weinzierl, B.: Vertical wind retrieved by airborne lidar and analysis of island induced gravity waves in combination with numerical models and in situ particle measurements, *Atmospheric Chemistry and Physics*, 16, 4675–4692, <https://doi.org/10.5194/acp-16-4675-2016>, 2016.
- Dixit, V., Nuijens, L., and Helfer, K.: Counter-gradient momentum transport through subtropical shallow convection in ICON-LEM simulations, *Journal of Advances in Modeling Earth Systems*, p. e2020MS002352, <https://doi.org/https://doi.org/10.1029/2020MS002352>,
370 e2020MS002352 2020MS002352, 2021.
- George, G., Halloran, G., Kumar, S., Indira Rani, S., Bushair, M., Jangid, B., George, J., and Maycock, A.: Impact of Aeolus horizontal line of sight wind observations in a global NWP system, *Atmospheric Research*, 261, 105742, <https://doi.org/https://doi.org/10.1016/j.atmosres.2021.105742>, 2021.
- Gisinger, S., Wagner, J., and Witschas, B.: Airborne measurements and large-eddy simulations of small-scale gravity waves at the tropopause inversion layer over Scandinavia, *Atmospheric Chemistry and Physics*, 20, 10091–10109, <https://doi.org/10.5194/acp-20-10091-2020>,
375 2020.
- Helfer, K. C. and Nuijens, L.: The Morphology of Simulated Trade-Wind Convection and Cold Pools Under Wind Shear, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035148, <https://doi.org/https://doi.org/10.1029/2021JD035148>, e2021JD035148 2021JD035148, 2021.
- 380 Horányi, A., Cardinali, C., Rennie, M., and Isaksen, L.: The assimilation of horizontal line-of-sight wind information into the ECMWF data assimilation and forecasting system. Part I: The assessment of wind impact, *Quarterly Journal of the Royal Meteorological Society*, 141, 1223–1232, <https://doi.org/https://doi.org/10.1002/qj.2430>, 2015.
- Jungo, G. V. and Porté-Agel, F.: Measurement procedures for characterization of wind turbine wakes with scanning Doppler wind LiDARs, *Adv. Sci. Res.*, p. 71–75, <https://doi.org/10.5194/asr-10-71-2013>, 2013.
- 385 Käsler, Y., Rahm, S., Simmet, R., and Kühn, M.: Wake Measurements of a Multi-MW Wind Turbine with Coherent Long-Range Pulsed Doppler Wind Lidar, *Journal of Atmospheric and Oceanic Technology*, 27, 1529 – 1532, <https://doi.org/10.1175/2010JTECHA1483.1>, 2010.

- Krishnamurthy, R., Choukulkar, A., Calhoun, R., Fine, J., Oliver, A., and Barr, K.: Coherent Doppler lidar for wind farm characterization, *Wind Energy*, 16, 189–206, <https://doi.org/https://doi.org/10.1002/we.539>, 2013.
- 390 LeMone, M. and Pennell, W.: The Relationship of Trade Wind Cumulus Distribution to Subcloud Layer Fluxes and Structure, *Monthly Weather Review*, 104, 524–539, [https://doi.org/10.1175/1520-0493\(1976\)104<0524:TROTWC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1976)104<0524:TROTWC>2.0.CO;2), 1976.
- Lenschow, D. and Stephens, P.: The role of thermals in the convective boundary layer, *Boundary-Layer Meteorology*, 19, 509–532, <https://doi.org/10.1007/BF00122351>, 1980.
- Lenschow, D. H., Mann, J., and Kristensen, L.: How Long Is Long Enough When Measuring Fluxes and Other Turbulence Statistics?, *Journal of Atmospheric and Oceanic Technology*, 11, 661 – 673, [https://doi.org/10.1175/1520-0426\(1994\)011<0661:HLILEW>2.0.CO;2](https://doi.org/10.1175/1520-0426(1994)011<0661:HLILEW>2.0.CO;2), 1994.
- 395 Li, Z., Zuidema, P., and Zhu, P.: Simulated Convective Invigoration Processes at Trade Wind Cumulus Cold Pool Boundaries, *Journal of the Atmospheric Sciences*, 71, 2823 – 2841, <https://doi.org/10.1175/JAS-D-13-0184.1>, 2014.
- Liu, Z., Barlow, J., Chan, P., Fung, J. C. H., Li, Y., Ren, C., Mak, H. W. L., and Ng, E.: A Review of Progress and Applications of Pulsed Doppler Wind LiDARs, *Remote Sensing*, 11, 2522, <https://doi.org/10.3390/rs11212522>, 2019.
- 400 Lux, O., Lemmerz, C., Weiler, F., Marksteiner, U., Witschas, B., Rahm, S., Geiß, A., and Reitebuch, O.: Intercomparison of wind observations from the European Space Agency’s Aeolus satellite mission and the ALADIN Airborne Demonstrator, *Atmospheric Measurement Techniques*, 13, 2075–2097, <https://doi.org/10.5194/amt-13-2075-2020>, 2020.
- Mallaun, C., Giez, A., and Baumann, R.: Calibration of 3-D wind measurements on a single-engine research aircraft, *Atmospheric Measurement Techniques*, 8, 3177–3196, <https://doi.org/10.5194/amt-8-3177-2015>, 2015.
- 405 Mann, J., Peña, A., Bingöl, F., Wagner, R., and Courtney, M. S.: Lidar Scanning of Momentum Flux in and above the Atmospheric Surface Layer, *Journal of Atmospheric and Oceanic Technology*, 27, 959 – 976, <https://doi.org/10.1175/2010JTECHA1389.1>, 2010.
- Nicholls, S. and LeMone, M.: The Fair Weather Boundary Layer in GATE: The Relationship of Subcloud Fluxes and Structure to the Distribution and Enhancement of Cumulus Clouds, *Journal of Atmospheric Sciences*, 37, 2051 – 2067, [https://doi.org/10.1175/1520-0469\(1980\)037<2051:TFWBLLI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2051:TFWBLLI>2.0.CO;2), 1980.
- 410 Pu, Z., Zhang, L., Zhang, S., Gentry, B., Emmitt, D., Demoz, B., and Atlas, R.: The Impact of Doppler Wind Lidar Measurements on High-Impact Weather Forecasting: Regional OSSE and Data Assimilation Studies, pp. 259–283, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-319-43415-5_12, 2017.
- Saggiorato, B., Nuijens, L., Siebesma, A. P., de Roode, S., Sandu, I., and Papritz, L.: The Influence of Convective Momentum Transport and Vertical Wind Shear on the Evolution of a Cold Air Outbreak, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001991, <https://doi.org/https://doi.org/10.1029/2019MS001991>, e2019MS001991 10.1029/2019MS001991, 2020.
- 415 Schaefer, A., Craig, G., Wernli, H., Arbogast, P., Doyle, J. D., McTaggart-Cowan, R., J., M., G., R., Ament, F., Boettcher, M., Bramberger, M., Cazenave, Q., Cotton, R., Crewell, S., Delanoë, J., Dörnbrack, A., Ehrlich, A., Ewald, F., Fix, A., Grams, C. M., Gray, S. L., Grob, H., Groß, S., Hagen, M., Harvey, B., Hirsch, L., Jacob, M., Kölling, T., Konow, H., Lemmerz, C., Lux, O., Magnusson, L., Mayer, B., Mech, M., Moore, R., Pelon, J., Quinting, J., Rahm, S., Rapp, M., Rautenhaus, M., Reitebuch, O., Reynolds, C. A., Sodemann, H., Spengler, T.,
- 420 Vaughan, G., Wendisch, M., Wirth, M., Witschas, B., Wolf, K., and Zinner, T.: The North Atlantic Waveguide and Downstream Impact Experiment, *Bulletin of the American Meteorological Society*, 99, 1607 – 1637, <https://doi.org/10.1175/BAMS-D-17-0003.1>, 2018.
- Schlemmer, L., Bechtold, P., Sandu, I., and Ahlgrimm, M.: Uncertainties related to the representation of momentum transport in shallow convection, *Journal of Advances in Modeling Earth Systems*, 9, 1269–1291, <https://doi.org/https://doi.org/10.1002/2017MS000915>, 2017.
- Schneemann, J., Theuer, F., Rott, A., Dörenkämper, M., and Kühn, M.: Offshore wind farm global blockage measured with scanning lidar, *Wind Energy Science*, 6, 521–538, <https://doi.org/10.5194/wes-6-521-2021>, 2021.
- 425

- Wagner, J., Dörnbrack, A., Rapp, M., Gisinger, S., Ehard, B., Bramberger, M., Witschas, B., Chouza, F., Rahm, S., Mallaun, C., Baumgarten, G., and Hoor, P.: Observed versus simulated mountain waves over Scandinavia – improvement of vertical winds, energy and momentum fluxes by enhanced model resolution?, *Atmospheric Chemistry and Physics*, 17, 4031–4052, <https://doi.org/10.5194/acp-17-4031-2017>, 2017.
- 430 Weissmann, M., Busen, R., Dörnbrack, A., Rahm, S., and Reitebuch, O.: Targeted Observations with an Airborne Wind Lidar, *Journal of Atmospheric and Oceanic Technology*, 22, 1706 – 1719, <https://doi.org/10.1175/JTECH1801.1>, 2005.
- Witschas, B., Rahm, S., Dörnbrack, A., Wagner, J., and Rapp, M.: Airborne Wind Lidar Measurements of Vertical and Horizontal Winds for the Investigation of Orographically Induced Gravity Waves, *Journal of Atmospheric and Oceanic Technology*, 34, 1371–1386, <https://doi.org/https://doi.org/10.1175/JTECH-D-17-0021.1>, 2017.
- 435 Witschas, B., Lemmerz, C., Geiß, A., Lux, O., Marksteiner, U., Rahm, S., Reitebuch, O., and Weiler, F.: First validation of Aeolus wind observations by airborne Doppler wind lidar measurements, *Atmospheric Measurement Techniques*, 13, 2381–2396, <https://doi.org/10.5194/amt-13-2381-2020>, 2020.
- Yuan, J., Xia, H., Wei, T., Wang, L., Yue, B., and Wu, Y.: Identifying cloud, precipitation, windshear, and turbulence by deep analysis of the power spectrum of coherent Doppler wind lidar, *Optics Express*, 28, 37 406–37 418, 2020.
- 440 Zhan, L., Letizia, S., and Valerio Iungo, G.: LiDAR measurements for an onshore wind farm: Wake variability for different incoming wind speeds and atmospheric stability regimes, *Wind Energy*, 23, 501–527, <https://doi.org/https://doi.org/10.1002/we.2430>, 2020.
- Zhu, P.: On the Mass-Flux Representation of Vertical Transport in Moist Convection, *Journal of the Atmospheric Sciences*, 72, 4445 – 4468, <https://doi.org/10.1175/JAS-D-14-0332.1>, 2015.
- Zuidema, P., Li, Z., Hill, R., Bariteau, L., Rilling, B., Fairall, C., Brewer, W., Albrecht, B., and Hare, J.: On Trade Wind Cumulus Cold Pools, 445 *Journal of the Atmospheric Sciences*, 69, 258 – 280, <https://doi.org/10.1175/JAS-D-11-0143.1>, 2012.