

# Influence of air mass origin on microphysical properties of low-level clouds in a subarctic environment.

Konstantinos M. Doulgeris<sup>1</sup>, Ville Vakkari<sup>1,2</sup>, Ewan J. O'Connor<sup>1</sup>, Veli-Matti Kerminen<sup>4</sup>  
Heikki Lihavainen<sup>1,3</sup> and David Brus<sup>1</sup>

<sup>1</sup>Finnish Meteorological Institute, Erik Palménin aukio 1, P.O. Box 503, FIN-00100 Helsinki, Finland

<sup>2</sup>Atmospheric Chemistry Research Group, Chemical Resource Beneficiation, North-West University, Potchefstroom, South Africa

<sup>3</sup>Svalbard Integrated Arctic Earth Observing System (SIOS), SIOS Knowledge Centre, Svalbard Science Centre, P.O. Box 156, 9171 Longyearbyen, Norway

<sup>4</sup>Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, Helsinki, Finland

Correspondence to: K.D (konstantinos.doulgeris@fmi.fi)

**Abstract:** In this work, an analysis was performed to investigate how different long-range transport air masses can affect ~~the occurrence and~~ the microphysical properties of low-level clouds in a clean subarctic environment. The cloud measurements ~~comprised~~ included in-situ and remote sensing ground-based techniques and were conducted during eight Pallas Cloud Experiments (PaCE) held in the autumn between 2004 and 2019. Each PaCE was carried out at the Pallas Atmosphere – Ecosystem Supersite, located in the Finnish subarctic region. Two cloud spectrometer ground setups were installed on the roof of the station to measure cloud microphysical properties: the Cloud, Aerosol and Precipitation Spectrometer probe (CAPS) and the Forward Scattering Spectrometer Probe (FSSP). Air mass histories were analyzed using the Lagrangian particle dispersion model FLEXPART in order to investigate the differences between five distinct source regions (Arctic, Eastern, Southern, Western and Local). We observed clear differences in the cloud microphysical properties for the air mass source regions. Arctic air masses were characterized by low liquid water content (LWC), low cloud droplet number concentration ( $N_c$ ), and comparatively large median volume and effective droplet diameter. Western region (marine North Atlantic) differed from Arctic by both higher  $N_c$  and LWC. Eastern region (continental Eurasia) had only a little higher LWC than Arctic, but substantially higher  $N_c$  and smaller droplet diameter. Southern region (continental Europe) had high  $N_c$  and LWC, and very similar droplet diameter to the Eastern region. Finally, the relationship between  $N_c$  and droplet size (i.e., the Twomey effect) was characterized for the different source regions, indicating that all region clouds were sensitive to increases in  $N_c$ . ~~The results showed that there was a connection between the microphysical properties of the clouds we sampled during PaCEs and the air mass source region. The occurrence of clouds at the measuring station was mainly connected with air masses coming from the continental regions. Higher values of cloud droplet number concentrations ( $N_c$ ) were related to air masses coming from the Southern and Eastern regions (continental), whereas the lowest values of  $N_c$  were related to Arctic and Western air masses (marine). In addition, the marine air masses were characterized by larger cloud droplets (median volume diameter approximately from 15 to 20  $\mu\text{m}$ ), in comparison to continental air masses~~

Formatted: Font: Italic

that were mainly characterized by cloud droplets with median volume diameters of approximately from 8 to 12  $\mu\text{m}$ . Finally, there was an indication that cloud droplets were more prone to grow in warm liquid clouds.

## 1. Introduction

Uncertainties in cloud processes and feedbacks are key challenges when developing climate projections (e.g., Boucher et al., 2013; Sherwood et al., 2020). Cloud microphysics and their dynamics are considered as a fundamental challenge (Morrison et al., 2020) due to their connection to the cloud radiative effect (e.g., Devenish et al., 2012; McFarquhar et al., 2020). Thus, it is necessary to distinguish between the effects of aerosol and varying meteorological conditions on clouds (Barthlott and Hoose, 2018) since the aerosol is influenced through meteorology by air mass history as well as cloud and precipitation processes (Rosenfeld et al., 2014). Long-range transport is significant when investigating the characteristics and the spatial distribution of aerosols (e.g., Raatz and Shaw 1984; Barrie 1986; Freud et al., 2017; Wang et al., 2020; Lee et al., 2022). Due to insufficient knowledge of the cloud dynamics and the interaction between aerosols and clouds (e.g., Painemal et al., 2014; Orbe et al., 2015a; Fuchs et al., 2017; Cho et al., 2021), it is important to understand how different air masses can influence the aerosols and the cloud microphysics (e.g., Painemal et al., 2014; Orbe et al., 2015a; Fuchs et al., 2017; Cho et al., 2021). Investigating subarctic clouds is of particularly high interest due to the Arctic amplification effect, since the Arctic surface energy budget and Arctic warming feedback are affected by cloud related radiative processes (e.g., Wendisch et al., 2019, Shupe et al., 2022).

Several observation efforts and experiments have been made to explore how air masses affect climate and the cloud macrophysical and microphysical properties (e.g., Hobbs and Rangno 1998; Gultepe et al., 2000; Orbe et al., 2015b; Solomon and Shupe 2019; Delgado et al., 2021). Hobbs and Rangno (1998) highlighted that air masses from the south resulted in the highest overall aerosol number concentration measured in altocumulus clouds over the Beaufort Sea. Gultepe et al. (2000) stated that Arctic clouds were affected by the air mass origin, which was strongly related to aerosol properties, dynamical and thermodynamical parameters. Gultepe and Isaac (2002) studied the cloud microphysics over the Arctic Ocean and found that there were differences in the number concentration, liquid water content and effective radius of Arctic clouds in air masses originating from the Arctic and Pacific Oceans. After investigating the air mass origin seasonality Orbe et al. (2015a) revealed that the northern hemisphere summer air-mass origin response to increases in greenhouse gases (Orbe et al., 2015b). Fuchs et al., (2017) highlighted the impact of air mass origin and dynamics on cloud property changes in the Southeast Atlantic during the biomass burning season based on a cluster analysis of 8 years of September data. Solomon and Shupe (2019) presented a case study of a sharp transition between high ice clouds and the formation of lower stratocumulus from Summit, Greenland, when a warm and moist air mass was advected to Greenland from lower latitudes. Iwamoto et al. (2021), using measurements

from a high mountain site located in Japan, showed that the cloud droplet number concentrations were significantly higher in continental air masses than in air masses from the Pacific Ocean. Patel and Jiang (2021) combined measurements of aerosol properties from a site located in Lamont, Oklahoma, with cluster analysis of back trajectories to study aerosol characteristics and their influences on cloud condensation nuclei (CCN) under various air mass environments and suggested that information on the aerosol chemical composition and mixing state are more crucial at lower supersaturations. Delgado et al. (2021), using aerosol and cloud measurements from a site in a tropical montane cloud forest on the Caribbean Island of Puerto Rico, suggested that air masses that arrived after passing over areas with anthropogenic emissions led to clouds with much higher cloud droplet concentrations. Cho et al. (2021) investigated wintertime cloud properties and radiative effects in connection with cold and warm air mass origins at Ny-Ålesund, Svalbard, using remote sensing measurements with cloud radar, ceilometer and microwave radiometer, and revealed that the effective radius of ~~ice~~-cloud particles in warm advection cases was approximately 5–10  $\mu\text{m}$  larger than that of cold advection cases at all altitudes.

One of the few sites that enables long-term in situ observations of cloud and aerosol properties in arctic and subarctic air masses is the Pallas Global Atmospheric Watch (GAW) station in northern Finland (e.g., Lihavainen et al., 2008; Hyvärinen et al., 2011; Anttila et al., 2012; Raatikainen et al., 2015; Gérard et al., 2019; Girdwood et al., 2020, 2022). However, after the initial case study (Lihavainen et al., 2008), which indicated a clear Twomey effect depending on air mass origin, no subsequent concerted study has investigated the effect of air mass origin on cloud microphysical properties at ~~Pallas~~. Pallas. In the Arctic, during autumn, the ultrafine aerosol number concentration and the occurrence of clean, natural Arctic background conditions is significantly increasing (Pernov et al., 2022). Subsequently, this allows us to ~~Here, we~~ focus in this work on quantifying the impact of air mass origin (e.g., clean arctic vs. long-range transported air from continental Europe) on the microphysical properties of low-level clouds and their patterns based on measurements at the Pallas GAW station. To our knowledge, this is the first study that was performed in a subarctic environment and connects extensive in situ cloud measurements to air mass origin. Size distribution is considered as one of the most important parameters of the cloud system due to its impact on the dynamics and microstructures within the cloud (Igel et al., 2017a, b). Measuring cloud microphysical properties, such as the median volume diameter and liquid water content, is of high importance for the identification and description of clouds (Pruppacher and Klett, 1977; Rosenfeld and Ulbrich, 2003; Donovan et al., 2015), cloud radiative properties and lifetime (Albrecht 1989; Small et al., 2009;), and the probability for which clouds precipitate (Rosenfeld and Ulbrich, 2003; Chang et al., 2019). We used *in situ* low-level cloud measurements from two ground-based cloud spectrometers from eight different years of campaigns to obtain the cloud droplet size distribution. A lidar ceilometer was used to monitor the cloud base. The FLEXible PARTicle (Flexpart) dispersion model was used to analyze the air mass history. A description of the measurement site and the instrumentation, as it was installed, is given in Sect. 2.1 and 2.2. Subsequently, in Sect. 2.3, a general overview of the campaigns is presented. In Sect. 2.4, we

present how the backward trajectories were calculated. In Sect. 3, the optimal threshold of travelling air masses within a region to represent an air mass type is identified, and a detailed analysis is made to find out to what extent the air mass type influences the microphysical properties of the low-level clouds. Finally, in Sect. 4, we summarize our main conclusions.


115

## 2. Methodology

### 2.1 Sampling Station

120 The measurements were conducted in subarctic Finland at the Pallas Atmosphere – Ecosystem Supersite ( $67^{\circ}580$  N,  $24^{\circ}070$  E), hosted by the Finnish Meteorological Institute. The site where the ground-based cloud spectrometers were installed was the Sammaltunturi station, located on a hilltop, 565 m above sea level (ASL) (Hatakka et al., 2003). The site where the ceilometer was installed was the Kenttäröva station, 347 m ASL, located at the foot of the same hill (Fig.1). A full description of the Pallas Atmosphere – Ecosystem Supersite can be found in Lohila et al. (2015).

125






Figure 1. a) Map of Finland showing the location of the Pallas Atmosphere – Ecosystem Supersite (yellow hut icon), and (b) map of the wider Pallas area showing the location of Sammallunturi (yellow hut icon) and Kenttäröva (yellow circle icon) stations (© Google Maps).

## 2.2 Instrumentation

The instruments that were used in this study are listed in Table 1, together with the measured and derived parameters, their uncertainties and their location. During PaCEs, we used ground-based in situ cloud spectrometers to monitor the cloud droplet size distribution, which has been recognized as a valid method for continuous cloud in situ measurements in the [Aerosol, Clouds and Trace Gases Research Infrastructure \(ACTRIS\)](#) network (Wandinger et al., 2018). Four microphysical parameters were derived from the measured size distribution (Droplet Measurement Technologies Manual, 2009; Doulgieris et al., 2020): the cloud droplet number concentration ( $N_c$ ,  $\text{cm}^{-3}$ ), the median volume diameter (MVD,  $\mu\text{m}$ ) and effective diameter (ED,  $\mu\text{m}$ ) of cloud droplets, and the cloud liquid water content (LWC,  $\text{g m}^{-3}$ ).

Two ground-based spectrometers were used installed on the roof of Sammallunturi station: the cloud, aerosol and precipitation spectrometer (CAPS) and the forward-scattering spectrometer probe

145 (FSSP-100, hereafter called FSSP for simplicity) (Fig. 2). CAPS was made by Droplet Measurement Technologies (DMT), Boulder, CO, USA). FSSP-100 (model SPP-100, DMT) was initially manufactured by Particle Measuring Systems (PMS Inc., Boulder CO, USA) and later acquired by DMT. The CAPS probe includes three instruments: the cloud and aerosol spectrometer (CAS), the cloud imaging probe (CIP), and the hot-wire liquid water content ( $LWC_{hw}$ ) sensor. [Only CAS data were used](#)  
150 [during this work](#). The size range of the CAS extends from 0.51 to 50  $\mu m$  and that of the FSSP from 0.5 to 47  $\mu m$  in diameter. In both CAPS and FSSP, the main measurement principle for the size detection is based on a conversion of the forward scattering of light into a size bin using Lorentz–Mie theory (Mie, 1908). Their main difference was that the CAPS was fixed and always heading to the main wind direction of the station (southwest,  $\sim 225^\circ$ ), whereas the FSSP-100 was deployed on a rotating platform to continuously face the wind. A description of both ground setups, installation, limitations and the methodology that was used is documented in Dougeris et al. (2020) and Dougeris et al. (2022). The  
155 instrument that monitored the cloud base was a lidar ceilometer [that was deployed at Kenttäröva site](#) (model CT25K, Vaisala Oyj, Vaisala users guide, 2002; Emeis et al., 2004), except in 2019 when it was replaced by a model CL31, Vaisala Oyj. The meteorological variables were monitored by an automatic  
160 weather station (model Milos 500, Vaisala Oyj) [that was deployed at Sammaltunturi site](#). All the weather sensors that were used in this work were described in Hatakka et al. (2003). The temperature was measured at 570 m ASL by a PT100 sensor, the horizontal visibility by a weather sensor (model FDP12P, Vaisala Oyj), the relative humidity by a HUMICAP (Vaisala Oyj), the barometric pressure by a BAROCAP (Vaisala Oyj) sensors, the wind direction by a heated wind vane and the wind speed by a  
165 heated cup (Vaisala Oyj).




Figure 2. a) CAPS and b) FSSP-100 ground setups as installed on the roof of Sammaltunturi station during PaCE 2017.

170 TABLE 1: Instrumentation that was used during PaCE along with measured and derived parameters, their sampling frequencies, accuracy of the instruments and the location that were installed.

Instrument	Measured, derived parameters	Sampling frequency	Accuracy	Location	References
CAS, DMT	Number size distribution of cloud droplets (0.51 to 50 $\mu\text{m}$ ); derived parameters $N_c$ , LWC, ED, MVD	1s	at ambient droplet concentrations of $500 \text{ cm}^{-3}$ , 27 %undercounting 20 %–30 % oversizing	Sammaltunturi	Doulgeris et al., 2020, 2022. Baumgardner 2001. Lance (2012)
FSSP, DMT	Number size distribution of cloud droplets (0.5 to 47 $\mu\text{m}$ ); derived parameters $N_c$ , LWC, ED, MVD	1s	$N_c$ accuracy: 16 % sizing accuracy: $\pm 3 \mu\text{m}$ LWC accuracy: 30 %–50 %	Sammaltunturi	Doulgeris et al., 2020, 2022 Brenguier 1989 Baumgardner et al. (2017) Baumgardner (1983)
Ceilometer CT25K, Vaisala	Cloud base altitude	60s	$\pm 2 \%$ $\pm 1/2 \times$ (resolution)	Kenttäröva	Vaisala users guide, 2002, Emeis et al., 2004
FD12P, Vaisala	Horizontal visibility	60s	$\pm 10 \%$ at 10 – 10.000 (m)	Sammaltunturi	Hatakka et., al 2003
PT100 sensor, Vaisala	Temperature	60s	$\pm 0.1(^{\circ}\text{C})$	Sammaltunturi	Hatakka et., al 2003
HUMICAP sensor, Vaisala	Relative humidity	60s	$\pm 0.8(\%) \text{ RH}$	Sammaltunturi	Hatakka et., al 2003
BAROCAP sensor, Vaisala	Barometric pressure	60s	$\pm 0.15(\%) \text{ (hPa)}$	Sammaltunturi	Hatakka et., al 2003
Wind vane, Vaisala	Wind direction	60s	$\pm 3 (^{\circ})$	Sammaltunturi	Hatakka et., al 2003
Heated cup, Vaisala	Wind speed	60s	$\pm 0.17 \text{ (ms}^{-1}\text{)}$	Sammaltunturi	Hatakka et., al 2003

### 2.3 Sampling campaigns

175

Measurements used in this study were conducted during the Pallas Cloud Experiments (PaCE). A description of the dataset (microphysical properties of clouds along with meteorological variables) that was obtained during PaCEs are available in Doulgeris et al. (2022). The PaCEs were approximately



two-month-long field campaigns conducted in the Finnish subarctic region at the Sammaltunturi station during autumn and lasted approximately from the beginning of September until the end of November. The reason for this choice was that during autumn the Sammaltunturi station is frequently inside a cloud, which allowed us to perform ground-based, continuous in situ cloud measurements (Hatakka et al., 2003). An overview of each campaign along with the availability of instruments and the hours of observations in cloud are presented in Table 2. During PaCEs, all measurements were performed with a 1 Hz acquisition frequency. For the data analysis, averages per minute from each instrument were calculated when the measuring site was inside a cloud. Each cloud event was inspected separately. Afterwards, in-situ cloud data were related to the air mass origin and classified accordingly. We only used measurements when the cloud spectrometers were facing the wind direction, as suggested by Doulgeris et al. (2020). Thus, since the CAS was fixed, data when the CAS was not facing the wind direction were disqualified from further analysis.

Fine particles at the Sammaltunturi are expected to be dominated by sulphate and particulate organic matter in continental air masses, while particulate organic matter, sodium and chlorine are the main components in marine air masses (Lihavainen et al., 2008, Brus et al., 2013a). Also, episodes of elevated concentrations of SO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> are possible in air masses arriving from the Kola Peninsula, which is a large source of SO<sub>2</sub> emissions (Kyro et al., 2014; Sipilä et al., 2020, Brus et al., 2013a, b). Elevated SO<sub>2</sub> concentrations and particle number concentrations in the accumulation-mode (0.1–1 µm in diameter) of the mass size distribution are also expected from air masses travelled over continental Europe. Total aerosol particle number concentrations at Sammaltunturi are typically low (average of 700 cm<sup>-3</sup>, in winter the daily averages may drop below 100 cm<sup>-3</sup>) (Hatakka et al., 2003; Komppula et al., 2003). Generally, in Finnish Lapland, aerosol particle number concentrations are expected to be the highest in air masses arriving from the Kola Peninsula (more than 1000 cm<sup>-3</sup>) and the lowest in marine air masses, especially in air originating from the Arctic Sea (often less than 100 cm<sup>-3</sup>) (Lihavainen et al., 2008). Higher particle number concentrations in the accumulation mode are also expected in air masses which have travelled over the continental Europe (Virkula et al., 1997). In Sammaltunturi, cloud condensation nuclei (CCN) concentrations are smaller than 100 cm<sup>-3</sup> for supersaturations from 0.1 to 0.5%. The aerosol particle population is dominated by the Aitken mode (30–100 nm in diameter) and a low hygroscopicity is expected (Paramonov et al., 2015).

TABLE 2: Overview of each campaign, including the starting and ending date, the availability of the ground cloud spectrometer probes and the ceilometer. ~~model.~~

Year	Starting date	Ending date	CAS	FSSP	Ceilometer	Cloud Observations CAPS (hours)	Cloud Observations FSSP (hours)
2004	25.10.	07.11.	Not available	On site	CT25K	-	42
2005	30.09.	05.10.	Not available	On site	CT25K	-	45.4




2009	11.09.	09.10.	Not available	On site	CT25K	-	34.2
2012	14.09.	30.10.	On site	On site from 09.10	CT25K	477.5	50
2013	14.09.	28.11.	On site from 15.10	On site	CT25K	483.5	492.6
2015	24.09.	02.12.	On site from 06.10	On site	CT25K	528.4	561.9
2017	18.09.	29.11.	On site	Not available	CT25K	681.8	-
2019	20.09.	24.11.	On site	Not available	CL31	479.6	-

## 2.4 Classification of air mass origin

Air mass origins were analyzed using the Lagrangian particle dispersion model FLEXPART version 10.4 (Seibert and Frank, 2004; Stohl et al., 2005; Piss0 et al., 2019). FLEXPART was run backward in time to calculate potential emission sensitivity (PES) fields. PES in a particular grid cell is proportional to the air mass residence time in that cell and was calculated in units of seconds (Seibert and Frank, 2004; Piss0 et al., 2019). ERA5 reanalysis by European Centre for Medium-Range Weather Forecasts (ECMWF) was used as meteorological input fields for FLEXPART at 1 hour temporal resolution and 0.25° resolution in latitude and longitude. In vertical, ERA5 levels 50 to 137 were used, which corresponds approximately to the lowest 20 km above surface. The model domain was from 125° W to 75° E and 10° N to 85° N, which was large enough to contain 96 h simulations backward in time. FLEXPART runs were initiated at an hourly time resolution for the in-cloud measurement periods at Sammaltunturi. The retro plume release height was set to 560-660 m ASL, as the terrain height in ERA5 at the site was approximately 300 m ASL. The PES output resolution was set to 0.2° latitude and longitude with a 250-m height resolution up to 5 km and two additional output levels at 10 km and 50 km.

The air mass source regions for the Sammaltunturi site were divided into five categories: Arctic, Eastern, Southern, Western and Local (Fig. 3). The division was based on previous studies that were conducted at Sammaltunturi (e.g., Aalto et al., 2003; Eneroth et al., 2005; Tunved et al., 2006; Asmi et al., 2011). Initially, the regions were classified using trajectories cluster analysis, following the method as Eneroth et al. (2003) proposed as proposed by Asmi et al. (2011). The choice of sectors represents roughly the characteristics of the region. The Arctic and Western regions represent marine areas, whereas the Eastern, Southern and Local regions represent continental areas. Figure 5a-4 illustrates an

235 example case for air masses arriving on 20.09.2012 at 11:00 UTC at Sammaltunturi. In Fig. 4a PES is  
summed up for the full duration of the 96-hour backward simulation and for all output heights at each  
latitude-longitude grid cell. Fig. 4b displays the vertical distribution of PES during the simulation.  
However, the information in Fig. 4b is not used in further analysis, but fraction of PES in each source  
region is calculated based on the integrated data presented in Fig. 4a. For the case in Fig. 4a, this results  
240 in PES fractions of 42% Local (area V), 35% Southern (area III) and 23% Western (area IV). Finally,  
Fig. 4c shows how the PES fractions evolve during 20.09.2012 at Sammaltunturi with the case in Fig.  
4a represented by the bar at 11 UTC. In Fig. 4c a clear change in air mass origin is observed at 18 UTC  
when the fraction of Arctic source region starts to increase, reaching up to 48% at 22 UTC.



245 Figure 3: Map of the air mass regions: I (Arctic), II (Eastern), III (Southern), IV (Western) and V  
(Local). Figure was adopted from Asmi et al., 2011.





Figure 4: a) Horizontal distribution of vertically integrated PES for air masses arriving at Sammaltunturi on 20.09.2012 11:00 UTC. Source regions are indicated with Roman numerals. b) Vertical distribution of PES in panel a as function of time before arrival at Sammaltunturi. c) Timeline of the PES fraction for each source region for 20.09.2012.

### 3.Results

#### 3.1 Local meteorological conditions

Figure 5 shows the daily average temperatures at 570 m observed at the Sammaltunturi measuring site for days with “cloud events”. [The seasonal range of temperatures from an average 4.5 °C \(SD 2.1 °C\) in September to -5.3 °C \(SD 1.8 °C\) in November and its interannual variability is revealed.](#) Days with “cloud events” were defined as the days when the station was at least 30 minutes immersed in a cloud. To identify the presence of a cloud at the station, four steps were followed. The droplet size distribution was checked from both cloud spectrometers; the relative humidity should be ~ 100 %; the horizontal visibility should be less than 1000 m, and a final inspection was performed visually using pictures

recorded by an automatic weather camera installed on the roof of the station. During days with no cloud  
 265 events, clouds could still exist at higher altitudes. Supercooled water droplets were expected at  
 temperatures  $<0^{\circ}\text{C}$  (usually during November and October of each campaign, in total 175 cloud events  
 with temperature  $<0^{\circ}\text{C}$  were sampled). Mixed-phase clouds, consisting of water vapor, ice particles  
 and supercooled liquid droplets, are frequent at temperatures from  $-10$  to  $-25^{\circ}\text{C}$  (Korolev et al., 2017,  
 Filioglou et al., 2019); however, they can be present up to temperatures of  $0^{\circ}\text{C}$  (Andronache, 2017).  
 270 During September, the average temperature was  $>0^{\circ}\text{C}$ , thus, the clouds were expected to consist of  
 liquid hydrometeors only (liquid droplets, drizzle drops and raindrops). Wind speed ranges during the  
 PaCEs were approximately from 0 to  $10\text{ m s}^{-1}$  and the average wind speed during each campaign was  
 around  $7\text{ m s}^{-1}$ . These values were lower in comparison to the probe air speed of both cloud ground-  
 based spectrometers (Doulgeris et al., 2022).







Figure 5: The daily averaged temperatures at the Sammaltunturi site for days with cloud events during all PaCE campaigns. The dotted line is used as a reference line for 0 °C temperature. The definition of a cloud event is provided in the text. The shaded area represents the corresponding standard deviations.

~~Figure 5: The daily averaged temperatures at the Sammaltunturi site for days with cloud events. The black solid line is used as a reference line for 0 °C temperature. The definition of a cloud event is provided in the text.~~

### 3.2. Identification of the air mass origin and its effect on the number concentration of cloud droplets.

First, we examined which was the optimal threshold of the PES fraction within one region that should be used for ~~this a particular~~ region to be representative of the air mass type.  $N_c$  was chosen to be used as a benchmark parameter. ~~The results are shown in Fig. 6.~~ Figure 6a shows hourly  $N_c$  values depending on the PES fraction of the Arctic region (~~in Arctic, during autumn, the ultrafine aerosol number concentration and the occurrence of clean, natural Arctic background conditions is significantly increasing, (Pernov et al., 2022))~~ ~~marine environment with no anthropogenic emissions~~). The lowest values of  $N_c$  ( $< 30 \text{ cm}^{-3}$ ) were related to PES fractions  $> 80\%$ . When the PES fraction was between 70 and 80%, the values of  $N_c$  were varying between 5 and  $80 \text{ cm}^{-3}$  and the highest values of  $N_c$  values ( $> 30 \text{ cm}^{-3}$ ) were related to PES fraction lower than 70%.

295



300

Figure 66: a) Hourly cloud droplet number concentration ( $N_c$ ) versus the different potential emission sensitivity (PES) fraction for the Arctic region. The shaded area represents the corresponding standard deviations, and b)  $N_c$  for each region and single PaCE campaign as they were measured by the cloud and aerosol spectrometer (CAS) and the forward scattering spectrometer probe (FSSP) where the PES fraction was within one region >80 % (black) and the PES fraction was within one region from 70 to 80 % (red).




Figure 7: Cloud droplet number concentration ( $N_c$ ) for each region and single PaCE campaign as they were measured by the cloud and aerosol spectrometer (CAS) and the forward-scattering spectrometer probe (FSSP) where the PES fraction was within one region >80 % (full symbols) and the PES fraction was within one region from 70 to 80 % (open symbols). Error bars indicate the corresponding standard deviation.

To achieve a generalization of the large-scale air mass influence on microphysical cloud properties, the characteristics of all regions were intercompared. In Fig. 6b-7 we summarize cloud  $N_c$  measurements from both CAPS and FSSP. Each point represents a single PaCE campaign for different regions according to the classification criteria that were introduced previously in this section. We present each campaign and instrument to demonstrate that there were no obvious changes through years or possible malfunction of the instruments that were used and could produce biased results. Two different levels of the PES fraction were chosen to be further investigated; i) when the PES fraction was within one region more than 80% and ii) when the PES fraction was within one region between 70 and 80 %. For the first level, the highest values of  $N_c$  (approximately 100-200  $\text{cm}^{-3}$ ) were clearly associated with Southern air masses, whereas the lowest ones (approximately 20  $\text{cm}^{-3}$ ) were observed in Arctic air masses. In general, marine air masses (Arctic, Western) arriving at Sammaltunturi resulted in lower values of  $N_c$  compared with continental air masses. However, there was also a difference (approximately by a factor of 2) between marine air masses:  $N_c$  in air masses travelling over the Atlantic Ocean, or the Norwegian Sea, were higher than those in air masses arriving from the Arctic Sea. For aerosol



populations, higher values of  $N_{CCN}$  are expected in more polluted air masses (Southern and Eastern air masses, due to emissions from Europe and Kola peninsula) (Virkula et al., 1997; Jaatinen et al., 2014; Sipila et al., 2021). Averaged temperatures at Sammaltunturi for each air mass were  $-3.1^{\circ}\text{C}$  (SD  $2.5^{\circ}\text{C}$ ),  $-2.2^{\circ}\text{C}$  (SD  $5.9^{\circ}\text{C}$ ),  $1.3^{\circ}\text{C}$  (SD  $3.9^{\circ}\text{C}$ ) and  $-2.8^{\circ}\text{C}$  (SD  $2.01^{\circ}\text{C}$ ) for the arctic, eastern, southern and western region respectively. Furthermore, there was no clear indication that there was any trend in  $N_c$  through different years of PaCEs. In this work,  $N_{CCN}$  measurements at different supersaturations were not conducted.

For PES fraction between 70 and 80 %, the impact of the air mass type on the  $N_c$  changed as the differences in  $N_c$  were less than for PES > 80%. ~~(Fig. 5b)~~. Clouds that were associated with Southern air masses had slightly higher values of  $N_c$  (approximately  $60\text{--}80\text{ cm}^{-3}$ ) in comparison with clouds from the other regions (approximately  $20\text{--}60\text{ cm}^{-3}$ ). As a result, for further analysis in this work, we decided to exclude measurements that were performed when the PES fraction was between 70 and 80 %. Thus, we considered that > 80 % of the PES fraction within a particular region would be the optimal threshold to represent an air mass type during PaCEs. Using the > 80% PES fraction from one source region as a criterion for further analysis left 492 hours of in-cloud measurements with the CAS and 214 hours of in-cloud measurements with the FSSP probe (from total 2004 hours of in situ cloud data 706 hours belong to non-mixed air mass origin), respectively, which ensured statistically robust results. Cloud observation related to Arctic, Eastern, Southern, Western and Local air masses were 118, 275, 152, 118 and 43 hours, respectively. The observation hours related to each region for each PaCE are presented in Table S2 of the supplementary material (SM). Clouds that were related with local air masses were excluded due to relatively small number of observations.


Based on the air mass origin classification, a statistical analysis was made to investigate the frequency of the air masses during cloud events at the measuring site. When the air masses were not mixed, the occurrence of clouds at the station related to continental and marine air masses in 31.9 % and 14.3 % of the cases, respectively. Focusing on each region separately, 29.6 % of the cloud's occurrence seemed to be related to Southern and Eastern air masses and 7.4 % were related to Arctic air masses, although the predominant air mass at Sammaltunturi was from the Arctic (Asmi et al., 2012).

### 3.3. Effect of the air mass origin on the cloud droplets size.

In this section, we focused on investigating the size distribution of the cloud droplets and the derived parameters ED, MVD and LWC. ED and MVD are strongly dependent on the shape of the cloud droplet size distribution, while LWC is a function of both  $N_c$  and sizes of the cloud droplets. To achieve a generalization, size distributions of cloud droplets related to each air mass origin for ~~each single~~ all PaCE campaigns are presented in Figure 7.8. Cloud droplet size distributions originating from marine regions (Arctic, Western) had a relatively broad shape with presence of large ( $10\text{--}20\text{ }\mu\text{m}$ ) droplets, whereas in continental air masses there was a clear absence of large cloud droplets. In general, the

Formatted: Font: Italic

average size distribution showed a spectrum with more droplets at small size ranges when the masses were continental and more droplets in larger size ranges when the air masses were marine. Cloud droplets larger than  $16.0\ \mu\text{m}$  started to appear in clouds that were characterizes by Arctic and western air masses. On the other hand, clouds that were characterized by Eastern and Southern air masses had cloud droplets mainly in the range from  $5$  to  $10\ \mu\text{m}$ . Values of  $N_c$  for different sizes of the cloud droplets suggest that higher aerosol loadings lead to higher number concentrations of cloud droplets and smaller cloud droplet effective diameters. This result is consistent with the Twomey effect (Twomey, 1977), reported in several *in situ* observations (e.g. Twomey et al., 2005; Freud et al., 2008; Goren and Rosenfeld 2014; Paramonov et al., 2015). In general, in a cloud system, it is expected that for a relatively constant LWC, the effective diameter of cloud droplets decreases as their number concentration increases.






Figure 78. Cloud droplet size distribution associated with the (a) Arctic, (b) Western, (c) Southern and (d) Eastern region as they were measured by the cloud and aerosol spectrometer (CAS) and the forward-scattering spectrometer probe (FSSP) during all PaCEs. Yearly average cloud droplet size distribution associated with the (a) Arctic, (b) Western, (c) Southern and (d) Eastern region as they were measured by the cloud and aerosol spectrometer (CAS) and the forward scattering spectrometer probe (FSSP).

We also investigated whether sizes of cloud droplets depend on the air temperature. For that reason, temperature bins of 4 °C range were created. Thus, the measurements were grouped at temperature bins of -10 to -6 °C, -6 to -2 °C, -2 to 2 °C and 2 to 6 °C. The mid temperature value of each bin was used to create Fig.-89 which shows that cloud droplets appeared to be more prone to grow at temperatures larger than -2 °C. A hypothesis to explain such growth could be the collision-coalescence procedures that can take place in warm clouds (e.g., Xue et al., 2008; Pruppacher and Klett 2010; Lohmann et al., 2016). In this study, all the sampled clouds are considered as warm clouds, however those at warmer air temperatures seem to consist of larger droplets. Both MVD and ED had a similar behavior. When the clouds were characterized by Arctic air masses, MVD and ED were approximately 15 μm within our temperature spectrum. The decrease of particle size for the 'Arctic' subsample in the FSSP data above 0 °C was due to the relatively low amount of observation in this temperature range (2 hours of observation). The observation hours related to each temperature bin for

each PaCE are presented in Table S3 of the SM. This could possibly be related to the fact that warm clouds that were associated with Arctic air masses had smaller lifetimes compared to the other regions within our dataset. For clouds related to eastern air masses MVD and ED were approximately 9  $\mu\text{m}$  when the temperature was below 0  $^{\circ}\text{C}$  and showed approximately 5–6  $\mu\text{m}$  larger hydrometeors in warmer clouds (above 0  $^{\circ}\text{C}$ ). However, more observations on existing and wider temperature ranges are needed to statistically ensure those results. In our case, we obtained approximately from 11 to 99 hours of observations for each temperature bin.




Figure 89. Hourly averages of median volume diameter (MVD) and effective diameter (ED) values versus temperature for all PaCE campaigns as they were measured by the cloud and aerosol spectrometer (CAS) and the forward-scattering spectrometer probe (FSSP) where the PES fraction was within one region >80 %. Solid lines were made to lead readers' eyes.

LWC of low-level clouds for the different air mass types are summarized for each PaCE campaign (Fig. 9a10a). The Arctic air masses were related to the lowest values of LWC (approximately 0.025  $\text{g m}^{-3}$ ), whereas the Southern air masses were related to the highest values of LWC (> 0.05  $\text{g m}^{-3}$ ). Western and Eastern air masses were related to LWC values of approximately 0.025 to 0.05  $\text{g m}^{-3}$ .

410 In this study, LWC of continental air masses were, on average, larger than those of marine air masses. This is [also](#) reflected in the higher  $N_c$  of continental air masses (Fig. [6b7](#)), as LWC is a function of both  $N_c$  and size of cloud droplets. In Figure [9b10b](#), the relation between the  $N_c$  and MVD is plotted. The points were divided into three different levels according to the measured LWC. The values of MVD were ranging from  $\sim 9$  to  $19 \mu\text{m}$ . MVD was larger for higher values of LWC and decreased with an

415 increasing cloud droplet number concentration for each LWC category. The LWC values of the clouds we sampled ( $\sim 0.03 \text{ g m}^{-3}$  for marine and  $\sim 0.06 \text{ g m}^{-3}$  for continental conditions) are comparable to those observed in several other *in situ* cloud studies (e.g., Gultepe and Isaac 1997; Zhao et al., 2012; Lu et al., 2014; Guyot et al., 2015; Dione et al., 2020).

Formatted: Justified




Figure 910: a) Statistical description of Liquid water content (LWC) for each region as they were it was measured by the cloud and aerosol spectrometer (CAS) and the forward-scattering spectrometer probe (FSSP) where PES was within one region  $>80 \%$ . The box plot for each region is also provided, b)

425 Median volume diameter (MVD) as a function of total cloud droplets number concentration ( $N_c$ ) for three different categories of LWC. Each point represents a single PaCE campaign for different regions.

### 3.4 Influence of the vertical position of probe on the derived parameters

In this section, we focus on investigating how the derived parameters change with changes in the vertical position (altitude) of the sampling probe.  $H$  (m) (e.g., Martins et al., 2011; McDonald et al., 2018; Alexandrov et al., 2020). Under theoretical adiabatic conditions, the vertical profile of LWC is expected to increase linearly with height above cloud base, with a constant gradient that is dependent on the temperature and pressure at cloud base (Brennguier, 1991).  $N_c$  is constant through the vertical profile of the cloud layer, while the size of the droplets increases with altitude (Pawlowska et al., 2006). Assuming homogenous mixing, this expectation of the cloud microphysical profile also holds for ‘scaled-adiabatic’ conditions which include the entrainment of drier air (Boers et al., 2000). In reality, there are more processes to consider, which lead to departures from this ideal condition, particularly towards cloud top (Pawlowska et al., 2006). As already discussed in Section 2.2., both ground-based spectrometers were fixed in one vertical position. Thus, there were cases that we sampled with different layers of a cloud in a range of 120 meters from the cloud base. The ground-based spectrometers were placed at the Sammaltunturi, 210 m above the ceilometer (installed at the Kenttäröva site). Kenttäröva is located 4.3 km to the east of hilltop Sammaltunturi station. Since the Sammaltunturi station is on a top of an Arctic fell, cloud formation and properties could also be influenced by the local topography via changes in turbulence or orographic flows. The ceilometer’s resolution in estimation of the cloud base was 30 meters.

In Fig. 1011, a statistical description of both-MVD and-ED-are is presented as they-it-were-was derived from both cloud spectrometers for-in-five different altitudes-ss-of-the-cloud-base-above-cloud-base. The distance of the cloud spectrometer was relative to the cloud base. From this analysis, it is apparent that there was no strong dependency between the vertical position of CAS and FSPP in the cloud and the derived sizing parameters. However, it is expected that number concentration provides a robust signal and can clearly be linked to air mass origin, whereas MVD and ED has some extra uncertainties depending on the altitude with respect to cloud base.

Formatted: English (United States)




Figure 11. Statistical description of hourly averages of median volume diameter ( $MVD$ ) as they were measured by the cloud and aerosol spectrometer (CAS) and the forward-scattering spectrometer probe (FSSP) where PES was within one region  $>80\%$  for five different levels of the position of the probes inside the cloud ( $H$ ) (relative distance of the cloud ground-based spectrometer). Cloud base was measured at the Kenttäröva station.

The cloud base height ( $h$ ) (relative distance of the cloud ground-based spectrometer) as it was measured at the Kenttäröva station versus hourly averages of median volume diameter ( $MVD$ ) and effective



diameter ( $ED$ ) values for all PaCEs as they were measured by the cloud and aerosol spectrometer (CAS) and the forward scattering spectrometer probe (FSSP) where PES was within one region >80 %.

#### 4. Summary and conclusions

Our main goal during this work was to quantify the effect of air mass origin on cloud microphysical properties in a clean subarctic environment. Thus, the impact of different air masses on cloud properties in the subarctic Finland was investigated based on data from 8 Pallas Cloud Experiments (PaCEs) made during 2004–2019. For measuring the cloud microphysical properties, we deployed two cloud ground-based spectrometer probes: the cloud and aerosol spectrometer and the forward-scattering spectrometer probe. For performing the air mass sources classification, the FLEXPART model was used with ERA5 meteorology. The air mass source regions were categorized into Arctic, Eastern, Southern, Western and Local sectors, with the Arctic and Western sectors representing marine air masses, and the Eastern, Southern and Local sectors representing continental airmasses.

Our analysis demonstrated that different air mass types had significant impacts on cloud microphysics. When 80 % of the potential emission sensitivity fraction was within a region, the observations were considered ~~to as~~ representative of that air mass type. ~~The occurrence of a cloud at the station was more frequent with air masses arriving from the Southern and Eastern regions than other source regions. Furthermore, C~~ontinental air masses led to the highest cloud droplet number concentrations ( $\sim 100\text{--}200\text{ cm}^{-3}$ ) and marine air masses to the lowest ones ( $\sim 20\text{ cm}^{-3}$ ). ~~The lowest values of cloud droplet concentration were related to clean arctic airmasses. Number concentration was expected to be a robust signal with no dependency from the vertical position of cloud spectrometer in the cloud. On the other hand, both effective radius and median volume diameter has some extra uncertainties depending on the altitude with respect to cloud base. We observed a clear relationship between air mass origin and cloud droplet number concentration. This connection is expected to be a robust signal as according to theoretical considerations (Brenquier 1991; Pawlowska et al., 2006), the measurements of cloud droplet number concentration do not depend on the vertical position of the cloud spectrometer within the cloud layer.~~ In general, the median volume diameter and effective ~~radius~~ diameter of cloud droplets was found to be influenced by the cloud droplet number concentration: Clouds associated with marine air masses had larger cloud droplets (ranging from 15 to 20  $\mu\text{m}$ ) in comparison with continental clouds (ranging from 8 to 12  $\mu\text{m}$ ). These results are in agreement with the Twomey effect (Twomey, 1977). ~~The above differences that were observed in cloud microphysical properties when the air masses were related to different regions show the need to investigate how the aerosol loading and meteorology of different airmasses along with local meteorological parameters change the cloud microphysics and to what scale.~~ Furthermore, there was an indication that cloud droplets in clouds in warmer air (from  $-2$  to  $6\text{ }^{\circ}\text{C}$ ) were more prone to grow. However, ~~more measurements are needed to confirm such temperature dependency of droplet sizes. all year round in~~

situ cloud measurements in the area are of high importance to confirm such temperature dependency of droplet sizes. Specifically, a large data set containing a wider temperature range needs to be obtained.

#### Data availability

The cloud probes and meteorological data used here are available in the Finnish Meteorological Institute (FMI) open data repository for each campaign and each cloud spectrometer ground setup individually (Doulgeris et al., 2021; Doulgeris et al., 2022). The FLEXPART simulations and the ceilometer dataset are available upon request to the corresponding author ([konstantinos.doulgeris@fmi.fi](mailto:konstantinos.doulgeris@fmi.fi)).

*Author contributions.* KD wrote the paper with contributions from all co-authors. HL planned and coordinated PaCE 2004, 2005, and 2009. HL and DB planned and coordinated PaCE 2012 and 2013. KMD and DB planned and coordinated PaCE 2015, 2017, and 2019. KMD and DB processed, analyzed, and quality controlled the data set. VV carried out the FLEXPART simulations. EOC provided the ceilometer data. VMK reviewed and edited the manuscript.

*Competing interests.* The authors declare no conflict of interest.

*Acknowledgements:* This work was supported by the Koneen Säätiö (grant no. 46-6817), Academy of Finland Flagship funding (grant no. 337552). This project has also received funding from the European Union, H2020 research and innovation program (ACTRIS-IMP, the European Research Infrastructure for the observation of Aerosol, Clouds, and Trace gases, no. 871115). The authors wish to acknowledge CSC – IT Center for Science, Finland, for computational resources. The authors would also like to thank all the people who help in PACE campaign measurements throughout the years.

#### References

Aalto, T., Hatakka, J. and Viisanen, Y.: Influence of air mass source sector on variations in CO<sub>2</sub> mixing ratio at a boreal site in northern Finland. *Boreal Env. Res.* 8, 285–393, 2003.

Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227–1230, 1989.

Alexandrov, M. D., Cairns, B., Emde, C., Ackerman, A. S., and van Diedenhoven, B. (2012): Accuracy assessments of cloud droplet size retrievals from polarized reflectance measurements by the research scanning polarimeter. *Remote Sensing of Environment*, 125, 92–111. <https://doi.org/10.1016/j.rse.2012.07.012>, 2012.

Andronache, C.: Mixed-phase Clouds: Observations and Modeling, Elsevier, Saint Louis, proQuest Ebook Central, available at: <https://ebookcentral.proquest.com/lib/fmi/detail.action?docID=5064425> (last access: 16 June 2022/2023), 2017.

- Anttila, T., Brus, D., Jaatinen, A., Hyvärinen, A.-P., Kivekäs, N., Romakkaniemi, S., Komppula, M., and Lihavainen, H.: Relationships between particles, cloud condensation nuclei and cloud droplet activation during the third Pallas Cloud Experiment, *Atmos. Chem. Phys.*, 12, 11435–11450, <https://doi.org/10.5194/acp-12-11435-2012>, 2012.
- Asmi, E., Kivekäs, N., Kerminen, V.-M., Komppula, M., Hyvärinen, A.-P., Hatakka, J., Viisanen, Y., and Lihavainen, H.: Secondary new particle formation in Northern Finland Pallas site between the years 2000 and 2010, *Atmos. Chem. Phys.*, 11, 12959–12972, <https://doi.org/10.5194/acp-11-12959-2011>, 2011.
- Barrie, L. A., 1986: Arctic air pollution: an overview of current knowledge, *Atmos. Environ.* 20, 643–663, 1986.
- Barthlott, C. and Hoose, C.: Aerosol effects on clouds and precipitation over central Europe in different weather regimes, *J. Atmos. Sci.*, 75, 4247–4264, <https://doi.org/10.1175/JAS-D-18-0110.1>, 2018.
- Baumgardner, D.: An analysis and comparison of five water droplet measuring instruments, *J. Appl. Meteorol.*, 22, 891–910, [https://doi.org/10.1175/1520-0450\(1983\)022<0891:AAACOF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1983)022<0891:AAACOF>2.0.CO;2), 1983.
- Baumgardner, D., Jonsson, H., Dawson, W., O'Connor D., and Newton R.: The cloud, aerosol and precipitation spectrometer (CAPS): A new instrument for cloud investigations, *Atmos. Res.*, 59–60, 251–264, [https://doi.org/10.1016/S0169-8095\(01\)00119-3](https://doi.org/10.1016/S0169-8095(01)00119-3), 2001.
- Baumgardner, D., Abel, S. J., Axisa, D., Cotton, R., Crosier, J., Field, P., Gurganus, C., Heymsfield, A., Korolev, A., Krämer, M., Lawson, P., McFarquhar, G., Ulanowski, Z., and Um, J.: Cloud Ice Properties: In Situ Measurement Challenges, *Meteor. Mon.*, 58, 9.1–9.23, <https://doi.org/10.1175/AMSMONOGRAPH-D-16-0011.1>, 2017.
- Boers, R., H. Russchenberg, J. Erkelens, V. Venema, A. van Lammeren, A. Apituley, and S. Jongen : Ground-based remote sensing of stratocumulus properties during CLARA, 1996. *Journal of Applied Meteorology*, 39(2), 169–181, 2000.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Doschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, 571–657, <https://doi.org/10.1017/CBO9781107415324.016>, 2013.
- Brenguier, J. L.: Coincidence and Dead-Time Corrections for Particles Counters. Part II: High Concentration Measurements with an FSSP, *J. Atmos. Ocean. Tech.*, 6, 585–598, [https://doi.org/10.1175/1520-0426\(1989\)006<0585:CADTCF>2.0.CO;2](https://doi.org/10.1175/1520-0426(1989)006<0585:CADTCF>2.0.CO;2), 1989.
- Brenguier, J. L.: Parameterization of the condensation process - A theoretical approach. *Journal of the Atmospheric Sciences*, 48(2), 264–282, 1991.
- Brus, D., Neitola, K., Asmi, E., Aurela, M., Makkonen, U., Svensson, J., Hyvärinen, A.-P., Hirsikko, A., Hakola, H., Hillamo, R., and Lihavainen, H.: Pallas cloud experiment, PACE 2012, AIP Conf. Proc., 1527, 964, [doi:10.1063/1.4803433](https://doi.org/10.1063/1.4803433), 2013a.
- Brus, D., Asmi, E., Raatikainen, T., Neitola, K., Aurela, M., Makkonen, U., Svensson, J., Hyvärinen, A.-P., Hirsikko, A., Hakola, H., Hillamo, R., and Lihavainen, H.: Ground-based observations of aerosol

- and cloud properties at sub-arctic Pallas GAW-station, Pallas cloud experiment (PACE 2012) REPORT  
 595 SERIES IN AEROSOL SCIENCE N:o 142 (2013), Proceedings of FCoE in 'Physics, Chemistry,  
 Biology and Meteorology of Atmospheric Composition and Climate Change' Annual Meeting,  
 Proceedings\_FCoE\_2013. 187-191, <http://www.faar.fi/wp-content/uploads/2020/04/RS142pdf>, 2013b  
 (Last access ~~16.08.2022~~ [16.08.2023](#)).
- 600 Chang, Y., Guo, X., Tang, J., and Lu, G.: Aircraft measurement campaign on summer cloud  
 microphysical properties over the Tibetan Plateau, *Sci. Rep.*, 9, 4912, , 2019.
- Cho, Y.; Park, S.-J.; Kim, J.-H.; Yeo, H.; Nam, J.; Jun, S.-Y.; Kim, B.-M.; Kim, S.-W. Investigating  
 Wintertime Cloud Microphysical Properties and Their Relationship to Air Mass Advection at Ny-  
 605 Ålesund, Svalbard Using the Synergy of a Cloud Radar–Ceilometer–Microwave Radiometer. *Remote*  
*Sens.* 2021, 13, 2529. <https://doi.org/10.3390/rs13132529>, [2021](#).
- Devenish, B. J., Bartello, P., Brenguier, J.-L., Collins, L. R., Grabowski, W. W., Jzermans, R. H. A.,  
 610 Malinowski, S. P., Reeks, M. W., Vassilicos, J. C., Wang, L.-P., and Warhaft, Z.: Droplet growth in  
 warm turbulent clouds, *Q. J. Roy. Meteor. Soc.*, 138, 1401–1429, <https://doi.org/10.1002/qj.1897>, 2012.
- Dionne, J., von Salzen, K., Cole, J., Mahmood, R., Leaitch, W. R., Lesins, G., Folkins, I., and Chang,  
 R. Y.-W.: Modelling the relationship between liquid water content and cloud droplet number  
 615 concentration observed in low clouds in the summer Arctic and its radiative effects, *Atmos. Chem.*  
*Phys.*, 20, 29–43, <https://doi.org/10.5194/acp-20-29-2020>, 2020.
- Doulgeris, K. M., Brus, D., Raatikainen, T., and Kerminen V.-M.: A Finnish Meteorological Institute–  
 Aerosol Cloud Interaction Tube (FMI–ACIT): Experimental setup and tests of proper operation, *J.*  
 620 *Chem. Phys.*, 149, 124201, <https://doi.org/10.1063/1.5037298>, 2018.
- Doulgeris, K.-M., Komppula, M., Romakkaniemi, S., Hyvärinen, A.-P., Kerminen, V.-M., and Brus,  
 D.: In situ cloud ground-based measurements in the Finnish sub-Arctic: intercomparison of three cloud  
 spectrometer setups, *Atmos. Meas. Tech.*, 13, 5129–5147, <https://doi.org/10.5194/amt-13-5129-2020>,  
 2020.
- 625 Doulgeris, K.-M., Lihavainen, H., Hyvärinen, A.-P., Kerminen, V.-M., and Brus, D.: Data set for  
 Doulgeris et al. 2021: In-situ microphysical characterization of low-level clouds in the Finnish sub-  
 Arctic, extensive dataset, Finnish Meteorological Institute  
 [dataset], <https://doi.org/10.23728/FMIB2SHARE.988739D21B824C709084E88ED6C6D54B>, 2021.
- Doulgeris, K. M., Lihavainen, H., Hyvärinen, A.-P., Kerminen, V.-M., and Brus, D.: An extensive data  
 630 set for in situ microphysical characterization of low-level clouds in a Finnish sub-Arctic site, *Earth Syst.*  
*Sci. Data*, 14, 637–649, <https://doi.org/10.5194/essd-14-637-2022>, 2022.
- Donovan, D. P., Klein Baltink, H., Henzing, J. S., de Roode, S. R., and Siebesma, A. P.: A  
 depolarisation lidar-based method for the determination of liquid-cloud microphysical properties,  
 635 *Atmos. Meas. Tech.*, 8, 237–266, <https://doi.org/10.5194/amt-8-237-2015>, 2015.
- Droplet Measurement Technologies Manual: Particle Analysis and Display System (PADS) Image  
 Probe Data Reference Manual DOC-0201 Rev A-2 PADS 2.5.6, DMT, Boulder, Colorado, USA, 2009.
- 640 Emeis, S.; Münkler, C.; Vogt, S.; Müller, W.J.; ~~and~~ Schäfer, K.: Atmospheric boundary-layer structure  
 from simultaneous SODAR, RASS and ceilometer measurements. *Atmos. Environ.* ~~2004~~, 38, 273–286,  
[2004](#).

- 645 [Eneroth, K., Kjellström, E. and Holmén, K.: A trajectory climatology for Svalbard: investigating how atmospheric flow patterns influence observed tracer concentrations. Phys. Chem. Earth 28, 1191–1203, doi: DOI: 10.1016/j.pce.2003.08.051., 2003](#)
- 650 [Eneroth, K., Aalto, T., Hatakka, J., Holmen, K., Laurila, T. and Viisanen Y.: Atmospheric transport of carbon dioxide to a baseline monitoring station in northern Finland. Tellus B, 57: 366– 374. <https://doi.org/10.1111/j.1600-0889.2005.00160.x>, 2005.](#)
- Filioglou, M., Mielonen, T., Balis, D., Giannakaki, E., Arola, A., Kokkola, H., Komppula, M., and Romakkaniemi, S.: Aerosol Effect on the 360 Cloud Phase of Low-Level Clouds Over the Arctic, J. Geophys. Res. Atmos., 124, 7886–7899, <https://doi.org/10.1029/2018JD030088>, 2019.
- 655 Freud, E., Krejci, R., Tunved, P., Leaitch, R., Nguyen, Q. T., Massling, A., Skov, H., and Barrie, L.: Pan-Arctic aerosol number size distributions: seasonality and transport patterns, Atmos. Chem. Phys., 17, 8101–8128, <https://doi.org/10.5194/acp-17-8101-2017>, 2017.
- 660 Fuchs, J., Cermak, J., Andersen, H., Hollmann, R., & Schwarz, K.: ~~(2017)~~. On the influence of air mass origin on low-cloud properties in the Southeast Atlantic. Journal of Geophysical Research: Atmospheres, 122, 11,076–11,091. <https://doi.org/10.1002/2017JD027184>, [2017](#).
- Gérard, V., Nozière, B., Fine, L., Ferronato, C., Singh, D., K., Frossard, A., Cohen, R. C. Asmi, E., Lihavainen, H., Kivekäs, N., Aurela, M., Brus, D., Frka, S., and Cvitešić Kušan, A.: Concentrations and Adsorption Isotherms for Amphiphilic Surfactants in PM1 Aerosols from Different Regions of Europe, Environ. Sci. Technol., 53, 12379–12388, <https://doi.org/10.1021/acs.est.9b03386>, 2019.
- 665 Girdwood, J., Smith, H., Stanley, W., Ulanowski, Z., Stopford, C., Chemel, C., Doulgieris, K.-M., Brus, D., Campbell, D., and Mackenzie, R.: Design and field campaign validation of a multi-rotor unmanned aerial vehicle and optical particle counter, Atmos. Meas. Tech., 13, 6613–6630, <https://doi.org/10.5194/amt-13-6613-2020>, 2020.
- Girdwood, J., Stanley, W., Stopford, C., and Brus, D.: Simulation and field campaign evaluation of an optical particle counter on a fixed-wing UAV, Atmos. Meas. Tech., 15, 2061–2076, <https://doi.org/10.5194/amt-15-2061-2022>, 2022.
- 675 Goren T., D. Rosenfeld :Decomposing aerosol cloud radiative effects into cloud cover, liquid water path and Twomey components in marine stratocumulus, Atmospheric Research, 138 ~~(2014)~~, pp. 378–393, 10.1016/j.atmosres.2013.12.008, [2014](#).
- 680 Gultepe, I., G. Isaac, D. Hudak, R. Nissen, and W. Strapp, Dynamical and microphysical characteristics of Arctic clouds during BASE, J. Clim., 13, 1225 – 1254, 2000.
- Gultepe, I. and Isaac, G. A.: Effects of air mass origin on Arctic cloud microphysical parameters for April 1998 during FIRE.ACE, J. Geophys. Res., 107, D21, <https://doi.org/10.1029/2000JC000440>, 2002.
- 685 Guyot, G., Goubeyre, C., Febvre, G., Shcherbakov, V., Burnet, F., Dupont, J.-C., Sellegri, K., and Jourdan, O.: Quantitative evaluation of seven optical sensors for cloud microphysical measurements at the Puy-de-Dôme Observatory, France, Atmos. Meas. Tech., 8, 4347–4367, <https://doi.org/10.5194/amt-8-4347-2015>, 2015.
- 690 Hatakka, J., Aalto, T., Aaltonen, V., Aurela, M., Hakola, H., Komppula, M., Laurila, T., Lihavainen, H., Paatero, J., Salminen, K., and Viisanen Y.: Overview of the atmospheric research activities and results at Pallas GAW station, Boreal Environ. Res., 8, 365–384, 2003.

Formatted: Finnish

Formatted: English (United States)

695 Hobs and Rango, ~~1998~~: Reply to “Comments by Alan M. Blyth and John Latham on ‘Cumulus  
glaciation papers by P. V. Hobbs and A. L. Rangno’” Quart. J. Roy. Meteor. Soc., 124, 1009–1011,  
[1998](#).

700 Hyvärinen, A.-P., Kolmonen, P., Kerminen, V.-M., Virkkula, A., Leskinen, A., Komppula, M.,  
Hatakka, J., Burkhardt, J., Stohl, A., Aalto, P., Kulmala, M., Lehtinen, K., Viisanen, Y., and Lihavainen,  
H.: Aerosol black carbon at five background measurement sites over Finland, a gateway to the Arctic,  
Atmos. Environ., 45, 4042–4050, 2011.

Igel, A. L., & van den Heever, S. C. (~~2017~~): The Importance of the Shape of Cloud Droplet Size  
705 Distributions in Shallow Cumulus Clouds. Part I: Bin Microphysics Simulations, Journal of the  
Atmospheric Sciences, 74(1), 249–258. Retrieved [January](#) ~~May~~ 27, 2023~~2~~,  
from <https://journals.ametsoc.org/view/journals/atsc/74/1/jas-d-15-0382.1.xml>, [2017](#).

Igel, A. L., & van den Heever, S. C. (2017). The Importance of the Shape of Cloud Droplet Size  
710 Distributions in Shallow Cumulus Clouds. Part II: Bulk Microphysics Simulations, Journal of the  
Atmospheric Sciences, 74(1), 259–273. Retrieved [January](#) 27, 2023~~May~~ ~~27~~, ~~2022~~,  
from <https://journals.ametsoc.org/view/journals/atsc/74/1/jas-d-15-0383.1.xml>, [2017](#).

Iwamoto, Y.; Watanabe, A.; Kataoka, R.; Uematsu, M. ~~and~~ Miura, K.: Aerosol–Cloud Interaction at  
the Summit of Mt. Fuji, Japan: Factors Influencing Cloud Droplet Number Concentrations. Appl. Sci.  
715 ~~2021~~, 11, 8439. <https://doi.org/10.3390/app11188439>, [2021](#).

Korolev, A., McFarquhar, G., Field, P. R., Franklin, C., Lawson, P., Wang, Z., Williams, E., Abel, S.  
J., Axisa, D., Borrmann, S., Crosier, J., Fugal, J., Krämer, M., Lohmann, U., Schlenczek, O., Schnaiter,  
M., ~~& and~~ Wendisch, M.: (~~2017~~) Mixed-Phase Clouds: Progress and Challenges, Meteorological  
720 Monographs, 58, 5.1–5.50, [2017](#).

Kreidenweis, S. M., Petters, M. D., and De Mott, P. J.: Single-parameter estimates of aerosol 930 water  
content, <https://doi.org/10.1088/1748-9326/3/3/035002>, 2008.

725 Kyrö, E.-M., Väänänen, R., Kerminen, V.-M., Virkkula, A., Petäjä, T., Asmi, A., Dal Maso, M.,  
Nieminen, T., Juhola, S., Shcherbinin, A., Riipinen, I., Lehtipalo, K., Keronen, P., Aalto, P. P., Hari,  
P., and Kulmala, M.: Trends in new particle formation in eastern Lapland, Finland: effect of decreasing  
sulfur emissions from Kola Peninsula, Atmos. Chem. Phys., 14, 4383–4396,  
<https://doi.org/10.5194/acp-14-4383-2014>, 2014.

730 Lance, S.: Coincidence Errors in a Cloud Droplet Probe (CDP) and a Cloud and Aerosol Spectrometer  
(CAS), and the Improved Performance of a Modified CDP, <https://doi.org/10.1175/JTECH-D-11-00208.1>, 2012.

735 Lee, H.J., Jo, Y.J., Kim, S., [Kim, D., Kim J., Choi, D., Jo, H., Bak, J., Park, S., Jeon, W. and Kim, C.-H. et al.](#)  
Transboundary aerosol transport process and its impact on aerosol-radiation-cloud feedbacks in  
springtime over Northeast Asia. Sci Rep 12, 4870. (~~2022~~) [https://doi.org/10.1038/s41598-022-08854-](https://doi.org/10.1038/s41598-022-08854-1)  
[1](#), 2022.

740 Lihavainen, H., Kerminen, V.-M., Komppula, M., Hyvärinen, A.-P., Laakia, J., Saarikoski, S.,  
Makkonen, U., Kivekäs, N., Hillamo, R., Kulmala, M., and Viisanen, Y.: Measurements of the relation  
between aerosol properties and microphysics and chemistry of low level liquid water clouds in Northern  
Finland, Atmos. Chem. Phys., 8, 6925–6938, <https://doi.org/10.5194/acp-8-6925-2008>, 2008.

745 Lohila, A., Penttilä, T., Jortikka, S., Aalto, T., Anttila, P., Asmi, E., Aurela, M., Hatakka, J., Hellén, H.,  
Henttonen, H., Hänninen, P., Kilkki, J., Kyllönen, K., Laurila, T., Lepistö, A., Lihavainen, H.,  
Makkonen, U., Paatero, J., Rask, M., Sutinen, R., Tuovinen, J.-P., Vuorenmaa, J., and Viisanen, Y.:

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Preface to the special issue on integrated research of atmosphere, ecosystems and environment at Pallas, Boreal Env. Res., 20, 431–454, 2015.

750

Lohmann, U., Lüönd, F., and Mahrt, F.: Microphysical processes in warm clouds. In *An Introduction to Clouds: From the Microscale to Climate* (pp. 186–217). Cambridge: Cambridge University Press. doi:10.1017/CBO9781139087513.008, 2016.

755 Lu, C., Liu, Y., Niu, S., and Endo, S.: Scale dependence of entrainment-mixing mechanisms in cumulus clouds, *J. Geophys. Res.-Atmos.*, 119, 13877–13890, 2014.

MacDonald, A.B., Dadashazar, H., Chuang, P.Y., Crosbie, E., Wang, H., Wang, Z., Jonsson, H.H., Flagan, R.C., Seinfeld, J.H., and Sorooshian, A., Characteristic Vertical Profiles of Cloud Water Composition in Marine Stratocumulus Clouds and Relationships With Precipitation. *J Geophys Res Atmos.* ~~2018~~<sup>Apr</sup>:123(7) 3704–3723. doi:10.1002/2017jd027900. PMID: 32025449; PMCID: PMC7002026, ~~2018~~.

760

Martins, J. V., Marshak, A., Remer, L. A., Rosenfeld, D., Kaufman, Y. J., Fernandez-Borda, R., Koren, I., Correia, A. L., Zubko, V., and Artaxo, P.: Remote sensing the vertical profile of cloud droplet effective radius, thermodynamic phase, and temperature, *Atmos. Chem. Phys.*, 11, 9485–9501, <https://doi.org/10.5194/acp-11-9485-2011>, 2011.

765

McFarquhar, G. M., Bretherton, C., Marchand, R., Protat, A., DeMott, P. J., Alexander, S. P., Roberts, G. C., Twohy, C. H., Toohey, D., Siems, S., Huang, Y., Wood, R., Rauber, R. M., Lasher-Trapp, S., Jensen, J., Stith, J., Mace, J., Um, J., Järvinen, E., Schnaiter, M., Gettelman, A., Sanchez, K. J., McCluskey, C. S., Russell, L. M., McCoy, I. L., Atlas, R., Bardeen, C. G., Moore, K. A., Hill, T. C. J., Humphries, R. S., Keywood, M. D., Ristovski, Z., Cravigan, L., Schofield, R., Fairall, C., Mallet, M. D., Kreidenweis, S. M., Rainwater, B., D'Alessandro, J., Wang, Y., Wu, W., Saliba, G., Levin, E. J. T., Ding, S., Lang, F., Truong, S. C., Wolff, C., Haggerty, J., Harvey, M. J., Klekociuk, A., and McDonald, A.: Observations of clouds, aerosols, precipitation, and surface radiation over the Southern Ocean: An overview of CAPRICORN, MARCUS, MICRE and SOCRATES, *B. Am. Meteorol. Soc.*, 102, E894–E928, , 2020.

775

780

Mie, G.: Beiträge zur Optik trüber Medien, speziell kolloidaler Metallösungen, *Ann. Phys.-Berlin*, 330, 377–445, 1908.

Morrison, H., van Lier-Walqui, M., Fridlind, A. M., Grabowski, W. W., Harrington, J. Y., Hoose, C., Korolov, A., Kumjian, M. R., Milbrandt, J. A., Pawlowska, H., Posselt, D. J., Prat, O. P., Reimel, K. J., Shima, S.-I., Van Diedenhoven, B., and Xue, L.: Confronting the challenge of modeling cloud and precipitation microphysics, *J. Adv. Model. Earth Sy.*, 12, e2019MS001689, <https://doi.org/10.1029/2019MS001689>, 2020.

785

Orbe, C., Newman, P. A., Waugh, D. W., Holzer, M., Oman, L. D., Li, F., & Polvani, L. M.: ~~(2015)~~. Air-mass Origin in the Arctic. Part I: Seasonality, *Journal of Climate*, 28(12), 4997–5014. Retrieved ~~May~~<sup>January</sup> ~~2023~~<sup>23</sup>, from <https://journals.ametsoc.org/view/journals/clim/28/12/jcli-d-14-00720.1.xml>, ~~2015~~.

790

Orbe, C., Newman, P. A., Waugh, D. W., Holzer, M., Oman, L. D., Li, F., & Polvani, L. M. ~~(2015)~~. Air-mass Origin in the Arctic. Part II: Response to Increases in Greenhouse Gases, *Journal of Climate*, 28(23), 9105–9120. Retrieved ~~May~~<sup>January</sup> ~~2023~~<sup>23</sup>, from <https://journals.ametsoc.org/view/journals/clim/28/23/jcli-d-15-0296.1.xml>, ~~2015~~.

795

Paramonov, M., Kerminen, V.-M., Gysel, M., Aalto, P. P., Andreae, M. O., Asmi, E., Baltensperger, U., Bougiatioti, A., Brus, D., Frank, G. P., Good, N., Gunthe, S. S., Hao, L., Irwin, M., Jaatinen, A.,



- 800 Jurányi, Z., King, S. M., Kortelainen, A., Kristensson, A., Lihavainen, H., Kulmala, M., Lohmann, U., Martin, S. T., McFiggans, G., Mihalopoulos, N., Nenes, A., O'Dowd, C. D., Ovadnevaite, J., Petäjä, T., Pöschl, U., Roberts, G. C., Rose, D., Svenningsson, B., Swietlicki, E., Weingartner, E., Whitehead, J., Wiedensohler, A., Wittbom, C., and Sierau, B.: A synthesis of cloud condensation nuclei counter (CCNC) measurements within the EUCAARI network, *Atmos. Chem. Phys.*, 15, 12211–12229, <https://doi.org/10.5194/acp-15-12211-2015>, 2015.
- 805 Painemal, D., Kato, S., and Minnis, P.: Boundary layer regulation in the southeast Atlantic cloud microphysics during the biomass 25 burning season as seen by the A-train satellite constellation, *Journal of Geophysical Research: Atmospheres*, 119, 11 288–11 302, doi:10.1002/2014JD022182, <http://doi.wiley.com/10.1002/2014JD022182>, 2014.
- 810 [Patel, P. and Hiang, J.: Cloud condensation nuclei characteristics at the Southern Great Plains site: role of particles size distribution and aerosol hygroscopicity: \*Environ. Res. Commun.\* 3 075002, 2021.](#)
- Pawlowska, H., W. W. Grabowski, and J. . L. Brenguier: Observations of the width of cloud droplet spectra in stratocumulus. *Geophysical Research Letters*, 33(19), L19810, doi: 10.1029/2006GL026841, 2006.
- 815 [Pernov, J. B., Beddows, D., Thomas, D. C., Dall'Osto, M., Harrison, R. M., Schmale, J., Skov, H., and Massling, A.: Increased aerosol concentrations in the High Arctic attributable to changing atmospheric transport patterns, \*npj Climate and Atmospheric Science\*, 5, 62, <https://doi.org/10.1038/s41612-022-00286-y>, 2022.](#)
- ~~Piyushkumar N Patel and Jonathan H Jiang 2021 *Environ. Res. Commun.* 3 075002~~
- 820 Pissu, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, *Geosci. Model Dev.*, 12, 4955–4997, <https://doi.org/10.5194/gmd-12-4955-2019>, 2019.
- 825 Pruppacher, H. R. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, Kluwer Academic Publishers, Dordrecht, 1997.
- Pruppacher, H. R. and Klett, J. D.: Growth of Cloud Drops by Collision, Coalescence and Breakup. In: *Microphysics of Clouds and Precipitation. Atmospheric and Oceanographic Sciences Library*, vol 18. Springer, Dordrecht. [https://doi.org/10.1007/978-0-306-48100-0\\_15](https://doi.org/10.1007/978-0-306-48100-0_15). 2010.
- 830 Raatikainen, T., Brus, D., Hyvärinen, A.-P., Svensson, J., Asmi, E., and Lihavainen, H.: Black carbon concentrations and mixing state in the Finnish Arctic, *Atmos. Chem. Phys.*, 15, 10057–10070, <https://doi.org/10.5194/acp-15-10057-2015>, 2015.
- ~~Raatz W.E. and Shaw G.E. -1984- -Long-range tropospheric transport of pollution aerosols into the Alaskan Arctic. *J. Climate App. Met.* 23: 1052–1064. 1984.~~
- 835 Rosenfeld, D. and Ulbrich, C. W.: Cloud Microphysical Properties, Processes, and Rainfall Estimation Opportunities, *Meteor. Mon.*, 30, 237–237, [https://doi.org/10.1175/0065-9401\(2003\)0302.0.CO;2](https://doi.org/10.1175/0065-9401(2003)0302.0.CO;2), 2003.
- 840 Rosenfeld, D., Sherwood, S., Wood R. and Donner, L.: ~~Atmospheric science-~~Climate effects of aerosol-cloud interactions. *Science*. 2014 Jan 24;343(6169):379-80. doi: 10.1126/science.1247490. PMID: 24458631. [2014.](#)

- Seibert, P. and Frank, A.: Source-receptor matrix calculation with a Lagrangian particle dispersion model in backward mode, *Atmos. Chem. Phys.*, 4, 51–63, <https://doi.org/10.5194/acp-4-51-2004>, 2004.
- ~~Seibert, P. and Frank, A.: Source-receptor matrix calculation with a Lagrangian particle dispersion model in backward mode, *Atmos. Chem. Phys.*, 4, 51–63, 2004, SRef ID: 1680-7324/acp/2004-4-51~~
- Sherwood, S., Webb, M.J., Annan, J.D., Armour, K.C., Forster, P.M., Hargreaves, J.C., Hegerl, G., Klein, S.A., Marvel, K.D., 605 Rohling, E.J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C.S., Foster, G.L., Hausfather, Z., von der Heydt, A.S., Knutti, R., Mauritsen, T., Norris, J.R., Proistosescu, C., Rugenstein, M., Schmidt, G.A., Tokarska, K.B., Zelinka, M.D.: An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics*, 58, <https://doi.org/10.1029/2019RG000678>, 2020.
- ~~Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., et al. (2020). An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics*, 58, e2019RG000678. <https://doi.org/10.1029/2019RG000678>.~~
- Shupe, MD, Rex, M, Blomquist, B, Persson, POG, Schmale, J, Uttal, T, Althausen, D, Angot, H, Archer, S, Bariteau, L, Beck, I, Bilberry, J, Bucci, S, Buck, C, Boyer, M, Brasseur, Z, Brooks, IM, Calmer, R, Cassano, J, Castro, V, Chu, D, Costa, D, Cox, CJ, Creamean, J, Crewell, S, Dahlke, S, Damm, E, de Boer, G, Deckelmann, H, Dethloff, K, Du`tsch, M, Ebell, K, Ehrlich, A, Ellis, J, Engelmann, R, Fong, AA, Frey, MM, Gallagher, MR, Ganzeveld, L, Gradinger, R, Graeser, J, Greenamyer, V, Griesche, H, Griffiths, S, Hamilton, J, Heinemann, G, Helmig, D, Herber, A, Heuze´, C, Hofer, J, Houchens, T, Howard, D, Inoue, J, Jacobi, H-W, Jaiser, R, Jokinen, T, Jourdan, O, Jozef, G, King, W, Kirchgaessner, A, Klingebiel, M, Krassovski, M, Krumpfen, T, Lampert, A, Landing, W, Laurila, T, Lawrence, D, Lonardi, M, Loose, B, Lu`pkes, C, Maahn, M, Macke, A, Maslowski, W, Marsay, C, Maturilli, M, Mech, M, Morris, S, Moser, M, Nicolaus, M, Ortega, P, Osborn, J, Pa`tzold, F, Perovich, DK, Peta`ja`, T, Pilz, C, Pirazzini, R, Posman, K, Powers, H, Pratt, KA, Preußer, A, Que`le´ ver, L, Radenz, M, Rabe, B, Rinke, A, Sachs, T, Schulz, A, Siebert, H, Silva, T, Solomon, A, Sommerfeld, A, Spreen, G, Stephens, M, Stohl, A, Svensson, G, Uin, J, Viegas, J, Voigt, C, von der Gathen, P, Wehner, B, Welker, JM, Wendisch, M, Werner, M, Xie, ZQ, Yue, F. 2022. Overview of the MOSAiC expedition: Atmosphere. *Elementa: Science of the Anthropocene* 10(1). DOI: <https://doi.org/10.1525/elementa.2021.00060>
- Sipilä, M., Sarnela, N., Neitola, K., Laitinen, T., Kempainen, D., Beck, L., Duplissy, E.-M., Kuittinen, S., Lehmuşjärvi, T., Lampilahti, J., Kerminen, V.-M., Lehtipalo, K., Aalto, P. P., Keronen, P., Siivola, E., Rantala, P. A., Worsnop, D. R., Kulmala, M., Jokinen, T., and Petäjä, T.: Wintertime subarctic new particle formation from Kola Peninsula sulfur emissions, *Atmos. Chem. Phys.*, 21, 17559–17576, <https://doi.org/10.5194/acp-21-17559-2021>, 2021.
- Small, J. D., Chuang, P. Y., Feingold, G., and Jiang, H.: Can aerosol decrease cloud lifetime?, *Geophys. Res. Lett.*, 36, L16806, <https://doi.org/10.1029/2009GL038888>, 2009.
- Solomon, A., & Shupe, M. D. (2019). A Case Study of Air mass Transformation and Cloud Formation at Summit, Greenland, *Journal of the Atmospheric Sciences*, 76(10), 3095–3113. Retrieved May 9, 2022, from <https://journals.ametsoc.org/view/journals/atms/76/10/jas-d-19-00561.xml>
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, <https://doi.org/10.5194/acp-5-2461-2005>, 2005.
- Torres-Delgado, E., Baumgardner, D., and Mayol-Bracero, O. L.: Measurement report: Impact of African aerosol particles on cloud evolution in a tropical montane cloud forest in the Caribbean, *Atmos. Chem. Phys.*, 21, 18011–18027, <https://doi.org/10.5194/acp-21-18011-2021>, 2021.

Formatted: Finnish

[Tunved, P., Hansson, H. C., Kerminen, V. M., Strom, J., Dal Maso, M., Lihavainen, H., Viisanen, Y., Aalto, P. P., Komppula, M., and Kulmala, M.: High natural aerosol loading over boreal forests, \*Science\*, 312, 261–263, 15.10.1126/science.1123052, 2006a.](#)

Formatted: Font: 11 pt

Twohy, C. H., M. D. Petters, J. R. Snider, B. Stevens, W. Tahnk, M. Wetzel, L. Russell, and F. Burnet (2005). Evaluation of the aerosol indirect effect in marine stratocumulus clouds: Droplet number, size, liquid water path, and radiative impact, *J. Geophys. Res.*, 110, D08203, doi:10.1029/2004JD005116, 2005.

Vaisala Oyj, Ceilometer CT25K: User's Guide; Vaisala Oyj: Vantaa, Finland, 2002.

Virkkula, A., Hillamo, R. E., Kerminen, V.-M. & and Stohl, A. :1997-. The influence of Kola Peninsula, continental European and marine sources on the number concentrations and scattering coefficients of the atmospheric aerosol in Finnish Lapland. *Boreal Env. Res.* 2: 317–336. ISSN 1239-6095, 1997.

Formatted: English (United States)

Wandinger, U., Apituley, A., Blumenstock, T., Bukowiecki, N., Cammas, J.-P., Connolly, P., De Mazière, M., Dils, B., Fiebig, M., Freney, E., Gallagher, M., Godin-Beekmann, S., Goloub, P., Gysel, M., Haeffelin, M., Hase, F., Hermann, M., Herrmann, H., Jokinen, T., Komppula, M., Kubistin, D., Langerock, B., Lihavainen, H., Mihalopoulos, N., Laj, P., Lund Myhre, C., Mahieu, E., Mertes, S., Möhler, O., Mona, L., Nicolae, D., O'Connor, E., Palm, M., Pappalardo, G., Pazmino, A., Petäjä, T., Philippin, S., Plass-Duelmer, C., Pospichal, B., Putaud, J.-P., Reimann, S., Rohrer, F., Russchenberg, H., Sauvage, S., Sellegri, K., Steinbrecher, R., Stratmann, F., Sussmann, R., Van Pinxteren, D., Van Roozendael M., Vigouroux C., Walden C., Wegene R., and Wiedensohler, A.: ACTRIS-PPP Deliverable D5.1: Documentation on technical concepts and requirements for ACTRIS Observational Platforms, available at: [https://www.actris.eu/Portals/46/Documentation/ACTRIS PPP/Deliverables/Public/WP5\\_D5.1\\_M18.pdf?ver=2018-06-28-125343-273](https://www.actris.eu/Portals/46/Documentation/ACTRIS_PPP/Deliverables/Public/WP5_D5.1_M18.pdf?ver=2018-06-28-125343-273) (last access: 21 April January 2023), 2018.

Wang, Y., Zheng, X., Dong, X., Xi, B., Wu, P., Logan, T., and Yung, Y. L.: Impacts of long-range transport of aerosols on marine-boundary-layer clouds in the eastern North Atlantic, *Atmos. Chem. Phys.*, 20, 14741–14755, <https://doi.org/10.5194/acp-20-14741-2020>, 2020.

Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B., Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrmann, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and Zeppenfeld, S.: The Arctic Cloud Puzzle: Using ACLOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *B. Am. Meteorol. Soc.*, 100, 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.

Jaatinen, A., Romakkaniemi, S., Anttila, T., Hyvärinen, A.-P., Hao, L. Q., Kortelainen, A., Miettinen, P., Mikkonen, S., Smith, J. N., Virtanen, A. & and Laaksonen, A. 2014: The third Pallas Cloud Experiment: Consistency between the aerosol hygroscopic growth and CCN activity. *Boreal Env. Res.* 19 (suppl. B): 368–382., 2014.

Xue, Y., Wang, L., and Grabowski, W. W. : Growth of Cloud Droplets by Turbulent Collision–Coalescence, *Journal of the Atmospheric Sciences*, 65(2), 331–356. <https://doi.org/10.1175/2007JAS2406.1>, 2008.

950 Zhao, C.F., Xie, S. C., Klein, S.A., Protat, A., Shupe, M.D., McFarlane, S.A., Comstock, J.M., Delanoe,  
J., Deng, M., Dunn, M., Hogan, R.J., Huang, D., Jensen, M.P., Mace, G.G., McCoy, R., O'Connor,  
E.J., Turner, D.D. and Wang, Z.E.: Toward understanding of differences in current cloud retrievals of  
ARM ground-based measurements. *J. Geophys. Res.-Atmos.*, 117, p. D10206, 10.1029/2011jd016792.  
2012.

955

960

965