

Responses to Referee #1:

This paper reports on measurements of spectrally resolved particle light absorption combined with aerosol chemical composition measured at the southeastern edge of the Tibetan Plateau. The particle light absorption data are used to infer BrC and BC levels. Based on the chemical signatures the aerosol was divided into two groups and the BrC optical properties were determined for each group. Overall radiative effects of the BrC relative to BC on a per particle mass basis showed that BrC has an important role. Back trajectory analysis was used to identify source regions of the BrC. The paper is well suited for publication in this journal. The authors consider uncertainty in their approach to determine BrC, which is a nice feature of this paper. A number of suggestions for further analysis and clarifications are noted below.

Response: We thank the reviewer for the helpful comments and providing us the opportunity to strengthen our research. We have carefully addressed the comments in point-by-point form as shown below. Detailed responses to each of the reviewer's comments are provided in blue, and the revised text is underlined. Attached please also find the marked-up manuscript to track the changes in the revised manuscript.

General Comments:

Comment (1): Some of the methods description may be too detailed, such as the HERM algorithm description. A plainer language description would be better. Eg, line 128 and on describing Eq (3). First, this is a classic inversion problem encountered for many instruments in aerosol science (and many other fields). My understanding is that X is the measured ACSM mass spectra of the organic species, (why call it the receptor site?), G is the source contributions, which is what is you are trying to determine, and F is the mass spectra of the specific sources. Normally one knows F and performs the inversion to solve for G. In this case F is unknown, so a modified approach is used. This can then be described.

Response: (a) Yes, as the reviewer points out, X is the measured ACSM mass spectra of the organic species. In the original manuscript, the receptor site refers to the sampling site. To make HERM algorithm more concise and clear, we have revised the relevant description of HERM following the reviewer's suggestion:

“HERM is an effective receptor model, which was performed to retrieve potential sources of OA in this study. The HERM algorithm groups the matrix X (measured mass spectra of organic fragments) into two nonnegative constant matrices G (source contribution) and F (mass spectra of specific sources), and the model residual matrix E, defined as:

$$X = G \times F + E \quad (3)$$

The model does not require prior mass spectra of sources, and the values of G and F can be obtained using an iterative conjugate gradient algorithm. The principle of HERM has been described in detail elsewhere (Chen and Cao, 2018).” (Page 5 Line 128–137)

(b) We moved the Section 2.3.4 of statistical metrics to the supplement as Text S1, and also simplified the description of calculation of optical parameters (Section 2.3.3) as follows:

“The MAC of OA component in this study was resolved by the multiple linear regression (MLR) model combined with $b_{\text{abs-BrC}}$ and OA source apportionment results, which is defined as follows:

$$b_{\text{abs-BrC}}(\lambda) = a_1(\lambda) \times [\text{BBOA}] + a_2(\lambda) \times [\text{po-OOA}] \quad (4)$$

Here, a_1 and a_2 denote the MAC of BBOA (MAC_{BBOA}) and po-OOA ($\text{MAC}_{\text{po-OOA}}$), respectively, in square meters per gram ($\text{m}^2 \text{g}^{-1}$); [BBOA] and [po-OOA] are the mass concentration of BBOA and po-OOA, respectively, in micrograms per cubic meter ($\mu\text{g m}^{-3}$). Tolerance (0.2) and variance inflation factor (4.7) for the ordinary least square fitting results indicated that there was no serious multicollinearity between two independent variables, however, heteroscedasticity existed according to “White Test” ($p < 0.05$). Thus, the weighted least squares method was used for parameter estimation in MLR model. The MAC of BC (MAC_{BC}) was directly calculated with $b_{\text{abs-BC}}$ divided by the mass concentration of BC, which was obtained by dividing b_{abs} (880 nm) by the default MAC (880 nm) used in the Model AE33 (Drinovec et al., 2015).

AAE describes the spectral dependence of light absorption by aerosols, and it can be calculated using a power law function with b_{abs} and MAC, respectively:

$$b_{\text{abs}}(\lambda) = k_1 \times \lambda^{-\text{AAE}} \quad (5)$$

$$\text{MAC}(\lambda) = k_2 \times \lambda^{-\text{AAE}} \quad (6)$$

Here, k_1 and k_2 are constants independent of wavelength.” (Page 6 Line 157–Page 7 Line 178)

Comment (2): A major possible issue is the characterization of the two sources identified by the source apportionment discussed above. Despite the complicated inversion, it seems the separation of source comes down to two things, the po-OOA source characterized by m/z 44 (which seems similar to MO-OOA in other studies using this instrument, see discussion below), and m/z 60 the fragments of levoglucosan and other carbohydrates known to be emitted primarily from biomass combustion (and cooking, but which is not discussed here). The issue is one way to interpret these source apportionment results is that both are from biomass burning, BBOA is the possibly fresher or less photochemically processed BBOA and po-OOA are more aged and chemically processed BBOA. To me, this clarifies the data interpretation and is supported by the idea that both have the same source region and that they are correlated (line 262, $r = 0.63$). However, arguing against this is that the MACs are higher for po-

OOA which one would not be expect if this was more aged and possibly more photochemically bleached relative to the fresher or less processed (less bleached) BBOA.

Response: In our study, po-OOA characterized by higher m/z 44 was more oxygenated than BBOA, which meant stronger photobleaching effect. We agree with the reviewer that more photobleaching would result in the lower MAC of OA (Lee et al., 2014). According to another reviewer’s suggestion, we recheck the accuracy of parameter estimation of the multiple linear regression (MLR) model. We find that the heteroscedasticity in MLR model existed when we use the ordinary least square fitting previously, and this may cause the overestimation of MAC_{po-OOA} .


To address this concern, we use the weighted least squares method to estimate parameter of MLR model. Figure 5 shows the latest results of MAC_{BBOA} and MAC_{po-OOA} . That is, the MAC_{BBOA} is higher than the MAC_{po-OOA} at wavelengths from 370 to 660 nm. In the revised manuscript, we first add the testing results and fitting method of MLR to the Sect. 2.3.3:

“Tolerance (0.2) and variance inflation factor (4.7) for the ordinary least square fitting results indicated that there was no serious multicollinearity between two independent variables, however, heteroscedasticity existed according to “White Test” ($p < 0.05$). Thus, the weighted least squares method was used for parameter estimation in MLR model.” (Page 6 Line 163–166)

We also update the MAC results and give the explanation that po-OOA has the lower MAC compared to BBOA in the Sect. 3.3:

“Compared with BBOA, more oxygenated po-OOA was possibly more photochemically bleached, which resulted in the lower MAC (Lee et al., 2014).” (Page 11 Line 296–297)

In the revised manuscript, Figure 5 shows:



“Figure 5. (a) The mass absorption cross section of BBOA and po-OOA (MAC_{BBOA} and MAC_{po-OOA} , respectively) at five wavelengths

($\lambda = 370, 470, 520, 590, \text{ and } 660 \text{ nm}$). The circle and shaded area represent the mean MAC values and the standard deviations, respectively. The dashed line is power-law fit. (b) Light absorption coefficient of BrC ($b_{\text{abs-BrC}}$) from BBOA and po-OOA and its fraction in the total reconstructed BrC absorption at different wavelengths.”

In the reference list, we add the new reference:

“Lee, H. J., Aiona, P. K., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Effect of solar radiation on the optical properties and molecular composition of laboratory proxies of atmospheric brown carbon, *Environ. Sci. Technol.*, 48(17), 10217–10226, <https://doi.org/10.1021/es502515r>, 2014.”

Specific Comments:

Comment (3): When discussing Eq (1), might want to say something about how well a power law fits data.

Response: AAE can be calculated using a pair of light absorption coefficients (b_{abs}) at two different wavelength:

$$\text{AAE}(\lambda_1 \lambda_2) = \frac{\ln\left(\frac{b_{\text{abs}}(\lambda_1)}{b_{\text{abs}}(\lambda_2)}\right)}{\ln\left(\frac{\lambda_1}{\lambda_2}\right)}$$

Eq (1) is derived from this method when b_{abs} at 880 nm and AAE of BC are known. Therefore, b_{abs} of BC obtained from Eq (1) always satisfies the power law relationship with wavelength.

Comment (4): Where does the term for one of the identified sources (po-OOA) come from. Has it been used before or is it being introduced here? Essentially, it seems to be driven by mass spectral peak at m/z 44, which is carboxylic acid fragment and which in past studies of ACSM/AMS is largely indicative of more aged oxygenated organic aerosol from various sources. Why not discuss this (ie, how that peak is determined in other studies and why a different name is used here).

Response: As noted by the reviewer, the fraction of m/z 44 in OA mass spectrum (f_{44}) is a surrogate of oxidation degree (Aiken et al., 2008). The photochemical-oxidation processed oxygenated OA (po-OOA) resolved in this study was characterized by the highest peak at m/z 44 (f_{44} , 27.8 %), which was quite similar to those of more-oxidized oxygenated OA (MO-OOA) ($f_{44} > 20\%$) identified frequently in previous ACSM and AMS studies (Tobler et al., 2021; Xu et al., 2018; Zhang et al., 2019). Here, since high O_3 was the driving factor of po-OOA formation, the term of po-OOA was introduced in this manuscript to stress the importance of photochemical processing in the Tibetan Plateau.

In the revised manuscript, to make it clear, we add:

“Another OA source was featured by the strong correlation with m/z 44 ($r = 0.97, p < 0.01$), which is a surrogate of oxidation degree (Aiken et al., 2008). The most abundant peak in mass spectrum of po-OOA was at m/z 44 ($f_{44}, 27.8\%$), similar to those in mass spectra of more-oxidized oxygenated OA (MO-OOA) ($f_{44} > 20\%$) identified frequently in previous studies (Tobler et al., 2021; Xu et al., 2018; Zhang et al., 2019). It indicated that this OA source was likely related to extensive secondary processes occurring during transport (Wang et al., 2017; Xu et al., 2017). Figure 4d shows that both po-OOA mass concentration and its fraction in OA increased with increasing O_3 ($R^2 = 0.79-0.87$), however, neither of them correlated with RH (Fig. S3). These results supported that high O_3 was the possible driving factor of po-OOA formation, thus the term of po-OOA was introduced in this study to stress the importance of photochemical-oxidation process in the TP.” (Page 10 Line 270–279)

In the reference list, we add the new reference:

“Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y. L., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prévôt, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with High-Resolution Time-of-Flight Aerosol Mass Spectrometry, *Environ. Sci. Technol.*, 42 (12), 4478–4485, <https://doi.org/10.1021/es703009q>, 2008.”

Comment (5): In the back trajectory analysis, one could test the sensitivity to the assumed starting height of 500 m by varying this parameter over some range and see if the predicted trajectories change much?

Response: We do some sensitivity tests for the starting height. Figure R1 and R2 show the 72-h backward trajectories and potential source regions for $b_{\text{abs-BrC}}$ based on the assumed starting height of 750 m and 1000 m, respectively. Although there are some difference in the 72-h backward trajectories under the assumption of height of 500 m, 750, and 1000 m, the spatial distributions of potential source for $b_{\text{abs-BrC}}$ from BBOA and po-OOA were similar. Therefore, the conclusion of “The source regions with the highest CWT values were located in the northern Myanmar and along the China-Myanmar border, while the CWT values in the areas surrounding Gaomeigu were relatively low.” in this study is robust and reliable.




Figure R1. (a) 72-h backward trajectories of Gaomeigu from 8:00 on 14 to 23:00 on 31 March, 2018. (b) and (c) Concentration weighted trajectory (CWT) maps of $b_{\text{abs-BrC}}$ at 370 nm (Mm^{-1}) from BBOA and po-OOA, respectively, based on the assumed starting height of 750 m.




Figure R2. (a) 72-h backward trajectories of Gaomeigu from 8:00 on 14 to 23:00 on 31 March, 2018. (b) and (c) Concentration weighted trajectory (CWT) maps of $b_{\text{abs-BrC}}$ at 370 nm (Mm^{-1}) from BBOA and po-OOA, respectively, based on the assumed starting height of 1000 m.

Comment (6): In the calculation of SFE, one might also do a sensitivity test on the assumed variables, possibly most importantly the albedo (also what type of ground cover does a surface albedo of 0.9 represent)?

Response: We are sorry that this is a writing error, and the surface albedo used in our study was 0.19. We have corrected this mistake:

“ τ_{atm} , F_c , and a_s are the atmospheric transmission (0.79), the cloud fraction (0.6), and the surface albedo (0.19), respectively, which are constants from the global average calculations” (Page 8 Line 218–219)

For SFE calculation, the relative uncertainty of SFE can be estimated as follows:

$$U_{\text{SFE}} = \sqrt{(2 \times U_{\tau_{\text{atm}}})^2 + U_{(1 - F_c)}^2 + U_{a_s}^2}$$

Here, $U_{\tau_{\text{atm}}}$, $U_{(1 - F_c)}$, and U_{a_s} represent the relative uncertainties of τ_{atm} , F_c , and a_s , respectively. Take an example, when τ_{atm} , F_c , and a_s have relative uncertainties of 10%,

$U_{\text{SFE}} = \sqrt{(2 \times 0.1)^2 + \left(\frac{0.6 \times 0.1}{1 - 0.6}\right)^2} + 0.1^2 \approx 0.27$. In this study, the atmospheric transmission, cloud fraction, and surface albedo are not measured values, we used the global average values in order to compare with other studies better.

Comment (7): Line 217, 18-41 fold is relative to what?

Response: 18–41 folds is the result of the maximum hourly b_{abs} value relative to the minimum hourly b_{abs} value during the sampling period. To make this point clear, we have made the following revisions:

“The hourly b_{abs} values at different wavelengths varied from minimum to maximum values by factors of 19–41 from 14 to 31 March 2018, reflecting that atmospheric environment at Gaomeigu is influenced by dynamic changes in emission sources and meteorological condition.” (Page 9 Line 228–230)

Comment (8): What are the bracketed variables in Table 1?

Response: The bracketed variables in Table 1 represented the minimum and maximum values of hourly b_{abs} during the sampling period. To make it clear, we have modified the relevant note in Table 1:

“**Table 1.** Submicron aerosol light absorption coefficient (b_{abs}) contributed by BrC ($b_{\text{abs-BrC}}$) and BC ($b_{\text{abs-BC}}$) at Gaomeigu during the sampling period (March 14th to 31th, 2018).”

Parameter*	Wavelength					
	370 nm	470 nm	520 nm	590 nm	660 nm	880 nm
b_{abs}	33.1 ± 24.4 (4.7–160.0)**	26.7 ± 19.7 (3.8–138.4)	20.3 ± 13.9 (2.6–93.0)	18.2 ± 12.5 (2.1–84.8)	13.7 ± 9.0 (1.7–56.9)	8.0 ± 4.9 (1.5–28.6)
$b_{\text{abs-BrC}}$	12.3 ± 13.8	10.7 ± 11.5	6.0 ± 6.0	5.8 ± 5.8	2.7 ± 2.6	0.0 ± 0.0
$b_{\text{abs-BC}}$	20.8 ± 12.8	16.0 ± 9.8	14.3 ± 8.8	12.4 ± 7.7	11.0 ± 6.8	8.0 ± 4.9

* $b_{\text{abs-BrC}}$ and $b_{\text{abs-BC}}$ were calculated based on the $\text{AAE}_{\text{BC}} = 1.1$.

**The measured hourly b_{abs} from minimum to maximum values.”

Comment (9): The AAE frequency distribution is stated to be normally distributed (line 225), but it looks possibly bimodal. What is the justification for stating it is normally distributed?

Response: Thank you for pointing this out. We did the Kolmogorov-Smirnov and Shapiro-Wilk tests for AAE values with p values less than 0.01. The results proved that the frequency distribution of AAE was not normally distributed. The relevant description has been revised as follows:

“Frequency histograms of hourly AAE values showed the media AAE value of 1.59 with interquartile range from 1.38 to 1.83 (Fig. S1). Over 72 % of AAE values were higher than 1.4 (Upper limit of AAE_{BC}), implying the presence of both BrC and BC absorption in the submicron aerosol at Gaomeigu.” (Page 9 Line 237–238)

Comment (10): Line 259 and on discussing po-OOA correlation with O₃ and RH. The logic here is not clear. Both O₃ and RH vary substantially both spatially and even diurnally. Given this, how can O₃ and RH at the measurement site be used to infer what the particles were exposed to over the time when transported from source region to where measured? It is the history of what the particles were exposed to that determines the properties at the measurements, the conditions at the measurement site may have only a small or minor impact.

Response: Yes. In fact, we are unable to elucidate what reactions occur in the secondary OA during atmospheric processes with the data measured in the sampling site alone. Establishing the relationship between indicators of oxidation (e.g., O₃, O_x, RH, and aerosol liquid water content) and secondary OA is the useful way to explore the potential formation mechanisms, but it is still not 100% confirmed whether these indicators are involved in secondary formation. In this study, both po-OOA mass concentration and its fraction in OA increased with increasing O₃ ($R^2 = 0.79-0.87$), however, neither of them correlated with RH. These results indicated, to some extent, that high O₃ was the possible driving factor of po-OOA formation. Meanwhile, the intense photochemical environment is an inherent feature of the TP. Thus, we believe that photochemical oxidation is an important potential pathway for the formation of po-OOA.

Comment (11): Line 261 to 264, seems that another explanation is that most of the po-OOA is processed BBOA, as discussed above.

Response: As addressed in the comment (2), in the revised manuscript, po-OOA was more oxygenated than BBOA, and had the lower MAC possibly due to more photobleaching effect.

Comment (12): The last statement of the Discussion (line 319) and in the conclusions is not clear. That is, how does knowledge of the secondary BrC help tackling climate change? Be more specific, this is too general a statement to be meaningful.

Response: Thank you for this comment. The revised discussion shows:

“The fractional radiative forcing by two OAs relative to BC was as high as 48.8 ± 15.5 %, in which the relative radiative forcing of po-OOA to BC (24.2 ± 13.2 %) was almost equal that of BBOA to BC (24.6 ± 9.1 %) (Fig. 7b). These results suggested that BrC emitted from biomass burning and formed by photochemical oxidation was an efficient radiative forcing agent, which, along with BC, can remarkably disturb the radiative balance over the TP. Thus, the inclusion of BrC in the climate models will provide a better understanding of climate change of the southeastern TP. It should also be noted that although BBOA and po-OOA had similar radiative effects, effective measures on tackling the impact of BrC are to reduce primary emission and volatile organic precursor of BrC from biomass burning in the future, since the intense photochemical environment is

an inherent feature of the TP.” (Page 12 Line 340–Page 13 Line 348)

Responses to Referee #2:

General comments:

This manuscript presents a comprehensive analysis of light-absorbing aerosols, especially organic aerosol, in the southeastern margin of Tibetan Plateau during the pre-monsoon season. Authors found light absorption of BrC was mainly contributed by biomass burning and secondary formation. Furthermore, radiative effect of BrC was found to be non-negligible compared to that of BC. Generally, in view of the concern about the melting of plateau glaciers caused by global warming, it is of great significance to evaluate the optical properties, sources and formation processes of BrC in the TP. The results of this study provide valuable dataset and analysis to illustrate the importance of BrC on climate change in the TP, which will be helpful in model simulation of radiative effect of aerosols. Overall, this manuscript is well written and results are interesting with novelty. I recommend this manuscript should be accepted for publication after addressing the following issues.

Response: We highly appreciate the thoughtful and valuable suggestions by the reviewer, which are helpful for us to improve the quality of our manuscript. We have carefully addressed the comments in point-by-point form as shown below. Detailed responses to each of the reviewer's comments are provided in blue, and the revised text is underlined. Attached please also find the marked-up manuscript to track the changes in the revised manuscript.

Major comments:

Comment (1): In this study, the mass spectra of po-OOA were characterized by high m/z 44 and low m/z 43. This indicated po-OOA was more-oxidized, which was similar to MO-OOA identified in AMS and ACSM studies. Generally, MO-OOA has lower MAC compared to primary OA due to photo-bleaching effect; however, in this study the estimated MAC of po-OOA was larger than that of BBOA. Could the authors explain it? On the other hand, I suggest authors could conduct the tests of collinearity and heteroscedasticity of independent variables considering the good correlation between BBOA and po-OOA, and this would guarantee the accuracy of multiple linear regression as shown in Eq. 5.

Response: We agree with the reviewer that the more-oxidized OOA might have the lower MAC compared to primary OA due to photo-bleaching effect. We thus have conducted multicollinearity and heteroscedasticity tests for the multiple linear regression (MLR) model, following the reviewer's suggestion. We find that there is no serious multicollinearity between two independent variables ([BBOA] and [po-OOA]), however, the heteroscedasticity existed when using the ordinary least square fitting.

To address this concern, we use the weighted least squares method to estimate parameter of MLR model. Figure 5 shows the latest results of MAC_{BBOA} and MAC_{po-OOA} . That is, the MAC_{BBOA} is higher than the MAC_{po-OOA} at wavelengths from 370 to 660 nm. The fact that po-OOA is more oxygenated than BBOA means more photobleaching on po-OOA. This might result in the lower MAC of po-OOA (Lee et


al., 2014). In the revised manuscript, we first add the testing results and fitting method of MLR to the Sect. 2.3.3:

“Tolerance (0.2) and variance inflation factor (4.7) for the ordinary least square fitting results indicated that there was no serious multicollinearity between two independent variables, however, heteroscedasticity existed according to “White Test” ($p < 0.05$). Thus, the weighted least squares method was used for parameter estimation in MLR model.” (Page 6 Line 163–166)

We also update the MAC results and give the explanation that po-OOA has the lower MAC compared to BBOA in the Sect. 3.3:

“Compared with BBOA, more oxygenated po-OOA was possibly more photochemically bleached, which resulted in the lower MAC (Lee et al., 2014).” (Page 11 Line 296–297)

In the revised manuscript, Figure 5 shows:



“Figure 5. (a) The mass absorption cross section of BBOA and po-OOA (MAC_{BBOA} and $\text{MAC}_{\text{po-OOA}}$, respectively) at five wavelengths ($\lambda = 370, 470, 520, 590,$ and 660 nm). The circle and shaded area represent the mean MAC values and the standard deviations, respectively. The dashed line is power-law fit. (b) Light absorption coefficient of BrC ($b_{\text{abs-BrC}}$) from BBOA and po-OOA and its fraction in the total reconstructed BrC absorption at different wavelengths.”

In the reference list, we add the new reference:

“Lee, H. J., Aiona, P. K., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Effect of solar radiation on the optical properties and molecular composition of laboratory proxies of atmospheric brown carbon, *Environ. Sci. Technol.*, 48(17), 10217–10226, <https://doi.org/10.1021/es502515r>, 2014.”

Comment (2): The amount of solar energy absorbed by BrC relative to BC ($f_{\text{BrC/BC}}$) was used to evaluate the radiative effect of BrC. In this study, the $f_{\text{BrC/BC}}$ was calculated by “simple forcing efficiency” and mass concentration. In fact, there are many studies have been conducted by $f_{\text{BrC/BC}}$ calculation according to the method reported by Kirillova et al. (2014, JGR-A). Have the authors compared these two approaches? Furthermore, the uncertainties of $f_{\text{BrC/BC}}$ need to be discussed. The uncertainties can be caused by e.g., the errors of assumption parameter, MAC and mass concentration.


Response: Following the Kirillova et al. (2014), the fraction of solar energy absorbed by BrC relative to BC ($f_{\text{BrC/BC}}$) can be calculated as follows:

$$f_{\text{BrC/BC}} = \frac{\int_{370}^{660} S_{\theta}(\lambda) \{1 - e^{-(\text{MAC}_{\text{BrC}}(\lambda) \times C_{\text{BrC}} \times h_{\text{ABL}})}\} d\lambda}{\int_{370}^{660} S_{\theta}(\lambda) \{1 - e^{-(\text{MAC}_{\text{BC}}(\lambda) \times C_{\text{BC}} \times h_{\text{ABL}})}\} d\lambda}$$

Here, S_{θ} is the solar irradiance based on the ASTM G173-03 reference spectra; MAC is the mass absorption cross section; C_{BrC} and C_{BC} are the mass concentrations of BrC and BC; h_{ABL} is the height of the atmospheric boundary layer, which is set to 1000 m. Based on the Kirillova’s method, the average fractional radiative forcing by two OAs relative to BC was 45.0 %, which is close to the result (48.8 %) calculated by “simple forcing efficiency” and mass concentration in the manuscript. This suggests that both methods are reasonable and valid in assessing the radiative effect of BrC relative to BC.

$f_{\text{BrC/BC}}$ is a relative value, and its uncertainties come from the errors of MAC and mass concentration. Considering that MAC with a 1 nm resolution is extrapolated using the power law fitting in this study, we gave the uncertainties in $f_{\text{BrC/BC}}$ caused by the variations in mass concentration as shown in Figure 7.

In the revised manuscript, Figure 7 shows:



“Figure 7. (a) Simple forcing efficiency (SFE) of light-absorbing particles from 370 to 660 nm and the integrated SFE over the entire

solar spectra (370–660 nm). (b) The fraction of solar radiation absorbed by OA components relative to BC. In each panel, the short line and square inside the boxes indicate the median and mean values, respectively. The lower and upper edges of the boxes denote the standard deviation. The vertical bars (“whiskers”) show the 5th and 95th percentiles. Scattered data points and normal distribution curve are also shown.”

Comment (3): From the results of this study, biomass burning emission and photochemical reactions both contributed to the formation of BrC. Since the intense photochemical environment is an inherent feature of the TP, it is difficult to control the photochemical production of BrC. Thus, effective control measures are to reduce the anthropogenic emissions of biomass burning. The authors should add some discussions at the end of the results to imply the significance of this work.

Response: Thank you for point this out. We add the relevant description in the revised discussion:

“The fractional radiative forcing by two OAs relative to BC was as high as 48.8 ± 15.5 %, in which the relative radiative forcing of po-OOA to BC (24.2 ± 13.2 %) was almost equal that of BBOA to BC (24.6 ± 9.1 %) (Fig. 7b). These results suggested that BrC emitted from biomass burning and formed by photochemical oxidation was an efficient radiative forcing agent, which, along with BC, can remarkably disturb the radiative balance over the TP. Thus, the inclusion of BrC in the climate models will provide a better understanding of climate change of the southeastern TP. It should also be noted that although BBOA and po-OOA had similar radiative effects, effective measures on tackling the impact of BrC are to reduce primary emission and volatile organic precursor of BrC from biomass burning in the future, since the intense photochemical environment is an inherent feature of the TP.” (Page 12 Line 340–Page 13 Line 348)

Specific Comments:

Comment (4): Line 12: Replace "as well as" with “compared to”.

Response: We have replaced “as well as” with “compared to” as follows:

“Brown carbon (BrC) in the atmosphere can greatly influence aerosol’s radiative forcing over the Tibetan Plateau (TP), because it has the non-negligible capacity of light absorption compared to black carbon (BC);” (Page 1 Line 11–12)

Comment (5): Line 25: Replace “BrC from biomass burning emissions” with “BrC of biomass burning origin”.

Response: We have replaced “BrC from biomass burning emissions” with “BrC of biomass burning origin” as follows:

“This study highlighted a significant influence of BrC of biomass burning origin and secondary formation on climate change over the TP region during the pre-monsoon season.” (Page 1 Line 26–28)

Comment (6): Line 38-39: Add citation to the statement of “In particular, the atmospheric heating of BrC is greater than that of BC in the tropical mid and upper troposphere.”.

Response: The statement of “In particular, the atmospheric heating of BrC is greater than that of BC in the tropical mid and upper troposphere.” is also cited from Zhang et al. (2020a). To make it clear, we revised the text as follows:

“Zhang et al. (2020a) estimated that globally BrC contributed more than 25% of BC RF, wherein the atmospheric heating of BrC is greater than that of BC in the tropical mid and upper troposphere.” (Page 2 Line 38–40)

Comment (7): Line 66-67: Replace “plateau” with “TP”. Full name of TP is already mentioned in Line 56.

Response: We have replaced “plateau” with “TP”. It now reads:

“For example, Zhang et al. (2015) reported that biomass burning from South Asia has the largest impact on BC in the central TP, while fossil fuel combustion contributed the most to BC burden in the northeast TP in all seasons and southeast TP in the summer.” (Page 3 Line 66–68)

Comment (8): Line 182: Check the publication year of Draxler and Hess.

Response: We have corrected this citation information.

“The trajectories were calculated with the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model developed by the National Oceanic and Atmospheric Administration (Draxler and Hess, 1998).” (Page 7 Line 192–194)

Comment (9): Line 216: Please consider to put Fig. S1 in the main figures of manuscript. Fig. S1 indicates the periods of high concentrations and b_{abs} that later discussed in this manuscript.

Response: We have moved Fig. S1 to the main manuscript and assigned as Fig. 2.

Comment (10): Line 256: Replace "as well as" with “similar to”.

Response: We have replaced “as well as” with “similar to” as follows:

“The most abundant peak in mass spectrum of po-OOA was at m/z 44 (f_{44} , 27.8 %), similar to those in mass spectra of more-oxidized oxygenated OA (MO-OOA) ($f_{44} > 20$ %) identified frequently in

previous studies (Tobler et al., 2021; Xu et al., 2018; Zhang et al., 2019).” (Page 10 Line 272–274)

Comment (11): Line 261: Replace "affect most" with “strongly affect”.

Response: We have deleted this sentence.

Comment (12): Line 283: How the $rb_{\text{abs-BrC}}$ was calculated? Is it reconstructed from the b_{abs} of BBOA and po-OOA? Have the authors compared the $rb_{\text{abs-BrC}}$ with the calculated $b_{\text{abs-BrC}}$ from Eq. 2?

Response: Yes, $rb_{\text{abs-BrC}}$ is calculated by the sum of the reconstructed b_{abs} from BBOA and po-OOA based on the MLR model. To express this point clear, the following sentence has been added in the revised manuscript:

“The total reconstructed light absorption coefficient of BrC ($rb_{\text{abs-BrC}}$) was the sum of $b_{\text{abs-BrC}}$ from BBOA and po-OOA obtained from the MLR model.” (Page 11 Line 301–302)

Impacts of biomass burning and photochemical processing on the light absorption of brown carbon in the southeastern Tibetan Plateau

Jie Tian^{1,2}, Qiyuan Wang^{1,2}, Yongyong Ma³, Jin Wang¹, Yongming Han^{1,2}, and Junji Cao^{1,4}

¹Key Laboratory of Aerosol Chemistry and Physics, State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710061, China

²CAS Center for Excellence in Quaternary Science and Global Change, Xi'an 710061, China

³Meteorological Institute of Shaanxi Province, Xi'an 710015, China

⁴Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

10 *Correspondence:* Qiyuan Wang (wangqy@ieecas.cn) and Junji Cao (jjcao@mail.iap.ac.cn)

Abstract. Brown carbon (BrC) in the atmosphere can greatly influence aerosol's radiative forcing over the Tibetan Plateau (TP), because it has the non-negligible capacity of light absorption ~~compared to as well as~~ black carbon (BC); however, our understanding of optical properties, sources, atmospheric processes of BrC in this region remains limited. In this study, a multiple-wavelength Aethalometer coupled with a quadrupole aerosol chemical speciation monitor were deployed to investigate the highly time resolved BrC in the submicron aerosol in the southeastern edge of the TP during the pre-monsoon season. The result showed that BrC had the substantial contributions (20.0–40.2 %) to the light absorption of submicron aerosol from 370 to 660 nm. Organic aerosol (OA), an alternative to BrC, was split into a biomass burning OA (BBOA) with aging process and a photochemical-oxidation processed oxygenated OA (po-OOA) by a hybrid environmental receptor model analysis. Combined with light absorption coefficient of BrC ($b_{\text{abs-BrC}}$), the source-specific mass absorption cross section of BBOA (~~0.6150–2.78175~~ $\text{m}^2 \text{g}^{-1}$) and po-OOA (~~0.308–1.43215~~ $\text{m}^2 \text{g}^{-1}$) at 370–660 nm were retrieved. On average, $b_{\text{abs-BrC}}$ from po-OOA (~~1.31–6.03~~ Mm^{-1}) was ~~comparable to higher than~~ that from BBOA (~~1.307–6.023~~ Mm^{-1}) at all wavelengths. The concentration weighted trajectory analysis showed that the most important potential source regions for $b_{\text{abs-BrC}}$ values from BBOA and po-OOA were located in the northern Myanmar and along the China-Myanmar border, indicating the cross-border transport of BrC from Southeast Asia. A “simple forcing efficiency” evaluation further illustrated the importance of BrC radiative effect with the high fractional radiative forcing by two OAs relative to BC (~~48.8506 ± 15.5187~~ %). This study highlighted a significant influence of BrC ~~of biomass burning origin from biomass burning emissions~~ and secondary formation on climate change over the TP region during the pre-monsoon season.

1 Introduction

30 Carbonaceous aerosols, a major component of atmospheric particles, play an important role in the global climate by directly absorbing and scattering solar and terrestrial radiation (Bellouin et al., 2013; IPCC, 2013; Yao et al., 2017). In the past, black carbon (BC) was often considered to be the only light-absorbing carbonaceous aerosol, and organic aerosol (OA) was thought to purely scatter light (Bond and Bergstrom, 2006; Koch et al., 2007). However, a fraction of OA has been found to absorb radiation efficiently in near-ultraviolet (UV) and visible spectral ranges with a strong
35 wavelength dependence (Kirchstetter et al., 2004). This light-absorbing OA, collectively known as brown carbon (BrC) (Andreae and Gelencsér, 2006), is receiving increasing attention due to its non-negligible radiative effect. Feng et al. (2013) and Lin et al. (2014) have reported that the radiative forcing (RF) caused by BrC absorption on a global scale can be up to + 0.25 and + 0.57 W m⁻², respectively. Zhang et al. (2020a) estimated that globally BrC contributed more than 25% of BC RF. ~~In particular,~~ wherein the atmospheric heating of BrC is greater than that of BC in the tropical mid and
40 upper troposphere. Zhang et al. (2017) also suggested that the clear-sky RFs of high- and low-altitude BrC were 0.35 ± 0.16 and 0.65 ± 0.34 W m⁻², corresponding to 34% and 24% of carbonaceous aerosol warming effect at the tropopause, respectively. The inclusion of BrC in climate models can reduce uncertainties in the global or regional RF assessment of aerosols. Therefore, a comprehensive understanding on light absorption properties of BrC is required.

In the atmosphere, primary BrC is mainly emitted from biomass burning and fossil fuel combustion (Olson et al., 2015; Washenfelder et al., 2015; Xie et al., 2017), and secondary BrC is commonly formed from photochemical-oxidation and aqueous reactions of biogenic or anthropogenic precursors (Hecobian et al., 2010; Nakayama et al., 2013). The complex sources and formation mechanisms of BrC lead to the spatial-temporal variations in its light absorption properties. Accurately quantifying the source-specific absorption capacity (i.e., mass absorption cross section (MAC)) of BrC is essential for modelling BrC climate effect. However, direct source apportionment of BrC is impossible with current
45 analytical method, since BrC constituents responsible for light absorption remain relatively unknown (Laskin et al., 2015). Recent studies have usually used OA as an alternative to BrC for two major reasons: (1) the definition of OA contains all BrC constituents, and (2) OA from either online monitoring or filter extraction can be apportioned to a few major primary and secondary sources with the development of mass spectrometry. This allows for establishing the relationship between primary and secondary OA types and BrC absorption, which provides better quantification of the
50 impact of BrC from different sources and formation mechanisms on regional and global climates (Kaskaoutis et al., 2021; Moschos et al., 2018; Qin et al., 2018; Wang et al., 2021).

The Tibetan Plateau (TP), often referred to as the “Third Pole”, is the largest and highest mountain region in the world and contains the most abundant ice outside of the polar regions (Yao et al., 2012). It has a huge impact on the large-scale atmospheric circulation and the hydrological cycle, which is the most sensitive and visible indicator of climate change

60 in the entire Asian continent (Chen and Bordoni, 2014; Immerzeel et al., 2010). In the recent decades, there has been the growing evidence of increased surface temperature in the Himalayas and the TP regions, accompanied by the accelerated glacier melt and retreat (Kehrwald et al., 2008; Liu and Chen, 2000; Wang et al., 2008). This rapid warming was firstly attributed to greenhouse gas warming; however, light-absorbing aerosols were found to be another major warming agent (Lau et al., 2010; Ramanathan et al., 2007). Previous studies paid a large amount of attention on BC due to its vital climatic implication in the TP. The sources of BC varied significantly with the receptor location and the season (Kopacz et al., 2011; Tan et al., 2021; Zhang et al., 2015). For example, Zhang et al. (2015) reported that biomass burning from South Asia has the largest impact on BC in the central **TPplateau**, while fossil fuel combustion contributed the most to BC burden in the northeast **TPplateau** in all seasons and southeast **TPplateau** in the summer. The direct RF of BC at the top of the atmosphere (+ 1.6–3.5 W m⁻²) induced atmospheric heating rates of 0.13–0.35 K day⁻¹ in the Himalayas and the TP regions (Liu et al., 2021; Panicker et al., 2020); meanwhile, BC deposited on the snow-covered areas can increase 1.0 °C of the surface temperature over the TP by reducing the snow albedo (Qian et al., 2011). It is clearly that there are primary OA emissions along with BC emitted from biomass burning and fossil fuel combustion, and secondary OA formation has been found in the TP (Xu et al., 2018; Zhang et al., 2019). However, the link between light absorption properties and sources of OA is less understood so far, which would lead to uncertainties in evaluating aerosol radiative effect of TP.

In this study, real-time measurements of both light absorption properties and chemical characteristics of submicron aerosol were conducted in the southeastern margin of the TP during the pre-monsoon season. The main objectives were to (1) characterize the light absorption properties of BrC, (2) quantify the source-specific MAC and absorption of BrC, and (3) evaluate the importance of BrC radiative effect from different sources. This study provides insights into light absorption properties of BrC, which is necessary for understanding the role of BrC in climate warming and revealing impacts of sources and atmospheric processes in the TP and surrounding areas.

2 Methodology

2.1 Sampling site and period

Submicron aerosol online measurements of optical and chemical properties were performed at the Lijiang Astronomical Station, Chinese Academy of Sciences, Gaomeigu County, Yunnan Province (26 °41' N, 100 °1' E; 3260 m a.s.l.) (Fig. 1). Continuous hourly O₃ and relative humidity (RH) were measured with the use of an ozone analyzer (EC9810, Ecotech Pty Ltd, Australia) and an automatic weather station (MAWS201, Vaisala, Helsinki, Finland). The monitoring station is situated in the southeastern edge of the TP, a natural channel for the transport of air pollutants from Southeast Asia to the TP (Tan et al., 2021). All instruments were placed on the rooftop of an office building (~10 m above the ground),

90 without any strong anthropogenic emission sources nearby. More detailed description about the sampling site can be found in Wang et al. (2019) and Liu et al. (2021). The sampling period lasted from 8:00 local stand time (LST, all time references that follow are given in LST) on 14 to 23:00 on 31 March, 2018, corresponding to the pre-monsoon season in the TP.

2.2 Submicron aerosol measurements

95 A newly developed Aethalometer (Model AE33, Magee Scientific, Berkeley, CA, USA) was used to measure aerosol light absorption coefficient (b_{abs}) at multiple wavelengths (i.e., 370, 470, 520, 590, 660, and 880 nm) with a 1 min time resolution. Briefly, the ambient air sampled at a flow rate of 5 L min⁻¹ was firstly selected by a PM₁ (particulate matter with an aerodynamic diameter $\leq 1.0 \mu\text{m}$) cyclone separator (BGI SCC 1.197, Mesa Labs, USA) to collect submicron aerosol on the filter. Light at different wavelengths emitted from diodes irradiated two parallel filter spots with deposition
100 rates of 3.85 and 1.15 L min⁻¹, respectively. Thereafter, the two light attenuations measured by optical detectors was used to calculate b_{abs} through a real-time loading effect compensation algorithm. This “dual-spot” technique for the Model AE33 can eliminate the nonlinearity effect caused by increasing amount of aerosol deposit on the filter (Weingartner et al., 2003). Additionally, the Model AE33 automatically used a factor of 2.14 to compensate the scattering effect of quartz filter. Detailed operating principles of the Model AE33 can be found in Drinovec et al. (2015).

105 OA in the non-refractory PM₁ was measured using a quadrupole aerosol chemical speciation monitor (Q-ACSM, Aerodyne Research Inc., Billerica, Massachusetts, USA) with a 30 min time resolution. The aerodynamic lens coupled with a 100 μm diameter critical aperture in the Q-ACSM created a beam of focused submicron aerosols (40–1000 nm aerodynamic diameter), which was vaporized at $\sim 600 \text{ }^\circ\text{C}$, ionized by a 70 eV electron impact, and subsequently characterized with a mass spectrometer. The details of the instrument have been described elsewhere (Ng et al., 2011b).
110 The measured Q-ACSM data was processed by the ACSM local tool version 1.5.3.5 compiled with Igor Pro 6.37 (Wavemetrics, Inc., Lake Oswego, OR, USA) to determine the mass concentration and ion-specified mass spectra of OA. For our study, the default collection efficiency (0.45) and relative ionization efficiency (1.4) were used to obtain OA mass concentration (Middlebrook et al., 2012). The mass concentration and error matrices of organic fragments from mass-to-charge (m/z) 12 to 120 were initialized following the method of Allan et al. (2003).

115 2.3 Data analysis

2.3.1 Separation of BrC and BC absorption

The extrapolation method based on Absorption Ångström exponent (AAE) is widely used to project the absorption at the longer wavelength to the shorter wavelength of the spectrum (Olson et al., 2015; Pokhrel et al., 2017). With an

120 assumption of b_{abs} at 880 nm ($b_{\text{abs}}(880 \text{ nm})$) solely from BC (Kirchstetter et al., 2004), the light absorption coefficient of BC ($b_{\text{abs-BC}}$) at wavelengths (λ) of 370, 470, 520, 590, and 660 nm can be obtained using the following formula:

$$b_{\text{abs-BC}}(\lambda) = b_{\text{abs}}(880 \text{ nm}) \times \left(\frac{880}{\lambda}\right)^{\text{AAE}_{\text{BC}}} \quad (1)$$

Here, b_{abs} and $b_{\text{abs-BC}}$ are given in inverse megameters (Mm^{-1}); AAE_{BC} is assumed to be 1.1 ± 0.3 , which represents the likely range of AAE for BC externally and internally mixed with non-absorbing material (Lack and Langridge, 2013). Then, BrC absorption is derived by subtracting BC absorption from the total submicron aerosol absorption via:

125
$$b_{\text{abs-BrC}}(\lambda) = b_{\text{abs}}(\lambda) - b_{\text{abs-BC}}(\lambda) \quad (2)$$

Here, $b_{\text{abs-BrC}}$ is the light absorption coefficient of BrC (Mm^{-1}).

2.3.2 Hybrid environmental receptor model (HERM) analysis

130 HERM is an effective receptor model, which analysis was performed to retrieve potential sources of OA in this study. The HERM algorithm groups the matrix X (measured mass spectra of organic fragments) into two nonnegative constant matrices G (source contribution) and F (mass spectra of specific sources), and the model residual matrix E, defined as using mass concentration and error matrices of organic fragments measured by the Q ACSM. The principle of HERM has been described elsewhere (Chen and Cao, 2018). Briefly, the HERM is a bilinear receptor model, which decomposes measured organic fragments matrix (X) at the receptor into matrices of the source contributions (G), source mass spectra (F), and the model residual (E):

135
$$X = G \times F + E \quad (3)$$

The model does not require prior mass spectra of sources, and the values of G and F can be obtained using an iterative conjugate gradient algorithm. The principle of HERM has been described in detail elsewhere (Chen and Cao, 2018). The HERM algorithm attempts to solve G and F by minimizing the object function Q, defined as:

140
$$Q = \frac{\sum_{i=1}^I \sum_{j=1}^J (x_{ij} - \sum_{k=1}^K g_{ik} f_{kj})^2}{\sigma_{x_{ij}}^2 + \sum_{k=1}^K (\sigma_{g_{ik}}^2 \sigma_{f_{kj}}^2 + \delta_{jk} \sigma_{x_{ij}}^2)} \quad (4)$$

145 Here, I, J, and K are the number of samples, m/z variables, and sources, respectively; x_{ij} is the measured concentration of the j th m/z in the i th sample; g_{ik} is the contribution of the j th source in the i th sample; f_{kj} is the j th m/z fraction of the total organic fragments in the k th source (so called mass spectrum); $\sigma_{x_{ij}}$ and $\sigma_{f_{kj}}$ represent the error in measured m/z concentration and the variability in constrained mass spectrum, respectively; δ_{jk} is set to 0 or 1 depending on whether the j th m/z in the k th mass spectrum is constrained or unconstrained, respectively.

Before HERM analysis, m/z from 12 to 120 with signal-to-noise between 0.2 and 2 and m/z 44 were down-weighted by increasing their errors by a factor of 2 (Ulbrich et al., 2009). HERM solutions from two to five factors with unconstrained mass spectrum were investigated to explore potential sources. The two-factor solution were chosen as the optimal solution, while greater number of factors (3–5) solutions existed many non-physical meaning factors dominated by individual m/z and do not further split new sources. Bootstrap (BS) method was adopted for two-factor solution (Brown et al., 2015). In 50 times BS, no mass spectrum was unmapped ($r < 0.6$) indicating the two-factor solution was robust. Therefore, a biomass burning OA (BBOA) and a photochemical-oxidation processed oxygenated OA (po-OOA) were finally identified. More detailed description of mass spectra, time series, and correlations with tracers of these two OA factors can be found in Sect. 3.2.

2.3.3 Calculation of optical parameters

~~MAC, expressed by normalized absorption cross sections to the mass of particles, is commonly used to describe the light absorption capacity of aerosols (Bond and Bergstrom, 2006).~~ The MAC of OA component in this study was resolved by the multiple linear regression (MLR) model combined with $b_{\text{abs-BrC}}$ and OA source apportionment results, which is defined. ~~The amount of $b_{\text{abs-BrC}}$ at different wavelengths can be estimated~~ as follows:

$$b_{\text{abs-BrC}}(\lambda) = a_1(\lambda) \times [\text{BBOA}] + a_2(\lambda) \times [\text{po-OOA}] \quad (54)$$

Here, a_1 and a_2 denote the MAC of BBOA (MAC_{BBOA}) and po-OOA ($\text{MAC}_{\text{po-OOA}}$), respectively, in square meters per gram ($\text{m}^2 \text{g}^{-1}$); $[\text{BBOA}]$ and $[\text{po-OOA}]$ are the mass concentration of BBOA and po-OOA, respectively, in micrograms per cubic meter ($\mu\text{g m}^{-3}$). Tolerance (0.2) and variance inflation factor (4.7) for the ordinary least square fitting results indicated that there was no serious multicollinearity between two independent variables, however, heteroscedasticity existed according to “White Test” ($p < 0.05$). Thus, the weighted least squares method was used for parameter estimation in MLR model. The MAC of BC (MAC_{BC}) was directly calculated with $b_{\text{abs-BC}}$ divided by the mass concentration of BC ~~[BC]:~~

$$\text{MAC}_{\text{BC}}(\lambda) = \frac{b_{\text{abs-BC}}(\lambda)}{[\text{BC}]}$$

(6), which

Here, ~~[BC]~~ was obtained by dividing b_{abs} (880 nm) by the default MAC (880 nm) used in the Model AE33 ($\mu\text{g m}^{-3}$) (Drinovec et al., 2015).

AAE describes the spectral dependence of light absorption by aerosols, and it ~~often reflect the composition of light absorbing components. Generally, the greater proportion of BrC relative to BC indicates the larger AAE (Lack and Cappa, 2010).~~ Equations (7) and (8) show the calculations of AAE can be calculated using a power law function with b_{abs} and MAC, respectively:

$$b_{\text{abs}}(\lambda) = k_1 \times \lambda^{-\text{AAE}} \quad (75)$$

$$\text{MAC}(\lambda) = k_2 \times \lambda^{-\text{AAE}} \quad (86)$$

Here, k_1 and k_2 are constants independent of wavelength.

2.3.4 Statistical metrics

180 ~~The uncentered correlation coefficient (UC) is a qualitative metric to characterize the similarity between mass spectra of sources, which is defined as follows (Ulbrich et al., 2009):~~

$$\text{UC} = \frac{x \cdot y}{\|x\| \|y\|} \quad (9)$$

Here, x and y represent a pair of mass spectra as vectors.

185 ~~The index of agreement (IOA) is used as an indicator to evaluate the performance of the simulated data from MLR model against the measured data (Willmott, 1981). The IOA varies between 0 (no agreement) and 1 (perfect agreement), and can be expressed as:~~

$$\text{IOA} = 1 - \frac{\sum_{i=1}^N (S_i - M_i)^2}{\sum_{i=1}^N (|S_i - M_{\text{ave}}| + |M_i - M_{\text{ave}}|)^2} \quad (10)$$

Here, N is the total number of the simulated data; S_i and M_i are the simulated and measured $b_{\text{abs-BrC}}$, respectively; and M_{ave} is the average measured $b_{\text{abs-BrC}}$.

190 2.4 Trajectory-related analysis

A geographic information system based software TrajStat was utilized to investigate the influences of regional transport on BrC absorption at Gaomeigu from 14 to 31 March, 2018 (Wang et al., 2009). The trajectories were calculated with the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model developed by the National Oceanic and Atmospheric Administration (Draxler and Hess, 1998). In this study, the model was set to run twenty-four times per day at starting times of 0:00–23:00 with 1 h step. 72-h backward trajectories at the height of 500 m above the ground level at Gaomeigu during the sampling period were produced based on the gridded meteorological data from Global Data Assimilation System (<ftp://arlftp.arlhq.noaa.gov/pub/archives/gdas1>, last access: 1 June, 2022).

195 The concentration weighted trajectory (CWT) method was further used to identify the potential source regions that likely affected the BrC absorption at Gaomeigu (Hsu et al., 2003). The geographic zone covered by the total number of backward trajectories (K) was divided into $I \times J$ grid cells with the resolution of $0.5^\circ \times 0.5^\circ$. The CWT value of each grid can be defined as follows:

$$b_{\text{abs-BrC-}ij} = \frac{\sum_{k=1}^K b_{\text{abs-BrC-}k} \tau_{ijk}}{\sum_{k=1}^K \tau_{ijk}} W_{ij} \quad (47)$$

Here, $b_{\text{abs-BrC-}ij}$ is the average weighted light absorption coefficient of BrC in the ij th cell; $b_{\text{abs-BrC-}k}$ is the light absorption coefficient of BrC observed on the arrival of trajectory k ; and τ_{ijk} is the time spent in the ij th cell by trajectory k . The weighting function of W_{ij} was applied to reduce the effect of the small number of back-trajectory segment endpoints that fall into the grid cell (Wang et al., 2006):

$$W_{ij} = \begin{cases} 1.00 & 135 < n_{ij} \\ 0.70 & 45 < n_{ij} \leq 135 \\ 0.42 & 15 < n_{ij} \leq 45 \\ 0.17 & n_{ij} \leq 15 \end{cases} \quad (428)$$

Here, n_{ij} is the total number of endpoints in the ij th cell. In this study, the total number of endpoints located in 72 cells of the geographic zone is 3268, so that the average number of endpoints in all cells is about 45.

2.5 Radiative effect calculation

The concept “simple forcing efficiency” (SFE) introduced by Bond and Bergstrom (2006) is a useful way to evaluate the radiative effect of atmospheric aerosols. Without consideration of aerosol scattering, a variant of wavelength-dependent SFE is given as follows:

$$\text{SFE}_i(\lambda) = \frac{S_0(\lambda)}{4} \times \tau_{\text{atm}}^2 \times (1 - F_c) \times [4a_s \times \text{MAC}_i(\lambda)] \quad (439)$$

Here, the subscript i represents BBOA, po-OOA, or BC; λ denotes the wavelength from 370 to 660 nm with 1 nm step; SFE is given in watts per gram (W g^{-1}), which represents the positive energy added to the Earth atmosphere system by a given mass of light-absorbing particles in the atmosphere; S_0 is the solar irradiance based on the ASTM G173-03 reference spectra in watts per square meters (W m^{-2}); τ_{atm} , F_c , and a_s are the atmospheric transmission (0.79), the cloud fraction (0.6), and the surface albedo (0.19), respectively, which are constants from the global average calculations; MAC with a 1 nm resolution is extrapolated using Eq. (86). And then, the fraction of solar radiation absorbed by OA component relative to BC ($f_{\text{OA/BC}}$) can be calculated as:

$$f_{\text{OA/BC}} = \frac{\sum_{\lambda=370}^{660} \text{SFE}_{\text{OA}}(\lambda) \times C_{\text{OA}}}{\sum_{\lambda=370}^{660} \text{SFE}_{\text{BC}}(\lambda) \times C_{\text{BC}}} \quad (441)$$

Here, the integrated SFE is the sum of the SFE from 370 to 660 nm; C_{OA} and C_{BC} are the average mass concentrations of OA component and BC during the sampling period.

3.1 Overview of BrC absorption

The temporal variation in submicron aerosol b_{abs} at wavelengths from 370 to 880 nm as well as the OA mass concentration during the entire campaign at Gaomeigu are depicted in Fig. [S1 in the Supplement](#). The hourly b_{abs} values at different wavelengths varied from minimum to maximum values by factors of 198–41 folds from 14 to 31 March 2018, reflecting that atmospheric environment at Gaomeigu is influenced by dynamic changes in emission sources and meteorological condition. Particularly, the larger variations in b_{abs} values at 370–660 nm, compared with that at 880 nm, highlighted the impact of non-BC light-absorbing materials. As shown in Table 1, the average b_{abs} values were $33.1 \pm 24.4 \text{ Mm}^{-1}$ (arithmetic mean \pm standard deviation) at 370 nm, $26.7 \pm 19.7 \text{ Mm}^{-1}$ at 470 nm, $20.3 \pm 13.9 \text{ Mm}^{-1}$ at 520 nm, $18.2 \pm 12.5 \text{ Mm}^{-1}$ at 590 nm, $13.7 \pm 9.0 \text{ Mm}^{-1}$ at 660 nm, and $8.0 \pm 4.9 \text{ Mm}^{-1}$ at 880 nm. The b_{abs} values obtained in this study were comparable with those reported previously at the sampling sites of the TP, where the major anthropogenic sources (i.e., industry, fossil fuel combustion, etc.) are limited locally (Niu et al., 2018; Zhao et al., 2019; Zhu et al., 2021). Frequency histograms of hourly AAE values showed the media AAE value of 1.59 with interquartile range from 1.38 to 1.83 ~~a typical normal distribution, with an average AAE value of 1.62 ± 0.28~~ (Fig. [S12](#)). Over 72 % of AAE values were higher than 1.4 (Upper limit of AAE_{BC}), implying the presence of both BrC and BC absorption in the submicron aerosol at Gaomeigu.

Based on Eqs. (1) and (2), $b_{\text{abs-BrC}}$ and $b_{\text{abs-BC}}$ were separated from the total absorption using the $\text{AAE}_{\text{BC}} = 1.1$. The average $b_{\text{abs-BrC}}$ values during the campaign were 12.3 ± 13.8 , 10.7 ± 11.4 , 6.0 ± 6.0 , 5.8 ± 5.7 , and $2.7 \pm 2.6 \text{ Mm}^{-1}$ at 370, 470, 520, 590, and 660 nm, respectively (Table 1). Figure [2–3](#) shows fractions of light absorption at specific wavelengths by BrC and BC in the submicron aerosol. BrC contributed substantially to b_{abs} , which accounted for 20.0–40.2 % from 370 to 660 nm. The average contributions of $b_{\text{abs-BrC}}$ to b_{abs} in the near-UV and blue spectral ranges (300–500 nm) were higher than those in other visible ranges (520–880 nm), indicating that BrC was a considerable absorbing material at short wavelengths in the atmosphere. It should be noted that the assumption for AAE_{BC} would lead to the biases in the BrC absorption calculation (Lack and Langridge, 2013). The uncertainties of AAE_{BC} (± 0.3) in this study caused variations of 14.3–16.6 %, 10.2–12.2 %, 10.3–11.5 %, 7.7–8.6 %, and 6.6–7.1 % in the estimation of BrC absorption contributions at 370, 470, 520, 590, and 660 nm, respectively (Fig. [23](#)).

With respect to the relationship between BrC absorption and OA components, $b_{\text{abs-BrC}}$ values at 370–660 nm were significantly correlated with OA mass concentrations ($r = 0.64\text{--}0.70$, $p < 0.01$) (Fig. [S3S2](#)), confirming a strong link between BrC-chromophores and OA in the southeastern margin of TP (Lack et al., 2013; Laskin et al., 2015).

3.2 OA source apportionment

255 HERM analysis identified two distinct OA sources, consisting of BBOA and po-OOA. Each of OA sources had unique characteristics on mass spectrum, temporary variation, and atmospheric processes. The detailed source apportionment results of OA are shown in Fig. 34.

The mass spectrum of BBOA resembled that of BBOA obtained in previous studies (Uncentered correlation coefficients $UC = 0.80-0.87$) (Crippa et al., 2013; Ng et al., 2011a; Wang et al., 2017). It was characterized by a prominent peak of m/z 60, and a strong positive correlation was found between BBOA and m/z 60 concentrations ($r = 0.72, p < 0.01$) (Figs. 3a-4a and b). We have known that m/z 60 is a typical molecular fragment of levoglucosan, mannosan, and galactosan, which are good organic tracers of biomass burning (Kalogridis et al., 2018; Kim et al., 2017; Reyes-Villegas et al., 2018). The fraction of m/z 60 in BBOA mass spectrum (f_{60} , 0.9 %) was higher than 0.3% (background level in absence of biomass burning), suggesting the impact of biomass burning at Gaomeigu (Cubison et al., 2011). Scatterplots of f_{44} vs. f_{60} was used to analyze aging degree of BBOA in the atmosphere (Fig. 3e4c). The f_{60} usually decreases from fresh to aged biomass burning emissions because of degradation and oxidation reactions during the atmospheric aging, while the f_{44} increases (Paglione et al., 2020). The f_{60} and f_{44} of BBOA resolved in this study (0.9 % and 6.3 %, respectively) indicates BBOA was partly aged, possibly caused by the long-distance regional transport. This is further demonstrated in our trajectory-related analysis (Sect. 3.3).

270 Another OA source was featured by the high correlation with m/z 44 ($r = 0.97, p < 0.01$), which is a surrogate of oxidation degree (Aiken et al., 2008), as well as previously reported OOA (Tobler et al., 2021; Xu et al., 2018; Zhang et al., 2019). The most abundant peak in mass spectrum of po-OOA was at m/z 44 (f_{44} , 27.8 %), similar to those in mass spectra of more-oxidized oxygenated OA (MO-OOA) ($f_{44} > 20$ %) identified frequently in previous studies (Tobler et al., 2021; Xu et al., 2018; Zhang et al., 2019).~~had a high f_{44} (27.8 %) and a low f_{43} (0.8 %);~~ It indicated that this OA source ~~that~~ was likely related to extensive secondary processes occurring during transport (Wang et al., 2017; Xu et al., 2017). Figure 3d-4d shows that both po-OOA mass concentration and its fraction in OA increased with increasing O_3 ($R^2 = 0.79-0.87$), however, neither of them correlated with RH (Fig. S4S3). These results supported that high O_3 was the possible driving factor of po-OOA formation, thus the term of po-OOA was introduced in this study to stress the importance of photochemical-oxidation process in the TP-~~appeared to affect most formation of po-OOA.~~ Moreover, the temporal variation of mass concentration in po-OOA significantly correlated with that in BBOA ($r = 0.63, p < 0.01$), indicating that a portion of po-OOA could be derived from oxidation of volatile organic precursor from biomass burning (Bruns et al., 2016; Posner et al., 2018).

3.3 Source-dependent characteristics of BrC absorption

To further quantify the contributions of OA sources to BrC light absorption, a MLR model was applied to retrieve MAC values from BBOA and po-OOA. As shown in Fig. 54a, the wavelength dependences of MAC_{BBOA} and MAC_{po-OOA} were generally described by the power-law relationship ($R^2 = 0.8477-0.87$), and the AAE of BrC values for BBOA and po-OOA were greater than 2.0 (Kirchstetter et al., 2004). The MAC_{BBOA} was $2.781.75 \pm 0.390.48 \text{ m}^2 \text{ g}^{-1}$ at 370 nm, and dropped to $0.610.50 \pm 0.078 \text{ m}^2 \text{ g}^{-1}$ at 660 nm. Taking the near-UV wavelength as the representative for discussion, the MAC_{BBOA} obtained in this study was within that range from biomass burning ($0.9-7.7 \text{ m}^2 \text{ g}^{-1}$) reported by laboratory experiments and field measurements studies (Kaskaoutis et al., 2021; Kumar et al., 2018; Lack et al., 2012; Qin et al., 2018; Wang et al., 2020; Washenfelder et al., 2015). The differences in light absorption capacity of OA from biomass burning may be partly associated with biomass types and combustion efficiencies (Budisulistiorini et al., 2017; Martinsson et al., 2015; Tian et al., 2019). In addition, the photobleaching effect of aerosol at different aging degree can also lead to the variation in MAC_{BBOA} (Sumlin et al., 2017; Zhong and Jang, 2014). The MAC_{po-OOA} was $1.432.15 \pm 0.234 \text{ m}^2 \text{ g}^{-1}$ at 370 nm, $1.2369 \pm 0.2019 \text{ m}^2 \text{ g}^{-1}$ at 470 nm, $0.6890 \pm 0.1009 \text{ m}^2 \text{ g}^{-1}$ at 520 nm, $0.6685 \pm 0.1009 \text{ m}^2 \text{ g}^{-1}$ at 590 nm, and $0.308 \pm 0.04 \text{ m}^2 \text{ g}^{-1}$ at 660 nm. Compared with BBOA, more oxygenated po-OOA was possibly more photochemically bleached, which resulted in the lower MAC (Lee et al., 2014). absorbed light more efficiently in the near UV, consisting with previous findings for photochemical production of BrC from biomass burning (Kumar et al., 2018; Moschos et al., 2018).

The MLR model reasonably reconstructed the temporal variation in the measured $b_{abs-BrC}$ values, with the index of agreement IOAs ranged from 0.73 to 0.79 for different wavelengths. The total reconstructed light absorption coefficient of BrC ($rb_{abs-BrC}$) was the sum of $b_{abs-BrC}$ from BBOA and po-OOA obtained from the MLR model. As shown in Fig. 4b5b, the average $b_{abs-BrC}$ values from BBOA were $6.02.3 \pm 3.60.3$, $5.02.2 \pm 3.00.3$, $2.81.3 \pm 1.70.4$, $2.61.2 \pm 1.60.4$, and $1.30.7 \pm 0.80.4 \text{ Mm}^{-1}$ at 370, 470, 520, 590, and 660 nm, respectively. The $b_{abs-BrC}$ from BBOA had considerable contributions contributed about half (48.729.5-5140.12 %) of to the total reconstructed light absorption coefficient of BrC ($rb_{abs-BrC}$) at 370-660 nm, indicating that biomass burning is an important primary source of BrC absorption at Gaomeigu. This was possibly related to transport of pollutants emitted from South and Southeast Asia during the pre-monsoon season, where biomass burning activities are intensive (Zhang et al., 2020b; Zhang et al., 2015). The po-OOA produced larger comparable $b_{abs-BrC}$ values ($6.03 \pm 4.20.7$, $5.14.9 \pm 3.60.7$, $2.86 \pm 2.00.6$, $2.85 \pm 1.90.6$, and $1.31 \pm 0.96 \text{ Mm}^{-1}$ at 370, 470, 520, 590, and 660 nm, respectively), suggesting the critical role of photochemical-oxidation processes in the BrC absorption at Gaomeigu. Four periods are marked in Fig. S12, characterized by the inclusion of obvious rising stages for both OA concentrations and b_{abs} values. The $b_{abs-BrC}$ at 370 nm from BBOA and po-OOA contributed a comparable fraction (42.1 % vs. 57.9 %) the most to $rb_{abs-BrC}$ (63.4 %) during the period I, while the contribution of b_{abs-}

BrC from po-OOA increased significantly during other periods (52.272.3–64.781.4 %) (Fig. S5S4). The rapid increases
315 in $b_{\text{abs-BrC}}$ from po-OOA were likely caused by photochemical-oxidation processes that were favored by relatively high
O₃ condition (75–84 ppb) during the periods II–IV; for comparison, the O₃ mixing ratio during period I was 52 ppb. We
noted that the largest $rb_{\text{abs-BrC}}$ occurred in period II when both primary source emissions and secondary formation were
strong. These results further highlighted the importance of biomass burning and photochemical-oxidation on light
absorption of BrC at Gaomeigu.

320 The air-mass trajectory and CWT analyses were used to identify whether local emission or regional transported air
pollution was the major source of $b_{\text{abs-BrC}}$ from OA components at Gaomeigu. Figure 5a–6a shows the 72-h backward
trajectories of the receptor site during the sampling period, and all those were originated from Myanmar. The percentage
of the trajectories with high OA concentration ($> 5.3 \mu\text{g m}^{-3}$, the median value of hourly OA concentration) exceed 50%.
From the CWT maps of $b_{\text{abs-BrC}}$ at 370 nm, the spatial distributions of potential source for $b_{\text{abs-BrC}}$ from BBOA and po-
325 OOA were similar (Figs. 5b–6b and c). The source regions with the highest CWT values were located in the northern
Myanmar and along the China-Myanmar border, while the CWT values in the areas surrounding Gaomeigu were
relatively low. This indicates that large $b_{\text{abs-BrC}}$ loadings at Gaomeigu were more likely resulted in strong cross-border
transport of BrC from biomass burning and secondary formation than local emission during the pre-monsoon season.

3.4 Radiative effect of BrC

330 As described in Sect. 2.5, a simple model was used to provide a first-order estimate of the radiative effect of light-
absorbing particles. Figure 76a shows SFEs of BBOA, po-OOA, and BC at wavelengths from 370 to 660 nm. The SFE
peaks of BBOA, po-OOA, and BC were observed at the boundary between the UV and blue spectra (i.e., ~450 nm), that
was mainly caused by the high MAC and strong solar irradiance at specific wavelength. The integrated SFEs of BBOA
and po-OOA over the entire solar spectra (370–660 nm) in this study were 24.2 W g⁻¹, 17.4 and 17.0 W g⁻¹, respectively,
335 comparable to that of primary OA (21 W g⁻¹, 300–1000 nm) reported in Lu et al. (2015). The integrated SFE of po-OOA
(12.5 W g⁻¹) was only half of that of BBOA due to the relative lower MAC. BC had a much higher integrated SFE (226.6
W g⁻¹) compared with OAs. This is consistent with the widely acknowledged view that BC is the strongest and most
important light-absorbing particle in the atmosphere (Bahadur et al., 2012; Bond et al., 2013). As the concentration of
OA in the atmosphere is generally greater than that of BC, the importance of BrC radiative effect was further evaluated
340 by calculating the fraction of solar radiation absorbed by OA relative to BC. The fractional radiative forcing by two OAs
relative to BC was as high as 48.850.6 ± 15.518.7 %, in which the relative radiative forcing of po-OOA to BC (24.232.9
± 13.218.0 %) was almost twice equal that of BBOA to BC (24.617.7 ± 9.16.6 %) (Fig. 76b). These results suggested
that BrC emitted from biomass burning and formed by photochemical oxidation was an efficient radiative forcing agent,
which, along with BC, can remarkably disturb the radiative balance over the TP. Thus, the inclusion of BrC in the climate

345 ~~models will provide a better understanding of climate change of the southeastern TP. It should also be noted that although BBOA and po-OOA had similar radiative effects, effective measures on tackling the impact of BrC are to reduce primary emission and volatile organic precursor of BrC from biomass burning in the future, since the intense photochemical environment is an inherent feature of the TP, the radiative effect of BrC, especially secondary BrC, should be given more consideration in tackling climate change of the southeastern TP.~~

350 **4 Conclusion**

This study conducted an intensive real-time measurement at Gaomeigu in the southeastern margin of the TP during the pre-monsoon season to investigate light absorption properties, sources, secondary formation, and radiative effect of BrC in the submicron aerosol. Based on the assumption of $AAE_{BC} = 1$, the average $b_{abs-BrC}$ values were calculated as 12.3 ± 13.8 , 10.7 ± 11.4 , 6.0 ± 6.0 , 5.8 ± 5.7 , and 2.7 ± 2.6 Mm^{-1} at 370, 470, 520, 590, and 660 nm, respectively, which
355 contributed 20.0–40.2 % of the total light absorption. OA was used as an alternative to BrC due to the significant correlation ($r = 0.64$ – 0.70 , $p < 0.01$) between its mass concentration and $b_{abs-BrC}$. The HERM analysis identified two OA sources, including a BBOA and a po-OOA. BBOA was partly aged as evidenced by the f_{60} (0.9 %) and f_{44} (6.3 %) of mass spectrum, while significant positive correlation between po-OOA and O_3 indicated that photochemical-oxidation process was possibly the main pathway for the formation of po-OOA. A MLR model combined with $b_{abs-BrC}$ and OA
360 concentration was used to estimate the MAC of OA. The result showed that po-OOA absorbed light ~~less~~ more efficiently ~~in the near-UV~~ compared with BBOA ~~due to more photobleaching effect~~. The MAC_{BBOA} and MAC_{po-OOA} was ~~2.781-75~~ ± 0.3948 and ~~1.432-15~~ ± 0.234 $m^2 g^{-1}$ at 370 nm, and dropped to ~~0.6150~~ ± 0.078 and ~~0.308~~ ± 0.04 $m^2 g^{-1}$ at 660 nm, respectively. $b_{abs-BrC}$ from BBOA contributed ~~about half (48.7–51.1 %)~~ ~~29.5–40.2 %~~ of the reconstructed $b_{abs-BrC}$ at 370–660 nm, while the rest was from po-OOA. All the 72-h backward trajectories of the Gaomeigu site came from
365 Myanmar. The spatial distributions of potential source regions for $b_{abs-BrC}$ showed the highest CWT values for BBOA and po-OOA were both in the northern Myanmar and along the China-Myanmar border, demonstrating that biomass burning emissions and secondary formation from the cross-border transport of Southeast Asia were the major source of $b_{abs-BrC}$ at Gaomeigu. According to the integrated SFEs of BBOA, po-OOA, and BC over the solar spectra (370–660 nm) (~~24.2-17.4~~, ~~12.5-17.0~~, and 226.6 $W g^{-1}$, respectively), the fractional radiative forcing by BBOA (~~24.6-17.7~~ $\pm 9.16-6$ %) and
370 po-OOA (~~24.2-32.9~~ $\pm 13.2-18.0$ %) relative to BC were obtained, highlighting the importance of BrC radiative effect. This study provides insights into light absorption properties of BrC and its potential impacts on climate change over the TP and surrounding areas.

Data availability. Data used to support the findings in this study are archived at the Institute of Earth Environment, Chinese Academy of Sciences, and are publicly available at <https://doi.org/10.5281/zenodo.74965047034650>.

375 *Competing interests.* The authors declare that they have no conflict of interest.

Author contributions. QW, YH, and JC designed the campaign. JT and JW conducted field measurements. QW, JT, and YM made data analysis and interpretation. JT wrote the paper with contributions from all co-authors.

Acknowledgments. The authors are grateful to Weikang Ran, Yonggang Zhang, and other staff at the sampling sites for their assistance with field sampling.

380 *Financial support.* This research was supported the National Natural Science Foundation of China (grant no. 41877391), the Second Tibetan Plateau Scientific Expedition and Research Program (STEP) (grant no. 2019QZKK0602), and the Youth Innovation Promotion Association of the Chinese Academy of Sciences (grant no. 2019402 and 2022416).

References

- 385 [Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y. L., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prévôt, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with High-Resolution Time-of-Flight Aerosol Mass Spectrometry, *Environ. Sci. Technol.*, 42 \(12\), 4478–4485, <https://doi.org/10.1021/es703009q>, 2008.](#)
- 390 Allan, J. D., Jimenez, J. L., Williams, P. I., Alfarra, M. R., Bower, K. N., Jayne, J. T., Coe, H., and Worsnop, D. R.: Quantitative sampling using an Aerodyne aerosol mass spectrometer – 1. Techniques of data interpretation and error analysis, *J. Geophys. Res.-Atmos.*, 108, 4090, <https://doi.org/10.1029/2002JD002358>, 2003.
- Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, <https://doi.org/10.5194/acp-6-3131-2006>, 2006.
- 395 Bahadur, R., Praveen, P. S., Xu, Y. Y., and Ramanathan, V.: Solar absorption by elemental and brown carbon determined from spectral observations, *P. Natl. Acad. Sci. USA.*, 109, 17366–17371, <https://doi.org/10.1073/pnas.120591010>, 2012.
- Bellouin, N., Quaas, J., Morcrette, J. J., and Boucher, O.: Estimates of aerosol radiative forcing from the MACC re-analysis, *Atmos. Chem. Phys.*, 13, 2045–2062, <https://doi.org/10.5194/acp-13-2045-2013>, 2013.
- 400 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- 405 Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Sci. Tech.*, 40, 27–67, <https://doi.org/10.1080/02786820500421521>, 2006.
- Brown, S. G., Eberly, S., Paatero, P., and Norris, G. A.: Methods for estimating uncertainty in PMF solutions: Examples with ambient air and water quality data and guidance on reporting PMF results, *Sci. Total Environ.*, 518–519, 626–635, <https://doi.org/10.1016/j.scitotenv.2015.01.022>, 2015.

- Bruns, E. A., El Haddad, I., Slowik, J. G., Kilic, D., Klein, F., Baltensperger, U., and Prévôt, A. S. H.: Identification of significant precursor gases of secondary organic aerosols from residential wood combustion, *Sci. Rep-UK.*, 6, 27881, <https://doi.org/10.1038/srep27881>, 2016.
- 415 Budisulistiorini, S. H., Riva, M., Williams, M., Chen, J., Itoh, M., Surratt, J. D., and Kuwata, M.: Light-absorbing brown carbon aerosol constituents from combustion of Indonesian peat and biomass, *Environ. Sci. Technol.*, 51, 4415–4423, <https://doi.org/10.1021/acs.est.7b00397>, 2017.
- Chen, J. Q. and Bordoni, S.: Orographic effects of the Tibetan Plateau on the East Asian Summer Monsoon: An energetic perspective, *J. Climate.*, 27, 3052–3072, <https://doi.org/10.1175/JCLI-D-13-00479.1>, 2014.
- 420 Chen, L. -W. A. and Cao, J. J.: PM_{2.5} source apportionment using a hybrid environmental receptor model, *Environ. Sci. Technol.*, 52, 6357–6369, <https://doi.org/10.1021/acs.est.8b00131>, 2018.
- Crippa, M., DeCarlo, P. F., Slowik, J. G., Mohr, C., Heringa, M. F., Chirico, R., Poulain, L., Freutel, F., Sciare, J., Cozic, J., Di Marco, C. F., Elsasser, M., Nicolas, J. B., Marchand, N., Abidi, E., Wiedensohler, A., Drewnick, F., Schneider, J., Borrmann, S., Nemitz, E., Zimmermann, R., Jaffrezo, J. L., Prévôt, A. S. H., and Baltensperger, U.: Wintertime aerosol chemical composition and source apportionment of the organic fraction in the metropolitan area of Paris, *Atmos. Chem. Phys.*, 13, 961–981, <https://doi.org/10.5194/acp-13-961-2013>, 2013.
- 425 Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., Diskin, G. S., Fisher, J. A., Fuelberg, H. E., Hecobian, A., Knapp, D. J., Mikoviny, T., Riemer, D., Sachse, G. W., Sessions, W., Weber, R. J., Weinheimer, A. J., and Jimenez, J. L.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049–12064, <https://doi.org/10.5194/acp-11-12049-2011>, 2011.
- 430 Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT_4 modelling system for trajectories, dispersion, and deposition, *Aust. Met. Mag.*, 47, 295–308, 1998.
- Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Müller, T., Wiedensohler, A., and Hansen, A. D. A.: The “dual-spot” Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, *Atmos. Meas. Tech.*, 8, 1965–1979, <https://doi.org/10.5194/amt-8-1965-2015>, 2015.
- 435 Du, W., Sun, Y. L., Xu, Y. S., Jiang, Q., Wang, Q. Q., Wang, W., Wang, F., Bai, Z. P., Zhao, X. D., and Yang, Y. C.: Chemical characterization of submicron aerosol and particle growth events at a national background site (3295 m a.s.l.) on the Tibetan Plateau, *Atmos. Chem. Phys.*, 15, 10811–10824, <https://doi.org/10.5194/acp-15-10811-2015>, 2015.
- 440 Duan, J., Huang, R. J., Gu, Y. F., Lin, C. S., Zhong, H. B., Xu, W., Liu, Q., You, Y., Ovadnevaite, J., Ceburnis, D., Hoffmann, T., and O’Dowd, C.: Measurement report: Large contribution of biomass burning and aqueous-phase processes to the wintertime secondary organic aerosol formation in Xi’an, Northwest China, *Atmos. Chem. Phys.*, 22, 10139–10153, <https://doi.org/10.5194/acp-22-10139-2022>, 2022.
- 445 Feng, Y., Ramanathan, V., and Kotamarthi, V. R.: Brown carbon: a significant atmospheric absorber of solar radiation? *Atmos. Chem. Phys.*, 13, 8607–8621, <https://doi.org/10.5194/acp-13-8607-2013>, 2013.
- Florou, K., Papanastasiou, D. K., Pikridas, M., Kaltsonoudis, C., Louvaris, E., Gkatzelis, G. I., Patoulias, D., Mihalopoulos, N., and Pandis, S. N.: The contribution of wood burning and other pollution sources to wintertime organic aerosol levels in two Greek cities, *Atmos. Chem. Phys.*, 17, 3145–3163, <https://doi.org/10.5194/acp-17-3145-2017>, 2017.
- 450

- Hecobian, A., Zhang, X., Zheng, M., Frank, N., Edgerton, E. S., and Weber, R. J.: Water-Soluble Organic Aerosol material and the light-absorption characteristics of aqueous extracts measured over the Southeastern United States, *Atmos. Chem. Phys.*, 10, 5965–5977, <https://doi.org/10.5194/acp-10-5965-2010>, 2010.
- 455 Hsu, Y. K., Holsen, T. M., and Hopke, P. K.: Comparison of hybrid receptor models to locate PCB sources in Chicago, *Atmos. Environ.*, 37, 545–562, [https://doi.org/10.1016/S1352-2310\(02\)00886-5](https://doi.org/10.1016/S1352-2310(02)00886-5), 2003.
- Huang, R. J., Wang, Y. C., Cao, J. J., Lin, C. S., Duan, J., Chen, Q., Li, Y. J., Gu, Y. F., Yan, J., Xu, W., Fröhlich, R., Canonaco, F., Bozzetti, C., Ovadnevaite, J., Ceburnis, D., Canagaratna, M. R., Jayne, J., Worsnop, D. R., El-Haddad, I., Prévôt, A. S. H., and O’Dowd, C. D.: Primary emissions versus secondary formation of fine particulate matter in the most polluted city (Shijiazhuang) in North China, *Atmos. Chem. Phys.*, 19, 2283–2298, <https://doi.org/10.5194/acp-19-2283-2019>, 2019.
- 460 Immerzeel, W. W., van Beek, L. P. H., and Bierkens, M. F. P.: Climate change will affect the Asian water towers, *Science*, 328, 1382–1385, <https://doi.org/10.1126/science.1183188>, 2010.
- IPCC: Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, edited by: Stocker, T. F., Qin, D., Plattner, G. -K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., ISBN 978-1-107-66182-0, 2013.
- 465 Kalogridis, A. C., Popovicheva, O. B., Engling, G., Diapouli, E., Kawamura, K., Tachibana, E., Ono, K., Kozlov, V. S., and Eleftheriadis, K.: Smoke aerosol chemistry and aging of Siberian biomass burning emissions in a large aerosol chamber, *Atmos. Environ.*, 185, 15–28, <https://doi.org/10.1016/j.atmosenv.2018.04.033>, 2018.
- 470 Kaskaoutis, D. G., Grivas, G., Stavroulas, I., Bougiatioti, A., Liakakou, E., Dumka, U. C., Gerasopoulos, E., and Mihalopoulos, N.: Apportionment of black and brown carbon spectral absorption sources in the urban environment of Athens, Greece, during winter, *Sci. Total Environ.*, 801, 149739, <https://doi.org/10.1016/j.scitotenv.2021.149739>, 2021.
- 475 Kehrwald, N. M., Thompson, L. G., Yao, T. D., Mosley-Thompson, E., Schotterer, U., Alfimov, V., Beer, J., Eikenberg, J., and Davis, M. E.: Mass loss on Himalayan glacier endangers water resources, *Geophys. Res. Lett.*, 35, L22503, <https://doi.org/10.1029/2008GL035556>, 2008.
- Kim, H., Zhang, Q., Bae, G. N., Kim, J. Y., and Lee, S. B.: Sources and atmospheric processing of winter aerosols in Seoul, Korea: Insights from real-time measurements using a high-resolution aerosol mass spectrometer, *Atmos. Chem. Phys.*, 17, 2009–2033, <https://doi.org/10.5194/acp-17-2009-2017>, 2017.
- 480 Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *J. Geophys. Res.*, 109, D21208, <https://doi.org/10.1029/2004JD004999>, 2004.
- Koch, D., Bond, T. C., Streets, D., Unger, N., and van der Werf, G. R.: Global impacts of aerosols from particular source regions and sectors, *J. Geophys. Res.-Atmos.*, 112, D02205, <https://doi.org/10.1029/2005JD007024>, 2007.
- 485 Kopacz, M., Mauzerall, D. L., Wang, J., Leibensperger, E. M., Henze, D. K., and Singh, K., 2011. Origin and radiative forcing of black carbon transported to the Himalayas and Tibetan Plateau, *Atmos. Chem. Phys.*, 11, 2837–2852, <https://doi.org/10.5194/acp-11-2837-2011>, 2011.
- Kumar, N. K., Corbin, J. C., Bruns, E. A., Massabó, D., Slowik, J. G., Drinovec, L., Močnik, G., Prati, P., Vlachou, A., Baltensperger, U., Gysel, M., El-Haddad, I., and Prévôt, A. S. H.: Production of particulate brown carbon during atmospheric aging of residential wood-burning emissions, *Atmos. Chem. Phys.*, 18, 17843–17861, <https://doi.org/10.5194/acp-18-17843-2018>, 2018.
- 490 Lack, D. A. and Langridge, J. M.: On the attribution of black and brown carbon light absorption using the Ångström exponent, *Atmos. Chem. Phys.*, 13, 10535–10543, <https://doi.org/10.5194/acp-13-10535-2013>, 2013.

- 495 Lack, D. A. and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, *Atmos. Chem. Phys.*, 10, 4207–4220, <https://doi.org/10.5194/acp-10-4207-2010>, 2010.
- Lack, D. A., Bahreini, R., Langridge, J. M., Gilman, J. B., and Middlebrook, A. M.: Brown carbon absorption linked to organic mass tracers in biomass burning particles, *Atmos. Chem. Phys.*, 13, 2415–2422, <https://doi.org/10.5194/acp-13-2415-2013>, 2013.
- 500 Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, *P. Natl. Acad. Sci. USA.*, 109, 14802–14807, <https://doi.org/10.1073/pnas.120657510>, 2012.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of atmospheric brown carbon, *Chem. Rev.*, 115, 4335–4382, <https://doi.org/10.1021/cr5006167>, 2015.
- 505 Lau, W. K. M., Kim, M. -K., Kim, K. -M., and Lee, W. -S.: Enhanced surface warming and accelerated snow melt in the Himalayas and Tibetan Plateau induced by absorbing aerosols, *Environ. Res. Lett.*, 5, 025204, <https://doi.org/10.1088/1748-9326/5/2/025204>, 2010.
- [Lee, H. J., Aiona, P. K., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Effect of solar radiation on the optical properties and molecular composition of laboratory proxies of atmospheric brown carbon, *Environ. Sci. Technol.*, 48\(17\), 10217–10226, <https://doi.org/10.1021/es502515r>, 2014.](https://doi.org/10.1021/es502515r)
- 510 Lin, G. X., Penner, J. E., Flanner, M. G., Sillman, S., Xu, L., and Zhou, C.: Radiative forcing of organic aerosol in the atmosphere and on snow: Effects of SOA and brown carbon, *J. Geophys. Res.-Atmos.*, 119, 7453–7476, <https://doi.org/10.1002/2013JD021186>, 2014.
- Liu, H. K., Wang, Q. Y., Xing, L., Zhang, Y., Zhang, T., Ran, W. K., and Cao, J. J.: Measurement report: quantifying source contribution of fossil fuels and biomass-burning black carbon aerosol in the southeastern margin of the Tibetan Plateau, *Atmos. Chem. Phys.*, 21, 973–987, <https://doi.org/10.5194/acp-21-973-2021>, 2021.
- 515 Liu, X. D. and Chen, B. D.: Climatic warming in the Tibetan Plateau during recent decades, *Int. J. Climatol.*, 20, 1729–1742, [https://doi.org/10.1002/1097-0088\(20001130\)20:14<1729::AID-JOC556>3.0.CO;2-Y](https://doi.org/10.1002/1097-0088(20001130)20:14<1729::AID-JOC556>3.0.CO;2-Y), 2000.
- Martinsson, J., Eriksson, A. C., Nielsen, I. E., Malmberg, V. B., Ahlberg, E., Andersen, C., Lindgren, R., Nyström, R., Nordin, E. Z., Brune, W. H., Svenningsson, B., Swietlicki, E., Boman, C., and Pagels, J. H.: Impacts of combustion conditions and photochemical processing on the light absorption of biomass combustion aerosol, *Environ. Sci. Technol.*, 49, 14663–14671, <https://doi.org/10.1021/acs.est.5b03205>, 2015.
- 520 Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of composition-dependent collection efficiencies for the aerodyne aerosol mass spectrometer using field data, *Aerosol Sci. Tech.*, 46, 258–271, <https://doi.org/10.1080/02786826.2011.620041>, 2012.
- 525 Moschos, V., Kumar, N. K., Daellenbach, K. R., Baltensperger, U., Prévôt, A. S. H., and El Haddad, I.: Source apportionment of brown carbon absorption by coupling UV/Vis spectroscopy with aerosol mass spectrometry, *Environ. Sci. Technol. Lett.*, 5, 302–308, <https://doi.org/10.1021/acs.estlett.8b00118>, 2018.
- Nakayama, T., Sato, K., Matsumi, Y., Imamura, T., Yamazaki, A., and Uchiyama, A.: Wavelength and NO_x dependent complex refractive index of SOAs generated from the photooxidation of toluene, *Atmos. Chem. Phys.*, 13, 531–545, <https://doi.org/10.5194/acp-13-531-2013>, 2013.
- 530 Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Zhang, Q., Ulbrich, I. M., and Worsnop, D. R.: Real-time methods for estimating organic component mass concentrations from aerosol mass spectrometer data, *Environ. Sci. Technol.*, 45, 910–916, <https://doi.org/10.1021/es102951k>, 2011a.

- 535 Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L., and Jayne, J. T.: An aerosol chemical speciation monitor (ACSM) for routine monitoring of the composition and mass concentrations of ambient aerosol, *Aerosol Sci. Tech.*, 45, 780–794, <https://doi.org/10.1080/02786826.2011.560211>, 2011b.
- 540 Niu, H. W., Kang, S. C., Wang, H. L., Zhang, R. D., Lu, X. X., Qian, Y., Paudyal, R., Wang, S. J., Shi, X. F., and Yan, X. G.: Seasonal variation and light absorption property of carbonaceous aerosol in a typical glacier region of the southeastern Tibetan Plateau, *Atmos. Chem. Phys.*, 18, 6441–6460, <https://doi.org/10.5194/acp-18-6441-2018>, 2018.
- 545 Olson, M. R., Garcia, M. V., Robinson, M. A., Rooy, P. V., Dietenberger, M. A., Bergin, M., and Schauer, J. J.: Investigation of black and brown carbon multiple-wavelength-dependent light absorption from biomass and fossil fuel combustion source emissions, *J. Geophys. Res.-Atmos.*, 120, 6682–6697, <https://doi.org/10.1002/2014JD022970>, 2015.
- 550 Paglione, M., Gilardoni, S., Rinaldi, M., Decesari, S., Zanca, N., Sandrini, S., Giulianelli, L., Bacco, D., Ferrari, S., Poluzzi, V., Scotto, F., Trentini, A., Poulain, L., Herrmann, H., Wiedensohler, A., Canonaco, F., Prévôt, A. S. H., Massoli, P., Carbone, C., Facchini, M. C., and Fuzzi, S.: The impact of biomass burning and aqueous-phase processing on air quality: a multi-year source apportionment study in the Po Valley, Italy, *Atmos. Chem. Phys.*, 20, 1233–1254, <https://doi.org/10.5194/acp-20-1233-2020>, 2020.
- 555 Panicker, A. S., Sandeep, K., Gautam, A. S., Trimbake, H. K., Nainwal, H. C., Beig, G., Bisht, D. S., and Das, S.: Black carbon over a central Himalayan Glacier (Satopanth): Pathways and direct radiative impacts, *Sci. Total Environ.*, 766, 144242, <https://doi.org/10.1016/j.scitotenv.2020.144242>, 2020.
- 560 Pokhrel, R. P., Beamesderfer, E. R., Wagner, N. L., Langridge, J. M., Lack, D. A., Jayarathne, T., Stone, E. A., Stockwell, C. E., Yokelson, R. J., and Murphy, S. M.: Relative importance of black carbon, brown carbon and absorption enhancement from clear coatings in biomass burning emissions, *Atmos. Chem. Phys.*, 17, 5063–5078, <https://doi.org/10.5194/acp-17-5063-2017>, 2017.
- 565 Posner, L. N., Theodoritsi, G., Robinson, A., Yarwood, G., Koo, B., Morris, R., Mavko, M., Moore, T., and Pandis, S. N.: Simulation of fresh and chemically-aged biomass burning organic aerosol. *Atmos. Environ.* 196, 27–37. <https://doi.org/10.1016/j.atmosenv.2018.09.055>, 2018.
- 570 Qian, Y., Flanner, M. G., Leung, L. R., and Wang, W.: Sensitivity studies on the impacts of Tibetan Plateau snowpack pollution on the Asian hydrological cycle and monsoon climate, *Atmos. Chem. Phys.*, 11, 1929–1948, <https://doi.org/10.5194/acp-11-1929-2011>, 2011.
- 575 Qin, Y. M., Tan, H. B., Li, Y. J., Li, Z. J., Schurman, M. I., Liu, L., Wu, C., and Chan, C. K.: Chemical characteristics of brown carbon in atmospheric particles at a suburban site near Guangzhou, China, *Atmos. Chem. Phys.*, 18, 16409–16418, <https://doi.org/10.5194/acp-18-16409-2018>, 2018.
- Ramanathan, V., Ramana, M. V., Roberts, G., Kim, D., Corrigan, C., Chung, C., and Winker, D.: Warming trends in Asia amplified by brown cloud solar absorption, *Nature*, 448, 575–578, <https://doi.org/10.1038/nature06019>, 2007.
- 570 Reyes-Villegas, E., Priestley, M., Ting, Y. C., Haslett, S., Bannan, T., Le Breton, M., Williams, P. I., Bacak, A., Flynn, M. J., Coe, H., Percival, C., and Allan, J. D.: Simultaneous aerosol mass spectrometry and chemical ionisation mass spectrometry measurements during a biomass burning event in the UK: Insights into nitrate chemistry, *Atmos. Chem. Phys.*, 18, 4093–4111, <https://doi.org/10.5194/acp-18-4093-2018>, 2018.
- 575 Sumlin, B. J., Pandey, A., Walker, M. J., Pattison, R. S., Williams, B. J., and Chakrabarty, R. K.: Atmospheric photooxidation diminishes light absorption by primary brown carbon aerosol from biomass burning, *Environ. Sci. Technol. Lett.*, 4, 540–545, <https://doi.org/10.1021/acs.estlett.7b00393>, 2017.

- Tan, T. Y., Hu, M., Du, Z. F., Zhao, G., Shang, D. J., Zheng, J., Qin, Y. H., Li, M. R., Wu, Y. S., Zeng, L. M., Guo, S., and Wu, Z. J.: Measurement report: Strong light absorption induced by aged biomass burning black carbon over the southeastern Tibetan Plateau in pre-monsoon season, *Atmos. Chem. Phys.*, 21, 8499–8510, <https://doi.org/10.5194/acp-21-8499-2021>, 2021.
- 580 Tian, J., Wang, Q. Y., Ni, H. Y., Wang, M., Zhou, Y. Q., Han, Y. M., Shen, Z. X., Pongpiachan, S., Zhang, N. N., Zhao, Z. Z., Zhang, Q., Zhang, Y., Long, X., and Cao, J. J.: Emission characteristics of primary brown carbon absorption from biomass and coal burning: Development of an optical emission inventory for China, *J. Geophys. Res.-Atmos.*, 124, 1879–1893, <https://doi.org/10.1029/2018jd029352>, 2019.
- 585 Tian, J., Wang, Q. Y., Zhang, Y., Yan, M. Y., Liu, H. K., Zhang, N. N., Ran, W. K., and Cao, J. J.: Impacts of primary emissions and secondary aerosol formation on air pollution in an urban area of China during the COVID-19 lockdown, *Environ. Int.*, 150, 106426, <https://doi.org/10.1016/j.envint.2021.106426>, 2021.
- 590 Tobler, A. K., Skiba, A., Canonaco, F., Močnik, G., Rai, P., Chen, G., Bartyzel, J., Zimnoch, M., Styszko, K., Necki, J., Furger, M., Róžański, K., Baltensperger, U., Slowik, J. G., and Prevot, A. S. H.: Characterization of non-refractory (NR) PM₁ and source apportionment of organic aerosol in Kraków, Poland, *Atmos. Chem. Phys.*, 21, 14893–14906, <https://doi.org/10.5194/acp-21-14893-2021>, 2021.
- Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, <https://doi.org/10.5194/acp-9-2891-2009>, 2009.
- 595 Wang, B., Bao, Q., Hoskins, B., Wu, G. X., and Liu, Y. M.: Tibetan Plateau warming and precipitation changes in East Asia, *Geophys. Res. Lett.*, 35, L14702, <https://doi.org/10.1029/2008GL034330>, 2008.
- 600 Wang, J. F., Ye, J. H., Zhang, Q., Zhao, J., Wu, Y. Z., Li, J. Y., Liu, D. T., Li, W. J., Zhang, Y. G., Wu, C., Xie, C. H., Qin, Y. M., Lei, Y. L., Huang, X. P., Guo, J. P., Liu, P. F., Fu, P. Q., Li, Y. J., Lee, H. C., Choi, H., Zhang, J., Liao, H., Chen, M. D., Sun, Y. L., Ge, X. L., Martin, S. T., and Jacob, D. J.: Aqueous production of secondary organic aerosol from fossil-fuel emissions in winter Beijing haze, *P. Natl. Acad. Sci. USA.*, 118, e2022179118, <https://doi.org/10.1073/pnas.2022179118>, 2021.
- Wang, Q. Y., Han, Y. M., Ye, J. H., Liu, S. X., Pongpiachan, S., Zhang, N. N., Han, Y. M., Tian, J., Wu, C., Long, X., Zhang, Q., Zhang, W. Y., Zhao, Z. Z., and Cao, J. J.: High contribution of secondary brown carbon to aerosol light absorption in the southerastern margin of Tibetan Plateau, *Geophys. Res. Lett.*, 46, 4962–4970, <https://doi.org/10.1029/2019GL082731>, 2019.
- 605 Wang, Q. Y., Liu, H. K., Wang, P., Dai, W. T., Zhang, T., Zhao, Y. Z., Tian, J., Zhang, W. Y., Han, Y. M., and Cao, J. J.: Optical source apportionment and radiative effect of light-absorbing carbonaceous aerosols in a tropical marine monsoon climate zone: the importance of ship emissions, *Atmos. Chem. Phys.*, 20, 15537–15549, <https://doi.org/10.5194/acp-20-15537-2020>, 2020.
- 610 Wang, Y. C., Huang, R. J., Ni, H. Y., Chen, Y., Wang, Q. Y., Li, G. H., Tie, X. X., Shen, Z. X., Huang, Y., Liu, S. X., Dong, W. M., Xue, P., Fröhlich, R., Canonaco, F., Elser, M., Daellenbach, K. R., Bozzetti, C., El Haddad, I., Prévôt, A. S. H., Canagaratna, M. R., Worsnop, D. R., and Cao, J. J.: Chemical composition, sources and secondary processes of aerosols in Baoji city of northwest China, *Atmos. Environ.*, 158, 128–137, <https://doi.org/10.1016/j.atmosenv.2017.03.026>, 2017.
- 615 Wang, Y. Q., Zhang, X. Y., and Arimoto, R.: The contribution from distant dust sources to the atmospheric particulate matter loadings at XiAn, China during spring, *Sci. Total Environ.*, 368, 875–883, <https://doi.org/10.1016/j.scitotenv.2006.03.040>, 2006.

- Wang, Y. Q., Zhang, X. Y., and Draxler, R. R.: TrajStat: GIS-based software that uses various trajectory statistical analysis methods to identify potential sources from long-term air pollution measurement data, *Environ. Modell. Softw.*, 24, 938–939, <https://doi.org/10.1016/j.envsoft.2009.01.004>, 2009.
- 620 Washenfelder, R. A., Attwood, A. R., Brock, C. A., Guo, H., Xu, L., Weber, R. J., Ng, N. L., Allen, H. M., Ayres, B. R., Baumann, K., Cohen, R. C., Draper, D. C., Duffey, K. C., Edgerton, E., Fry, J. L., Hu, W. W., Jimenez, J. L., Palm, B. B., Romer, P., Stone, E. A., Wooldridge, P. J., and Brown, S. S.: Biomass burning dominates brown carbon absorption in the rural southeastern United States, *Geophys. Res. Lett.*, 42, 653–664, <https://doi.org/10.1002/2014GL062444>, 2015.
- 625 Weingartner, E., Saathoff, H., Schnaiter, M., Streit, N., Bitnar, B., and Baltensperger, U.: Absorption of light by soot particles: determination of the absorption coefficient by means of aethalometers, *J. Aerosol Sci.*, 34, 1445–1463, [https://doi.org/10.1016/S0021-8502\(03\)00359-8](https://doi.org/10.1016/S0021-8502(03)00359-8), 2003.
- ~~Willmott, C. J.: On the validation of models. *Phys. Geogr.*, 2, 184–194, <https://doi.org/10.1080/02723646.1981.10642213>, 1981.~~
- 630 Xie, M. J., Hays, M. D., and Holder, A. L.: Light-absorbing organic carbon from prescribed and laboratory biomass burning and gasoline vehicle emissions, *Sci. Rep-UK.*, 7, 7318, <https://doi.org/10.1038/s41598-017-06981-8>, 2017.
- Xu, J. Z., Zhang, Q., Shi, J. S., Ge, X. L., Xie, C. H., Wang, J. F., Kang, S. C., Zhang, R. X., and Wang, Y. H.: Chemical characteristics of submicron particles at the central Tibetan Plateau: insights from aerosol mass spectrometry, *Atmos. Chem. Phys.*, 18, 427–443, <https://doi.org/10.5194/acp-18-427-2018>, 2018.
- 635 Xu, W. Q., Han, T. T., Du, W., Wang, Q. Q., Chen, C., Zhao, J., Zhang, Y. J., Li, J., Fu, P. Q., Wang, Z. F., Worsnop, D. R., and Sun, Y. L.: Effects of aqueous-phase and photochemical processing on secondary organic aerosol formation and evolution in Beijing, China, *Environ. Sci. Technol.*, 51, 762–770, <https://doi.org/10.1021/acs.est.6b04498>, 2017.
- 640 Yao, T. D., Thompson, L. G., Mosbrugger, V., Zhang, F., Ma, Y. M., Luo, T. X., Xu, B. Q., Yang, X. X., Joswiak, D. R., Wang, W. C., Joswiak, M. E., Devkota, L. P., Tayal, S., Jilani, R., and Fayziev, R.: Third Pole Environment (TPE), *Environ. Dev.*, 3, 52–64, <http://doi.org/10.1016/j.envdev.2012.04.002>, 2012.
- 645 Yao, H., Song, Y., Liu, M. X., Archer-Nicholls, S., Lowe, D., McFiggans, G., Xu, T. T., Du, P., Li, J. F., Wu, Y. S., Hu, M., Zhao, C., and Zhu, T.: Direct radiative effect of carbonaceous aerosols from crop residue burning during the summer harvest season in East China, *Atmos. Chem. Phys.*, 17, 5205–5219, <https://doi.org/10.5194/acp-17-5205-2017>, 2017.
- Zhang, A. X., Wang, Y. H., Zhang, Y. Z., Weber, R. J., Song, Y. J., Ke, Z. M., and Zou, Y. F.: Modeling global radiative effect of brown carbon: A larger heating source in the tropical free troposphere than black carbon, *Atmos. Chem. Phys.*, 20(4), 1901–1920, <https://doi.org/10.5194/acp-20-1901-2020>, 2020a.
- 650 Zhang, M. X., Zhao, C., Cong, Z. Y., Du, Q. Y., Xu, M. Y., Chen, Y., Chen, M., Li, R., Fu, Y. F., Lei, Z., Kang, S. C., Zhao, D. L., and Yang, Y.: Impact of topography on black carbon transport to the southern Tibetan Plateau during the pre-monsoon season and its climatic implication, *Atmos. Chem. Phys.*, 20, 5923–5943, <https://doi.org/10.5194/acp-20-5923-2020>, 2020b.
- 655 Zhang, R., Wang, H., Qian, Y., Rasch, P. J., Easter, R. C., Ma, P. L., Singh, B., Huang, J., and Fu, Q.: Quantifying sources, transport, deposition, and radiative forcing of black carbon over the Himalayas and Tibetan Plateau, *Atmos. Chem. Phys.*, 15, 6205–6223, <https://doi.org/10.5194/acp-15-6205-2015>, 2015.
- Zhang, X. H., Xu, J. Z., Kang, S. C., Zhang, Q., and Sun, J. Y.: Chemical characterization and sources of submicron aerosols in the northeastern Qinghai–Tibet Plateau: insights from high-resolution mass spectrometry, *Atmos. Chem. Phys.*, 19, 7897–7911, <https://doi.org/10.5194/acp-19-7897-2019>, 2019.

- 660 Zhang, Y. Z., Forrister, H., Liu, J. M., Dibb, J., Anderson, B., Schwarz, J. P., Perring, A. E., Jimenez, J. L., Campuzano-Jost, P., and Wang, Y. H.: Top-of-atmosphere radiative forcing affected by brown carbon in the upper troposphere, *Nat. Geosci.*, 10, 486–489, <https://doi.org/10.1038/NGEO2960>, 2017.
- 665 Zhao, Z. Z., Cao, J. J., Chow, J. C., Watson, J. G., Chen, L. -W. A., Wang, X. L., Wang, Q. Y., Tian, J., Shen, Z. X., Zhu, C. S., Liu, S. X., Tao, J., Ye, Z. L., Zhang, T., Zhou, J. M., and Tian, R. X.: Multi-wavelength light absorption of black and brown carbon at a high-altitude site on the Southeastern margin of the Tibetan Plateau, China, *Atmos. Environ.*, 212, 54–64, <https://doi.org/10.1016/j.atmosenv.2019.05.035>, 2019.
- Zhong, M. and Jang, M.: Dynamic light absorption of biomass-burning organic carbon photochemically aged under natural sunlight, *Atmos. Chem. Phys.*, 14, 1517–1525, <https://doi.org/10.5194/acp-14-1517-2014>, 2014.
- 670 Zhu, C. S., Qu, Y., Huang, H., Chen, J., Dai, W. T., Huang, R. J., and Cao, J. J.: Black carbon and secondary brown carbon, the dominant light absorption and direct radiative forcing contributors of the atmospheric aerosols over the Tibetan Plateau, *Geophys. Res. Lett.*, 48, e2021GL092524, <https://doi.org/10.1029/2021GL092524>, 2021.

Table 1. Submicron aerosol light absorption coefficient (b_{abs}) contributed by BrC ($b_{\text{abs-BrC}}$) and BC ($b_{\text{abs-BC}}$) at Gaomeigu during the sampling period (March 14th to 31th, 2018).

Parameter* (Mm ⁻¹)	Wavelength					
	370 nm	470 nm	520 nm	590 nm	660 nm	880 nm
b_{abs}	33.1 ± 24.4 (4.7–160.0)**	26.7 ± 19.7 (3.8–138.4)	20.3 ± 13.9 (2.6–93.0)	18.2 ± 12.5 (2.1–84.8)	13.7 ± 9.0 (1.7–56.9)	8.0 ± 4.9 (1.5–28.6)
$b_{\text{abs-BrC}}$	12.3 ± 13.8	10.7 ± 11.5	6.0 ± 6.0	5.8 ± 5.8	2.7 ± 2.6	0.0 ± 0.0
$b_{\text{abs-BC}}$	20.8 ± 12.8	16.0 ± 9.8	14.3 ± 8.8	12.4 ± 7.7	11.0 ± 6.8	8.0 ± 4.9

* $b_{\text{abs-BrC}}$ and $b_{\text{abs-BC}}$ were calculated based on the $\text{AAE}_{\text{BC}} = 1.1$.

** ~~The Variations of the~~ measured hourly b_{abs} from minimum to maximum values.

Figure 1. Topography map of the Tibetan Plateau and the location of the sampling site at Gaomeigu.

Figure 2. Hourly variations in (a) OA mass concentrations and (b) submicron aerosol light absorption coefficients (b_{abs}) at different wavelengths (370, 470, 520, 590, 660 and 880 nm) at Gaomeigu from 14 to 31 March, 2018.

680 **Figure 23.** Light absorption fractions at specific wavelengths contributed by BrC and BC under different absorption Ångström exponent of BC (AAE_{BC}) assumptions. The red, blue, and green lines were the dividing lines between BrC and BC light absorption fractions when $AAE_{\text{BC}} = 1.1, 0.8,$ and $1.4,$ respectively. The grey filled area represents variations in the BrC absorption fraction caused by the uncertainties of $AAE_{\text{BC}} (\pm 0.3).$

685 **Figure 34.** (a) Mass spectra of BBOA and po-OOA. (b) Pearson correlations between mass concentrations of OA components and the tracer molecular fragments. (c) Scatterplots of f_{44} vs. f_{60} for BBOA resolved in this study and reported by previous literatures. (d) Variations of po-OOA mass concentration and its fraction in OA as a function of O_3 . The data are grouped in O_3 bins (10 ppb increment).

690 **Figure 45.** (a) The mass absorption cross section of BBOA and po-OOA (MAC_{BBOA} and $\text{MAC}_{\text{po-OOA}}$, respectively) at five wavelengths ($\lambda = 370, 470, 520, 590,$ and 660 nm). The circle and shaded area represent the mean MAC values and the standard deviations, respectively. The dashed line is power-law fit. (b) Light absorption coefficient of BrC ($b_{\text{abs-BrC}}$) from BBOA and po-OOA and its fraction in the total reconstructed BrC absorption at different wavelengths.

Figure 56. (a) 72-h backward trajectories of Gaomeigu from 8:00 on 14 to 23:00 on 31 March, 2018. (b) and (c) Concentration weighted trajectory (CWT) maps of $b_{\text{abs-BrC}}$ at 370 nm (Mm^{-1}) from BBOA and po-OOA, respectively.

695 **Figure 67.** (a) Simple forcing efficiency (SFE) of light-absorbing particles from 370 to 660 nm and the integrated SFE over the entire solar spectra (370–660 nm). (b) The fraction of solar radiation absorbed by OA components relative to BC. In each panel, the short line and square inside the boxes indicate the median and mean values, respectively. The lower and upper edges of the boxes denote the standard deviation. The vertical bars (“whiskers”) show the 5th and 95th percentiles. Scattered data points and normal distribution curve are also shown.




Figure 1. Topography map of the Tibetan Plateau and the location of the sampling site at Gaomeigu.






Figure 2. Hourly variations in (a) OA mass concentrations and (b) submicron aerosol light absorption coefficients (b_{abs}) at different wavelengths (370, 470, 520, 590, 660 and 880 nm) at Gaomeigu from 14 to 31 March, 2018.



705 **Figure 23.** Light absorption fractions at specific wavelengths contributed by BrC and BC under different absorption Ångström exponent of BC (AAE_{BC}) assumptions. The red, blue, and green lines were the dividing lines between BrC and BC light absorption fractions when $AAE_{BC} = 1.1, 0.8,$ and $1.4,$ respectively. The grey filled area represents variations in the BrC absorption fraction caused by the uncertainties of $AAE_{BC} (\pm 0.3).$



710 **Figure 34.** (a) Mass spectra of BBOA and po-OOA. (b) Pearson correlations between mass concentrations of OA components and the tracer molecular fragments. (c) Scatterplots of f_{44} vs. f_{60} for BBOA resolved in this study and reported by previous literatures. (d) Variations of po-OOA mass concentration and its fraction in OA as a function of O_3 . The data are grouped in O_3 bins (10 ppb increment).




Figure 45. (a) The mass absorption cross section of BBOA and po-OOA (MAC_{BBOA} and $\text{MAC}_{\text{po-OOA}}$, respectively) at five wavelengths ($\lambda = 370, 470, 520, 590,$ and 660 nm). The circle and shaded area represent the mean MAC values and the standard deviations, respectively. The dashed line is power-law fit. (b) Light absorption coefficient of BrC ($b_{\text{abs-BrC}}$) from BBOA and po-OOA and its fraction in the total reconstructed BrC absorption at different wavelengths.






Figure 56. (a) 72-h backward trajectories of Gaomeigu from 8:00 on 14 to 23:00 on 31 March, 2018. (b) and (c) Concentration weighted trajectory (CWT) maps of $b_{\text{abs-BrC}}$ at 370 nm (Mm^{-1}) from BBOA and po-OOA, respectively.





725

Figure 67. (a) Simple forcing efficiency (SFE) of light-absorbing particles from 370 to 660 nm and the integrated SFE over the entire solar spectra (370–660 nm). (b) The fraction of solar radiation absorbed by OA components relative to BC. In each panel, the short line and square inside the boxes indicate the median and mean values, respectively. The lower and upper edges of the boxes denote the standard deviation. The vertical bars (“whiskers”) show the 5th and 95th percentiles. Scattered data points and normal distribution curve are also shown.

730