

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Supplementary information

Yields and molecular composition of gas phase and secondary organic aerosol from the photooxidation of benzyl alcohol: formation of highly oxygenated and hydroxy nitroaromatic compounds

Mohammed Jaoui¹, Kenneth S. Docherty², Michael Lewandowski¹, Tadeusz E. Kleindienst¹

¹Center for Environmental Measurement & Modeling, U.S. Environmental Protection Agency, Research Triangle Park, NC, 27711, USA

²Jacobs Technology, Inc., Research Triangle Park, NC, 27709, USA




Correspondence: Mohammed Jaoui (Jaoui.mohammed@epa.gov)


29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

Section 1. Purity of benzyl alcohol


Benzyl alcohol (BnOH) was purchased at the highest purity commercially available (Sigma aldrich, anhydrous, 99.8%), and kept in the refrigerator when not in use. BnOH oxidizes slowly to one of the main toxic oxidation product benzaldehyde (BnAld) when exposed to air (Urakami et al., 2000). To a lesser extent, additional impurities were reported including benzoic acid, benzyl benzoate and dibenzyl ether (Ferri et al., 2006; Abend et al., 2004). In this study, the purity of BnOH was investigated by GC-MS analysis. A stock solution of BnOH at 345.86 $\mu\text{g/ml}$ was prepared by dissolving 10 μl of standard BnOH in 30 ml acetonitrile. Two working solutions at 6.78 and 3.39 $\mu\text{g/ml}$, respectively, were prepared from the stock solution by diluting 10 and 5 μl in 0.5 ml methanol. The three solutions were analyzed using the same GC-MS parameters reported in the main manuscript for underivatized extracts (2 μl was injected from each solution), with an on-column injected mass of 692, 14 and 7 ng, respectively.

Two peaks associated with BnOH (isomers) were detected at 10.6 (small peak) and 10.7 (large peak) minutes, in addition to a small peak of BnAld eluted at 9.5 min (Figure S1). Figure S1 shows the TIC chromatogram obtained from analysing the stock solution (692 ng on column: highest amount injected). Figure S1 also shows the mass spectra associated with BnOH and BnAld in EI and CI mode. No benzoic acid or other impurities were observed when underivatized sample was analyzed. To check for benzoic acid (BnCOOH) impurity, BSTFA derivatization was conducted using the same procedure described in the main paper. Figure S2 shows chromatograms and mass spectra associated with BnOH and BnCOOH in EI and CI mode as BSTFA derivatives.





54




55

56


Figure S1. GC-MS chromatograms of benzyl alcohol standard, and EI and methane-CI mass spectra of benzylalcohol and benzaldehyde (underivatized).

58


59







60



61





66 **Figure S2.** Artefacts associated with BnOH when exposed to chamber air, sonication, and/or BSTFA derivatization. (a)
67 GC-MS chromatograms originated from BSTFA derivatization of stock solution. (b) GC-MS chromatogram originated
68 from BSTFA derivatization of GP chamber background extract (light off). (c) GC-MS chromatogram originated from
69 underivatized GP chamber background extract (light off). Mass spectra in EI and CH₄-CI associated with BnOH and
70 benzoic acid (BnCOOH).

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89


90

91

92

93

94



96 **Figure S3.** Selected EI mass spectra associated with peaks observed in the BnOH/H₂O₂ and/or BnOH/NO_x underivatized
 97 denuder extracts eluting at retention time of 9.3 min (benzaldehyde), 11.3 min; 12.0 min; and 12.8 min.

98

99

100

101

102

103


104

105

106


107

108




109 **Figure S4.** Selected EI mass spectra associated with peaks observed in the BnOH/H₂O₂ and/or BnOH/NO_x BSTFA
 110 derivatized denuder extracts.

111
 112
 113
 114
 115
 116
 117




118 **Figure S5.** Selected EI mass spectra associated with peaks observed in the BnOH/H₂O₂ and/or BnOH/NO_x BSTFA
 119 derivatized filter extracts.

120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131




132

133



134



135




136

137


138

139

140



144
145
146 **Figure S6.** EI mass spectra associated with BSTFA derivatives of meso-tartaric acid, *L*-tartaric acid, d-tartaric acid
147 standards, 4-nitrocatechol, 2-nitrophenloglycinol (isomer), and a peak eluted at 38.18 min.
148



150

151 **Scheme S1.** Proposed reaction pathways leading to some intermediate products from the reaction of R1-23OO and R2-

152 16OO.

153 **References (SI)**

154 Abend, A. M., Chung, L., Bibart, R. T., Brooks, M., and McCollum, D. G.: Concerning the stability of benzyl alcohol:
155 formation of benzaldehyde dibenzyl acetal under aerobic conditions, *J. Pharm. Biomed. Anal.*, 34, 5, 957-962,
156 doi:10.1016/j.jpba.2003.11.007, 2004.

157
158 Ferri, D., Mondelli, C., Krumeich, F., and Baiker, A.: Discrimination of active palladium sites in catalytic liquid-phase
159 oxidation of benzyl alcohol, *J. Phys. Chem. B.*, 110, 46, 22982-22986, doi:10.1021/jp065779z, 2006.

160
161 Urakami, K., Kobayashi, C., Miyazaki, Y., Nishijima, K., and Yoshimura, Y.: Degradation products generated by
162 sonication of benzyl alcohol, a sample preparation solvent for the determination of residual solvents in pharmaceutical
163 bulks, on capillary gas chromatography, *Chem. Pharm. Bull.*, 48, 1299-1303, 2000.

164

165

166

167