

Indicators of the ozone recovery for selected sites in the Northern Hemisphere mid-latitudes derived from various total column ozone datasets (1980–2020)

Janusz Krzyścin¹

5 ¹Institute of Geophysics, Polish Academy of Sciences, Warsaw, 01-452, Poland

Correspondence to: Janusz Krzyścin (jkrzys@igf.edu.pl)

Abstract. We propose a method to examine the current status of the ozone recovery attributed to ozone-depleting substances (ODS) changes in the stratosphere. The total column ozone (TCO₃) datasets used are based on the ground-based (by the Dobson and/or Brewer spectrophotometer) measurements, satellite observations (from the Solar Backscatter Ultraviolet, SBUV, and Ozone Mapping and Profiler Suite, OMPS, instrument), and output of reanalyses (Multi-Sensor Reanalysis version 2, MSR2, and Modern-Era Retrospective Analysis for Research and Applications, version 2, MERRA2). The TCO₃ time series are calculated for selected sites in the NH mid-latitudes (35°N–60°N), which are station locations with long-term TCO₃ observations archived at the World Ozone and Ultraviolet Radiation Data Centre (WOUDC). The TCO₃ monthly means (1980–2020) are averaged over the April–September period to ~~obtained~~obtain TCO₃ time series for the warm sub-period of the year. Two types of the averaged TCO₃ time series are considered: the original one and ~~that non-proxy time series~~ with removed natural variability by a standard multiple regression model. TCO₃ time series were smoothed by the ~~Locally Weighted Scatterplot Smoother (LOWESS)~~locally weighted scatterplot smoother and the ~~Super Smoother (SS)~~super smoother. The smoothed TCO₃ values in 1980, 1988, 1997, and 2020 are used to build Ozone Recovery Indices (ORIs) in 2020. These are key years in the equivalent effective stratospheric chlorine (EESC) time series ~~for the period 1980–2020~~, i.e., ~~the EESC minimum year when~~the stratosphere was only slightly contaminated by ODS; ~~in 1980, 1988 is~~ the year ~~(before the EESC maximum) within which~~ the EESC value is equal to ~~that its value~~ at the end ~~of the TCO₃ data, the EESC maximum year in the Northern Hemisphere mid-latitudes,~~(2020), and the ~~end of TCO₃ data~~EESC maximum was in 1997 in mid-latitude stratosphere. The first proposed ORI, ORI₁, is the normalized difference between the TCO₃ values in 2020 and 1988. The second one, ORI₂, is the percentage of the recovered TCO₃ in 2020 since the ODS maximum. Following these definitions, the corresponding reference ~~values, ranges from~~ -0.5% to 1% for ORI₁ and ~~51.7~~from 40% to 60% for ORI₂; are obtained ~~from the analyzing a set of possible~~ EESC time series ~~simulated via the Goddard automailer~~. The ozone recovery phases are classified comparing the current ORI values and their uncertainty ranges (by the bootstrapping) with ~~the~~these reference ~~values~~ranges. In the analyzed TCO₃ time series, for specific combinations of datasets, data types, and the smoother used, we find faster (~~for~~ ORI₁ $>0\%$ or ORI₂ $>51.7\%$), ~~above the reference range~~, slower (~~for~~ ORI₁ $<0\%$ or ORI₂ $<51.7\%$) ~~below the reference range~~ recovery in 2020 than that inferred from the EESC change, and a continuation of TCO₃ decline after the EESC peak (ORI₂ $<0\%$). ~~Using the original datasets, the strong~~Strong signal of the slower TCO₃ recovery ~~can be~~is found in ~~five stations: Nashville, Toronto, Hohenpeissenberg, Hradec Kralove, and Belsk. This is also in the series with removed proxy effects for those stations excluding Nashville.~~ A continuation of ozone decline after the turnaround in ODS concentration is found both in the original and non-proxy time series from WOUDC (Toronto), SBUV&OMPS (Toronto, Arosa, Hohenpeissenberg, Uccle, Hradec Kralove, and Belsk), and MERRA2 data (~~Arosa, Hohenpeissenberg, Hradec Kralove, and Belsk~~).

1 Introduction

Unexpected low total column ozone (TCO_3) values observed in the early 1980s over Antarctica alarmed both scientists and public because of anticipated increase of ultraviolet radiation (UVR) reaching the ~~Earth~~Earth's surface (Chubachi, 1984; Farman et al., 1985; Solomon et al., 1986). Widespread threats of thinning of the stratospheric ozone layer and corresponding danger for the ~~Earth~~Earth's environment led to signing the Montreal Protocol (MP) in 1987 to phaseout of the man-made ozone depleting substances (ODS). Overturning of the ODS concentration in the stratosphere (from large increase since the early 1980s to slight decrease beginning in the mid-1990s) was an evident sign of the success of MP and its subsequent amendments. The ODS turnaround in the stratosphere was observed around the middle 1990s in the mid-latitudes and at the beginning of the 2000s in Antarctica. This also triggered numerous studies to reveal the corresponding change point also in TCO_3 trends to support impact of the MP and its later amendments on the ozone layer (e.g., Reinsel et al., 2005; Mäder et al., 2007; Harris et al., 2008; Weber et al., 2018).

Trends in ozone were usually calculated using multiple linear regression (MLR) with a number of proxies to eliminate TCO_3 variations related to dynamical oscillations in the atmosphere, ~~and driven by~~ the solar 11-year activity, and volcanic eruptions. The anthropogenic component of the trend term was usually modelled as independent (disjoint) or dependent (joint) two lines drawn for the periods of increasing and decreasing ODS concentration in the stratosphere (e.g., Weber et al., 2018). There were only a few papers using a non-linear smoothed trend pattern based on dynamical linear model (Laine et al., 2014 ~~and~~ Maillard Barras et al., 2022), Fourier series (Bozhkova et al., 2019), and other smoothers, e.g., locally weighted scatterplot smoothing (LOWESS) by Krzyścin and Rajewska-Więch (2016) and wavelets by Delage et al. (2022).

Coldewey-Egbers et al. (2022) calculated ~~longitude~~latitude/longitude linear TCO_3 trends since 1995 over the entire globe based on the satellite data record. The trends in the NH mid-latitudes were mostly insignificant. In some isolated regions ~~there~~, they found statistically significant trends, e.g., negative in East Europe and positive in the northern part of the North Atlantic. This pattern was linked with the opposite trends in the tropopause height. A continuation of the TCO_3 declining tendency since the ODS turnaround was surprisingly revealed in the NH lower stratosphere (Ball et al., 2018). Rather positive trend was expected due to a strengthening of the Brewer–Dobson circulation that was suggested by chemistry–climate models (e.g., Dietmüller et al. 2021). The negative trends in the NH lower stratosphere might be attributed to enhanced horizontal air mass exchange with the tropical region less abundant with ozone in the lower stratosphere (Thompson et al. 2021). However, other studies using observational data did not reveal such decline in the NH lower stratospheric ozone mostly due to large uncertainty in the satellite observations (e.g., Arosio et al. 2019).

~~NASA~~NOAA proposed the Ozone Depleting Gas Index (ODGI) to keep track on changes in ODS concentration in the mid-latitudes and Antarctica (Montzka et al., 2022). Equivalent effective stratospheric chlorine (EESC) was used as a measure of the stratospheric halogen loading, which is weighted by the ozone destruction potential of each gas depleting stratospheric ozone. ODGI was defined as an indicator of the EESC current decline since its peak that is expressed in percentage of the corresponding decline needed to reach the EESC level in 1980, when ODS concentration in the stratosphere was only slightly affected by man-made substances. The EESC peak year in the mid-latitude stratosphere was found in 1997 being three years later than the ODS peak in the troposphere (Montzka et al., 2022). The corresponding ODGI value in 2020 was 51.7 % providing almost half reduction of EESC necessary to reach its undisturbed level existing in 1980. This also provided an estimate of the recovery time around 2045 in the mid-latitudes if factors affecting ODS changes were the same as those in the period 1997–2020. In addition, the EESC level in 2020 corresponds to ~~that of the~~ 1988 level (Fig. 1).

Following ODGI concept, we propose indices to monitor ozone recovery attributed to the EESC changes using various TCO_3 dataset from ground-based and satellite observations, and two reanalyzes. These indices will be calculated for selected NH mid-latitude sites corresponding to locations of the ground-based stations with long-term time series archived in the World Ozone and Ultraviolet Radiation Data Centre (WOUDC). From the smoothed pattern of long-term TCO_3 changes, we will extract TCO_3 values in key years: 1980, 1988, 1997 and 2020 (Fig. 1). These four values will be used to calculate the

80 proposed ozone recovery indices (ORIs) showing the current stage of the stratospheric ozone recovery. [The Goddard automailer \(http://acdb-ext.gsfc.nasa.gov/Data_services/automailer/index.html\)](http://acdb-ext.gsfc.nasa.gov/Data_services/automailer/index.html) will be implemented to simulate various possible EESC patterns to estimate variability of the EESC peak year and the year when the EESC value is equal to its value in 2020.

85 In Sect. 2, the TCO₃ datasets are presented. In Sect. 3, the data preparation, ORI definition, and a method to obtain the uncertainty range of the ORI estimate are described. Section 4 presents ORI values for selected mid-latitudinal sites for various datasets using [combinationcombinations](#) of the data smoother and data [categoriescategory](#). In Sect. 5, the discussion and conclusions are presented.




Fig.

90 **Figure 1:** The EESC time series with marked key years: the EESC maximum in 1997 and 1988 when the EESC value was the same as in 2020 (the end of total column [ozone data used in the paper](#)) based on the EESC pattern proposed by Montzka et al. (2022).

2 Total column ozone datasets

95 Four TCO₃ data sets are used in the study based on the ground-based and satellite observations as well as from two reanalyses. The TCO₃ time series are calculated for selected sites in the NH mid-latitudes (35°N–60°N) corresponding to locations of stations that archived results of the ground-based observations by the Dobson and/or Brewer spectrophotometer at WOUDC. Arosa and Davos data sets [arewere](#) combined as the ozone monitoring at Arosa was moved to nearby station Davos in 2014. 16 stations with long-term and continuous observations (starting at least before 1980 and ending after 2020) are selected (Table 1). The monthly mean TCO₃ values for these stations are taken from the WOUDC web site.

100 ~~The NASA Merged Ozone Data (MOD) version 8.7 is used for a comparison with the WOUDC data. Overpass subset of MOD provides daily means of ozone content in various stratospheric layers and column ozone over the WOUDC sites including also those listed in Table 1. MOD time series are build using the homogenized spectral measurements of the solar backscattered UV on various satellite platforms Nimbus 4, Nimbus 7, NOAA 9, 11, 14, 16, and 17-19, and Suomi National Polar orbiting Partnership (NPP) (Weber et al., 2022).~~

105 **Table 1:** Selected ground-based total column ozone observing stations in the NH [midlatitudesmid-latitudes](#) with the data record archived in WOUDC.

No.	WOUDC No.	Station	Lat.	Lon.	Elevation
Japan					
1	14	Tateno	36.0°N	140.1°E	31 m
2	12	Sapporo	43.1°N	141.3°E	19 m

North America						
3	106	Nashville	36.3°N	86.6°W	182 m	
4	67	Boulder	40.1°N	105.3°W	1689 m	
5	65	Toronto	43.8°N	79.5°W	198 m	
6	19	Bismarck	46.8°N	100.1°W	511 m	
7	76	Goose Bay	53.3°N	60.4°W	40 m	
8	21	Edmonton	53.6°N	114.1°W	766 m	
9	77	Churchill	58.8°N	94.1°W	35 m	
Europe						
10	35	Arosa	46.8°N	9.7°E	1840 m	
11	99	Hohenpeissenberg	47.8°N	11.0°E	975 m	
12	96	Hradec Kralove	50.2°N	15.9°E	285 m	
13	53	Uccle	50.8°N	4.4°E	100 m	
14	68	Belsk	51.8°N	20.8°E	180 m	
15	165	Oslo	59.9°N	10.7°E	90 m	
16	43	Lerwick	60.0°N	1.2°W	80 m	

The NASA Merged Ozone Data (MOD) version 8.7 is used for a comparison with the WOUDC data. Overpass subset of MOD provides daily means of ozone content in various stratospheric layers and column ozone over the WOUDC sites including also those listed in Table 1. MOD time series are build using the homogenized spectral measurements of the solar backscattered UV on various satellite platforms Nimbus 4, Nimbus 7, NOAA 9, 11, 14, 16, and 17–19, and Suomi National Polar-orbiting Partnership (NPP) (Frith et al., 2014; Weber et al., 2022).

Other two data sets represent the category of reanalyzed data, i.e., global TCO₃ field taken from output of 3-D chemistry and transport model (CTM). Here we examine time series for the selected sites interpolated from the Multi-Sensor Reanalysis, version 2 (MSR2) global data with the 0.5° x 0.5° resolution (van der A et al., 2015) and Modern-Era Retrospective analysis for Research and Applications, version 2 (MERRA2) with the 0.5° (latitude)x 0.625°(longitude) resolution (Wargan et al., 2017). Table 2 present the sources of all data sets used in the study.

Table 2: Source of total column ozone datasets.

Datasets	Time Resolution	Data Start	WEB Page
WOUDC	Day	1926 (Arosa)	https://woudc.org/archive/Summaries/TotalOzone/Daily_Summary/
MOD	Day	1970	https://acd-ext.gsfc.nasa.gov/anonftp/toms/sbuv/MERGED/
MSR2	Month	1979	http://climexp.knmi.nl/select.cgi?field=o3col
MERRA2	Month	1980	https://disc.gsfc.nasa.gov/datasets/M2TMNXCHM_5.12.4/summary

3 Methods/Method

The TCO₃ monthly means from all data sets described in Sect. 2 are averaged over warm sub-period of the year (April–September) to build seasonal TCO₃ time series (1980–2020) for all NH mid-latitude selected sites listed in Table 1. The dynamic variability of ozone in these months is much smaller than in the cold-subperiod of the year, so the variability of TCO₃ is lower and chemical changes are potentially more pronounced. Moreover, ground-based TCO₃ observations by the spectrometers are the most accurate in this part of the year because of due to the Sun's high solar elevations and better weather conditions, allowing AD the most reliable ozone observations by the Dobson instrument with spectrophotometers (Komhyr, 1980).

During the warm sub-period of the year, UVR is much higher comparing to the cold sub-period. UV overexposures having detrimental effect on the Earth ecosystems are most frequent in the warm sub-period of the year. Therefore, analysis of the long-term TCO₃ changes in this period of the year is of special importance to discuss detrimental biological effects of the

130 ozone changes. In the mid-1980s, the anticipated risks of UV overexposure led to international initiatives to protect the ozone layer. In addition, the results based on for the entire year (January–December) and the cold sub-period (October–next year March) data are presented in Appendix A.

3.1 Data smoother

135 Two data smoother types are examined in the study: LOWESS by Cleveland (1979) and super smoother (SS) by Friedman (1984). Here, for LOWESS application, the smoothness level is pre-defined to have up to two turning points in the smoothed time series, i.e., the smoothing parameter f is set equal to 0.5. The first turning point corresponds to the ODS turning point in 1997 and next point is to reveal a change of the recovery rate after 1997 (if it exists). For SS, we use an option with not pre-defined smoothness level. The smoothed curve is obtained by means of a bivariate regression smoother based on local linear regression with adaptive bandwidths. Figure 2 illustrates the performance of both smoothers applied to the time series of seasonal (April–September) TCO₃ means for Tateno and Toronto from the ground-based observations.




Fig.

Figure-2. Examples of the long-term time series of total column ozone from ground-based observations in the warm sub-period (April–September) of the year by locally weighted scatterplot smoother (Lowess–blue curve) and super smoother (red curve): Tateno (a), Toronto (b).

3.2 Removal of the ozone natural variability

155 The smoothed pattern of TCO₃ time series is used to discuss long-term variability in the series comprising 41 yearly values. The smoothers described in Sect. 3.1 are applied to both original time series and the series with removed variability due to various dynamical/chemical processes not directly involved with the anthropogenic emission of chemicals affecting the stratospheric ozone. Various proxies (explanatory variables in MLR regressions) for these “natural” processes have been proposed parameterizing dynamical/chemical forcing on the ozone layer. Frequently in MLR, the solar activity cycle (e.g., 10.7 cm solar radio flux), indices of internal atmospheric fluctuations (Quasi Biennial Oscillations, El-NiñoNiño Southern OscillationOscillations, and Arctic OscillationOscillations), optical depth of the stratospheric aerosols, and the eddy heat flux in the stratosphere (to parameterize the intensity in the Brewer–Dobson circulation) were used. The proxy set proposed by

160 Weber et al. (2022) is used here with the sources listed in their Table 2. The only differences between our and Weber et. (2022) proxy set is using the eddy heat flux at 100-hPa from the NASA Atmospheric Chemistry and Dynamics Laboratory data base (https://acd-ext.gsfc.nasa.gov/Data_services/met/ann_data.html) and the stratospheric aerosol optical depth updated from Sato et al. (1993) for the entire time data (i.e., no different aerosol data sources after 1990).

165 The standard MLR with two independent linear trend terms (before and after the year of ODS turnaround) is used to extract the trend pattern and the combined signal due to all proxies. The MLR used here is identical to the one labeled as full MLR in Weber et. (2022). Finally, the ~~explained part of TCO₃ variability related to the combined time series combining proxy effects/signals~~ is subtracted from the original time series to obtain the TCO₃ series ~~comprising/containing~~ only the anthropogenic long-term component and noise. Further in the text, this time series is called non-proxy times series.

3.3 Ozone Recovery Index

170 The proposed ozone recovery indices (ORIs) follow the ODGI concept (Montzka et al., 2022) of using selected values from the EESC series to define the recovery status. Key TCO₃ values are taken at the beginning (1980), at the end of the data (2020), in the year of TCO₃ trend ~~overturing~~ overturning (1997), and in the year (1988) when the EESC value in the NH mid-latitudes was the same as ~~at~~ in 2020. Last two years are taken according the ~~end-year~~. Namely, smoothed EESC pattern shown in Fig.1. Smoothed TCO₃ values in the selected key years T , denoted as $\langle TCO_3(T) \rangle$, $T=\{1980, 1988, 1997, 2020\}$, are used for calculation of the following dimensionless ORIs in 2020:

$$\begin{aligned}
 & ORI_1(1988, 2020) \\
 & = 100\% \frac{\langle TCO_3(2020) \rangle - \langle TCO_3(1988) \rangle}{\langle TCO_3(1980) \rangle} \frac{\langle TCO_3(2020) \rangle - \langle TCO_3(1988) \rangle}{\langle TCO_3(1980) \rangle} \\
 & ORI_2(2020) = 100\% \frac{\langle TCO_3(2020) \rangle - \langle TCO_3(1997) \rangle}{\langle TCO_3(1980) \rangle - \langle TCO_3(1997) \rangle} (1997, 2020) \\
 & = 100\% \frac{\langle TCO_3(2020) \rangle - \langle TCO_3(1997) \rangle}{\langle TCO_3(1980) \rangle - \langle TCO_3(1997) \rangle} \quad (2)
 \end{aligned}$$

180 If the ozone recovery follows ODS changes, $ORI_1(1988, 2020)$ will be equal to 0 % but negative (positive) for slower (faster) TCO₃ recovery than that found in the EESC pattern. Correspondingly, $ORI_2(1997, 2020)$ value in the Northern Hemisphere will be equal to 48.3 % (i.e. 100 % - ODGI(2020), see ODGI(2020)= 51.7 % in Montzka et al. 2022) if the ozone recovery in the period 1980-2020 follows EESC changes and lower (higher) than this reference value if the ozone recovery is slower (faster) comparing to that ~~found~~ existing in the EESC. ~~If $ORI_2(2020)$ is less than 0 % and the ozone is declining before the ODS overturning year (usual case of the NH mid-latitudinal TCO₃), the ozone depletion will continue throughout the period 1980-2020.~~

185 To calculate $ORI_2(T1)$ in the NH mid-latitudes in any year $T1 (>1997)$, the year $T2 (<1997)$ when $ODS(T2) = ODS(T1)$ needs to be delineated from ODS pattern. In this case, the general formulas for ORIs are as follows:

$$ORI_1(T1) = 100\% \frac{\langle TCO_3(T1) \rangle - \langle TCO_3(T2) \rangle}{\langle TCO_3(1980) \rangle} \quad (3)$$

$$ORI_2(T1) = 100\% \frac{\langle TCO_3(T1) \rangle - \langle TCO_3(1997) \rangle}{\langle TCO_3(1980) \rangle - \langle TCO_3(1997) \rangle} \quad (4)$$

190 The ozone recovery will happen in the year TR when $\langle TCO_3(TR) \rangle = \langle TCO_3(1980) \rangle$. This means that the ozone recovered to its initial level when the stratosphere was not contaminated by man-made compounds (in 1980). In this case, $ORI_2(1997, TR)=100\%$.

If ORI_2 (Negative $ORI_2(T1)$ values will be found when 1997, 2020) is less than 0 % and the ozone decline, which was observed in declining before the period 1980–1997 (i.e., $\langle ODS \text{ overturning year (usual case for the NH mid-latitudinal } TCO_3(1980) \rangle - \langle TCO_3(1997) \rangle$ is always positive), continues), the ozone depletion will continue also after 1997–the EESC overturning. It is worth mentioning that for a searching of the TCO_3 recovery over Antarctica in 2020, the smoothed TCO_3 values in 1980, 1993, 2001, and 2020 need to be selected as these years correspond with key years in the EESC pattern for this region (Montzka et al., 2022).

3.4 ORI reference range

The specific EESC pattern, which is shown in Fig.1, was used in the calculation of ODGI and the year when EESC was equal its value in 2020. Various shapes of the EESC pattern can be provided via the Goddard automailer depending on parameters characterizing ODS in the stratosphere: mean age of air, age of air spectrum width, Bromine scaling factor, and the fractional release type. For mid-latitudes, the default values in the automailer are 3 yr and 60 for the mean age of the air and the Bromine scaling factor, respectively. We examine additional two options, i.e., 3.3 yr & 2.7 yr and 45 & 75, based on the uncertainties (1σ) of these parameters discussed by Velders and Daniel (2014) for mid-latitudes (see their Table 2). Moreover, two options are assumed for age of air spectrum width, i.e., 1.5 yr (default in the automailer) and 3 yr (arbitrarily selected). Moreover, three types of the fractional release type are implemented: Newman et al. (2006), Laube et al. (2010), and default type denoted as WMO 2010. In total, 54 EESC patterns have been obtained. Figure 3a shows the EESC's peak values and the corresponding years at the EESC maxima derived from the automailer together with the pair (1936 ppt and 1997) used in ODGI calculations by Montzka et al. (2022). The range of $100\% - ODGI(2020)$ values (equivalent to $ORI_2(1997, 2020)$ obtained with the corresponding EESC values instead of TCO_3 in Eq. (2)) is from 40 % to 60 % using all simulated EESC curves (Fig. 3b). This gives the reference range for $ORI_2(1997, 2020)$ when the ozone recovery is similar to that of the EESC.




Fig. 3. The EESC characteristics derived via the Goddard automailer (open circles) and those (full circles) taken from Montzka et al. (2022): EESC maximum value versus the year at the peak (a), $ORI_2(1997, 2020)$ value calculated on the basis of the EESC values in Eq. (2) (equivalent to $100\% - ODGI(2020)$) versus the year when EESC value is equal to that in 2020 (b).

The year (before the EESC peak) T when EESC was equal to its value in 2020 changes from 1986 to 1989 (Fig. 3b). The mean total ozone value, $\langle TCO_3(T) \rangle$, for this year can be calculated assuming the ozone linear trend with rate A (in Dobson Unit per year) as almost linear depletion of TCO_3 existed in mid-latitudes in the 1980s (e.g. Hudson et al., 2006):

$$\langle TCO_3(T) \rangle = \langle TCO_3(1988) \rangle + A(T - 1988) \quad (3)$$

For the earliest (1986) and latest year (1989) shown in Fig. 3b using the definition of $T, ORI_1(T, 2020) \stackrel{\text{def}}{=} 0$, we obtain after simple mathematical manipulations with Eq. (1) and Eq. (3) that $ORI_1(1988, 2020)$ equals to $-2 \text{ yr } A'$ and $1 \text{ yr } A'$,

respectively, when $A' = 100 \% A < TCO_3(1980) >^{-1}$ is the ozone trend (in percentage) in the 1980s. Earlier studies estimated A' of a few percent per decade in the NH mid-latitudes (e.g. WMO, 1999; Hudson et al., 2006). Finally, taking A' of -0.5% per year, the reference range for $ORI_1(1988, 2020)$ can be estimated from -0.5% up to 1% .

3.5 Uncertainty of ORI estimates

The original TCO_3 time series ~~in year t , $TCO_3(t)$~~ , $TCO_3(t)$ is divided into two parts to account for the long-term variability, ~~$<TCO_3(t)>$~~ , $< TCO_3(t) >$, which is extracted by the smoother, and the residual parts, $Resid_TCO_3(t)$:

$$TCO_3(t) = < TCO_3(t) > + Resid_TCO_3(t) \quad (5) \quad (4)$$

For each selected WOUDC station and data ~~type~~type (original ~~and/or~~ without proxy effects), two smoothers (LOWESS and SS) are applied to extract the long-term variability and the residual part of TCO_3 variability comprising dynamical/chemical part of variations and noise. The uncertainty range (between the 5th and the 95th percentile) of each ORI estimate is calculated from the set of ordered (lowest to highest) synthetic ORI values derived by bootstrapping. This method of calculations uncertainties of the trend estimates has been applied in our previous studies (e.g., Krzyścin et al., 2015).

First step of bootstrapping is building synthetic n -th time series, ~~$Resid_TCO_3(t)$~~ , $Resid_TCO_3^n(t)$, to mimic the residual term in Eq. (5),

$$Resid_TCO_3^n(t) = Resid_TCO_3(t_x), \quad t_x \in \{1980, \dots, 2020\}, \quad n = \{1, \dots, N\} \quad (65)$$

Random year ~~t_x~~ t_x is obtained by drawing with replacement any year between 1980 and 2020. The total number of the bootstrapped time series used, $N=10000$, is derived experimentally to have stable estimates of ORIs. The use of the drawing with replacement in the construction of ~~analytical~~synthetic residuals was allowed because the original residuals were random, which was confirmed by the Wald-Wolfowitz one sample run test (Wald and Wolfowitz, 1940).

The original and synthetic residuals should be identical in nature, i.e., have almost similar cumulative distribution. The Wald-Wolfowitz two samples run test is applied to check the difference between the original and synthetic residuals. If the time series of residuals derived by the drawing with replacement was found to be significantly different (~~at 5 % significance level with the 95 % confidence~~) from the original ones, such the residuals were not used in the bootstrapping. This means that the number of draws was greater than the total number of the series (N) used in the bootstrap sample.

Next step of the bootstrapping is adding $Resid_TCO_3^n$ term to the smoothed part of the original TCO_3 series to build n -th synthetic ~~TCO_3~~ $TCO_3(t)$ time series:

$$TCO_3^n(t) = < TCO_3(t) > + Resid_TCO_3^n(t) \quad (76)$$

The smoothed n -th time series, $< TCO_3^n(t) >$ is obtained applying LOWESS (or SS) to synthetic $TCO_3^n(t)$ time series. Repeating this procedure, a set of synthetic series, $TCO_3^n(t)$, $n = \{1, \dots, N\}$, is constructed.

Finally, for each $< TCO_3^n(t) >$ series, $ORI_1^n(2020)(1988, 2020)$ and $ORI_2^n(2020)(1997, 2020)$ are calculated using Eq. (1) and Eq. (2), respectively. From the ordered set of $ORI_1^n(2020)(1988, 2020)$ and $ORI_2^n(2020)(1997, 2020)$ values, the uncertainty range (5th percentile–95th percentile) is obtained. This allows to discuss if ORIs values are significantly

different than the reference ORIs values ORI ranges (Sect. 3.4) defining the recovery stage, i.e., -0.5 % to 1.0 % and 51.740 % to 60 % for $ORI_1(2020)(1988, 2020)$ and $ORI_2(1997, 2020)$, respectively.

4 Results

Figures 3–4–5 illustrate the median and uncertainty range of $ORI_1(1988, 2020)$ and $ORI_2(2020)(1997, 2020)$, respectively, for all WOUDC stations listed in Table 1. These values are calculated using TCO₃ monthly mean values averaged over the warm sub-period of the year. Results are shown in Figs. 3–4–5 (a–d) for WOUDC, MOD, MSR2, and MERRA2 data, respectively. The order of the stations on “x” axis in these Figures is the same as in Table 1, i.e., first there are two Japanese stations, then seven North American, and finally seven European stations. For each region, (between the vertical dashed lines), the results are arranged from the southernmost to the northernmost station.





Figure 3

Fig. 4. Median of the ozone recovery index, $ORI_1(1988, 2020)$, estimate (triangle) based on Eq. (1) and its uncertainty range (line) for 16 NH mid-latitude stations (Table 1) and various combination of data smoothers and datasets: WOUDC data (a), MOD data (b), MSR2 (c), and MERRA2 (d). Results in red are for the original data and in blue for the data with removed natural variability. The hatched area marks the reference range, -0.5 % to 1.0 %, calculated in Sect. 3.4.

The location of the uncertainty range of ORIs in Figs. 3–4–5 relative to the horizontal reference lines range defined in Sect. 3.4 provides the stage of the ozone recovery at each selected station. Namely, if the uncertainty range of ORI is entirely above (below) the horizontal reference lines, i.e., 0 % for $ORI_1(2020)$ and 51.7 % for $ORI_2(2020)$, range, the ozone recovery is faster (slower) than that found for existing in the EESC change in for the period 1980–2020. This means that the TCO₃ recovery rate is significantly different (at least at the 95 % confidence level) when compared with corresponding the EESC values. If the uncertainty range crosses the horizontal reference line range, the hypothesis of the TCO₃ recovery rate similar to

that in following the EESC change changes cannot be rejected. The depletion of TCO_3 will continue even after the EESC maximum if the uncertainty range for $\text{ORI}_2(2020)(1997, 2020)$ is completely below 0% reference line (dashed line in



% Fig. 4). In this case, there is no TCO_3 recovery between for the entire period (1980 and 2020).

Figure 4. The same as Fig. 3 but for the ozone recovery index, $\text{ORI}_2(2020)$, based on Eq. (2).

The ozone recovery is discussed both for the original data (results in red in Figs. 3–4) and for the data non-proxy time series (with removed combined proxy signal (results in blue). Further in the text, the later data type will be called as the non-proxy data from the original data). A strong signal that the TCO_3 recovery at a selected station is slower (faster) than that found in the EESC pattern if the uncertainty ranges of both ORIs are below (above) the reference lines (i.e., 0% for Figs. 3 and 51.7% for Figs. 4) ranges for all data types used (WOUDC, MOD, MSR2, and MERRA2), regardless of the smoother type used.

For strong support for the original times series, the strong signal of slower TCO_3 recovery can be found for five three European stations: Nashville, Toronto, Hohenpeissenberg, Hradec Kralove, and Belsk. This is also in the non-proxy series for those stations excluding Nashville as the $\text{ORI}_1(2020)$ uncertainty range crosses the zero line when SS is used to smooth WOUDC data (For Fig. 3a). In Arosa and Uccle, using $\text{ORI}_2(1997, 2020)$, the slower ozone recovery is found in all datasets except WOUDC. This may suggest a problem with the homogeneity of WOUDC data at least for Arosa, as the results are different than those obtained at the nearby Hohenpeissenberg station. The slower TCO_3 recovery from the original data is identified in Oslo using all datasets except WOUDC. For Toronto and Nashville, using $\text{ORI}_2(1997, 2020)$ derived from the non-proxy time series, the slower ozone recovery can be suggested.




Fig. 5. The same as Fig. The TCO₃ recovery faster than in the ESSC pattern only occurs at one station (Bismark). It can be found using $ORI_1(2020)$ from all original datasets except MOD. In addition, there is a faster recovery for the ozone recovery index, $ORI_2(1997, 2020)$, based on Eq. (2). The reference range (hatched area) is between 40 % and 60 % according estimates in Sect. 3.4.

There are many individual cases where the localization of the uncertainty range of ORI estimate suggests significant difference between the TCO₃ and EESC recovery rates in at least one or two datasets. For example, this is the case for Sapporo/Nashville for both ORIs derived from MOD data (Fig. 3b4b and Fig. 4b5b). The differences between ORIs values derived from the original and non-proxy series sometimes appear, e.g., both ORIs for Goose based on the original data show the non-proxy time series for Oslo shows slower recovery but the non-proxy data supports this only if using $ORI_2(2020)(1997, 2020)$ for all data types excluding WOUDC but this is obtained from MOD (Fig. 4b) not found in the original time series.

A consistent pattern of the ORIs variability is found for all data records. Lower ORIs for Nashville and Toronto were calculated among the North American stations. Moreover, ORIs below the reference range appeared in Europe (except two northernmost stations). This was not found in Arosa and Uccle using ground-based data probably due to instrumental problems. ORIs values indicated that the TCO₃ recovery in Japan follows the EESC change. For this region, slower recovery was found only in the MOD original data for Sapporo using $ORI_2(1997, 2020)$. Usually, uncertainty of ORIs estimates in MOD time series were the smallest that allows to identify additional sites with slower recovery (e.g. Nashville for the original and non-proxy time series, Fig. 5b). In all cases, the TCO₃ recovery was never faster than that in the EESC pattern.

A continuation of ozone decline after the turnaround in ODS concentration, which appears when the uncertainty range of $ORI_2(2020)(1997, 2020)$ is entirely below 0 regardless of the smoother type applied, is found both in the original and non-proxy time series from WOUDC (Toronto, see corresponding also Fig. 2b for the TCO₃ time series in Fig. 2b), MOD (Toronto, Arosa, Hohenpeissenberg, Uccle, Hradec Kralove, and Belsk), and MERRA2 data (Arosa, Hohenpeissenberg, Hradec Kralove, and Belsk).

5 Discussion and conclusions

We proposed a novel tool to examine the stage of ozone recovery attributed to ~~EESC changes-the EESC's change~~. We introduce two ORIs. The first ORI is a difference between the TCO₃ value in 2020 and that in the year before the EESC maximum with the same EESC value as that in 2020. The second ORI is the percentage of the recovered ozone since the EESC maximum. The following ozone recovery phases can be identified when compared with the EESC change, i.e., faster, slower, ~~nonconclusive~~ no conclusive (the hypothesis of the TCO₃ recovery driven entirely by EESC change cannot be rejected), and a continuation of TCO₃ decline after the EESC peak. For NH mid-latitudes in 2020, first three categories are found from the location of the ORI uncertainty ranges relative to the reference ~~lines, ranges, from -0.5 % to 1 % for~~ ORI₁(2020)=0%(1988, 2020) or from 40 % to 60 % for ORI₂(1997, 2020)=51.7 %, which. The reference ranges were obtained from simulations of the 1980-2020 mid-latitude EESC time series-via the Goddard automailer. The last category appears when the uncertainty range of ORI₂ is completely below its 0 % reference line.

The stage of ozone recovery level was usually discussed comparing linear trend values before and after the year of the EESC overturning in the mid-1990s for the NH ~~midlatitudes-mid-latitudes~~. The trends were obtained using MLR applied to various TCO₃ datasets and the slopes of the trend lines, which can be joint (Reinsel et al., 2005) or disjoint (Weber et al., 2018) at the EESC turning year, were compared. Weber et al. (2022) found that for the increasing rate of near global ozone (60°S–60°N) after 1995 was roughly a third of the decreasing rate calculated in the period 1978–1995 that corresponds with the ratio of the EESC linear change after and before the EESC maximum. This ~~supports~~ supported the success of MP and its further amendments. For this case, corresponding ORI₂(2020) of ~47 % ~~will be roughly~~ (i.e., inside the reference range) is estimated from the ratio between the TCO₃ trends (1/3) multiplied by the ratio between duration of the period after (24 yr→) and before (17 yr→) the EESC maximum. This estimate is close to the corresponding 100 %–ODGI(2020) value in-equal to 48.3 % when ODGI(2020-of)=51.7% calculated for the NH mid-latitudes by % is taken according to Montzka et al. (2022) calculations (see ODGI(2020) in their Table 2). Then, two approaches, which are based on the TCO₃ trends and ORI₂(2020), are almost equivalent and provide the TCO₃ recovery ~~following~~ corresponding with the EESC changes-change.

In our approach, to disclose phase of the ozone recovery, we compare ORIs, which are derived from the smoothed pattern of the ozone time series, with corresponding indicators of the mid-latitude EESC recovery. ORIs are calculated for each selected mid-latitude WOUDC station with the long time TCO₃ series. 16 stations are selected and various datasets for these stations based on the ground-based measurements, satellite overpasses, and two reanalyzes, are examined. For almost all stations and data types, the ozone recovery follows EESC pattern. Only in few cases it is slower. The firm sign of the slower ozone recovery is identified for a group of neighborhood central/western Europe sites (Hohenpeissenberg, Hradec Kralove, and Belsk) ~~and for Toronto~~ as all ORI uncertainty ranges, regardless of the smoother and data type, are below the reference ~~values-obtained~~ ORI ranges. This means that TCO₃ trends are less than expected from the EESC change. The smallest uncertainty ranges of ORIs estimates are for the MOD dataset that result in the ~~largest~~ larger number of sites with slower TCO₃ recovery ~~comparing with the EESC change~~.

A negative ORI₂(1997, 2020) value means that the mean TCO₃ level in 2020 is lower than that in 1997 as the denominator in Eq. (2) is positive because ~~the~~ TCO₃ usually declined ~~was declining~~ in NH mid-latitudes ~~in~~ before the ~~period-1980-1997~~ EESC overturning. When the uncertainty range of ORI₂(1997, 2020) is completely below zero, this means that the decline in ozone continues even after the EESC peak. This case is found for the group of central/western Europe stations from MOD and MERRA2 data. The continuation of ozone depletion in this region was previously discussed by Coldewey-Egbers et al. (2022) using merged satellite data.

Negative trends in TCO₃ since the EESC peak provides lower TCO₃ values at the end of time series (2020) and also negative ORIs thereafter. Szelag et al. (2020) reveals negative trends in the ozone vertical profile ranging from -1 % per decade to -2 %

per decade in the lower/middle stratosphere at 30–60° N for the period 2000–2018 during the summer (June–July–August). Moreover, according to WMO (2022), declining tendency in the tropospheric ozone due to air quality improvement in some regions can provide additional source for TCO₃ depletion for sites where the troposphere was cleaned from the ozone precursors. This is probable for the central/western European sites with $ORI_2(1997, 2020) < 0\%$. Superposition of the negative ozone trend in the lower/middle stratosphere over NH mid-latitudes in summer and the troposphere cleaning supports possibility of negative $ORI_2(1997, 2020)$ value that also means no ozone recovery for the entire period (1980–2020).

Recent studies have focused on anthropogenic trends extracted by MLR to assess changes in the ozone layer by halogens, which implies parameterization of both trend and the natural ozone variability. In this study, two data types, the original and with the combined proxy effects removed, are under interests. The former is used to quantify changes in the UV radiation reaching the Earth's surface caused by ozone and the latter one is to delineate anthropogenic changes (due to man-made halogens) in the ozone layer. ~~We only~~In the main text, we examine the TCO₃ means for the warm period of the year, since UVR levels are naturally high during this period, ~~and if~~. If excessive UV exposure occurs due to lower TCO₃ values, it will cause serious consequences for health and the environment (Barnes et al., 2019). ORIs from the original and non-proxy datasets can sometimes be different, e.g., $ORI_2(1997, 2020)$ based on the ~~Belsk's original non-proxy~~ MSR2 data for Belsk and Hradec Kralove shows a continuation of ozone decline, but the slower recovery stage only comes from the ~~non-proxy~~original data (Fig. 4e5c).

~~The declining TCO₃ tendency in the original data (MOD and MERRA2) is revealed for the group of the central/western European stations for the entire period 1980–2020 using the original MOD and MERRA 2 dataset. In addition, the lower TCO₃ recovery rate appears in all datasets for this region. The ozone issue, which was raised in the early 1980s due to the anticipated UVR increase due to man-made halogens impact on the ozone layer, is still worth considering.~~

Similar analysis for the monthly mean TCO₃ data averaged of the entire year (January–December, Figs. A1–A2) and the cold sub-period of the year (October–next year March, , Figs. A3–A4) are shown in Appendix A. For only a few cases, a faster TCO₃ recovery rate in the period 1980–2020 than that in the EESC series is found using the ground-based data for the cold sub-period of the year. This was identified in Arosa, Uccle, and Oslo when using $ORI_1(1988, 2020)$ after removal proxy effects from the original data (Fig. A3a). In Oslo, the faster TCO₃ recovery is also found in the original WOUDC data. Uccle is only one station where $ORI_2(1997, 2020)$ is above the reference range using the non-proxy TCO₃ time series (Fig. A4a) that suggests a faster TCO₃ recovery. These results for the cold-sub-period of the year should be treated with caution as the ground-based TCO₃ observations in this part of the year are usually less precise than those taken in the warm sub-period because of weather conditions (many cloudy days) and low solar elevation near (Komhyr, 1980). In the cold sub-period, slower TCO₃ recovery than that in the EESC time series was found only in Sapporo (Fig. A4b and Fig. A4d). Using the yearly mean TCO₃, slower recovery was revealed at fewer sites compared to data for the warm sub-period. In addition, this was also identified in Sapporo (Fig. A2b and Fig. A2d). However, a continuation of the ozone decline after the EESC peak for the European stations cannot be revealed. The differences between the European stations and other stations were not as pronounced as in the case of data from the warm-subperiod.

The variability of the ozone layer caused by the gradual removal of long-lived halogens from the stratosphere and changes in dynamic processes in the atmosphere, which are to some extent related to climate change, continue to pose a threat to the environment and health. The ozone recovery could be delayed (if it ever happens) in some isolated areas. The ozone issue, which was raised in the early 1980s due to the anticipated UVR increase is still worth considering. This study is a kind of an introduction to use ORIs, which are based on the smoothed TCO₃ values in key EESC years, to monitor the current state of the ozone layer and its relationship with changes in man-made halogen loading in the stratosphere. We plan to use the ORI concept to discuss the ozone recovery globally from various gridded datasets.

Appendix A

These supporting figures provide further insight into ORIs performances for 16 mid-latitudinal stations using averaged monthly mean TCO_3 values over the entire year (January–December) and cold sub-period of the year (October– next year March). The same data and smoother type were examined as those for ORIs based on the TCO_3 values averaged over the warm sub-period of the year (Figs. 4–5).




Fig. A1. The same as Fig. 4 but for the ozone recovery index, $\text{ORI}_1(1997, 2020)$, based on the TCO_3 monthly means averaged over the entire year (January--December).




Fig. A2. The same as Fig. A1 but for the ozone recovery index, ORI_2 (1997, 2020).




Fig. A3 The same as Fig. A1 but for $ORI_1(1997, 2020)$ based on the TCO_3 monthly means averaged over the cold sub-period of the year (October—next year March).




Fig. A4. The same as Fig. A3 but for the ozone recovery index, $ORI_2(1997, 2020)$.

Data availability. The TCO_3 data used in this study are taken from the WEB sources, listed in Table 2, with non-limited access. “Full model” version of MLR by Weber et al. (2022) is applied to remove “natural” variability from the original TCO_3 data. The MLR proxy set is taken from the WEB sources listed in their Table 2.

Author contributions. JK contributed to all aspects of the data analysis, manuscript preparation, and interpretation of the results.

Competing interests. The author declares that he has no conflict of interest.

Acknowledgments. We thank the Chief Inspectorate of Environmental Protection, Poland, for funding the ozone measurements at Belsk.

Funding. This work was supported by the Ministry of Science and Higher Education of Poland under Grant number 3841/E-41/S/2022.

References:

Arosio, C., Rozanov, A., Malinina, E., Weber, M., and Burrows, J.P.: Merging of ozone profiles from SCIAMACHY, OMPS and SAGE II observations to study stratospheric ozone changes, *Atmos. Meas. Tech.*, 12, 2423–2444, <https://doi.org/10.5194/amt-12-2423-2019>, 2019.

- Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R., and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmos. Chem. Phys.*, 18, 1379–1394, <https://doi.org/10.5194/aep18acp-18-1379-2018>, 2018.
- 440 Barnes, P.W., Williamson, C.E., Lucas, R.M., Robinson, S.A., Madronich, S., Paul, N.D., Janet F. Bornman, J.F., Bais, A.F., Sulzberger, B., Wilson, S.R., Andrady, A.L., McKenzie, R.L., Neale, P.J., Austin, A.T., Bernhard, G.H., Solomon, K.R., Neale, R.E., Young, P.J., Norval, M., Rhodes, L.E., Hylander, S., Rose, K.C., Longstreth, J., Aucamp, P.J., Ballaré, C.L., Cory, R.M., Flint, S.D., de Gruijl, Häder, D.P., Heikkilä, A.M., Jansen, M.A., Pandey, K.K., Robson, T.M., Sinclair, C.A., Wängberg, S., Worrest, R.C., Seyhan Yazar, S., Antony R. Young, A.R., and Richard G. Zepp, R.G.: Ozone depletion, ultraviolet radiation, climate change and prospects for a sustainable future, *Nat Sustain.*, 2, 569–579, <https://doi.org/10.1038/s41893-019-0314-2>, 2019.
- 445 Bozhkova, V., Liudchik, A., and Umreiko, S.: Long-term trends of total ozone content over mid-latitudes of the Northern Hemisphere, *Int. J. Remote Sens.*, 40, 5216–5229, <https://doi.org/10.1080/01431161.2019.1579384>, 2019.
- ~~Chehade, W., Weber, M., and Burrows, J. P.: Total ozone trends and variability during 1979–2012 from merged data sets of various satellites, *Atmos. Chem. Phys.*, 14, 7059–7074, <https://doi.org/10.5194/acp-14-7059-2014>, 2014.~~
- 450 Chubachi, S.: Preliminary results of ozone observations at Syowa Station from February 1982 to January 1983, in: Proc. Sixth Symposium on Polar Meteorology and Glaciology, edited by: Kusunoki, K., vol. 34 of Mem. National Institute of Polar Research Special Issue, 13–19, 1984.
- Cleveland, W. S.: Robust locally weighted regression and smoothing scatterplots. *J. Am. Stat. Assoc.*, 74, 829–836, <https://doi.org/10.2307/2286407>, 1979.
- 455 Coldewey-Egbers, M., Loyola, D. G., Lerot, C., and Van Roozendael, M.: Global, regional and seasonal analysis of total ozone trends derived from the 1995–2020 GTO-ECV climate data record, *Atmos. Chem. Phys.*, 22, 6861–6878, <https://doi.org/10.5194/acp-22-6861-2022>, 2022.
- Delage, O., Portafaix, T., Bencherif, H., Bourdier, A., and Lagracie, E.: The Empirical Adaptive Wavelet Decomposition (EAWD): An adaptive decomposition for the variability analysis of observation time series in atmospheric science, *Nonlin. Processes Geophys.*, 29, 265–277, <https://doi.org/10.5194/npg-29-265-2022>, 2022.
- 460 Dietmüller, S., Garny, H., Eichinger, R., and Ball, W. T.: Analysis of recent lower-stratospheric ozone trends in chemistry climate models, *Atmos. Chem. Phys.*, 21, 6811–6837, <https://doi.org/10.5194/acp-21-6811-2021>, 2021.
- Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica reveal seasonal ClO_x / NO_x interaction, *Nature*, 315, 207–210, <https://doi.org/10.1038/315207a0>, 1985.
- 465 Friedman, J. H. 1984. A variable span smoother. Laboratory for Computational Statistics, Department of Statistics, Stanford University: Technical Report, SLAC-PUB-3477, <https://doi.org/10.2172/1447470>, 1984.
- ~~Frith, S.M., Kramarova, N.A., Stolarski, R.S., McPeters, R.D., Bhartia, P.K., and Labow, G.J.: Recent changes in total column ozone based on the SBUV Version 8.6 Merged Ozone Data Set, *J. Geophys. Res. Atmos.*, 119, 9735–9751, <https://doi.org/10.1002/2014JD021889>, 2014.~~
- 470 Harris, N. R. P., Kyrö, E., Staehelin, J., Brunner, D., Andersen, S.- B., Godin-Beekmann, S., Dhomse, S., Hadjinicolaou, P., Hansen, G., Isaksen, I., Jrrar, A., Karpetchko, A., Kivi, R., Knudsen, B., Krizan, P., Lastovicka, J., Maeder, J., Orsolini, Y., Pyle, J. A., Rex, M., Vanicek, K., Weber, M., Wohltmann, I., Zanis, P., and Zerefos, C.: Ozone trends at northern mid- and high latitudes – a European perspective, *Ann. Geophys.*, 26, 1207–1220, <https://doi.org/10.5194/angeo-26-1207-2008>, 2008.
- 475 ~~Hudson, R. D., Andrade, M. F., Follette, M. B., and Frolov, A. D.: The total ozone field separated into meteorological regimes – Part II: Northern Hemisphere mid-latitude total ozone trends, *Atmos. Chem. Phys.*, 6, 5183–5191, <https://doi.org/10.5194/acp-6-5183-2006>, 2006.~~

Komhyr, W.: Dobson spectrophotometer systematic total ozone measurement error, *Geophys. Res. Lett.*, 7(2), 161–163, 1980.

<https://doi.org/10.1029/GL007i002p00161>, 1980.

Krzyścin, J.W., Rajewska-Więch, B. and Pawlak, H.: Long-Term Ozone Changes Over the Northern Hemisphere Mid-Latitudes for the 1979–2012 Period, *Atmosphere-Ocean*, 53:1, 153–160, <https://doi.org/10.1080/07055900.2014.990869>, 2015.

Krzyścin, J., Rajewska-Więch, B.: Specific variability of total ozone over Central Europe during spring and summer in the period 1979–2014. *International Journal of Climatology, Int. J. Climatol.*, 36, 3539–3549, <https://doi.org/10.1002/joc.4574>, 2016.

Laine, M., Latva-Pukkila, N., and Kyrölä, E.: Analysing time-varying trends in stratospheric ozone time series using the state space approach, *Atmos. Chem. Phys.*, 14, 9707–9725, <https://doi.org/10.5194/acp-14-9707-2014>, 2014.

[Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Sturges, W. T., Braß, M., and Röckmann, T.: Fractional release factors of long-lived halogenated organic compounds in the tropical stratosphere. *Atmos. Chem. Phys.*, 10, 1093–1103, <https://doi.org/10.5194/acp-10-1093-2010>, 2010.](https://doi.org/10.5194/acp-10-1093-2010)

Mäder, J. A., Staehelin, J., Brunner, D., Stahel, W. A., Wohltmann, I., and Peter, T.: Statistical modeling of total ozone: Selection of appropriate explanatory variables, *J. Geophys. Res.*, 112, D11108, <https://doi.org/10.1029/2006JD007694>, 2007.

Maillard Barras, E., Haeferle, A., Stübi, R., Jouberton, A., Schill, H., Petropavlovskikh, I., Miyagawa, K., Stanek, M., and Froidevaux, L.: ~~DLMD~~[Dynamical linear modeling](https://doi.org/10.5194/acp-22-14283-2022) estimates of long-term ~~Ozone~~ trends from ~~homogenized~~ Dobson ~~and Brewer~~ Umkehr profiles ~~at Arosa/Davos, Switzerland~~, *Atmos. Chem. Phys. Discuss.*, 22, 14283–14302, <https://doi.org/10.5194/acp-22-14283-2022>, in review, 2022.

Montzka, S.A., Dutton, G., and Vimont, I.: The NOAA Ozone Depleting Gas Index: Guiding Recovery of the Ozone Layer, <https://gml.noaa.gov/odgi/>, 2022. (last accessed ~~15 September 2022~~ 19 January 2023)

[Newman, P. A., Nash, E. R., Kawa, S. R., Montzka, S. A., and Schauffler, S. M.: When will the Antarctic ozone hole recover?, *Geophys. Res. Lett.*, 33, L12814, doi:10.1029/2005GL025232, 2006.](https://doi.org/10.1029/2005GL025232)

Reinsel, G. C., Miller, A. J., Weatherhead, E. C., Flynn, L. E., Nagatani, R. M., Tiao, G. C., and Wuebbles, D. J.: Trend analysis of total ozone data for turnaround and dynamical contributions, *J. Geophys. Res.*, 110, D16306, <https://doi.org/10.1029/2004JD004662>, 2005.

Sato, M., Hansen, J.E., McCormick, M.P. and Pollack, J.B.: Stratospheric aerosol optical depths, 1850–1990. *J. Geophys. Res.*, 98, 22987–22994, <https://doi.org/10.1029/93JD02553>, 1993.

Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, *Nature*, 321, 755–758, <https://doi.org/10.1038/321755a0>, 1986.

[Szeląg, M. E., Sofieva, V. F., Degenstein, D., Roth, C., Davis, S., and Froidevaux, L.: Seasonal stratospheric ozone trends over 2000–2018 derived from several merged data sets, *Atmos. Chem. Phys.*, 20, 7035–7047, <https://doi.org/10.5194/acp-20-7035-2020>, 2020.](https://doi.org/10.5194/acp-20-7035-2020)

Thompson, A. M., Stauffer, R. M., Wargan, K., Witte, J. C., Kollonige, D. E., and Ziemke, J. R.: Regional and seasonal trends in tropical ozone from SHADOZ profiles: Reference for models and satellite products, *J. Geophys. Res.-Atmos.*, 126, e2021JD034691, <https://doi.org/10.1029/2021JD034691>, 2021.

van der A, R. J., Allaart, M. A. F., and Eskes, H. J.: Extended and refined multi sensor reanalysis of total ozone for the period 1970–2012, *Atmos. Meas. Tech.*, 8, 3021–3035, <https://doi.org/10.5194/amt-8-3021-2015>, 2015.

[Velders, G. J. M., and Daniel, J. S.: Uncertainty analysis of projections of ozone-depleting substances: mixing ratios, EESC, ODPs, and GWPs, *Atmos. Chem. Phys.*, 14, 2757–2776, <https://doi.org/10.5194/acp-14-2757-2014>, 2014.](https://doi.org/10.5194/acp-14-2757-2014)

- 520 Wald, A., and Wolfowitz, J.: On a test whether two samples are from the same population, *Ann. Math. Stat.*, 11, 147–162,
1940. <https://doi.org/10.1214/aoms/1177731909>, 1940.
- Wargan, K., Labow, G.J., Frith, S.M., Pawson, S., Livesey, N.J., and Partyka, G.S.: Evaluation of the Ozone Fields in NASA's
MERRA-2 Reanalysis. *J. Clim.*, 30, 2961–2988. <https://doi.org/10.1175/JCLI-D-16-0699.1>, 2017.
- 525 Weber, M., Coldewey-Egbers, M., Fioletov, V. E., Frith, S. M., Wild, J. D., Burrows, J. P., Long, C. S., and Loyola, D.:
Total ozone trends from 1979 to 2016 derived from five merged observational datasets – the emergence into ozone
recovery, *Atmos. Chem. Phys.*, 18, 2097–2117, <https://doi.org/10.5194/acp-18-2097-2018>, 2018.
- Weber, M., Arosio, C., Coldewey-Egbers, M., Fioletov, V. E., Frith, S. M., Wild, J. D., Tourpali, K., Burrows, J. P., and
Loyola, D.: Global total ozone recovery trends attributed to ozone-depleting substance (ODS) changes derived from five
merged ozone datasets, *Atmos. Chem. Phys.*, 22, 6843–6859, <https://doi.org/10.5194/acp-22-6843-2022>, 2022.
- 530 [WMO: Scientific Assessment of Ozone Depletion: 1998, 496 pp., Global Ozone Research and Monitoring Project - Rep. No. 44, World Meteorological Organization, Geneva, Switzerland, 1999.](#)
- [WMO: Executive Summary. Scientific Assessment of Ozone Depletion: 2022, 56 pp., GAW Report No. 278, World Meteorological Organization, Geneva, Switzerland, 2022.](#)