



Technical note: Sublimation of frozen CsCl solutions in ESEM: determining the number and size of salt particles relevant to sea-salt aerosols

5 Ľubica Vetráková¹, Vilém Neděla¹, Kamila Závacká¹, Xin Yang², Dominik Heger³

¹Environmental Electron Microscopy Group, Institute of Scientific Instruments of the Czech Academy of Sciences, Brno, Czech Republic

²British Antarctic Survey, Natural Environment Research Council, Cambridge, UK

³Department of Chemistry, Faculty of Science, Masaryk University, Brno, Czech Republic

10 *Correspondence to:* Ľubica Vetráková (vetrakova@isibrno.cz) and Dominik Heger (hegerd@chemi.muni.cz)

Abstract. Here we present a novel technique that enlightens the mechanism of formation of small aerosolizable salt particles from salty frozen samples. We demonstrated that CsCl may be a suitable probe for the sea salt due to their similar subzero properties and sublimation outcomes: Using CsCl substantially increased the visibility of the salt both during and after ice sublimation. Hence, we identified the factors that, during the sublimation of a frozen salty solution, are important in generating fine salt particles as a possible source of salt aerosol. The number, size, and structure of the particles that remain after ice sublimation were investigated with respect to the concentration of the salt in the sample, the freezing method, and the sublimation temperature. The last-named aspect is evidently of primary importance for the preference of fine salt crystals over a large compact piece of salt: We showed that the formation of the small salt particles is generally restricted if the brine is liquid during the ice sublimation, i.e., at temperatures higher than the eutectic temperature (T_{eu}). Small salt particles that might be a source of atmospheric aerosols were formed predominantly at the temperatures below the T_{eu} , and their structures strongly depended on the concentration of the salt. For example, the sublimation of those samples that exhibited less than 8 psu (0.05 M) often produced small aerosolizable isolated particles readily able to be windblown. Conversely, the sublimation of 78 psu (0.5 M) samples led to the formation of relatively stable and largely interconnected salt structures. Our findings are in a good agreement with other laboratory studies unsuccessfully seeking for salt aerosols, e.g., from the frost flowers, at temperatures above the T_{eu} . This study offers an explanation of this previously unexplained behaviour.

1 Introduction

Periods of low tropospheric ozone concentrations (e.g., < 10 ppbv) in springtime were repeatedly observed in the polar tropospheric boundary layer (Barrie et al., 1988; Bottenheim et al., 2002; Richter et al., 1998). These ozone depletion events are attributed to the inorganic halogens in the polar atmosphere (Abbatt et al., 2012; Richter et al., 1998). Bromine radicals destroy ozone very efficiently; moreover, the number of reactive gaseous bromine can increase markedly during “bromine



explosions”, in which every Br atom of a gaseous HOBr molecule can cause an extra bromide release from saline sea salt aerosol or other salty particles (Kaleschke et al., 2004). Thus, all saline crystals in sea ice zones are potential sources of reactive bromine: the proposed candidates include frost flowers (Kaleschke et al., 2004), first-year sea ice (Simpson et al., 2007), sea salt aerosol (SSA) from blowing snow (Frey et al., 2020; Yang et al., 2008, 2019), SSA from open leads (Peterson et al., 2017), and the photochemistry within snowpack (Pratt et al., 2013). However, the exact sources of bromine in polar spring are still disputed (Abbatt et al., 2012).

Chemical analyses of SSAs in inland Antarctica show that the aerosols are sulphate-depleted relative to sodium, which points to a sea-ice source of the SSA (Jourdain et al., 2008; Legrand et al., 2017; Rankin and Wolff, 2003; Wagenbach et al., 1998). The suggested mechanism of the sulphate-depletion of the SSA is mirabilite ($\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$) precipitation from brine below -6.4°C (Butler et al., 2016), a fractionation not plausible in sea spray particles generated directly from open oceans. The windblown snow particles of relatively low salinity were suggested to be an efficient source of the SSA (Yang et al., 2008, 2019). The proposed blowing snow mechanism relating to SSA production was supported by the Antarctic Weddell Sea data (Frey et al., 2020). Previously, highly saline frost flowers on thin sea ice were also regarded as a significant source of the SSA (Kaleschke et al., 2004; Rankin and Wolff, 2003; Wolff et al., 2003); this was contradicted by both the wind tunnel experiment (Roscoe et al., 2011) and the *in situ* sublimation of frost flowers in an environmental scanning microscope (ESEM) (Yang et al., 2017). However, these two laboratory studies were performed in a relatively warm temperature range, between -5 to -18°C ; these values are above the eutectic temperature, $T_{eu} = -21^\circ\text{C}$, of a NaCl solution (Rodebush, 1918). Apart from our recent study that complements this paper (Závacká et al., 2022), we are not aware of any laboratory research on the formation of the SSA at temperatures below T_{eu} , which often occur in polar regions in winter and early spring; moreover, most of the sea ice is covered by saline ice crystals, including either high salinity formations such as frost flowers and the basal part of a snowpack or low salinity ices comprising the snow at the surface layer of a snowpack and on thick multiyear sea ice. The investigation of the impact exerted by the temperature and saline concentration on SSA formation is critical in understanding the implications to polar atmospheric chemistry and climate. This is because SSAs not only are a reservoir of various chemical compounds but also function as cloud condensation nuclei (e.g. O’Dowd et al., 1999) or even ice nucleating particles (DeMott et al., 2016; Wise et al., 2012).

In this study, we utilize an ESEM to visualize the process of ice sublimation *in-situ* as well as the structures of the CsCl salt particles formed upon the sublimation. We inspect the potential of the frozen solutions to generate fine salt particles that may become a source of salt aerosol when airborne. The number, size, and structure of those salt particles that remain after ice sublimation were analyzed as the function of the salt concentration in the sample, the freezing method, and the sublimation temperature. Similarly, the effect of liquid sample evaporation slightly above the freezing point was inspected. This article follows on our previous paper in which the morphologies of salty frozen solutions were detailed (Vetráková et al., 2019), and complements our recent letter publication that briefly introduces this topic by imaging sea salt solutions in the ESEM (Závacká



et al., 2022). Using CsCl instead of sea salt enlightens the mechanism of formation of the individual salt structures, therefore it provides a missing piece of information, mediating detailed knowledge not obtainable from sea salts.

65 2 Methods

2.1 Description of the ESEM

The microscopic images were recorded using a non-commercial ESEM AQUASEM II constructed from a Tescan VEGA SEM at the Institute of Scientific Instruments of the Czech Academy of Sciences (Neděla, 2007). Compared to a conventional SEM, the ESEM does not require high vacuum conditions in the specimen chamber. The ESEM is suitable for the direct imaging of
70 wet, electrically non-conductive and electron beam-sensitive samples (Michaloudi et al., 2018) at chamber pressures up to 2,000 Pa; the samples do not require conductive coating (Schenk Mayerová et al., 2014). Moreover, the device allows in-situ observation under dynamically changing environmental conditions (Neděla et al., 2020). The samples are placed on the silicon pad of a Peltier cooling stage, which can reach temperatures down to -27 °C ; this ESEM facilitates the imaging of frozen samples. The electron beam energy to enable the imaging amounted to 20 keV. The low beam current (100 pA) and short dwell
75 time (14 μs) minimise the radiation damage and local heating of frozen samples. The YAG:Ce³⁺ scintillation detector of backscattered electrons (BSE) is sensitive to signal from atoms with a large atomic number (Neděla et al., 2018); therefore, very good material contrast between the ice and the CsCl salt is obtained.

2.2 CsCl as a proxy for sea salt

We observed the sublimation, evaporation, and structure of the residua of the CsCl solutions under various conditions (the
80 concentration of the salt, the freezing method, and the sublimation temperature) in the ESEM. Although the project focused on identifying the parameters important for the formation of SSAs, the experiments utilized CsCl salt instead of sea salt of a complex composition in order to increase the visibility of the salt in the BSE detector (Figure S1). In the images, both the CsCl crystals and the brine are represented in white, the ice is usually black, and the silicon surface of the cooling stage is grey. Due to this contrast, we were able to easily distinguish liquid brine and solidified salt from the ice crystals and cooling stage and to
85 monitor the whole process of sublimation, not only the structure of the residua. For example, we were able to assign the original location of various salty structures within frozen samples.

Besides the benefit of superior contrast in the ESEM, CsCl was chosen as a proxy for the sea salt due to the similarity in relevant physical properties with NaCl, which is by far the most abundant salt present in seawater. The T_{eu} of the CsCl solution (the published values range from -24.83 to -22.3 °C (Chen et al., 2005; Cohen-Adad and Lorimer, 1991; Dubois et al., 1993; Fujiwara and Nishimoto, 1998; Gao et al., 2017; Monnin and Dubois, 1999)) is very close to that of the NaCl (-21 °C (Rodebush, 1918)). The eutectic solubilities of CsCl and NaCl are also similar (7.75 vs. 5.11 mol kg^{-1} (Gao et al., 2017; Swenne, 1983)). Both salts can therefore be expected to exhibit a similar tendency to crystallize. Others (Hullar and Anastasio,



2016) and we (Vetráková et al., 2019) have previously applied CsCl to understand the freezing of salty solutions. Seawater contains also other salts besides NaCl; these crystallize at various temperatures (e. g. ikaite and mirabilite have much higher eutectic point than NaCl) and may possibly perform certain secondary tasks that are not covered in this study. We tested the applicability of CsCl as the proxy for the sea salt by performing several analogous experiments using the proper sea salt. The overall sublimation behaviour was very similar to the one presented in the current study (Závacká et al., 2022). However, CsCl grant much better contrast (Figure S1) allowing the observation of subtle features otherwise hardly discernible. Thus, we are of opinion that CsCl can be a suitable probe for the sea salt when enhanced contrast is required, and the important factors affecting the structures of the sublimation residua and their potential to form salt aerosols revealed in this study might be applied also to the formation of natural SSAs.

2.3 Freezing and sublimation

CsCl solutions with the concentrations of 0.005, 0.05, and 0.5 M (which approximates to 0.084, 0.83, and 7.8 wt%, or 0.84, 8.3, and 78 psu, respectively) were prepared via dissolving appropriate amounts of CsCl in MilliQ H₂O. The droplets with a diameter of ~ 4 μm were frozen at atmospheric pressure via three distinct methods:

- (I) Spontaneous freezing of a droplet without seeding. The droplet was deposited onto the cooling stage of the ESEM. The temperature of the cooling stage was gradually lowered from room temperature (about 23 °C) down to the desired sublimation temperature (T_{sub}) at a cooling rate of ~ 0.5 °C/s; the sample froze spontaneously at -10 to -12 °C in the replicate experiments. Within this article, the samples are referred to as *non-seeded*. The term spontaneous freezing was used in this text because no additional nucleation agents were introduced into the samples on purpose; it must not be mistaken for homogenous ice nucleation which can occur at temperatures as low as -40 °C (Mason, 1958). Presumably, the nucleation is still initialized heterogeneously, e.g. by dust particles or on the surface of the cooling pad.
- (II) Controlled ice nucleation (seeding). The droplet was placed onto the cooling stage of the ESEM; subsequently, the stage was cooled down to -3 °C and, after the thermal equilibration, several small ice crystals were added to the edge of the sample to initiate the nucleation process. Ice started to form, we could observe the movement of the freezing front with the naked eye. Eventually, the temperature was lowered to the T_{sub} at a cooling rate of ~ 0.5 °C/s. In the text, these samples are referred to as *seeded*.
- (III) Freezing in liquid nitrogen (LN). A droplet of the sample was deposited on a piece of aluminium foil and immersed into the LN. The foil was used to ensure a flat bottom of the sample, guaranteeing a better thermal contact with the cooling stage of the ESEM. Then, the frozen sample was transferred onto the stage precooled to the T_{sub} . Within the article, these samples are denoted as *LN-frozen*. In some of the experiments, small frozen spheres (a diameter of approx. 100 to 200 μm) were prepared (by spraying a solution into the LN) to evaluate the effect of the sample size.

The specimen chamber of the ESEM was closed and evacuated. The ambient pressure inside the chamber was maintained at 500 Pa. The gaseous environment comprised mostly blown-in (dry) air, although a small amount of water vapour was present too due to ice sublimation. T_{sub} for all the above-mentioned methods was either -20 or -25 °C; the frozen samples were held at this temperature during whole sublimation. The chosen temperatures allow us to observe the differences in the sublimation just above and just below the T_{eu} .



The freezing methods I and II are identical with those described in our previous paper (Vetráková et al., 2019), where the
130 freezing rates were estimated to be (154 ± 13) mm s⁻¹ in the non-seeded sample (method I) and ~ 0.2 mm s⁻¹ in the seeded one
(method II). The rate of the LN freezing (method III) was not determined. Apart from the freezing rate, the three freezing
methods differ in the direction of freezing (a vector perpendicular to the progressing freezing front). Estimated directions
incident to these freezing methods are outlined in Figure 1.

135 2.4 Evaporation

Droplets (~ 4 mm in diameter) of the liquid 0.005 and 0.05 M CsCl solutions were evaporated in the evacuated specimen
chamber of the ESEM at the ambient air pressure of 650 Pa and the temperature (T_{evap}) of 2 °C.

3 Results and discussion

This paper describes the ice sublimation process above and below the T_{eu} and the morphology of the resulting residua of the
140 non-seeded, seeded, and LN-frozen samples. The outcomes are compared to the residua after the evaporation of the salty water.

3.1 Sublimation above the T_{eu}

When the temperature during the sublimation reached -20 °C, which is above the T_{eu} , the CsCl brine was still in the liquid
state; no salt crystals were observed on the surface of the ice (Figure S2a, c, e). Following pattern was observed for the seeded
and non-seeded samples in all of the concentrations (Figures 3-5, left and middle columns) and in the 0.005 M LN-frozen
145 samples (Figure 3c). The liquid brine accumulated at the edge of the sample on the cooling stage. As the sublimation proceeded,
the diameter of the sample decreased, and the brine slowly shifted towards the retreating sample edge. At some point, the brine
at the edge crystallized. The frozen sample continued reducing its size due to the sublimation, and soon the edge was separated
from the crystallized brine. New liquid brine kept leaking out of the sample, and the situation recurred incessantly. The
crystallization sequence at the edge is illustrated in Figure 2. The amount of the crystallized brine diminished progressively
150 towards the centre of the sublimated sample because of the preceding drainage (Figures 3-5). After the sublimation, the
crystalline CsCl was arranged in the form of concentric circles (Figure 6). The salt rings were usually thick and compact close
to the periphery, but they became thinner towards the centre and, especially in the samples with the lowest salt concentration,
even small isolated particles occurred at the very centre of the samples (Figure 3). There are two concurrent processes that
may run during the described sequence: ice sublimation and evaporation of water from the brine. The tendency of the sample
155 to sublimate or evaporate can be expressed in terms of saturated vapour pressures above the ice and the brine, respectively. At
 -20 °C, the saturated vapour pressures above ice and brine are 103 (Wexler, 1977) and 99 Pa (Text S1). The vapour pressure
above the ice at -20 °C is then slightly higher than that above the brine, and the sublimation of ice is expected to be preferred



slightly more than the evaporation from the brine. Nevertheless, as the difference is small, both processes occur simultaneously. The sublimation of the ice only shrinks the sample, without disturbing the equilibrium, while the evaporation of water from the brine causes its supersaturation. The brine that is in close contact with the ice can compensate for the water loss through ice melting to sustain the equilibrium brine concentration, and thus the brine remains liquid. However, the ice melting cannot supply the water lost by evaporation from the brine at the very edge of the sample. Therefore, only the brine at the edge of the sample, far from the ice, becomes crystalline above the T_{eu} . In due time, the sublimation and evaporation lead to a shrinkage of the sample and the subsequent accumulation of the brine beyond the edge, followed by crystallization of the brine; these processes continually iterate slightly closer to the centre of the sample. As a result, the pieces of crystallized salt are arranged in the observed pattern of concentric (semi)circles (Figures 2-6); the crystalline salt firmly adheres to the silicon pad of the cooling stage and cannot be blown away in our experimental conditions. These patterns of the sublimation residua (formed above the T_{eu}) resemble those of the “stick-slip” mechanism observed in TiO_2 nanoparticles from evaporating ethanol (Moffat et al., 2009).

In contrast to the output described in the preceding paragraph, the 0.05 and 0.5 M LN-frozen samples exhibited some differences (Figures 4-5, right columns). As in the previous cases, the liquid brine accumulated at the edge of the sample and, at a certain point, started to crystallize. However, the crystallization continued all over the surface of the frozen sample, and a fibrous, shell-like structure was formed on the surface. Despite its delicate appearance, the structure retained its shape even when the ice had completely sublimated.

The obtained results seem to indicate that the formation of salt aerosols at temperatures above the T_{eu} is generally improbable due to the crystallization of the liquid brine into large pieces of salt unless the salt concentration is very low, and due to the adhesion of the crystalline salt to the ground (however, we admit the base of the saline ice is different in nature and so the adhesion to the ground may vary in comparison with our experiments). These results are in accordance with our previous study where we applied sea salt instead of CsCl (Závacká et al., 2022). Therefore, we believe the observed crystallization of the salt into large chunks above the T_{eu} is transferrable also to natural samples and prevents the ice to become an effective source of aerosolizable particles under these conditions. Still, a portion of fine salt particles (with the size of few micrometres) may form above the T_{eu} due to the brine leakage beyond the original sample position; this process is described in chapter 3.3.

3.2 Sublimation below the T_{eu}

For the sublimation to proceed below the T_{eu} , the temperature in our experiments was set to -25 °C ; due to limitation of the cooling stage, we were not able to achieve lower temperature. Although the temperature of the surface may slightly deviate from that of the cooling stage (the sample temperature estimated with a Pt1000 sensor was $\sim 2\text{ °C}$ higher than the pre-set one), solidified CsCl brine was visibly present on the surfaces of the frozen samples under the pre-set conditions in all of the concentrations and freezing methods. In a few replicates, however, the surface brine was still liquid at this temperature due to closeness to the eutectic point; these results are not included in the article.



190 3.2.1 Concentrations of 0.005 and 0.05 M

The sublimation residua of the 0.005 and 0.05 M CsCl are displayed in Figures 7 and 8, respectively. The non-seeded and seeded samples of the two concentrations appear to be largely similar. Three types of residual salt structures were observed:

1. A flat rim at the location of the circumference of the frozen sample (Figures 7a, 7c, 8a, 8c); this rim supposedly formed from the brine that had leaked to the sample's edge before the T_{eu} had been reached during the freezing processes. This formation mechanism is supported by the fact that, if the temperature temporarily rises above the T_{eu} (and the brine liquifies) and then drops again, the salt rim becomes wider (analogous situation is shown in Figure S3). During these processes, surplus liquid brine was observed to leak to the edge of the sample.

2. Fine salt particles that appeared from beneath the samples and remained lying on the stage (Figures 7a, 7c, 7d, 8a, 8d, 9) as the samples were sublimating; typically, their widths and lengths were approximately 1 μm and several micrometres, respectively. In the seeded samples, the microparticles were often arranged in straight lines (Figures 7c, 8c, 8d), while a random arrangement dominated in the non-seeded samples (Figures 7a, 8a). In the 0.05 M samples, they even interconnected into web-like structures instead of producing separated crystals (Figure 8c). Based on the adhesion of the fine particles to the silicon pad (Text S2), we suppose the particles were formed by the brine crystallizing at the bottom of the frozen sample, which is in contact with the cooling stage. The well-preserved regular lines of the small salt crystals on the surface of the silicon pad in the non-seeded and seeded droplets at the two lower concentrations enable us to grasp the original lamellar or cellular arrangement of ice and brine. The morphology of ice is closely related to particular freezing temperatures (Schremb and Tropea, 2016; Shibkov et al., 2003). The spacing of the lamellae depends on the freezing rate and solute concentration (Maus, 2007; Rohatgi and Adams, 1967; Wettlaufer, 1992).

3. Larger flakes (tens of micrometres in size) of an irregular shape appeared on the surface of the frozen samples during the sublimation (Figures 7b, 10, and 11). When the ice underneath these flakes sublimed, they detached from the surface but remained slightly fixed to the actual sample. As the frozen sample was retreating due to the sublimation, the flakes kept approaching to the centre of the sample until they fell down onto the stage (Figures 10 and 11). Thus, the flakes were scarcely found within the periphery, but their abundance increased towards the centre of the sublimated sample. A very similar behaviour was described and explained previously in the sublimation of a frozen colloidal suspension (Jambon-Puillet, 2019). Although we observed the flakes right on the ice surface, the crystallized brine had supposedly originated also from inside the frozen sample, as the sublimating ice continuously exposed deeper layers of the sample (such as the veins previously hidden in the bulk). Small flakes and aggregates were not affixed to the stage and could be readily blown away or split, possibly forming aerosol particles (Text S2). In the 0.05 M samples, large tufts of aggregated lichen-like structures, formed probably by the brine crystallizing in the veins between the ice grains, were abundant in the central part of the sublimated samples (Figures 8b, d). Having sizes of tens to hundreds of micrometres, the tufts were usually much larger than the salt flakes.

We processed the micrographs by using Mountains® software and estimated the abundance and size of the salt particles on the pad (Texts S3, S4). We detected approximately 3,400 and 13,000 particles mm^{-2} in the 0.005 M non-seeded and seeded



225 samples, and approximately 39,000 and 60,000 particles mm^{-2} in the non-seeded and the seeded 0.05 M samples, respectively (Tables 1, S1). The majority of the particles were less than 10 μm in maximum diameter (Figure 12). Although the particles were spread non-homogeneously over the pad, and the numerical estimates can be biased, the seeded samples evidently produced markedly more fine particles than the non-seeded ones at both the concentrations. Therefore, seeding favours the formation of aerosolable salt particles significantly more than spontaneous freezing.

230 Compared to both the non-seeded and the seeded samples, the structures of the sublimation residua from the LN-frozen sample were completely different (Figures 7e, 7f, 8e, 8f), forming large fluffy tufts of salt having a lichen-like appearance. The smaller aggregates exhibited a size of about 100 μm , but the large ones exceeded the dimension of our field of view, which is about 400 μm . Unlike the other two freezing methods, the LN technique did not produce any salt rim on the original circumference of the sample, as the sample had already been much below the T_{eu} when transferred onto the cold stage of the ESEM. We did not find any smaller ($< 100 \mu\text{m}$) crystals in the sublimation residua of the LN-frozen samples. This effect is supposedly due to the fact that during the LN-freezing the sample freezes from the outside, and the freeze-concentrated brine is expelled by the growing ice towards the interior of the sample. Such sub-surface brine inclusions were revealed in LN-frozen spheres of aqueous solutions already previously (Vetráková et al., 2019). As an excess of the salt forms inside the sample, and the ice crystals emanated during the fast freezing are small, the veins between the ice crystals in the frozen sample may be more interconnected, and thus the resulting sublimation residua form structures larger than those delivered by the other two freezing methods.

240 3.2.2 Concentration of 0.5 M

High solute concentrations yield a large amount of the brine, which is sufficiently interconnected throughout the 0.5 M samples; such a scenario then leads to the formation of relatively stable and large self-supporting structures that differed significantly from those revealed at lower concentrations. All of the sublimation residua retained the original shapes of the frozen samples, forming salty casts in which the solidified brine delineates the shapes of the ice crystals (Figure 13). The freezing method was a major factor determining the structures of the sublimation residua. The variation of the outcomes of the three freezing methods is striking: The surface of the sublimated non-seeded sample was covered with a fibrous film of salt, through which hollows of various sizes were visible (Figure 13a); however, most of the structure underneath the film remained hidden. This effect corresponds to the results of our previous studies (Vetráková et al., 2019, 2020) in terms of the accumulation of the solutes on the surface of the non-seeded sample, as in a globally supercooled sample cooled from the bottom the freezing likely proceeds from the bottom upwards (Figure 1a). The observed surface “skin” restricts to a large extent the access to the interior of the sample, thus preventing the more subtle and possibly fragile parts from being easily withdrawn by the blowing winds. The situation differed in the sublimated seeded sample (Figure 13b); the seeding likely promoted freezing downwards from the surface, where the nucleating crystal had been placed (Figure 1b). Almost no film of salt was present on the surface, allowing the inner structure to be clearly discernible. The well-resolved fibrous ice casts formed after the ice had sublimated;



255 these hollows in the salt matrix apparently retained the original sizes and shapes of the sublimed ice crystals that were very oblong, with strong parallel arrangement (Figure 13b). This well-preserved structure resembling the cells of a honeycomb illustrates that not only the veins (triple junctions, i.e. places where three ice crystals meet) but also the whole grain walls were filled with an amount of brine sufficient to support the structure. In the LN-frozen samples, a sponge-like structure appeared after the sublimation of both the LN-frozen droplet (Figure 13c) and the LN-frozen microspheres (Figure 13d). The pores in
260 the structure were numerous but much smaller than in the samples prepared via the other freezing methods, as abrupt freezing in LN induces the formation of small ice crystals. A comparison of the images in Figures 13c and 13d will reveal that the size of the sample does not determine the sizes of the ice crystals: The submillimetre spheres and millimetre-sized droplet exhibit similar dimensions of the pores in the residual structures. The structures hold together in large pieces even after the ice sublimation, with no collapse observed. By inspecting the sponge-like features in the detailed image of the LN-frozen residua
265 provided in the bottom-left corner of Figure 13c, we established that the structure of the residua delineates only the veins between the ice crystals, not the whole grain boundaries, as in the seeded sample. Moreover, the internal structures of the sublimated LN-frozen samples appear denser than those of the seeded sample, due to the very different freezing rates and directionalities: Freezing from the outside expels the brine towards the interior of the sample, as demonstrated in Figure 1c. Although 0.5 M samples form relatively stable and large self-supporting structures that are not eligible for aerosol formation,
270 these fine structures are expected to exhibit poor mechanical stability against external forces (including, for example, wind-cropping or physical collision with large mobile particles), and their potential breakdown could also lead to the production of small aerosol-sized particles.

3.3 Supercooled brine beyond the edge of a frozen sample

During our experiments, we often detected tiny brine droplets beyond the salt rim, where no observable frozen sample had
275 been previously present. These microdroplets did not crystallize at the eutectic temperature and stayed supercooled (Figure 14a), even though the brine in the sample (in the left part of the panel in Figure 14a) had already fully solidified. Evaporation of water from the microdroplets resulted in salt crystallization and formation of a large number of (often) rectangular crystals (Figure 14b) having typical sizes of less than 20 μm (Figure S4). Such supercooled brine microdroplets (and corresponding salt microcrystals) behind the circumference of the original sample were detected routinely in the non-seeded and the seeded
280 samples of all tested concentrations at both temperatures and in the LN-frozen samples above the T_{eu} ; however, they were absent from the LN-frozen samples below the T_{eu} . Thus, the generation of small microdroplets/crystals behind the perimeter of the visible sample was noticed in all samples whose temperature exceeded the T_{eu} during the experiments (Figures 4, 5, 14): the non-seeded and the seeded samples had been loaded on the cooling stage in the liquid state and subsequently frozen, while the LN-frozen samples were treated externally and had already fully solidified before being loaded on the stage. Therefore, we
285 assume that the observed effect relates to the interaction of the liquid brine with the silicon surface. Presumably, a small portion of a sample solution spread over the silicon pad and formed a thin film (invisible for the ESEM) due to surface wetting during



the freezing while the temperature was above the T_{eu} . Due to the evaporation, micrometric droplets were produced from the film; enhanced salt concentration in the microdroplets enabled their visualization in the ESEM. Further evaporation led to salt crystallization.

290 Previously, wetting the upper molecular layers of a silicon (1,0,0) surface with water was simulated (Barisik and Beskok, 2013; Ozcelik et al., 2020); the researchers observed the water spread beyond the drop boundary. The nanometre-sized crystals observed centimetres away from the creeping front were considered a proof of precursor nanofilms (De Gennes, 1985; Qazi et al., 2019). Our findings not only confirm such nanofilms in CsCl solutions but also reveal their formation around salty ices at
295 sub-zero centigrade temperatures can embody a viable mechanism of spreading the salts to the previously pristine snow in polar conditions. By this way micrometric salt crystals might form on a surface of ice or a rock.

3.4 Evaporation

We took a step further and observed the evaporation of the aqueous solution at +2 °C. Such conditions resemble the behaviour of sea water at solid supports on coastlines and grounds after the melting of sea ice (Keene et al., 2007). As the diameter of the
300 observed drops is about 4 mm, they are much larger than the jet drops from bubbles bursting on the water surface (Spiel, 1995), which embody the most common sources of the sea aerosols. The evaporation of the liquid CsCl solution in the ESEM was very quick: The droplet evaporated before we evacuated the specimen chamber and made the ESEM ready for the imaging. The evaporated sample exhibited a wide salt ring at the periphery (Figure 15a, c) and mostly dendritic, sword-like salt crystals in the central part (Figure 15b, d). The evaporation residua resembled to some extent the sublimation ones above the T_{eu} . This
305 is not surprising, because the evaporation-crystallization mechanisms of both the processes are similar (the salt in the frozen sample above T_{eu} is dissolved in the liquid brine and evaporation of water from the brine induces salt crystallization).

3.5 Implications to polar atmosphere and environment

This study identifies the conditions at which the sublimation of salty frozen samples generates small particles of salt that may potentially become a source of salt aerosols, and presents CsCl as a suitable probe for the sea salt in terms of their similar
310 properties and outcomes of the freeze-drying process. It complements our recent letter that introduces this topic by imaging sea salt solutions in the ESEM (Závacká et al., 2022). There, an artificially prepared (non-seeded) sea ice sublimated at -16, -30, and -40 °C. The results show that the sizes and amount of the salt particles are comparable in the sea salt and CsCl sublimation residues for analogous sublimation temperatures and salt concentrations. The greatest disadvantage in using sea salt was the poor contrast between the salt and the ice; all we could recognize were the final sublimation residue. On the
315 opposite, the liquid CsCl brine and the CsCl crystals were well visible on the surface of the ice (Figure S1), and this allowed us to observe the sublimation process, understand the sublimation mechanism, detect the origin of the fine particles, and correlate the absence of liquid brine with the formation of fine salt particles. The experiments were performed in the narrow



temperature range between -20 and -25 °C, i.e., close to the eutectic temperature, to show that it is the eutectic solidification that makes a difference. Without using contrasting CsCl, such information would not be acquired.

320 Denoted molar concentrations of 0.005, 0.05, and 0.5 M CsCl equal to 0.84, 8.3, and 78‰ (psu), respectively. These concentration values were applied to mimic the broad range of salinities in the environment, from low salinity surface snows in Arctic coastal regions to high salinity sea ice (formed from approx. 0.5 M seawater). We showed that the formation of the small particles is restricted if the brine is liquid during the ice sublimation, i.e., at temperatures higher than the T_{eu} . The sublimation temperature appears to be the most critical parameter in this respect; the concentration and freezing method seem

325 to affect the resulting structures of the sublimation residua above the T_{eu} only marginally. In the given context, we can comprehend the fact that previous studies did not succeed in seeking the SSA production from the frost flowers in a wind tunnel (Roscoe et al., 2011) and in an ESEM (Yang et al., 2017). In the former article, although the cold chamber air temperatures in most of the experiments were -30 °C or lower, the measured ice temperature reached -5 °C (Howard Roscoe and Eric Wolff, personal communications), and the temperature of the sampling line was about -10 °C (Roscoe et al., 2011).

330 In the latter one, the sublimation of the brine-covered frost flowers at the temperatures of -5 and -17 °C yielded a large chunk of salt (Yang et al., 2017). In both of the studies, the temperature of the frost flowers was well above the T_{eu} ; in that respect, the lack of aerosol-forming particles is in good agreement with the results of our study. A large amount of small salt particles and fine structures that might act as a source of salt aerosols can form only at temperatures below the T_{eu} . The structures of the sublimation residua that formed at this temperature strongly depended on the concentration: Lower concentrations produced

335 small, isolated particles, while high ones resulted in large aggregates and self-supporting structures (Table 2, Figure 16). At ≤ 0.05 M solute concentrations, the amount of the brine was too small to allow the veins in a frozen sample to interconnect sufficiently, and therefore the structures of the veins broke apart in the process of the ice sublimation (Figure 7); fine salt particles, small salt flakes, or small lichen-like tufts formed. At these concentrations and $T_{sub} < T_{eu}$, the residua of the non-seeded and the seeded samples appeared very similar, so the dependency on the freezing method may not be very strong. Based on

340 the appearance of the residua, we deduce that the lowest salt concentration (0.005 M) exhibits the highest proportion of fine particles over larger aggregates, and the ratio decreases with increasing concentration. From this perspective, the lowest salinity samples appear to be the best generators of small aerosolizable particles. Conversely, the absolute number of fine salt particles generated per mm^2 of the frozen sample substantially increased when the concentration rose from 0.005 to 0.05 M: In our experiments, the estimated number of the salt particles rose 11 and 5 times in non-seeded and seeded samples, respectively

345 (Tables 1, S1). However, the number of the particles then abruptly decreased at 0.5 M, as the highly concentrated sample did not yield any smaller salt particles and a self-supporting structure was formed instead. Thus, low to middle salt concentrations (0.005 to 0.05 M) are required in order to generate small aerosol-sized particles below the T_{eu} , and samples with the middle salt concentration can be even more effective in producing aerosolable particles than those having the low concentration. As we have not acquired enough reliable data on the relationship between salt concentration and number of small particles, we do



350 not attempt to extrapolate or estimate the number of small particles from samples having lower concentrations than the tested ones. Neither can we estimate the concentration at which the maximum number of small particles is formed.

We wish to show that CsCl serves as a good proxy for the sea salt for the performed experiments. Certainly, seawater freezing and sublimation exhibit some aspects where the nature of the individual salts is of central importance, however, the performed experiments with sea salt and CsCl showed a lot of similarities in terms of particles sizes and numerical densities. Based on
355 the similar physical properties of CsCl and NaCl salts near their eutectic points (chapter 2.2) and the consistency of the experimental results concerning the sublimation residues of the sea salt (Závacká et al., 2022) and CsCl (this work), we are convinced that CsCl can be a suitable probe for the sea salt when enhanced contrast is required, mediating detailed knowledge not obtainable from sea salts. We acknowledge that our system embodies merely a model of natural sea ice, but we demonstrate that there is a behaviour, previously undescribed, that is bifurcated by eutectic temperature. Indeed, the kinetics of the freezing
360 and the shape of ice crystals can vary greatly to deviate from our model observations. However, as the outputs in the two “mild” freezing methods (spontaneous freezing and seeding) are largely similar under the conditions when aerosolizable particles are detected ($T_{\text{sub}} < T_{\text{eu}}$; conc. ≤ 0.05 M), we believe the concept presented in this study is, to some extent, applicable also for freezing in natural conditions.

3.6 Seeking for a source of SSA

365 The argument for the frost flowers being a source of the SSA was threefold: (1) whether they are sulphate-depleted; (2) whether they break and form small particles; and, finally, (3) whether they cover enough area to be relevant. When the ice temperature ranges below -6.4°C , which is a typical situation in a polar winter and spring, mirabilite ($\text{Na}_2\text{SO}_4 \cdot 10 \text{H}_2\text{O}$) precipitates out from the brine, and therefore the frost flowers growing from the brine are sulphate-depleted, as seen in the observations (e.g. Rankin and Wolff, 2003). The temperature gradients above the thin ice are strong, but it is unlikely that the temperature of
370 newly grown frost flowers is below the T_{eu} , except for the tips of the flowers. However, when sea ice becomes thicker, ice surface temperature will approach that of the air and may decrease below the T_{eu} , making aged frost flowers more prone to generating SSAs, on condition that the brine concentration is low enough. Nevertheless, the aged frost flowers are more likely to have been buried by precipitation of snowfall, reducing the opportunity for the frost flowers to serve as a source of SSA.

Both the recent field data (Frey et al., 2020) and the modelling (e.g., Huang and Jaeglé, 2017; Levine et al., 2014; Rhodes et al., 2017; Yang et al., 2019) have demonstrated the importance of airborne saline snow particles as a source of the SSA in
375 polar regions. The salty snow lying on the sea ice has also the potential to generate the SSA at temperatures below the T_{eu} . On average, the snow salinity is several orders of magnitude lower than that of sea water and frost flowers (Frey et al., 2020). Thus, according to the results presented herein, the snowpack is supposed to yield smaller salt particles and fine structures of salts upon sublimation at very low temperatures, and, therefore, to be potentially an effective source of the SSA. However,
380 there might be significant differences depending on whether the snowpack lies on young sea ice or multi-year sea ice, and whether there is flooding, which depends on snow thickness vs. ice thickness. Normally, young sea ice is thin and thus



relatively warmer than thick ice, and a temperature as low as the T_{eu} could be difficult to reach on very thin young ice. Therefore, the snowpack on multi-year ice is more likely to form fine salt structures and SSA than the snow on relatively thin sea ice. There are many other factors that may affect the structure of the residua – e. g. salt crystallographic system, activity of the brine solution, or geometry of the cooling environment. Finally, we should also note that the formation of the sublimation
385 residua will probably be accompanied by acidity changes during both the freezing and the sublimation (Vetráková et al., 2017) which may have major consequences as regards the heterogeneous reactivity of the aerosols thus formed (Pratt et al., 2013). The implications of our findings to the polar atmosphere and environment are significant; however, further data (field, laboratory and modelling) are still necessary to confirm the conclusions.

390 3.7 Connection to the morphology and properties of the original frozen samples

The observed morphologies of the sublimation residua formed below the T_{eu} bear important information about the internal environment of the original frozen samples. We previously studied the properties of the frozen samples from which the residua had formed (Vetráková et al., 2019). In the relevant article, the differences in the surface morphologies were well visible, but the observation of the interiors was insufficient. The brine structures visible on the ice surfaces and the sublimation residua of the spontaneously frozen samples sublimed at -25 °C are compared in Figure 17.
395

The sublimation below the T_{eu} in the research reported herein allowed us to inspect also the sub-surface morphology by the ESEM more thoroughly. Another method for the direct visualization of solute locations was microtomography (μ CT) (Hullar and Anastasio, 2016). The technique provided a 3D scan of the location of impurities in ice, but the resolution was much coarser than that achieved with the ESEM (~ 16 and $2\text{ }\mu\text{m}$ in the μ CT vs ~ 500 and 50 nm in the ESEM for an overview and
400 details, respectively). Thus, the researchers were able to visualize larger inclusions of brine in the ice, but the brine in the veins (65 to 88 wt% of the salt) was beyond the detection ability of the μ CT method. The deduction of the morphology of a frozen sample from the morphology of a sublimation residuum can now supply the missing piece of the puzzle.

There appear to be several characteristic domains where impurities can accumulate within ice – on the surface, at the grain boundaries, in liquid inclusions, or within the ice crystals (Bartels-Rausch et al., 2014; Blackford et al., 2007; Dominé et al.,
405 2003; Hullar and Anastasio, 2016; Light et al., 2009). The existence of various microenvironments in the vicinity of guest molecules within ice was also observed spectroscopically (Heger et al., 2011; Heger and Klán, 2007; Krausko et al., 2014; Ondrušková et al., 2018). The location of impurities in ice and snow is an important factor for their reactivity and potential release to the atmosphere, due to considerable variations in the accessibility of gas-phase oxidants and photons in the domains (Hullar and Anastasio, 2016 and the references within). For example, our experimental results indicate that the LN-frozen
410 microspheres contain most of the brine inside the ice, not on the surface (Vetráková et al., 2020). It remains an open question if the molecules on the surface of the frozen microsphere are exposed to the incoming radiation or reactive gasses to a larger extent than the molecules inside and if and how the accessibility of these compartments depend on the temperature (Ray et al., 2011).



When comparing the sublimation residua formed below the T_{eu} from the 0.5 M frozen samples (Figure 13), it is obvious that
415 an abundance of the salt on the surface markedly depends on the freezing method. While the non-seeded sample is completely
covered with salt, there is no excessive surface salt in the seeded sample and LN-frozen samples. This corroborates our previous
observations of the frozen (unsublimated) samples (Vetráková et al., 2019), where the amount of salt on the ice surface was
related to the directionality of freezing. Higher amounts of surface salt in the frozen non-seeded sample can thus lead to
enhanced oxidation of halogens by atmospheric gases and their release to the atmosphere. The sublimation of ice from the
420 concentrated frozen samples leads to the formation of large salt chunks (above T_{eu} , Figure 5) or highly porous structures (below
the T_{eu} , Figure 13). Therefore, the sublimation temperature can significantly affect the accessibility of gases and photons to the
salt and thus alter the reactivity within these compartments.

The inspection of the residua formed by the sublimation of the 0.5 M frozen samples below the T_{eu} (Figure 13) also indicated
substantial differences in the thickness of the brine compartments in the interior of the seeded and the LN-frozen samples.
425 Regrettably, we could not evaluate this parameter in the non-seeded sample, as its surface was covered with a salt film to a
large extent and the internal morphology was hidden. In the seeded sample, the salt crystallized in the form of relatively thick
walls surrounding individual ice crystals. In the LN-frozen sample, the salt crystallized as very thin tortuous veins – the overall
shape of the residuum resembled a sponge. The most important differences between these two preparation methods consisted
in the directionality and freezing rate. Seeding is a method of slow freezing from the upper surface (Figure 13b), generating
430 large ice crystals, while LN-freezing proceeds under fast freezing from all sides (Figure 13c), generating small ice crystals
(Vetráková et al., 2019). Thus, the freezing rate and/or directionality affect the thickness of the brine compartments in a frozen
sample. It is intuitively obvious that the brine surrounding small ice crystals with large specific surface areas needs to spread
more, and therefore the grain boundaries and veins containing the salt ought to be thinner than in the sample of identical
concentration containing larger ice crystals. In this study, this relationship is proved experimentally (Figure 13b vs. 13c).
435 Previously, slow cooling rates were related to a larger extent of aggregation of impurities in ice, (Heger et al., 2005) while fast
cooling rates led to partial vitrification of the brine (Imrichová et al., 2019; Ondrušková et al., 2020), presumably due to
confined space and more efficient cooling of thin veins. The subtle differences in the ice-impurity morphology can possibly
alter the immediate molecular environment and consequently variate the absorption (Bononi et al., 2020), aggregation (Kania
et al., 2014), pH changes, (Heger et al., 2006) and, therefore, also the reactivity of compounds and their photochemical quantum
440 yields (Hullar et al., 2018; Kahan and Donaldson, 2007; Klánová et al., 2003).

Herein we observed several characteristic units among the sublimation residua. The residua formed above the T_{eu} cannot be
assigned to particular domains in the ice, because liquid brine from the individual domains pours together after the ice has
sublimated. However, the flow of the crystallized salt below the T_{eu} is restricted, and the shape of the residua (fine particles,
salt flakes, and lichen and sponge-shaped tufts) partially mirrors the one of the domains in the ice. From our observations of
445 the sublimation processes, we deduce the genesis of the individual salt features as follows: Fine salt particles are likely formed
by crystallization of the brine in the veins at the bottom of the sample. Arguments supporting this opinion were already outlined



above (adhesion of the particles to the cooling pad, their visual absence on the gradually sublimated ice surface). However, we cannot exclude that a portion of the fine particles comes from thin or separated veins inside the ice body that broke to small pieces after ice had sublimated. Remnants of the crystalline salt from larger and more interconnected veins can be presumably found in the form of lichen-like tufts. Lastly, crystallization of the brine in pools on the surface and liquid inclusions in the ice body likely led to formation of 2-dimensional salty flakes. The knowledge of the morphologies brought up in this paper and their relationship to the properties of original frozen samples should be borne in mind when designing laboratory experiments to simulate natural ice and snow. As the properties of frozen samples largely depend on the sample preparation method, artificial samples must be prepared with care in order to mimic natural ice and snow.

455 5 Summary

In this study we present a novel technique that facilitates identification of the most suitable conditions for formation of salt aerosols. We used CsCl as a proxy for natural sea salts due to its excellent visibility within the ice by the ESEM, and observed the transformation of frozen salty solutions into residual salt particles. By inspecting the morphology and number of the resulting salt particles based on the salt concentration, freezing method and sublimation temperature, we demonstrated that the sublimation temperature is the most important factor for the formation of aerosolable particles upon sublimation of various frozen salty ices. When the sublimation takes place at temperatures above the T_{eu} , the formation of large compact pieces of salt with very little aerosol-forming potential is preferred for all the concentrations and freezing methods tested herein; the concentration and the freezing method seem to play a less important role in the structure of the sublimation residua. Similar effect was observed when a liquid salty solution evaporated. Conversely, the structures of the residua formed at temperatures below the T_{eu} strongly depend on the concentration: Low salinity samples yield small isolated particles and small aggregates ($< 10 \mu\text{m}$) that are directly available to be windblown and become salt aerosols; high salinity samples transform into large aggregates with fine structures. Regarding the size and abundance of the fine particles in various residua, the 0.05 M seeded sample yielded the largest number of fine salt particles and thus would be the most promising source of salt aerosols. The outcomes of this article clearly indicate that the sublimation process on highly saline ices, such as frost flowers lying on young sea ice, does not directly induce the formation of SSAs unless the interconnected salt structures are potentially brought down by external forces, such as physical collision with large mobile particles. On the other hand, less saline snow lying on aged ice is more likely to directly generate separated fine salt particles during the sublimation process at temperatures below the T_{eu} . Our data emphasize the requirement of very low temperatures for the formation of SSAs. This condition favours salty snow over frost flowers as an efficient source of the SSA, provided that no flooding has occurred.

475 Inspecting CsCl-containing frozen and sublimated samples in the ESEM generally led to very similar outcomes in comparison with the sea salt (Závacká et al., 2022), but provided us with many details in the samples' morphology that would otherwise remain unseen. We are aware of the limitations in using artificial samples to mimic natural processes. However, such



480 systematic study where we vary parameters (sublimation temperature, concentration, freezing method) and monitor the
outcome are very difficult (and nearly impossible) to perform with natural samples, as there are many collateral effects in
nature that we are not able to fully distinguish and interpret. Nevertheless, evaluation of these effects step by step in a laboratory
may be very helpful when interpreting the behaviour of complex natural samples.

Author contributions

485 L.V. conducted the experiments, analyzed the data, and prepared the manuscript; K.Z. performed auxiliary experiments; D.H.
and X.Y. developed the idea of the project; and V.N., D.H., and X.Y. contributed to the discussions and writing of the final
paper.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared
to influence the work reported in this paper.

Acknowledgement

490 This work was supported by Czech Science Foundation via project GA22-25799S. The study was supported by Strategy AV21
– programme #23 “City as a laboratory of change; construction, historical heritage and place for safe and quality life”. We
thank Martin Olbert for the particle size evaluation and Přemysl Dohnal for the language corrections.

References

- 495 Abbatt, J. P. D., Thomas, J. L., Abrahamsson, K., Boxe, C., Granfors, A., Jones, A. E., King, M. D., Saiz-Lopez, A., Shepson,
P. B., Sodeau, J., Toohey, D. W., Toubin, C., Von Glasow, R., Wren, S. N. and Yang, X.: Halogen activation via interactions
with environmental ice and snow in the polar lower troposphere and other regions, *Atmos. Chem. Phys.*, 12(14), 6237–6271,
doi:10.5194/acp-12-6237-2012, 2012.
- Barisik, M. and Beskok, A.: Wetting characterisation of silicon (1,0,0) surface, *Mol. Simul.*, 39(9), 700–709,
doi:10.1080/08927022.2012.758854, 2013.
- 500 Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J. and Rasmussen, R. A.: Ozone destruction and photochemical
reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334(6178), 138–141, doi:10.1038/334138a0, 1988.
- Bartels-Rausch, T., Jacobi, H. W., Kahan, T. F., Thomas, J. L., Thomson, E. S., Abbatt, J. P. D., Ammann, M., Blackford, J.
R., Bluhm, H., Boxe, C., Domine, F., Frey, M. M., Gladich, I., Guzmán, M. I., Heger, D., Huthwelker, T., Klán, P., Kuhs, W.



- F., Kuo, M. H., Maus, S., Moussa, S. G., McNeill, V. F., Newberg, J. T., Pettersson, J. B. C., Roeselová, M. and Sodeau, J.
505 R.: A review of air–ice chemical and physical interactions (AICI): liquids, quasi-liquids, and solids in snow, *Atmos. Chem. Phys.*, 14(3), 1587–1633, doi:10.5194/acp-14-1587-2014, 2014.
- Blackford, J. R., Jeffree, C. E., Noake, D. F. J. and Marmo, B. A.: Microstructural evolution in sintered ice particles containing NaCl observed by low-temperature scanning electron microscope, *Proc. Inst. Mech. Eng. Part L-Journal Mater. Appl.*, 221(L3), 151–156, doi:10.1243/14644207jmda134, 2007.
- 510 Bononi, F. C., Chen, Z., Rocca, D., Andreussi, O., Hullar, T., Anastasio, C. and Donadio, D.: Bathochromic Shift in the UV-Visible Absorption Spectra of Phenols at Ice Surfaces: Insights from First-Principles Calculations, *J Phys Chem A*, doi:10.1021/acs.jpca.0c07038, 2020.
- Bottenheim, J. W., Fuentes, J. D., Tarasick, D. W. and Anlauf, K. G.: Ozone in the Arctic lower troposphere during winter and spring 2000 (ALERT2000), *Atmos. Environ.*, 36(15–16), 2535–2544, doi:10.1016/S1352-2310(02)00121-8, 2002.
- 515 Butler, B. M., Papadimitriou, S. and Kennedy, H.: The effect of mirabilite precipitation on the absolute and practical salinities of sea ice brines, *Mar. Chem.*, 184, 21–31, doi:10.1016/j.marchem.2016.06.003, 2016.
- DeMott, P. J., Hill, T. C. J., McCluskey, C. S., Prather, K. A., Collins, D. B., Sullivan, R. C., Ruppel, M. J., Mason, R. H., Irish, V. E., Lee, T., Hwang, C. Y., Rhee, T. S., Snider, J. R., McMeeking, G. R., Dhaniyala, S., Lewis, E. R., Wentzell, J. J. B., Abbatt, J., Lee, C., Sultana, C. M., Ault, A. P., Axson, J. L., Diaz Martinez, M., Venero, I., Santos-Figueroa, G., Stokes,
520 M. D., Deane, G. B., Mayol-Bracero, O. L., Grassian, V. H., Bertram, T. H., Bertram, A. K., Moffett, B. F. and Franc, G. D.: Sea spray aerosol as a unique source of ice nucleating particles, *Proc. Natl. Acad. Sci.*, 113(21), 5797–5803, doi:10.1073/pnas.1514034112, 2016.
- Dominé, F., Lauzier, T., Cabanes, A., Legagneux, L., Kuhs, W. F., Techmer, K. and Heinrichs, T.: Snow metamorphism as revealed by scanning electron microscopy, *Microsc. Res. Tech.*, 62(1), 33–48, doi:10.1002/jemt.10384, 2003.
- 525 Frey, M. M., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Yang, X., Jones, A. E., Nerentorp Mastromonaco, M. G., Jones, D. H. and Wolff, E. W.: First direct observation of sea salt aerosol production from blowing snow above sea ice, *Atmos. Chem. Phys.*, 20(4), 2549–2578, doi:10.5194/acp-20-2549-2020, 2020.
- De Gennes, P. G.: Wetting: statics and dynamics, *Rev. Mod. Phys.*, 57(3), 827, doi:10.1103/RevModPhys.57.827, 1985.
- Heger, D. and Klán, P.: Interactions of organic molecules at grain boundaries in ice: A solvatochromic analysis, *J. Photochem. Photobiol. A Chem.*, 187(2–3), doi:10.1016/j.jphotochem.2006.10.012, 2007.
- 530 Heger, D., Jirkovský, J. and Klán, P.: Aggregation of Methylene Blue in Frozen Aqueous Solutions Studied by Absorption Spectroscopy, *J. Phys. Chem. A*, 109(30), 6702–6709, doi:10.1021/jp050439j, 2005.
- Heger, D., Klanova, J. and Klan, P.: Enhanced protonation of cresol red in acidic aqueous solutions caused by freezing, *J. Phys. Chem. B*, 110(3), 1277–1287, doi:10.1021/jp0553683, 2006.
- 535 Heger, D., Nachtigallová, D., Surman, F., Krausko, J., Magyarová, B., Brumovský, M., Rubeš, M., Gladich, I. and Klán, P.: Self-organization of 1-methylnaphthalene on the surface of artificial snow grains: A combined experimental-computational



- approach, *J. Phys. Chem. A*, 115(41), doi:10.1021/jp205627a, 2011.
- Huang, J. and Jaeglé, L.: Wintertime enhancements of sea salt aerosol in polar regions consistent with a sea ice source from blowing snow, *Atmos. Chem. Phys.*, 17(5), 3699–3712, doi:10.5194/acp-17-3699-2017, 2017.
- 540 Hullar, T. and Anastasio, C.: Direct visualization of solute locations in laboratory ice samples, *Cryosphere*, 10(5), 2057–2068, doi:10.5194/tc-10-2057-2016, 2016.
- Hullar, T., Magadia, D. and Anastasio, C.: Photodegradation Rate Constants for Anthracene and Pyrene Are Similar in/on Ice and in Aqueous Solution, *Environ. Sci. Technol.*, doi:10.1021/acs.est.8b02350, 2018.
- Imrichová, K., Veselý, L., Gasser, T. M., Loerting, T., Neděla, V. and Heger, D.: Vittrification and increase of basicity in
545 between ice I h crystals in rapidly frozen dilute NaCl aqueous solutions, *J. Chem. Phys.*, 151(1), 014503, doi:10.1063/1.5100852, 2019.
- Jourdain, B., Preunkert, S., Cerri, O., Castebrunet, H., Udisti, R. and Legrand, M.: Year-round record of size-segregated aerosol composition in central Antarctica (Concordia station): Implications for the degree of fractionation of sea-salt particles, *J. Geophys. Res.*, 113(D14), D14308, doi:10.1029/2007JD009584, 2008.
- 550 Kahan, T. F. and Donaldson, D. J.: Photolysis of Polycyclic Aromatic Hydrocarbons on Water and Ice Surfaces, *J. Phys. Chem. A*, 111(7), 1277–1285, doi:10.1021/jp066660t, 2007.
- Kaleschke, L., Richter, A., Burrows, J., Afe, O., Heygster, G., Notholt, J., Rankin, A. M., Roscoe, H. K., Hollwedel, J., Wagner, T. and Jacobi, H. W.: Frost flowers on sea ice as a source of sea salt and their influence on tropospheric halogen chemistry, *Geophys. Res. Lett.*, 31(16), L16114, doi:10.1029/2004GL020655, 2004.
- 555 Kania, R., Malongwe, J. K., Nachtigallová, D., Krausko, J., Gladich, I., Roeselová, M., Heger, D. and Klán, P.: Spectroscopic properties of benzene at the air-ice interface: A combined experimental-computational approach, *J. Phys. Chem. A*, 118(35), doi:10.1021/jp501094n, 2014.
- Keene, W. C., Maring, H., Maben, J. R., Kieber, D. J., Pszenny, A. A. P. P., Dahl, E. E., Izaguirre, M. A., Davis, A. J., Long, M. S., Zhou, X., Smoydzin, L. and Sander, R.: Chemical and physical characteristics of nascent aerosols produced by bursting
560 bubbles at a model air-sea interface, *J. Geophys. Res.*, 112(D21), D21202, doi:10.1029/2007JD008464, 2007.
- Klánová, J., Klán, P., Heger, D. and Holoubek, I.: Comparison of the effects of UV, H₂O₂/UV and γ -irradiation processes on frozen and liquid water solutions of monochlorophenols, *Photochem. Photobiol. Sci.*, 2(10), doi:10.1039/b303483F, 2003.
- Krausko, J., Runštuk, J., Neděla, V., Klán, P. and Heger, D.: Observation of a brine layer on an ice surface with an
565 environmental scanning electron microscope at higher pressures and temperatures, *Langmuir*, 30(19), 5441–5447, doi:10.1021/la500334e, 2014.
- Legrand, M., Preunkert, S., Wolff, E., Weller, R., Jourdain, B. and Wagenbach, D.: Year-round records of bulk and size-segregated aerosol composition in central Antarctica (Concordia site) – Part 1: Fractionation of sea-salt particles, *Atmos. Chem. Phys.*, 17(22), 14039–14054, doi:10.5194/acp-17-14039-2017, 2017.




- 570 Levine, J. G., Yang, X., Jones, A. E. and Wolff, E. W.: Sea salt as an ice core proxy for past sea ice extent: A process-based model study, *J. Geophys. Res. Atmos.*, 119(9), 5737–5756, doi:10.1002/2013JD020925, 2014.
- Light, B., Brandt, R. E. and Warren, S. G.: Hydrohalite in cold sea ice: laboratory observations of single crystals, surface accumulations, and migration rates under a temperature gradient, with application to “snowball earth,” *J. Geophys. Res. Ocean.*, 114(7), 1–17, doi:10.1029/2008JC005211, 2009.
- 575 Mason, B. J.: The supercooling and nucleation of water, *Adv. Phys.*, 7(26), 221–234, doi:10.1080/00018735800101237, 1958.
- Maus, S.: Prediction of the cellular microstructure of sea ice by morphological stability theory, in *Physics and Chemistry of Ice*, edited by W. F. Kuhs, pp. 371–382, RSC Publishing., 2007.
- Michaloudi, E., Papakostas, S., Stamou, G., Neděla, V., Tihlaříková, E., Zhang, W. and Declerck, S. A. J.: Reverse taxonomy applied to the brachionus calyciflorus cryptic species complex: Morphometric analysis confirms species delimitations revealed
- 580 by molecular phylogenetic analysis and allows the (re) description of four species, *PLoS One*, doi:10.1371/journal.pone.0203168, 2018.
- Moffat, J. R., Sefiane, K. and Shanahan, M. E. R.: Effect of TiO₂ Nanoparticles on Contact Line Stick–Slip Behavior of Volatile Drops, *J. Phys. Chem. B*, 113(26), 8860–8866, doi:10.1021/jp902062z, 2009.
- Neděla, V.: Methods for additive hydration allowing observation of fully hydrated state of wet samples in environmental SEM,
- 585 *Microsc. Res. Tech.*, 70(2), 95–100, doi:10.1002/jemt.20390, 2007.
- Neděla, V., Tihlaříková, E., Runštuk, J. and Hudec, J.: High-efficiency detector of secondary and backscattered electrons for low-dose imaging in the ESEM, *Ultramicroscopy*, 184, 1–11, doi:10.1016/j.ultramic.2017.08.003, 2018.
- Neděla, V., Tihlaříková, E., Maxa, J., Imrichová, K., Bučko, M. and Gemeiner, P.: Simulation-based optimisation of thermodynamic conditions in the ESEM for dynamical in-situ study of spherical polyelectrolyte complex particles in their
- 590 native state, *Ultramicroscopy*, 211, 112954, doi:10.1016/j.ultramic.2020.112954, 2020.
- O’Dowd, C. D., Lowe, J. A., Smith, M. H. and Kaye, A. D.: The relative importance of non-sea-salt sulphate and sea-salt aerosol to the marine cloud condensation nuclei population: An improved multi-component aerosol-cloud droplet parametrization, *Q. J. R. Meteorol. Soc.*, 125(556), 1295–1313, doi:10.1002/qj.1999.49712555610, 1999.
- Ondrušková, G., Krausko, J., Stern, J. N., Hauptmann, A., Loerting, T. and Heger, D.: Distinct Speciation of Naphthalene
- 595 Vapor Deposited on Ice Surfaces at 253 or 77 K: Formation of Submicrometer-Sized Crystals or an Amorphous Layer, *J. Phys. Chem. C*, 122(22), doi:10.1021/acs.jpcc.8b03972, 2018.
- Ondrušková, G., Veselý, L., Zezula, J., Bachler, J., Loerting, T. and Heger, D.: Using Excimeric Fluorescence to Study How the Cooling Rate Determines the Behavior of Naphthalenes in Freeze-Concentrated Solutions: Vitrification and Crystallization, *J. Phys. Chem. B*, 124(46), 10556–10566, doi:10.1021/acs.jpcc.0c07817, 2020.
- 600 Ozcelik, H. G., Ozdemir, A. C., Kim, B. and Barisik, M.: Wetting of single crystalline and amorphous silicon surfaces: effective range of intermolecular forces for wetting, *Mol. Simul.*, 46(3), 224–234, doi:10.1080/08927022.2019.1690145, 2020.
- Peterson, P. K., Pöhler, D., Sihler, H., Zielcke, J., General, S., Frieß, U., Platt, U., Simpson, W. R., Nghiem, S. V., Shepson,




- P. B., Stirm, B. H., Dhaniyala, S., Wagner, T., Caulton, D. R., Fuentes, J. D. and Pratt, K. A.: Observations of bromine monoxide transport in the Arctic sustained on aerosol particles, *Atmos. Chem. Phys.*, doi:10.5194/acp-17-7567-2017, 2017.
- 605 Pratt, K. A., Custard, K. D., Shepson, P. B., Douglas, T. A., Pöhler, D., General, S., Zielcke, J., Simpson, W. R., Platt, U., Tanner, D. J., Gregory Huey, L., Carlsen, M. and Stirm, B. H.: Photochemical production of molecular bromine in Arctic surface snowpacks, *Nat. Geosci.*, 6(5), 351–356, doi:10.1038/ngeo1779, 2013.
- Qazi, M. J., Salim, H., Doorman, C. A. W., Jambon-Puillet, E. and Shahidzadeh, N.: Salt creeping as a self-amplifying crystallization process, *Sci. Adv.*, 5(12), doi:10.1126/SCIADV.AAX1853/SUPPL_FILE/AAX1853_SM.PDF, 2019.
- 610 Rankin, A. M. and Wolff, E. W.: A year-long record of size-segregated aerosol composition at Halley, Antarctica, *J. Geophys. Res. D Atmos.*, 108(D24), AAC 9-1-AAC 9-12, doi:10.1029/2003jd003993, 2003.
- Ray, D., Kurková, R., Hovorková, I. and Klán, P.: Determination of the Specific Surface Area of Snow Using Ozonation of 1,1-Diphenylethylene, *Environ. Sci. Technol.*, 45(23), 10061–10067, doi:10.1021/es202922k, 2011.
- Rhodes, R. H., Yang, X., Wolff, E. W., McConnell, J. R. and Frey, M. M.: Sea ice as a source of sea salt aerosol to Greenland ice cores: A model-based study, *Atmos. Chem. Phys.*, 17(15), 9417–9433, doi:10.5194/acp-17-9417-2017, 2017.
- 615 Richter, A., Wittrock, F., Eisinger, M. and Burrows, J. P.: GOME observations of tropospheric BrO in northern hemispheric spring and summer 1997, *Geophys. Res. Lett.*, 25(14), 2683–2686, doi:10.1029/98GL52016, 1998.
- Rodebush, W. H.: The freezing points of concentrated solutions and the free energy of solution of salts, *J. Am. Chem. Soc.*, 40(8), 1204–1213, doi:10.1021/ja02241a008, 1918.
- 620 Rohatgi, P. K. and Adams, C. M.: Ice–Brine Dendritic Aggregate formed on Freezing of Aqueous Solutions, *J. Glaciol.*, 6(47), 663–679, doi:10.1017/s0022143000019936, 1967.
- Roscoe, H. K., Brooks, B., Jackson, A. V., Smith, M. H., Walker, S. J., Obbard, R. W. and Wolff, E. W.: Frost flowers in the laboratory: Growth, characteristics, aerosol, and the underlying sea ice, *J. Geophys. Res.*, 116(D12), D12301, doi:10.1029/2010JD015144, 2011.
- 625 Schenk Mayerová, A., Bučko, M., Gemeiner, P., Treľová, D., Lacić, I., Chorvát, D., Ačai, P., Polakovič, M., Lipták, L., Rebroš, M., Rosenberg, M., Štefuca, V., Neděla, V. and Tihlaříková, E.: Physical and Bioengineering Properties of Polyvinyl Alcohol Lens-Shaped Particles Versus Spherical Polyelectrolyte Complex Microcapsules as Immobilisation Matrices for a Whole-Cell Baeyer–Villiger Monooxygenase, *Appl. Biochem. Biotechnol.*, 174(5), 1834–1849, doi:10.1007/s12010-014-1174-x, 2014.
- Schremb, M. and Tropea, C.: Solidification of supercooled water in the vicinity of a solid wall, *Phys Rev E*, 94(5–1), 52804, doi:10.1103/PhysRevE.94.052804, 2016.
- 630 Shirkov, A. A., Golovin, Y. I., Zheltov, M. A., Korolev, A. A. and Leonov, A. A.: Morphology diagram of nonequilibrium patterns of ice crystals growing in supercooled water, *Phys. A Stat. Mech. its Appl.*, 319, 65–79, doi:https://doi.org/10.1016/S0378-4371(02)01517-0, 2003.
- Simpson, W. R., Carlson, D., Hönninger, G., Douglas, T. A., Sturm, M., Perovich, D. and Platt, U.: First-year sea-ice contact predicts bromine monoxide (BrO) levels at Barrow, Alaska better than potential frost flower contact, *Atmos. Chem. Phys.*,
- 635



- 7(3), 621–627, doi:10.5194/acp-7-621-2007, 2007.
- Spiel, D. E.: On the births of jet drops from bubbles bursting on water surfaces, *J. Geophys. Res.*, 100(C3), 4995, doi:10.1029/94JC03055, 1995.
- Vetráková, L., Vykoukal, V. and Heger, D.: Comparing the acidities of aqueous, frozen, and freeze-dried phosphate buffers: Is there a “pH memory” effect?, *Int. J. Pharm.*, 530(1–2), doi:10.1016/j.ijpharm.2017.08.005, 2017.
- Vetráková, L., Neděla, V., Runštuk, J. and Heger, D.: The morphology of ice and liquid brine in an environmental scanning electron microscope: a study of the freezing methods, *Cryosph.*, 13(9), 2385–2405, doi:10.5194/tc-13-2385-2019, 2019.
- Vetráková, L., Neděla, V., Runštuk, J., Tihlaříková, E., Heger, D. and Shalaev, E.: Dynamical in-situ observation of the lyophilization and vacuum-drying processes of a model biopharmaceutical system by an environmental scanning electron microscope, *Int. J. Pharm.*, 585, 119448, doi:10.1016/j.ijpharm.2020.119448, 2020.
- Wagenbach, D., Ducroz, F., Mulvaney, R., Keck, L., Minikin, A., Legrand, M., Hall, J. S. and Wolff, E. W.: Sea-salt aerosol in coastal Antarctic regions, *J. Geophys. Res. Atmos.*, 103(D9), 10961–10974, doi:10.1029/97JD01804, 1998.
- Wettlaufer, J. S.: Directional solidification of salt water: deep and shallow cells, *Eur. Lett.*, 19(4), 337–342, doi:10.1209/0295-5075/19/4/015, 1992.
- Wexler, A.: Vapor pressure formulation for ice, *J. Res. Natl. Bur. Stand.*, 81A(February), 5, doi:10.6028/jres.081A.003, 1977.
- Wise, M. E., Baustian, K. J., Koop, T., Freedman, M. A., Jensen, E. J. and Tolbert, M. A.: Depositional ice nucleation onto crystalline hydrated NaCl particles: a new mechanism for ice formation in the troposphere, *Atmos. Chem. Phys.*, 12(2), 1121–1134, doi:10.5194/acp-12-1121-2012, 2012.
- Wolff, E. W., Rankin, A. M. and Röthlisberger, R.: An ice core indicator of Antarctic sea ice production?, *Geophys. Res. Lett.*, 30(22), CLM 4-1-CLM 4-4, doi:10.1029/2003GL018454, 2003.
- Yang, X., Pyle, J. A. and Cox, R. A.: Sea salt aerosol production and bromine release: Role of snow on sea ice, *Geophys. Res. Lett.*, 35(16), 1–5, doi:10.1029/2008GL034536, 2008.
- Yang, X., Neděla, V., Runštuk, J., Ondrušková, G., Krausko, J., Vetráková, L. and Heger, D.: Evaporating brine from frost flowers with electron microscopy and implications for atmospheric chemistry and sea-salt aerosol formation, *Atmos. Chem. Phys.*, 17(10), 6291–6303, doi:10.5194/acp-17-6291-2017, 2017.
- Yang, X., Frey, M. M., Rhodes, R. H., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Jones, A. E. and Wolff, E. W.: Sea salt aerosol production via sublimating wind-blown saline snow particles over sea ice: parameterizations and relevant microphysical mechanisms, *Atmos. Chem. Phys.*, 19(13), 8407–8424, doi:10.5194/acp-19-8407-2019, 2019.
- Závacká, K., Neděla, V., Olbert, M., Tihlaříková, E., Vetráková, L., Yang, X. and Heger, D.: Temperature and Concentration Affect Particle Size Upon Sublimation of Saline Ice: Implications for Sea Salt Aerosol Production in Polar Regions, *Geophys. Res. Lett.*, 49(8), 1–10, doi:10.1029/2021gl097098, 2022.



670 **Figure 1:** The directionalities of the three freezing methods: (a) Spontaneous freezing (non-seeded sample); (b) controlled ice nucleation (seeding); and (c) freezing in LN. The arrows show the expected directions of the progress of the ice crystal growth front.





675 **Figure 2:** The sublimation process in the 0.5 M non-seeded sample at $-20\text{ }^{\circ}\text{C}$. The images were recorded at the edge of the sample, within intervals of 10 s.






Figure 3: The salt residua after the sublimation of the frozen samples prepared from 0.005 M CsCl via the indicated freezing methods. The samples sublimated at $-20\text{ }^{\circ}\text{C}$ (above the T_{eu}). The scale in panel c applies to all of the images.



680




Figure 4: The salt residua after the sublimation of the frozen samples prepared from 0.05 M CsCl via the indicated freezing methods. The center and the periphery of the samples are displayed. The samples sublimated at $-20\text{ }^{\circ}\text{C}$ (above T_{eu}). The scale in panel f applies to all of the images.




685 **Figure 5:** The salt residua after the sublimation of the frozen samples prepared from 0.5 M CsCl via the indicated freezing methods. The center and the periphery of the samples are displayed. The samples sublimated at $-20\text{ }^{\circ}\text{C}$ (above the T_{eu}). The scale in panel i applies to all of the images.




690 **Figure 6:** A photograph of the cooling stage of the ESEM covered with the crystallized CsCl salt after the sublimation of the 0.5 M non-seeded sample at $-20\text{ }^{\circ}\text{C}$. The diameter of the stage is about 6 mm. The pattern of the concentric circles is nicely visible.




695 **Figure 7:** The salt residues after the sublimation of the frozen samples containing 0.005 M CsCl; the sublimation temperature was $-25\text{ }^{\circ}\text{C}$ (below the T_{eu}). Each row represents one freezing method, named on the left-hand side. The images in the left- and right-hand columns were taken at the periphery and close to the centre of the original sample, respectively. The scale in panel *e* applies to all the images. The examples of the salt rim (blue rectangles), fine salt particles (yellow circles), salt flakes (red dotted circles), and lichen-like tufts (green squares) are highlighted.



700 **Figure 8:** The salt residues after the sublimation of the frozen samples containing 0.05 M CsCl; the sublimation temperature was -25°C (below the T_{eu}). Each row represents one freezing method, named on the left-hand side. The images in the left- and right-hand columns were taken at the periphery and close to the centre of the original sample, respectively. The scale in panel *e* applies to all of the images.



705 **Figure 9:** The retreating ice (black) in the 0.05 M seeded sample reveals tiny salt particles (the white particles encircled in yellow) lying on the cooling stage (grey). These microparticles were most likely formed by the brine crystallizing at the bottom of the sample, in contact with the stage. Conversely, the larger salt flakes (pointed to by the red arrows), which emerge on the surface of the sample before falling down onto the stage, originate from the body and the surface of the sample.



710 **Figure 10.** The retreating ice (black, upper part) in the 0.005 M non-seeded sample reveals salt particles (white) lying on the cooling stage (grey, lower part). The sublimation occurs at $-25\text{ }^{\circ}\text{C}$. The encircled objects are the salt flakes on the ice surface.






Figure 11: The salt flakes (white) falling down the surface of the retreating ice (black, left upper part) onto the cooling stage (grey, right part) in the 0.005 M seeded sample.








Figure 12: The histograms showing the distribution of the maximum particle diameters of the residual salt particles in the non-seeded and seeded samples sublimated at -25°C . The smallest applied interval is $1\ \mu\text{m}$. Each histogram embraces the particles from 3 representative images of the sublimated samples. The data were recorded with the instrumental resolution of $0.5\ \mu\text{m}$.




720


Figure 13: The salt residues after the sublimation of the frozen samples containing $0.5\ \text{M CsCl}$: (a) a non-seeded droplet, (b) a seeded droplet, (c) an LN-frozen droplet, (d) LN-frozen microspheres. The samples sublimated at -25°C (below the T_{ew}). The scale in panel *d* applies to all of the images. The corresponding morphologies are detailed in panels *a* and *c*, where the black bar represents $10\ \mu\text{m}$.




725 **Figure 14:** The supercooled brine beyond the edge of the 0.05 M non-seeded sample (a) and the salt crystals formed via the eventual crystallization of the supercooled brine (b). The original frozen sample had been located in the left side of the panels a and b, its edge is represented by the broader white line (the salt rim).



730 **Figure 15:** The structures of the CsCl salt residua after the evaporation of the 0.005 and 0.05 M liquid samples at 2 °C. We imaged spots close to the periphery and the centre of the original sample, respectively.



735 Figure 16: The structure of the salt residua after the sublimation of the ice, influenced by the sublimation temperature, salt concentration, and freezing method. According to their ability to be windblown and to become aerosols, the salt residua are categorized as follows: readily available (the green box with dotted border), potentially available after their mechanical breakdown (the yellow boxes with dashed borders), improbable (the red box with dash-dotted border).



740 **Figure 17.** The structure of the frozen samples (before the sublimation) and the CsCl salt residues after the sublimation of the 0.005, 0.05, and 0.5 M spontaneously frozen samples at $-25\text{ }^{\circ}\text{C}$, i.e. below T_{eu} .

| | | | | |
|--|-------------------------------------|---------------------------------|------------------------------------|--------------------------------|
| | 0.005 M CsCl, non-seeded | 0.005 M CsCl, seeded | 0.05 M CsCl, non-seeded | 0.05 M CsCl, seeded |
|--|-------------------------------------|---------------------------------|------------------------------------|--------------------------------|



| | | | | |
|---|-------------|---------------|---------------|----------------|
| Surface number density of salt particles / mm ⁻² | 3405 ± 3033 | 12758 ± 7017 | 39200 ± 25032 | 59625 ± 11885 |
| Volume number density of salt particles / mm ⁻³ | 6300 ± 5613 | 23604 ± 12983 | 72528 ± 46314 | 110319 ± 21989 |

Table 1. The calculated surface and volume number densities (mm⁻³) with sample standard deviations under different conditions. The method of the calculation is described in SI (Text S3, Table S1).

745

| Concentration | Freezing method | The occurrence of residual salt particles according to their sizes | | | | |
|---------------|-----------------|--|--------------|---------------|---------------|----------|
| | | 1 to 10 μm | 10 to 100 μm | 100 to 200 μm | 200 to 400 μm | > 400 μm |
| 0.005 M | Non-seeded | *** | *** | | | |
| | Seeded | *** | ** | | | |
| | LN-frozen | | | ** | * | *** |
| 0.05 M | Non-seeded | *** | *** | ** | * | |
| | Seeded | *** | *** | *** | *** | |
| | LN-frozen | | | * | ** | *** |
| 0.5 M | Non-seeded | | | | | *** |
| | Seeded | | | | | *** |
| | LN-frozen | | | | | *** |

Table 2: The distribution of the sizes of the residual salt particles in the samples sublimated at -25 °C. The occurrence of the particles according to their sizes is expressed as follows: *** often; ** occasionally; * seldom.