

Tropical tropospheric ozone and carbon monoxide distributions: characteristics, origins and control factors, as seen by IAGOS and IASI

Maria Tsivlidou¹, Bastien Sauvage¹, Yasmine Bennouna¹, Romain Blot¹, Damien Boulanger², Hannah Clark³, Eric Le Flochmoën¹, Philippe Nédélec¹, Valérie Thouret¹, Pawel Wolff², and Brice Barret¹

¹Laboratoire d'Aérologie (LAERO), Université Toulouse III – Paul Sabatier, CNRS, Toulouse, France

²Observatoire Midi-Pyrénées (OMP-SEDOO), Université Toulouse III - Paul Sabatier, CNRS, Toulouse, France

³IAGOS-AISBL, 98 Rue du Trône, Brussels, Belgium

Correspondence: Maria Tsivlidou (mtsivl@gmail.com)

Abstract. The characteristics and seasonal variability of the tropical tropospheric distributions of ozone (O₃) and carbon monoxide (CO) were analysed based on in situ measurements provided by the In-service Aircraft for a Global Observing System (IAGOS) program since 1994 and 2002 respectively, combined with observations from the Infrared Atmospheric Sounding (IASI) instrument on board the Met-op A satellite since 2008. The SOFT-IO model, which couples back trajectories with CO emission inventories, was used to explore the origins and sources of the tropical CO observed by IAGOS.

The highest O₃ and CO mixing ratios occur over Western Africa in the low troposphere (LT: surface to 750 hPa) during the fire season (75 ppb of O₃ at 2.5 km and 850 ppb of CO at 0.3 km over Lagos in January), mainly due to anthropogenic (AN) emissions, and a major contribution from fires. The secondary maxima are observed in Asia, in mid (MT: 750–300 hPa) and upper (UT: 300–200 hPa) troposphere in April for O₃ and in LT in January for CO, with larger contributions from AN emissions. The lowest O₃ and CO mixing ratios occur over Caracas.

In the tropical LT, the majority of the location clusters are affected by local and regional AN emissions. The highest AN impact is found over Asia, Arabia and Eastern Africa, and South America (> 75 % of CO). Biomass burning (BB) emissions also originate from local or regional sources but with stronger seasonal dependence. The highest BB impact is found over Southern Tropical Africa (57–90 %), except in April, mostly due to local fires, but also from Northern Hemisphere Africa in January (45–73 %) and Southern Hemisphere South America in October (29 % over Windhoek).

In the MT and UT, AN emissions are more important and dominate in the Eastern part of the Tropics (from Middle East to Asia). BB contributions are more important than in the LT, especially from the African fires in January and July, and from South-East and Equatorial Asia in April and October.

The overall highest amount of CO is exported from Africa, with main transport pathway from the dry-season African regions towards the wet-season ones. In contrast, the impact of the Asian emissions in the LT and MT is limited on a local or regional scale. The transport of polluted Asian air masses is important in the UT during the Asian summer monsoon and post-monsoon seasons, when convection is active.

1 Introduction

25 Tropospheric ozone (O_3) and carbon monoxide (CO) are key components in the atmosphere. O_3 has a significant impact on human health close to the surface (Curtis et al., 2006; Jerrett et al., 2009) and on climate by being a powerful greenhouse gas (Gauss et al., 2003; IPCC, 2021). O_3 is a secondary pollutant produced by photochemical oxidation of precursors such as CO and volatile organic compounds (VOCs) in the presence of nitrogen oxides (NO_x) (Logan et al., 1981). Its distribution is controlled by: transport processes at intercontinental and hemispheric scale (Wild et al., 2004); emissions of precursors (natural and anthropogenic); destruction processes (photochemical and depositional) (Monks et al., 2015) and stratospheric transport (Stevenson et al., 2013). Due to its longer lifetime, CO is considered a powerful pollution tracer of combustion products at a hemispheric level (Logan et al., 1981). CO impacts the oxidation capacity of the atmosphere by being the major sink of OH radicals in non polluted atmosphere (Lelieveld et al., 2016), and the climate by producing greenhouse gases, such as CO_2 and O_3 , during its oxidation (Myhre et al., 2013). CO is primarily emitted by incomplete combustion, thus by anthropogenic (AN) (including biogenic fuels) and biomass burning (BB) (wildfires) sources (Galanter et al., 2000; Granier et al., 2011), with contributions between 450–600 and 350–600 Tg CO yr⁻¹ respectively (Lamarque et al., 2010; van der Werf et al., 2006). Its secondary sources include oxidation of VOCs and methane (450–1200 and 600– 1000 Tg CO y⁻¹ resp.; Stein et al., 2014).

Recent studies (Gaudel et al., 2018, 2020; Zhang et al., 2016) have shown increasing tropospheric O_3 burden in the second half of the 20th century mostly due to increase of precursors in the Tropics. Based on aircraft observations, Bourgeois et al. (2020) recently presented a global-scale distribution of O_3 in the remote troposphere. However, uncertainties still remain in the global O_3 distribution and sources of precursors due to paucity of observations in the free troposphere, especially over developing countries in the Tropics (Gaudel et al., 2018; Tarasick et al., 2019).

The Tropics are of particular interest regarding tropospheric O_3 and CO. They combine: i) intense photochemistry due to high UV radiation and humidity, ii) large active natural sources of CO and other O_3 precursors through BB (Ziemke et al., 2009), biogenic (Aghedo et al., 2007) and lightning emissions (Sauvage et al., 2007b, c), iii) increasing AN due to rapid industrialisation (Duncan et al., 2016; Granier et al., 2011), iv) large O_3 net production potential because deep convection can transport surface emissions to higher altitudes, where their lifetime is increased due to lack of surface deposition and dilution with unpolluted background (Pickering et al., 1995) and v) dynamic processes capable of redistributing chemical species in a regional and global scale (Zhang et al., 2016). Thus, the Tropics are a region where O_3 production is favoured.

50 Satellite observations from the OMI and MLS sensors (Ziemke et al., 2019) and simulations from the GEOS-Chem chemical transport model (Zhang et al., 2016, 2021) display the highest O_3 burden increase in the Tropics - mostly over India, East Asia and SouthEast Asia. Most studies tend to confirm an increase of O_3 in the Tropics but they are mostly based on model simulations, sparse ground observations or satellite data with little consistency, and it is not clear what can cause such an increase. Indeed the trends are attributed to different factors such as BB (Heue et al., 2016), dynamics (Lu et al., 2019; Thompson et al., 2021), or AN (Zhang et al., 2016; Gaudel et al., 2020). Thus, further investigation based on in situ observations is required in

order to better constrain models, validate satellite retrievals, and reduce the uncertainties in the quantification of O₃ and CO trends and source attribution over the Tropics.

Measurements of tropical O₃ and CO are available by satellite observations, but they have a coarse vertical resolution (e.g. Barret et al., 2008; Thompson et al., 2001). On short time scales, several field campaigns have been carried out in the Tropics, yielding measurements of various species over Africa (from TROPOZ 1987 to CAFE-Africa), Asia (from INDOEX to EMerge-Asia), South America (Cite-1/2/3, TROCCiNOX) and the tropical Pacific (from PEM-WEST-A/B to CAST/CON-TRAST/ATTREX and Atom). These campaigns have provided invaluable insights on the atmospheric chemistry and dynamics of the Tropics. On greater timescales, the Southern Hemisphere ADditional OZone Sounding (SHADOZ) program (Thompson et al., 2003a) provides long-term O₃ observations over the Tropics using ozonesondes since 1998. These measurements have allowed a better understanding on the vertical distribution and trends of tropical O₃ (e.g. Thompson et al., 2021).

In a complementary way to these datasets, the IAGOS (In-service Aircraft for a Global Observing System) (Marenco et al., 1998; Petzold et al., 2015; Thouret et al., 2022) program has provided O₃ and CO measurements over the Tropics since 1994 and 2002 respectively. Using equipped commercial aircraft, IAGOS samples vertical profiles at take off and landing, along with the lower part of the upper tropical troposphere at cruise altitude between 300 and 185 hPa (UT_{cruise}). Previous studies have documented the tropical composition over Africa (Lannuque et al., 2021; Sauvage et al., 2005, 2007a, d), South America (Yamasoe et al., 2015) and South Asia (Sahu et al., 2014; Sheel et al., 2014) using IAGOS data. However, they are focused on specific regions of the Tropics and have limited temporal coverage, especially for CO as fewer measurements were available at this time. Thus, the O₃ and CO distributions and their interlocking in the entire Tropics are still not well documented.

The SOFT-IO model (Sauvage et al., 2017) has been developed to supplement the analysis of the IAGOS dataset by estimating AN and BB contributions to the observed CO measurements. These measurements, along with the SOFT-IO output allow us to trace the CO origin over the Tropics. Further, global distributions provided by Infrared Atmospheric Sounding Interferometer (IASI)-Software for a Fast Retrieval of IASI Data (SOFRID) (Barret et al., 2011; De Wachter et al., 2012) retrievals since 2008 complement the O₃ and CO distributions provided by IAGOS. They allow us to understand the spatial extent of pollution plumes, and explore intercontinental transport patterns.

In this article we take advantage of the unique IAGOS database to (i) document the characteristics and seasonal variability of these two atmospheric species over the whole tropical band for the last decade, (ii) explore the origin of the observed CO anomalies, (iii) investigate transport processes driving the CO and O₃ distributions in the Tropics.

The observational (IAGOS and IASI) and model based (SOFT-IO) datasets, and methodology are introduced in Sect. 2. In Section 3, the IAGOS observations are analysed to document O₃/CO vertical profiles, along with the upper tropospheric composition over the Tropics. In addition, the sources of observed CO are explored with SOFT-IO.

2 Data and Methods

2.1 IAGOS observations

The Research Infrastructure IAGOS (Petzold et al., 2015; Thouret et al., 2022) provides in situ measurements of trace gases (O₃, CO, water vapour, NO_y between 2001 and 2005 (e.g. Gressent et al., 2014), and more recently NO_x, CH₄, CO₂ and cloud particles, see <https://www.iagos.org/iagos-data/>) and meteorological parameters (temperature and winds), using equipped commercial aircraft. Full description of the instruments can be found in Nédélec et al. (2015). O₃ (resp. CO) is measured using a dual-beam ultraviolet absorption monitor (resp. infrared analyser) with an overall uncertainty of ± 2 ppbv ± 2 % (resp. ± 5 ppbv ± 5 %) and a time resolution of 4 (resp. 30) seconds (Nédélec et al., 2015). IAGOS measures vertical profiles during ascend and descend phases, and the UT_{cruise} during cruise phases. Considering the aircraft speed (7–8 m s⁻¹ during ascent/descent; 900 km h⁻¹ during cruise), the time resolution of the instruments corresponds to a vertical resolution of 30 m (resp. 225 m), and a horizontal resolution of 1 km (resp. 7.5 km) for O₃ (resp. CO).

O₃ (resp. CO) observations have been collected since 1994 (resp. 2002) in the frame of the IAGOS Research Infrastructure and its predecessor MOZAIC (Marengo et al., 1998) program, based on the same instrument technologies. Good consistency in the measurements between the two programs (hereafter referred to as IAGOS) (Blot et al., 2021; Nédélec et al., 2015) leads to IAGOS temporal coverage of 26 (resp. almost 20) years for O₃ (resp. CO), depending on the availability of the flights. IAGOS data provides robust O₃ and CO climatologies, allowing studies of long-term trends (e.g. Cohen et al., 2018) along with validation of chemistry transport models (e.g. Gressent et al., 2016; Sauvage et al., 2007b) and satellite data retrievals (e.g. De Wachter et al., 2012) on a global scale. To complement the IAGOS observations, we use the potential vorticity (PV) field, which is part of the ancillary data (<https://doi.org/10.25326/3>) from the IAGOS database. The PV is calculated from the European Centre for Medium-Range Weather Forecast's (ECMWF) operational fields (horizontal resolution 1°, time resolution 3 hours), interpolated along IAGOS trajectories.

2.1.1 Data treatment

The tropical zone can be defined in several ways, such as by meteorological characteristics (e.g. location of the subtropical jets), climatic elements (e.g. precipitation rates) or by the geographical extent. Following the latter way, in the Tropospheric Ozone Assessment Report, Phase II (TOAR-II; <https://igacproject.org/activities/TOAR/TOAR-II>) Ozone and Precursors in the Tropics working group, the Tropics are defined between 20° S and 20° N. In our study, we consider the extended area between 25° S and 25° N, in order to investigate interactions of pollution and the transport of air masses between the Tropics and the Subtropics. Only tropospheric measurements are taken into account, by applying a PV filter of 2 PV units (pvu) for each measurement during cruise phase, and for the measurements between 20–25° N/S during ascend/descent. The UT_{cruise} climatologies are derived by averaging the cruise data on a 2 x 2.5° grid, for the period 1994–2020 (resp. 2002–2020) for O₃ (resp. CO). For the same time periods, the climatologies over the vertical are derived by averaging the data into 10 hPa pressure bins from the surface up to 200 hPa. Data are selected within a 300km radius circle centered on the airport location, similar

to Petetin et al. (2016). This way we reduce uncertainties due to possible horizontal heterogeneity in the measurements, as the aircraft keeps moving in the horizontal plane during ascent and descent.

120 Table 1 shows the total number of profiles taken into account for this study for each location for the whole IAGOS period. The temporal availability of the measurements differ for each location and cluster, as it depends on the flight schedule of the aircraft (see Fig. S1 and Fig. S2). For this reason, to determine a reliable climatological profile, we need to assess the statistical significance of the data. Similar to Logan (1999) and Sauvage et al. (2005), we compute the relative standard error (RSE) of the O₃ (CO) monthly mean, versus the number of flights per month. The RSE is defined as the ratio between the standard error ($SE = \frac{\sigma}{\sqrt{N}}$, with σ the square root of the sample variance and N the number of flights) and the O₃ (CO) monthly mean. The minimum number of flights required for statistical significance corresponds to the number above which $RSE \leq 10\%$. We choose RSE less than 10 %, because RSE depends not only on the number of measurements, but also on the O₃ (CO) variability which is high over the Tropics (Thompson et al., 2003b). For each location with an adequate number of flights per month, we consider an individual profile of O₃ (CO). Otherwise, we combine locations in clusters, in order to increase the amount of data and get statistically significant climatological profiles. Besides, the clusters can be useful for validation of models with a coarse horizontal resolution, because they represent a wider area as resolved by the models, which are not expected to capture small-scale variations in the ozone field (e.g. Emmons et al., 2010). For clustering, the locations should be: i) in relatively close distance from each other, ii) governed by similar meteorological conditions, and iii) display similar characteristics in the vertical distribution of O₃ (CO) (see Sect. 3.1).

135 The meteorological conditions in the Tropics are peculiar, with different seasonal patterns depending on the region. For instance, in Africa the main seasons are two (dry and wet) with two intermediate seasons passing from wet to dry and vice versa (Lannuque et al., 2021; Sauvage et al., 2005). On the other hand, in Asia the seasons are defined by the Asian monsoon phases: Asian summer monsoon (wet season); Asian winter monsoon (dry season) and post monsoon. Thus, we considered it more appropriate for our analysis to deviate from the classical definition of the seasons, which fits better to studies concerning higher latitudes. Instead, we analyse the O₃/CO profiles and horizontal distributions for months during the peak tropical seasons (January, April, July and October), to highlight seasonal patterns.

2.2 SOFT-IO model

The SOFT-IO (SOft attribution using FlexparT and carbon monoxide emission inventories for In-situ Observation database) tool (Sauvage et al., 2017; <http://dx.doi.org/10.25326/2>) has been developed to investigate the origin of the observed IAGOS-
145 CO, by coupling FLEXPART 20-days backward transport simulations with emission inventories. For each point of IAGOS trajectory, SOFT-IO estimates the CO contribution coming from 14 different geographical regions (see Fig. 1), for AN and BB origin separately. We use Community Emissions Data System (CEDS2) AN emissions (McDuffie et al., 2020) and the Global Fire Assimilation System (GFAS) BB emissions (Kaiser et al., 2012) which include fire injection heights, to discriminate sources of CO over different regions of interest. For the calculations, the AN (resp. BB) emissions are updated on a monthly
150 (resp. daily) basis. The meteorological fields are based on 1° x 1° ECMWF analysis and forecast with a time resolution of 6 and 3h respectively.

Table 1. Description of individual locations and clusters used in this study. The locations are displayed in Fig. 1. P O₃ and P CO indicate the total number of profiles for the IAGOS period.

Individual locations/Clusters		IAGOS locations	P O ₃	P CO
South America	South Brazil	Rio de Janeiro (Brazil), São Paulo (Brazil)	518	62
	Caracas	Caracas (Venezuela)	414	248
	Bogota	Bogota (Colombia)	190	142
Western Africa	Lagos	Lagos (Nigeria)	311	199
	Sahel	Abuja (Nigeria), Ouagadougou (Uganda), Niamey (Niger)	193	202
	Gulf of Guinea	Lome (Togo), Yaounde (Cameroon), Douala (Cameroon), Libreville (Gabon), Accra (Ghana), Abidjan (Ivory Coast), Malabo (Equatorial Guinea) Cotonou (Benin), Port Harcourt (Nigeria)	414	302
Southern Tropical Africa	Central Africa	Luanda (Angola), Brazzaville (Congo) Kinshasa (Democratic Republic of Congo)	169	89
	Windhoek	Windhoek (Namibia)	651	692
Arabia and Eastern Africa	AbuDhabi	Abu Dhabi (United Arab Emirates), Muscat (Oman)	118	56
	Khartoum	Khartoum (Sudan)	157	116
	Addis Ababa	Addis Ababa (Ethiopia)	121	110
	Jeddah	Jeddah (Saudi Arabia)	154	108
Asia	South China	Hong Kong (China), Guangzhou (China), Xiamen (China)	337	562
	Gulf of Thailand	Kuala Lumpur (Malaysia), Singapore (Singapore)	162	140
	Madras	Madras (India)	239	253
	Hyderabad	Hyderabad (India)	159	170
	Mumbai	Mumbai (India)	61	29
	Manila	Manila (Philippines)	66	101
	Bangkok	Bangkok (Thailand)	526	336
	Ho Chi Minh City	Ho Chi Minh City (Vietnam)	113	90

SOFT-IO estimates the contribution to anomalies in CO mixing ratios i.e. CO emitted by primary sources during the last 20 days, while it does not calculate the background CO. The background CO can be emitted by primary sources older than 20 days, and by secondary sources such as oxidation of methane and non-methane volatile organic compounds. In our study, CO anomalies (observed by IAGOS) are defined as the positive difference between the observed and the background CO mixing ratio. The background CO mixing ratio represents a reference value, not affected by surface emission or pollution events. For this reason, it is computed as the monthly climatological median CO of a remote area away from polluted regions, in the UT_{cruise} (during the whole study period 2002–2020). Overall in the Tropics, depending of the month and region, a 80 ppb to 100 ppb background has to be added to SOFT-IO for direct comparison with IAGOS observations.

Several studies (e.g. Cussac et al., 2020; Lannuque et al., 2021; Petetin et al., 2018b) have used SOFT-IO to track back the sources of CO measured by IAGOS. Sauvage et al. (2017) validated SOFT-IO performance against IAGOS CO observations for several regions and tropospheric levels. As detailed in their study, SOFT-IO uncertainties and biases are mostly due to uncertainties in emission inventories, and to a lesser extent to uncertainties concerning the meteorological fields and FLEXPART transport parameterizations (turbulence, convection). Their results show that SOFT-IO can simulate 95 % of the observed number of anomalies, without any strong dependence on altitude or region of the CO plume. SOFT-IO captures the intensity of CO anomalies with a bias lower than 10-15 ppb with respect to observed anomalies for most of the regions and tropospheric layers. The bias is higher in extreme pollution events and might be related to uncertainties in the emissions inventories.

We performed an evaluation of SOFT-IO for the low (LT: surface–750 hPa), mid (MT: 750–350 hPa), and upper (UT: 300–200 hPa) troposphere. Our simulations detect CO anomalies at the same rates as Sauvage et al. (2017). On average, SOFT-IO underestimates the observed CO anomalies by 10 ppb in the MT and UT, and by 45 ppb in the LT. When looking at the differences between IAGOS and SOFT-IO anomalies (taking into account the background not simulated by the model), SOFT-IO reproduces on average more than 93% of the observed anomalies for the 3 layers and 87% in the LT. April is the month where the model gives the largest underestimation, with 78% of the anomalies simulated on average over the 20 clusters (13% standard deviation). The lowest performance is in the LT of Sahel in January, with 52% of the anomaly simulated by SOFT-IO. Another source of uncertainty comes from the definition of background CO. To assess this source of uncertainty, we used an alternative definition for the background calculated as the median CO mixing ratio between 600 and 300 hPa for each location. The differences between this background and the one used in the study are within 2.5-60 ppbv. Nevertheless, using the alternative background did not make any difference in the anomaly source attribution and in the relative contributions.

2.3 IASI-SOFRID observations

The IASI sensor onboard MetOp-A (launched in 2006) has a 12 km footprint at nadir and a 2200 km swath allowing an overpass twice daily at 9:30 and 21:30 local solar time. IASI provides information for the atmospheric composition e.g. content of trace gases such as O_3 (Barret et al., 2011; Boynard et al., 2016; Eremenko et al., 2008), CO (De Wachter et al., 2012; George et al., 2009) and N_2O (Barret et al., 2021). We use O_3 (v3.5) and CO (v2.1 and v2.2 after 2014) IASI retrievals performed with SOFRID (Barret et al., 2020; De Wachter et al., 2012).




Figure 1. Tropical locations served by IAGOS, and geographical source regions used in SOFT-IO model. BONA: Boreal North America; TENA: Temperate North America; CEAM: Central America; NHSA: Northern Hemisphere South America; SHSA: Southern Hemisphere South America; EURO: Europe; MIDE: Middle East; NHAf: Northern Hemisphere Africa; SHAF: Southern Hemisphere Africa; BOAS: Boreal Asia; CEAS: Central Asia; SEAS: SouthEast Asia; EQAS: Equatorial Asia; AUST: Australia and New Zealand.

185 SOFRID- O_3 v3.5 retrievals use a dynamical a priori profile based on latitude, season and the tropopause height (Barret
 et al., 2020). In the Tropics, where the surface temperature, thermal contrast and tropopause height are the highest, SOFRID- O_3
 retrievals allow two independent pieces of information, one in the troposphere and one in the UTLS (Barret et al., 2020).
 Comparisons with ozonesonde measurements for the period 2008–2017, showed that SOFRID- O_3 is biased low in the tropical
 troposphere and UTLS, by $3 \pm 16 \%$ and $12 \pm 33 \%$ respectively in the Northern Tropics ($0-30^\circ N$), and by $8 \pm 14 \%$ and 21
 190 $\pm 30 \%$ in the Southern Tropics ($0-30^\circ S$) (Barret et al., 2020). The measurements above Northern Africa are erroneous, due
 to retrieval problems in the presence of desert ground with sand emissivity interfering with the O_3 signature (Boynard et al.,
 2018). The stripes along the 10° latitude bands in IASI O_3 maps (Fig. 2 e–h and q–t) are due to the use of a dynamical a priori
 profile, resulting in discontinuities between adjacent latitude bands with different a priori profiles. Nevertheless, the use of a
 dynamical a priori profile largely improves the retrieved O_3 profiles in terms of biases, variability and correlation relative to
 195 the previous version based on a single a priori profile (Barret et al., 2020).

For SOFRID-CO (v2.1 and 2.2 after 2014), two independent pieces of information are provided in the low and upper tropo-
 sphere (De Wachter et al., 2012). IASI correctly captures the seasonal variability of CO over southern Africa (Windhoek) and
 European mid-latitudes (Frankfurt) in the low (resp. upper) troposphere relative to IAGOS data (resp. correlation coefficients
 of 0.85 (0.70)). At Windhoek, SOFRID-CO is biased low in the low (resp. upper) troposphere by $13 \pm 20 \%$ (resp. $4 \pm 12 \%$).

200 We use monthly averaged SOFRID-CO and O₃ retrievals on a 1° x 1° grid from 2008–2019. We focus on daytime measurements when larger thermal contrast between the surface and the atmosphere results in increased sensitivity of the instrument (Clerbaux et al., 2009) and on pressure levels corresponding approximately to the independent pieces of information: low troposphere defined between 900–700 hPa (LT_{IASI}), mid troposphere between 600–400 hPa (MT_{IASI}) and upper troposphere between 290–220 hPa (UT_{IASI}).

205 3 Results

3.1 Regional characteristics of tropical O₃ and CO

Figure 2 displays the horizontal distributions of CO in LT_{IASI} (a–d), UT_{IASI} (i–l) and UT_{cruise} (q–t), and of O₃ in the MT_{IASI} (e–h), UT_{IASI} (m–p) and UT_{cruise} (u–x). The results shown in Fig. 2 motivated our choice in combining IAGOS locations in clusters when it is necessary to increase the number of measurements.

210 Figures 3, 5, 6, 7 (panels 1 and 2) display the monthly average vertical distributions of O₃ and CO based on IAGOS data, for the African, Asian, South American, Arabian and Eastern African clusters. Panels 3 to 5 represent the mean recent contribution to CO mixing ratios anomalies (above the background) from AN and BB emissions as estimated by SOFT-IO, with information about their geographical origin (see Sect. 2.2 and Fig. 1). To better summarise O₃ and CO anomalies, Fig. 4 displays the CO contributions for the most representative clusters in three tropospheric layers related to different dynamical regimes: LT corre-
215 sponding roughly to planetary boundary layer; MT above, and UT corresponding to the beginning of convective detrainment.

3.1.1 Africa

The striking feature of CO and O₃ over the African tropical clusters (Sahel, Gulf of Guinea, Lagos and Central Africa) is the low tropospheric maxima during the dry season (January for the Northern Hemisphere and July for the Southern Hemisphere) below 4km (Fig. 3 panels 1–2 and Fig. S3 panels 1–2). Above CO and O₃ display mixing ratio close to the background.
220 Enhanced O₃ and CO are confined in the low troposphere due to the stability of dynamical circulations which prevent vertical mixing (Sauvage et al., 2005). CO maximises close to the surface, with larger mixing ratios over Lagos (850 ppb) than Sahel (500 ppb), Gulf of Guinea or Central Africa (400 ppb). O₃ displays a vertical gradient from the minimum at the surface, likely related to surface deposition and titration by highly concentrated nitrogen oxide (Monks, 2005), to the maximum near 2.5km, likely indicating chemically processed air masses. At Windhoek, O₃ and CO have different vertical distributions, with maxima
225 in October at the end of the Southern Hemisphere dry season (Fig. 3 panels 1c and 2c). Windhoek CO enhancement has the smallest magnitude among the African clusters (150 ppb at 4 km), while the O₃ peak is among the largest, reaching 80 ppb in the UT (11.5 km).

The annual minima of CO and O₃ over the African clusters occur during the transition from the wet to the dry season (April and October), when the local fires are minimum (Fig. S4b and d).




Figure 2. Monthly mean low tropospheric IASI CO distributions (900–700 hPa; a–d), mid tropospheric IASI O₃ distributions (600–400 hPa; e–h), upper tropospheric CO and O₃ distributions based on IASI (290–220 hPa; i–l and m–p resp.) and IAGOS (300–185 hPa; q–t and u–x resp.).




Figure 3. IAGOS monthly mean O₃ (panel 1) and CO (panel 2) vertical distributions. The annotated numbers correspond to the total number of flights per month for the IAGOS period, given in the same colour as in the legend. The shadowed part corresponds to ± 1 standard deviation. Vertical distribution of CO contributions (in ppb) (panels 3 to 5) with the hatched part showing AN contribution, and the non-hatched part BB contribution. For clarity reasons the CO contribution for Sahel and Gulf of Guinea are displayed in Fig. S6.




Figure 4. Mean SOFT-IO contributions (in ppb), averaged over all the positive CO anomalies for the tropical locations (Caracas, Lagos, Central Africa, South China and Bangkok) for LT (a), MT (b) and UT (c) for the AN (hatched) and BB (unhatched) contributions. Each pie corresponds to a different month and each group of four pies refer to a different location (see panel b).

230 Regional fires generally dominate CO contributions in the troposphere of Central Africa and Windhoek (Fig. 3 panel 4 and 5; Fig. 4 and Fig. S5). For the clusters located over Western Africa (Sahel, Gulf of Guinea and Lagos), anthropogenic emissions dominate, but during fire seasons, the contribution from fires can be close to that from anthropogenic sources (for instance in January for the 3 clusters, in July for Lagos and Sahel) (Fig. 3 panel 3; Fig. 4 and Fig. S6 panels 1 and 2).

235 In January the Northern Hemisphere African fires contribution reaches 53 % over the Gulf of Guinea and also influences Central Africa with maximum contribution between 2 and 4km. During transport, the air masses impacted by BB are chemically processed resulting in the formation of an O₃ secondary maximum of 50 ppb coincident with the CO maximum (Fig. 3 panels 1d and 2d), as described in Sauvage et al. (2005). During this month, the AN contribution dominates over Lagos (58%) and Sahel (57%).

240 In July the Southern Hemisphere African fires are responsible for the CO anomalies over Central Africa in the LT and MT by up to 90% (Fig. 4a and b), and also influence Sahel, Gulf of Guinea and Lagos, accounting for up to 68 ppb in the LT of Lagos (Fig. S6 panels 1c and 2c; Fig. 4a and Fig. 3 panel 3c). O₃ LT mixing ratio is slightly higher over Central Africa (85 ppb) in July than over Northern Africa (in January) likely indicating rapid photochemical O₃ production by BB precursors (Singh et al., 1996) during the Southern Hemisphere fires.

245 During each previous dry season, BB air masses transport from the fire hemisphere (Fig. S4a and c) (where the highest CO is measured) to the opposite hemisphere is allowed either by the north-easterly Harmattan flow (January) or the south easterly winds and monsoon flow (July) (Sauvage et al., 2005). This transport is visible with IASI LT_{IASI} CO (Fig. 2a and c) and seems to be more effective in the south-to-north direction (in July) than in the opposite direction (in January).

250 In October CO peaks over Windhoek because fires are localized further south in Africa during this month, contributing to 68 % versus 12 % for AN (Fig. S5a), but also because of fire contribution from South America (20 %) (Fig. 3 panel 5d and Fig. S5a). These high CO mixing ratios in October are detected by IASI in the LT_{IASI} (Fig. 2d) over the whole of Southern Africa reflecting the large extent of the impact of these emissions. The South American fires contribution increases with height (30 % contribution in MT and 50 % in UT) contributing to the O₃ maximum observed in the UT (Fig. S5b and c) (Sauvage et al., 2006), also observed at 8km over Central Africa (80 ppb) (Fig. 3 panel 1b), both part of the South Atlantic maximum visible on IASI UT_{IASI} (Fig. 2p). This maximum is also explained by intense lightning activity (Sauvage et al., 2007b, c).

255 In April, during the transition from dry to wet season, when the fires are reduced (Fig. S4b), secondary CO and O₃ maxima (Fig. 3 panels 1 and 2 and Fig. S3) are observed below 4 km over the Western African clusters (Lagos, Sahel and Gulf of Guinea). These high CO mixing ratios in April are detected by IASI in the LT_{IASI} (Fig. 2b) over the whole of Western Africa. The annual CO surface maximum in Central Africa occurs also in April. LT CO is attributed by more than 85% to local AN emissions (Fig. 3 panel 4b and Fig. 4a) when few fires are detected by MODIS (Yamasoe et al. (2015); their Fig.7). The measured CO anomaly reaches +200 ppb (after removing the background from the observed CO), while SOFT-IO attributes 40 to 80 ppb to AN. This indicates that African AN emissions might be underestimated. The enhanced LT O₃ indicates possible O₃ formation during the transport of the aforementioned emissions towards Sahel and the Gulf of Guinea, but could also be explained by enhanced NO_x mixing ratios, possibly due to soil emissions when rains start (Saunois et al., 2009), especially

over Sahel (Fig. 3 panels 1a–c) because of higher NO₂ mixing ratios above dry savannas than over wet savannas and forests
265 (Adon et al., 2010).

3.1.2 Asia

Similarly to Africa, Asian clusters present CO maximum in the LT, but below 2km, and with lower O₃ enhancement. Above
2km, CO is characteristic of the background, while O₃ presents a strong seasonal variability (Fig. 5 panels 1 and 2 and Fig.
S3).

270 Overall, AN emissions largely dominate CO contributions over fire ones, throughout the troposphere (Fig. 5 panels 3–5 and
Fig. S6 panels 3–7). The dominant AN region is South-East Asia, followed by Central-East Asia which has a great influence
over the South China cluster. Biomass burning from South-East and Equatorial Asia bring a substantial contribution in April
and October respectively, but less important than the AN one.

In January, the annual CO maximum observed close to the surface for all the clusters (except over the Gulf of Thailand, Fig.
275 5 panel 2 and Fig. S3 panel 2) are related to local AN emissions (Fig. 5 panels 3a–5a and Fig. S6 panels 3a–7a) due to the low
boundary layer height in winter. CO maxima are coincident with O₃ enhancements also observed by IASI (Fig. 2f), and due to
the chemical ageing favoured by the confinement of the CO-rich air masses due to the large-scale subsidence (Lelieveld et al.,
2001) and by the cloud free conditions. Central-East Asian emissions mainly control LT CO anomalies for South China, Gulf
of Thailand and Manila, from 52 % to 95 % (over Gulf of Thailand and South China resp., Fig. 4a; Fig. S5a and Fig. S7a),
280 due to their advection by the northeasterly trade winds. South-East Asian emissions dominate over the Indian clusters with
an 85% contribution (Fig. S7a). As in Africa, the CO-rich air masses accumulated in the LT are accompanied by a secondary
LT O₃ maximum, significantly lower (40–60 ppb) than the Western African ones (65–75 ppb). It might be because the O₃
enhancement over Asia is caused by AN-polluted air, while in Africa by mixed (AN and BB) polluted air. During the Atom
campaign, Bourgeois et al. (2021) found that O₃ levels are more enhanced in mixed air pollution, because they are associated
285 with greater NO_x and peroxy acyl nitrates (a NO_x reservoir compound), and thus increased O₃ production, in comparison to
BB- or AN-polluted air alone.

In April during the pre-monsoon season, CO and O₃ are both enhanced above the boundary layer and below 4 to 6 km over
most locations (Fig. 5 panels 1 and 2 and Fig. S3). Local South-East Asian AN emissions dominate CO anomalies over the
majority of the locations (Fig. 5 panels 3–5 and Fig. S6). South-East Asian fires, especially located in the region of Myanmar,
290 Northern Thailand and Laos (Fig. S4b), also contribute over South China, Bangkok, Ho Chi Minh City and Manila by at
least 20–30 % in the LT and MT (Fig. 4 and Fig. S7), leading to large CO mixing ratios captured by IASI (Fig. 2b). The BB
outflow towards South China and the Pacific Ocean is due to westward low and mid tropospheric winds (Fig. S8b and f). The
enhanced MT O₃ is attributable to the intense solar radiation associated with precursors from AN and BB emissions which
were previously evidenced. This is in agreement with the observed O₃ maximum in spring over South China (Dufour et al.,
295 2010) and Bangkok (Sahu et al., 2013). Mumbai is the only location showing a more distant influence of AN emissions, from
the Middle East (36%) in the LT, Northern Hemisphere Africa (30%) in the MT and Northern Hemisphere Africa (AN and




Figure 5. Same as Fig. 3 for the Asian clusters.

BB) in the UT (54 %) (Fig. S7). This transport from Africa towards the Arabian Sea and South India is captured by IASI (Fig. 2j and n), suggesting O₃ photochemical production during the transport.

In October, during the post-monsoon season, either BB or AN can dominate depending on the cluster and the altitude. South-East Asian AN emissions dominate the entire troposphere of the Indian clusters (Fig. S7), while Central-East Asia AN emissions dominate over South China (Fig. 5 panel 4d and Fig. 4). A mix of South-East and Central Asian AN emissions have the largest influence over Bangkok, Ho Chi Minh City (LT and MT) and Manila (LT) (Fig. 4 and Fig. S7). Equatorial Asian BB emissions, principally from Indonesia (Fig. S4d), dominate CO anomalies over the entire troposphere of the Gulf of Thailand (Fig. S5), and also over Madras and Ho Chi Minh City (UT) and Manila (MT and UT) (Fig. S7). The same BB has a substantial influence (10 % on average) over the entire troposphere of South China and Bangkok (Fig. 4), and in the LT of Hyderabad and Manila (Fig. S7). BB CO-rich air masses advection towards the South East Asian coastal clusters is captured by IASI in the LT (Fig. 2d) and wind fields (Fig. S8d). The collocated O₃ enhancement (below 2 km) over these clusters (Fig. 5 panel 1) indicates O₃ production during transport. The Equatorial Asian BB contribution generally intensifies in the UT (40–57 %) (Fig. 4; Fig. S5 and Fig. S7), and is also captured by IASI and IAGOS (Fig. 2). Livesey et al. (2013) also found an upper tropospheric CO maximum over Indonesia and attributed it to episodically strong convection, in agreement with the low OLR in Fig. 2p. In contrast, the UT_{IASI} and MT_{IASI} O₃ distributions show a SE-NW gradient (Fig. 2) with lower O₃-levels over the Maritime continent and the southern Indian Ocean and higher ones over India and the Arabian Sea, as a result of convection over the first region and subsidence of precursor over the second one (Barret et al., 2011).

In July, during the monsoon season, LT O₃ and CO mixing ratios are minima for all the Asian clusters (Fig. 5 panels 1 and 2 and Fig. S3). The reversal of the north-easterly trades to the monsoon flow (Fig. S8c) results in the advection of O₃- and CO-poor air masses from the Indian Ocean towards Asia. Furthermore, cloudy conditions reducing solar radiation and promoting the scavenging of O₃ precursors result in lower O₃ production than in clear sky conditions (Mari et al., 2000; Safieddine et al., 2016). The steep CO gradient close to the surface (below 1 km) indicates the convective uplift of polluted boundary layer air masses towards the UT. The resulting enhancement of CO in the UT within the Asian monsoon anticyclone (AMA) analysed in Park et al. (2008) and Barret et al. (2016) is clear from IASI (Fig. 2). In contrast, the positive South-North O₃ gradient between the Maritime continent and north South Asia and the Middle East (Fig. 2) is likely associated with the photochemical ageing of air masses while they are recirculating towards Middle East (Lawrence and Lelieveld, 2010), and with the high insolation over the Middle East (Barret et al., 2016).

3.1.3 South America

South American clusters present CO maximum in the LT, similarly to African and Asian ones (Fig. 6 panel 2). O₃ is increasing from the surface to the MT/UT (Fig. 6 panel 1), probably due to the lack of depositional and chemical sinks above the LT, in combination with lower water vapour mixing ratios, and lightning emissions, as discussed in Sec. 3.1.1. Local AN emissions generally dominate the CO anomalies, either from North Hemisphere South America (Caracas and Bogota) or from South Hemisphere South America emissions (SBrazil) (Fig. 6 panels 3–5). Local (South America) and intercontinental (Africa) emissions due to fires make a significant contribution, which can reach or exceed the one of AN emissions.

In April CO is maximum in the LT (Fig. 6 panel 2), reaching 350 ppb below 1.5 km over SBrazil, 400 ppb below 1km over Bogota and 200 ppb over Caracas below 2 km. For the last two locations CO anomalies are related to a mix of AN (35 %) and BB (32 %) emissions from Northern Hemisphere South America (Fig. 4 and Fig. 6 panels 3b and 4b), corroborated by the elevated IASI LT_{IASI} CO (Fig. 2b) collocated with the strong AN emissions above Colombia and Venezuela, and active
335 fires above the latter (Fig. S4b and Fig. S9b). 20 % of the anomalies are caused by transport of BB Northern Hemisphere African emissions (Fig. 4a). The O_3 maximum collocated with the CO one at 2 km over Caracas (Fig. 6 panel 1), indicates O_3 production during transport of Northern Hemisphere African air masses impacted by BB, while the second one above 5 km is attributed by Yamasoe et al. (2015) to local AN sources followed by lightning. Contrary to Caracas and Bogota, SBrazil annual CO maximum is mostly due to local AN emissions located over the southern part of Brazil (Fig. S9b). Only 65 ppb of
340 the 260 ppb observed CO anomalies are simulated (Fig. 6 panel 5b), suggesting that Southern Hemisphere South American AN emissions are underestimated. The CO enhancement at 1.5 km (Fig. 6 panel 2c) not reported by Yamasoe et al. (2015) is related to increased local AN contributions for March–April–May 2014 relative to the previous years. This is in agreement with the CEDS2 inventory, which shows a peak in AN emissions over South Brazil in 2014.

In January CO is minimum over the three clusters with CO below 300 ppb (resp. 180 ppb) over Bogota (resp. Caracas and
345 SBrazil) below 1km (Fig. 6 panel 2).

In October, two CO maxima are observed over SBrazil, between 2–4 km (150–200 ppb), and above 8 km (200–250 ppb) (Fig. 6 panel 2c). IASI detects the MT_{IASI} and UT_{IASI} maxima over most of tropical South America (Fig. 2). At the same time O_3 peaks above 2km over Bogota and SBrazil, reaching 45 and 60 ppb in the UT (Fig. 6 panel 2). While CO enhancement below 1 km of SBrazil is caused by local AN (52 %) and BB (44 %) emissions (Fig. 6), BB contribution exceeds 80 % in the
350 MT and UT because of the strong convection moving over the BB regions (Liu et al., 2010), documented by low OLR (Fig. 2l). The uplifted BB products are trapped in an anticyclonic circulation developed over Central South America (Fig. S8l). CO from Southern Hemisphere South American fires are transported over Bogota, at the edge of the anticyclone, but does not reach Caracas which lies outside of the anticyclone (Fig. 6 panels 3d and 4d). Their photochemical processing contributes to the seasonal O_3 enhancement over South America (Thompson et al., 2003b; Sauvage et al., 2006), highlighted by the collocation
355 of IASI UT_{IASI} CO (Fig. 2l) and O_3 (Fig. 2p) maxima.

3.1.4 Arabia and Eastern Africa (AEA)

AEA clusters present CO maximum in the LT (Fig. 7 panel 2 and Fig. S3 panel 2e), similar to other tropical locations mostly due to AN local (Middle East or Africa) emissions, except Addis Ababa and Khartoum in July, showing substantial African BB influence (Fig. 7 panels 3–5 and Fig. S6 panel 8).

360 In January, strong AN emissions from Northern Hemisphere Africa control the LT CO anomalies over Addis Ababa with contribution of 71 % in the LT and 58 % in the MT (Fig. S5). Influence from the Northern Hemisphere African fires is also evident (12 % in the LT and 20 % in the MT). The impact from these fires intensifies over Khartoum and Jeddah between 2 and 4 km with contributions of around 55 % (Fig. 7 panels 4a and 5a).




Figure 6. Same as Fig. 3 for the South American clusters.

In July, the CO LT maximum is again caused by local AN emissions, except over Khartoum where air masses from Southern Hemisphere African fires are the dominant source of CO (Fig. 7 panel 5c). The combination of local AN (70 %) and Southern Hemisphere African BB (23 %) emissions is responsible for the annual CO maximum at the surface over Addis Ababa (Fig. S6 panel 8c and Fig. S5).

The striking feature is the elevated O₃ around 8 km (70 ppb on average) for all the clusters during April, and for Jeddah and Abu Dhabi during July (Fig. 7 panel 1 and Fig. S3 panel 1e). Low CO accompanying the O₃ enhancements (Fig. 7 panel 2 and Fig. S3 panel 2e) points to a dynamical origin of O₃. In July these features are also detected by IASI in the MT_{IASI} and UT_{IASI} (Fig. 2j, k, n and o) and can be due to either i) tropopause foldings in the vicinity of the subtropical jet stream associated with downward transport of stratospheric ozone (Stohl et al., 2003; Lelieveld et al., 2009; Safieddine et al., 2014; Tang et al., 2011) or ii) emissions from South-East Asia (50 %), with smaller contributions from the Middle East and Africa (30 %) (Jia et al., 2017) trapped in the AMA (Barret et al., 2016; Park et al., 2007). This last influence is in agreement with SOFT-IO, showing a significant contribution from South-East Asia over Jeddah (29 % to 78 %), Khartoum (60 %) and Addis Ababa (46 %) (Fig. S5), affected by the tropical easterly jet transport. SOFT-IO also attributes large contributions from Northern Hemisphere African AN and BB emissions, and also from American sources over Abu Dhabi (Fig. 7 panel 3a), related to eastward transport by the subtropical westerly jet in the UT, not present in the rest of the AEA clusters.

In July O₃ is minimum over Addis Ababa (45–50 ppb) (Fig. S3 panel 1e), related to the Inter Tropical Convergence Zone located between 5° N and 10° N (Lannuque et al., 2021).

In October, in the LT the long-range transport from Asia (South-East AN and Equatorial BB) plays a significant role in CO anomalies over the AEA locations (Fig. 7 panels 3d–5d and Fig. S6 panel 8d), especially over Addis Ababa and Jeddah. In the LT_{IASI}, IASI captures the northeasterlies (Fig. S8d) transport of CO-rich air masses from Asia towards eastern Africa (Fig. 2d). Above 4 km, O₃ enhancements are observed over the AEA locations, especially over Abu Dhabi which is the easternmost location of the AEA region (Fig. 7 panel 1). IASI detects a MT_{IASI} O₃ increase above the Arabian Sea and Northern India (Fig. 2h), previously detected with ozone soundings during the INDOEX campaign (1999–2000) and attributed to Indian sources uplifted over the marine boundary layer by the sea breeze circulation (Lawrence and Lelieveld, 2010; Barret et al., 2011). The O₃-rich air masses are further transported towards Eastern Africa by the prevailing northeasterlies (Fig. S8h) as documented by the predominant South-East Asian origin of MT and UT CO over the AEA locations (Fig. 7).

3.2 Control factors of tropical O₃ and CO

First, we give an overview of the Tropics, focusing on the extrema of O₃ and CO of all clusters, then on AN and BB contributions in the upper troposphere. Finally, we look at tropical connections between emission source regions and CO and O₃ distributions. Figure 8 displays the annual maxima/minima of O₃ (a) and CO (b) mixing ratios and their corresponding mean height in terms of absolute altitude. The annual maxima/minima are calculated based on monthly averaged mixing ratios over vertical layers with 40 hPa thickness.

Overall, the CO profiles display an annual maximum close to the surface, approximately at 0.5 km (Fig. 8b) with strong magnitude variations. The minima are generally located in the free troposphere (MT or UT), in the absence of recent pollution




Figure 7. Same as Fig. 3 for the Arabian and Eastern African clusters.




Figure 8. O₃ (a) and CO (b) annual maximum (higher bar) and minimum (lower bar) mixing ratio observed over the tropical clusters. The annotated number on top of each bar indicates the absolute altitude (in km) of the observed annual maximum/minimum mixing ratio. The colour in the bar indicates the month of the maximum/minimum.

and where CO is chemically destroyed, and are uniform in terms of mixing ratios, close to the background levels. Both maxima and minima present strong seasonality related to surface emissions and meteorological conditions.

400 For the majority of the Northern Tropical locations the CO maxima are mainly attributed to local AN emissions. However, for African locations and Caracas, BB contribution is almost similar to or greater than AN (Fig. 3 panel 3; Fig. S6 panels 1 and 2 and Fig. 6 panel 3). This confirms the key role of AN emissions in the Northern Hemisphere, related to a larger population compared to the Southern Hemisphere, and enhanced AN urban and industrial activity. The surface-layer pollution of the Southern Tropics is predominantly caused by BB over Africa (Central Africa and Windhoek) during the dry season, and by

405 AN over SBrazil (Fig. 3 panels 4 and 5 and Fig. 6 panel 5). The CO maxima over the latter occur before the burning season, in accordance with previous studies suggesting fossil fuels as the main CO source over Sao Paulo and Rio de Janeiro (Alonso et al., 2010), and decreasing BB over South America (Andela et al., 2017; Deeter et al., 2018) by more than 40 % due to the long-term declining deforestation rates, especially over forested areas and over savanna and shrublands (Naus et al., 2022).

Because of its complex chemistry, the situation for O₃ is more complicated. Africa is the only region where the annual O₃

410 maximum occurs in the LT (at 2.5 km) during the dry season (Fig. 8a Sahel, Guinea Gulf, Lagos and Central Africa). The

Table 2. Total (AN + BB) CO emission rates (in $10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$) based on CEDS and GFAS emission inventories over West Africa ($10^{\circ} \text{ W} - 12.5^{\circ} \text{ E}$; $0 - 12.5^{\circ} \text{ N}$), Central Africa ($10 - 35^{\circ} \text{ W}$; $2.5 - 20^{\circ} \text{ S}$), East Asia ($92.5 - 110^{\circ} \text{ E}$; $10 - 27^{\circ} \text{ N}$), Maritime Continent ($93 - 121^{\circ} \text{ E}$; $10^{\circ} \text{ S} - 10^{\circ} \text{ N}$), South Brazil ($35 - 50^{\circ} \text{ W}$; $0 - 20^{\circ} \text{ S}$) and Arabia and Eastern Africa ($30 - 60^{\circ} \text{ E}$; $5 - 25^{\circ} \text{ N}$).

	West Africa	Central Africa	East Asia	India	Maritime Continent	South Brazil	AEA
January	6	1	4	3	1.5	1	1.5
April	3	1	11	3.5	1.5	0.5	1
July	2.5	10	3	3	2	1	0.5
October	2.5	3	3	3	6	4	1
Annual	3.5	3.7	5.5	3.1	3	1.5	1

co-occurrence of maximum O_3 with enhanced CO over Africa indicates the stronger dependency of O_3 on the surface AN and BB CO emissions for these regions, in agreement with Sauvage et al. (2007b). South China is the only Asian cluster where the annual O_3 maximum is observed in the LT (at 2.6 km) during the active local fires (April). In contrast, over the other regions, the annual O_3 maximum is observed above 6 km (Fig. 8a). This indicates that O_3 is likely associated with larger O_3 production efficiency in the MT and UT (Sauvage et al., 2007c). In contrast, in regions such SBrazil and Windhoek in October, the co-occurrence of O_3 and CO enhancement in the MT and UT indicates O_3 production from surface sources with LiNOx emissions (Secs. 3.1.1 and 3.1.3).

The annual O_3 minima for all the tropical clusters are observed close to the surface (below 0.5 km on average), likely related to deposition and titration by NO, or because of lower O_3 production in the presence of convective clouds relative to clear sky conditions (Sect. 3.1.2).

Throughout the Tropics, the highest CO and O_3 maxima occur over Western Africa in the LT (Fig. 8) during the dry season (January). Table 2 displays the total (AN and BB) CO emissions rates over several regions of interest based on the sum of CEDS2 and GFAS emission inventories. The NO_x -limited O_3 production regime over Western Africa (Saunio et al., 2009; Zhang et al., 2016) likely explains the O_3 maxima when the local emissions, and thus the NO_2 mixing ratios increase in the region (Jaeglé et al., 2004). The largest O_3 and CO mixing ratio over Lagos is due to the strong Nigerian AN emissions and its location downwind of fire plumes through Harmattan, as confirmed by SOFT-IO (see Sect. 3.1.1).

The second highest CO and O_3 maxima over the Tropics are observed over Asia (Fig. 8) in January, mostly due to AN emissions. According to Table 2, the CO emissions over East Asia and India are lower than the ones over West Africa in January, explaining the lower CO mixing ratios over Asia than over Lagos. SOFT-IO represents better the Asian contributions than the African ones. As mentioned in Sect. 3.1.1, the emissions over Africa are likely underestimated by the SOFT-IO computations. Previous studies have found mixing ratios of pollutants in West Africa comparable to those observed over Asian megacities (Assamoi and Liousse, 2010; Adon et al., 2016; Sauvage et al., 2007b). The rapid growth of African megacities is indeed responsible for increasing emissions from diffuse and inefficient combustion sources (Marais and Wiedinmyer, 2016), such as residential sources mainly for cooking and heating (Zheng et al., 2019), and traffic emissions (related to a large number

435 of two-stroke vehicles, poor fuel quality and poorly-maintained engines) (Assamoi and Liousse, 2010). In contrast, Eastern
China has had one of the largest decreases in CO emissions (Hedelius et al., 2021) due to technological changes with improved
combustion efficiency (Zheng et al., 2018), such as replacing residential coal use with electricity and natural gas (Buchholz
et al., 2021), reduction of NO_x emissions from coal power plants since 2010 (Wang et al., 2017), and implementation of Clean
Air Policies (van der A et al., 2017) around 2010. Over South China, CO emissions have also been reduced in the last years,
440 but the O₃ levels remain relatively high (Fig. 8a). This is because the O₃ production regime over South China is VOCs-limited
(Li et al., 2013), and the total NMVOCs emissions increased by a factor of 3.5 (1997–2017) because of activity increases
in the solvent, energy, and industry sectors (Zheng et al., 2018). In India, on the other hand, the lack of efficient pollution
regulations explains the highest CO mixing ratios among the Asian clusters (Fig. 8). Previous studies have already reported
increasing CO emissions over India from 1996–2015, due to several factors such as increases in residential and agricultural
445 sources (Pandey et al., 2014), power production and transport activities (Sadavarte and Venkataraman, 2014). Furthermore,
Indian emissions are mostly associated with incomplete combustion processes from biofuel burning (Kumar et al., 2012) and
are therefore characterised by higher NMVOCs to NO_x ratios than other Asian regions leading to a NO_x-limited O₃ production
regime (Lawrence and Lelieveld, 2010).

The smallest LT CO maxima over the Northern Tropics are observed over the AEA clusters and South America (Fig. 8)
450 because of the smallest emissions rates among the Tropics (Table 2). The Middle East CO emissions are mainly related to
electricity generation, water desalination, and industry supplied by oil and gas deposits with cheap but relatively clean fuels
(Krotkov et al., 2016). O₃ maxima in the MT and UT of the AEA and South American clusters are related respectively to
pollution transport and LiNO_x emissions at higher altitudes (Sects. 3.1.4 and 3.1.3).

From the previous analysis, all the tropical clusters are primarily influenced by local sources of CO. However, AN and BB
455 emissions can be redistributed vertically and then carried horizontally aloft by zonal winds to influence the tropical upper
troposphere. Figure 9 displays the AN and BB contributions to the CO anomalies (in ppb) over the tropical UT_{cruise}. AN
emissions have the largest influence (Fig. 9) especially in the Eastern part, from the Middle East to Asia in July and October.
BB has a marked seasonal influence, less intense than the AN one. Fires strongly influence deep convection outflow regions,
above the Inter Tropical Convergence Zone over Africa in January and July, above the South Atlantic Convergence over South
460 America and the South Pacific Convergence Zone above Asia in October. For the latter, the influence of fire is of the same
order as that of AN emissions (24 ppb simulated).

In addition to the local influence, long-range transport plays an important role in the redistribution of these AN and BB emis-
sions and, consequently, in the distribution of CO in the Tropics. To illustrate this, Figure 10 displays the transport pathways
of CO emissions from the African, South American and Asian source regions, towards the 20 tropical IAGOS locations in the
465 LT (a), MT (b) and UT (c). Only the source regions and the months corresponding to the largest amounts of transported CO (in
ppb) are depicted.

CO sources located over Africa show the largest influence on the regional tropical CO. Large amounts of CO are indeed
transported inter-hemispherically from the dry-season region towards the wet-season region (Fig. 9 and Fig. 10) from the LT
with advection by the trades winds to the UT with the Hadley cells after vertical convective uplift in the convergence zone. As




Figure 9. Mean AN (a–d) and BB (e–h) contributions in ppb between 300–185 hPa from 2002–2019.




Figure 10. Transport of CO (AN+BB) emissions from the African, South American and Asian source regions towards the 20 tropical locations taken into account for this study. The colorbar shows the amount of CO transported in ppb.

470 a result, CO contributions of 45–50 ppb (resp. over 50 ppb) from Northern Hemisphere (resp. Southern Hemisphere) Africa
are found over Southern (resp. Northern) Africa during the respective dry season in the entire troposphere. In the UT this
explains why the Southern (resp. Northern) Hemisphere African emissions are dominant in the wet-season hemisphere during
July (resp. January) (Fig. 9 and resp. Fig. S12 and Fig. S10). During the dry season, the impact of the Northern Hemisphere
African emissions is stronger and extends to a wider area over South America, the Middle East and South Asia (Fig. S10
475 NHAf). Also, Northern Hemisphere African emissions are transported towards South America (10–15 ppb in MT; 5–10 ppb
in the rest) in April (Fig. not shown) and significantly contribute to the local South American annual maximum (30 % and 50
% of CO anomalies over Caracas, LT and MT resp.) (Fig. 10 and Fig. 4a–b).

The contribution of Asian emissions in the tropical LT is limited to a regional scale, as they are mostly impacting neighbour
Asian regions (Fig. 10a). CO export from Asia is enhanced during the Asian summer monsoon and post-monsoon (July and
480 October) in the UT, where transport is favoured due to stronger winds relative to the surface (Fig. S3). During the Asian
summer monsoon, the CO-rich (and O₃-poor) air masses from the boundary layer (Fig. 2c and g) are convectively uplifted
to the UT_{cruise} (Fig. 9c and Fig. S12), and trapped in the AMA circulation (see Sect. 3.1.2). Asian emissions are transported
towards Arabia (25–30 ppb) and Eastern Africa (10–15 ppb) in the UT_{cruise} (Fig. 9c; Fig. 10 panel 3c and Fig. S12 SEAS and
CEAS). Subsidence of air masses from AMA above Arabia is contributing to an O₃ maximum above AEA (Fig. 8a) (Sect.
485 3.1.4). During the post-monsoon season (October), sporadic convection uplifts CO emitted by the Equatorial Asian fires in the
UT_{cruise} (Fig. 9g and Fig. S13 EQAS) (see Sect. 3.1.2). At the same time, convection over continental Asia uplifts South-East
Asian AN emissions (Fig. 9d and Fig. S13 SEAS) to the UT_{cruise}. As a result, Asian emissions impact CO anomalies in the
UT over Eastern Africa (15–20 ppb from Equatorial Asia and 5–10 ppb from South-East Asia; Fig. not shown).

South American emissions mostly influence the troposphere over the South Atlantic and South America in October (Fig. 10)
490 through the South Atlantic Anticyclone circulation, accounting for around 25 % of the observed CO.

4 Summary and conclusions

IAGOS O₃ and CO observations since 1994 and 2002 respectively, were used to analyse tropospheric profiles over 20 tropi-
cal locations, along with the UT_{cruise} distributions. Despite their sparse spatial coverage, these observations provide a good
overview of the O₃ and CO distributions in most tropical regions. They are indeed representative of both the urban background
495 close to the ground and of the regional scale starting a few hundred meters above the surface (Petetin et al., 2018a). Further-
more, global IASI-SOFRID O₃ and CO distributions allow us to complement IAGOS observations over the whole tropical
band.

The results of the study indicate that the highest O₃ and CO mixing ratios are observed over Africa during the fire season in
January, with anomalies located in the LT (75 ppb at 2.5 km for O₃ and 800 ppb at 0.3 km for CO over Lagos), and explained
500 mainly by anthropogenic emissions, but with a strong contribution from fires. The secondary O₃ and CO maxima are observed
over Asia, in the MT-UT in April for the first and in the LT in January for the second, with a larger contribution from AN

emissions. The Asian distributions are largely controlled by meteorological conditions associated with the Asian monsoon phases. The lowest anomalies are measured over Caracas.

505 More specifically in the tropical LT, where the CO is maximum, the majority of the clusters are mostly influenced by local and regional AN contributions all year long. In particular, AN emissions have the greatest impact over Asia, Arabia and Eastern Africa and South America, where they account for more than 75 % of the CO. These results highlight the importance of the AN emissions over the tropical locations, in accordance with the global decreasing trends of BB and the increasing AN emissions. The impact of BB in the LT has also a predominantly local and regional influence with a strong seasonal dependence. BB contributions increase over South China (35 % in April) and dominate over the Gulf of Thailand (90 % in October) during the
510 local fire season (South-East and Equatorial Asian resp.). In contrast to the other continents, African LT with the highest O₃ and CO measured, is more impacted by BB during the dry seasons (January and July) with an influence of the same order as AN. This is also the case over Caracas (April) and SBrazil (October). The highest BB impact is found over Southern Hemisphere Africa all year long (57–90%), except in April, with BB emissions mostly originating from local fires, but also from Northern Hemisphere Africa in January (45-73 %) and Southern Hemisphere Southern America in October (29 % over Windhoek).

515 In the MT and UT, AN contributions are more important and dominate in the Eastern part of the Tropics, from the Middle East to Asia. BB contributions are increased compared to the LT and are higher in the outflow of the deep convection regions, especially African fires over Africa in January and July, and South-East and Equatorial Asian fires over the South East Asian coast (South China, Manila, Ho Chi Minh City) in April or October. Also, the contribution of long-range transport is more important than in the LT and allows connections between source regions and other continents.

520 Africa is the leading tropical region in terms of emissions export to the tropical troposphere. The main export pathway is the inter-hemispheric transport of O₃ and CO from the dry-season African regions to the wet-season ones. Northern (resp. Southern) Hemisphere African fires are also the dominant source of CO over Arabia and Eastern Africa (resp. Khartoum and Jeddah) in the MT and UT, and they also reach the MT and UT of India, accounting for 20% of the observed CO. Transport of African emissions from Africa occurs all year round towards northern South America in the entire troposphere, with the largest
525 impact over Caracas in the MT and UT (30 % on average).

In contrast, the impact of Asian emissions is mostly limited on a regional or local scale, especially in the LT and MT. Nevertheless, long-range transport of Asian emissions is responsible for 60 to 80% of the CO anomalies in the UT over AEA in July and October. During the Asian summer monsoon CO from South Asia is transported westwards within the AMA while in October, the tropical easterly jet transports air masses impacted by Indonesian fires towards Africa.

530 Last, over South America the local CO contributions at the surface level are as low as over AEA. During the dry season (October), when the convection moves over the South American fires, CO and precursors are trapped in an anticyclonic circulation developed over Central South America, resulting in the annual local maxima of O₃ and CO. As a result, Northern and Southern Hemisphere South American emissions contributed to 30 % of the CO observed in the MT over Windhoek, after transport in the South Atlantic anticyclone.

535 *Data availability.* The IAGOS data are available on the IAGOS data portal (<https://doi.org/10.25326/20>). The SOFT-IO v1.0 products are part of the ancillary products of IAGOS central database (<https://doi.org/10.25326/2>; <https://doi.org/10.25326/3>). The SOFRID-O₃ data are freely available on the IASI-SOFRID website (<http://thredds.sedoo.fr/iasi-sofrid-o3-co/>, last access: 8 June 2022; SEDOO, 2014).

Author contributions. MT, BS and BB designed the research. All the co-authors contributed to acquisition of data. MT analysed the data. MT, BS and BB interpreted the data. MT drafted the article. MT, BS and BB revised the article. VT and HC commented the article. ELF and
540 BB are responsible for the SOFRID retrieval software. ELF is in charge of the production and quality control of SOFRID data.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We acknowledge the strong support of the European Commission, Airbus and the airlines (Deutsche Lufthansa, Air France, Austrian, Air Namibia, Cathay Pacific, Iberia, China Airlines and Hawaiian Airlines) that have carried the MOZAIC or IAGOS equipment and performed the maintenance since 1994. IAGOS has been funded by the European Union projects IAGOS–DS and IA-
545 GOS–ERI. Additionally, IAGOS has been funded by INSU-CNRS (France), Météo-France, Université Paul Sabatier (Toulouse, France) and Research Center Jülich (FZJ, Jülich, Germany). The IAGOS database is supported in France by AERIS (<https://www.aeris-data.fr>). IASI is a joint mission of EUMETSAT and the Centre National d’Etudes Spatiales (CNES, France). The authors acknowledge the CNES for financial support for the IASI activities.

Financial support. This research has been supported by Bonus Stratégique programme at Université Paul Sabatier Toulouse III who funded
550 the first author’s doctoral position.

References

- Adon, M., Galy-Lacaux, C., Yoboué, V., Delon, C., Lacaux, J., Castera, P., Gardrat, E., Pienaar, J., Al Ourabi, H., Laouali, D., et al.: Long term measurements of sulfur dioxide, nitrogen dioxide, ammonia, nitric acid and ozone in Africa using passive samplers, *Atmospheric Chemistry and Physics*, 10, 7467–7487, <https://doi.org/10.5194/acp-10-7467-2010>, 2010.
- 555 Adon, M., Yoboué, V., Galy-Lacaux, C., Lioussé, C., Diop, B., Gardrat, E., Ndiaye, S. A., Jarnot, C., et al.: Measurements of NO₂, SO₂, NH₃, HNO₃ and O₃ in West African urban environments, *Atmospheric Environment*, 135, 31–40, <https://doi.org/10.1016/j.atmosenv.2016.03.050>, 2016.
- Aghedo, A. M., Schultz, M. G., and Rast, S.: The influence of African air pollution on regional and global tropospheric ozone, *Atmospheric Chemistry and Physics*, 7, 1193–1212, <https://doi.org/10.1016/j.atmosenv.2016.03.050>, 2007.
- 560 Alonso, M. F., Longo, K. M., Freitas, S. R., da Fonseca, R. M., Marécal, V., Pirre, M., and Klenner, L. G.: An urban emissions inventory for South America and its application in numerical modeling of atmospheric chemical composition at local and regional scales, *Atmospheric Environment*, 44, 5072–5083, <https://doi.org/10.1016/j.atmosenv.2010.09.013>, 2010.
- Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz, G., Hantson, S., Kloster, S., et al.: A human-driven decline in global burned area, *Science*, 356, 1356–1362, <https://doi.org/10.1126/science.aal4108>, 2017.
- 565 Assamoi, E.-M. and Lioussé, C.: A new inventory for two-wheel vehicle emissions in West Africa for 2002, *Atmospheric Environment*, 44, 3985–3996, <https://doi.org/10.1016/j.atmosenv.2010.06.048>, 2010.
- Barret, B., Ricaud, P., Mari, C., Attié, J.-L., Boussez, N., Josse, B., Le Flochmoën, E., Livesey, N., Massart, S., Peuch, V.-H., et al.: Transport pathways of CO in the African upper troposphere during the monsoon season: a study based upon the assimilation of spaceborne observations, *Atmospheric Chemistry and Physics*, 8, 3231–3246, <https://doi.org/10.5194/acp-8-3231-2008>, 2008.
- 570 Barret, B., Le Flochmoën, E., Sauvage, B., Pavelin, E., Matricardi, M., and Cammas, J.-P.: The detection of post-monsoon tropospheric ozone variability over south Asia using IASI data, *Atmospheric Chemistry and Physics*, 11, 9533–9548, <https://doi.org/10.5194/acp-11-9533-2011>, 2011.
- Barret, B., Sauvage, B., Bennouna, Y., and Le Flochmoën, E.: Upper-tropospheric CO and O₃ budget during the Asian summer monsoon, *Atmospheric Chemistry and Physics*, 16, 9129–9147, <https://doi.org/10.5194/acp-16-9129-2016>, 2016.
- 575 Barret, B., Emili, E., and Le Flochmoën, E.: A tropopause-related climatological a priori profile for IASI-SOFRID ozone retrievals: improvements and validation, *Atmospheric Measurement Techniques*, 13, 5237–5257, <https://doi.org/10.5194/amt-13-5237-2020>, 2020.
- Barret, B., Gouzenes, Y., Le Flochmoën, E., and Ferrant, S.: Retrieval of Metop-A/IASI N₂O profiles and validation with NDACC FTIR data, *Atmosphere*, 12, 219, <https://doi.org/10.3390/atmos12020219>, 2021.
- Blot, R., Nedelec, P., Boulanger, D., Wolff, P., Sauvage, B., Cousin, J.-M., Athier, G., Zahn, A., Obersteiner, F., Scharffe, D., et al.: Internal consistency of the IAGOS ozone and carbon monoxide measurements for the last 25 years, *Atmospheric Measurement Techniques*, 14, 3935–3951, <https://doi.org/10.5194/amt-14-3935-2021>, 2021.
- 580 Bourgeois, I., Peischl, J., Thompson, C. R., Aikin, K. C., Campos, T., Clark, H., Commane, R., Daube, B., Diskin, G. W., Elkins, J. W., et al.: Global-scale distribution of ozone in the remote troposphere from the ATom and HIPPO airborne field missions, *Atmospheric Chemistry and Physics*, 20, 10611–10635, <https://doi.org/10.5194/acp-20-10611-2020>, 2020.
- 585 Bourgeois, I., Peischl, J., Neuman, J. A., Brown, S. S., Thompson, C. R., Aikin, K. C., Allen, H. M., Angot, H., Apel, E. C., Baublitz, C. B., et al.: Large contribution of biomass burning emissions to ozone throughout the global remote troposphere, *Proceedings of the National Academy of Sciences*, 118, e2109628 118, <https://doi.org/10.1073/pnas.2109628118>, 2021.

- Boynard, A., Hurtmans, D., Koukouli, M. E., Goutail, F., Bureau, J., Safieddine, S., Lerot, C., Hadji-Lazaro, J., Wespes, C., Pommereau, J.-P., et al.: Seven years of IASI ozone retrievals from FORLI: validation with independent total column and vertical profile measurements, *Atmospheric Measurement Techniques*, 9, 4327–4353, <https://doi.org/10.5194/amt-9-4327-2016>, 2016.
- 590 Boynard, A., Hurtmans, D., Garane, K., Goutail, F., Hadji-Lazaro, J., Koukouli, M. E., Wespes, C., Vigouroux, C., Keppens, A., Pommereau, J.-P., et al.: Validation of the IASI FORLI/EUMETSAT ozone products using satellite (GOME-2), ground-based (Brewer–Dobson, SAOZ, FTIR) and ozonesonde measurements, *Atmospheric Measurement Techniques*, 11, 5125–5152, <https://doi.org/10.5194/amt-11-5125-2018>, 2018.
- 595 Buchholz, R. R., Worden, H. M., Park, M., Francis, G., Deeter, M. N., Edwards, D. P., Emmons, L. K., Gaubert, B., Gille, J., Martínez-Alonso, S., et al.: Air pollution trends measured from Terra: CO and AOD over industrial, fire-prone, and background regions, *Remote Sensing of Environment*, 256, 112 275, <https://doi.org/10.1016/j.rse.2020.112275>, 2021.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., et al.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmospheric Chemistry and Physics*, 9, 6041–6054, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- 600 Cohen, Y., Petetin, H., Thouret, V., Marécal, V., Josse, B., Clark, H., Sauvage, B., Fontaine, A., Athier, G., Blot, R., et al.: Climatology and long-term evolution of ozone and carbon monoxide in the upper troposphere–lower stratosphere (UTLS) at northern midlatitudes, as seen by IAGOS from 1995 to 2013, *Atmospheric Chemistry and Physics*, 18, 5415–5453, <https://doi.org/10.5194/acp-18-5415-2018>, 2018.
- Curtis, L., Rea, W., Smith-Willis, P., Fenyves, E., and Pan, Y.: Adverse health effects of outdoor air pollutants, *Environment international*, 32, 815–830, <https://doi.org/10.1016/j.envint.2006.03.012>, 2006.
- 605 Cussac, M., Marécal, V., Thouret, V., Josse, B., and Sauvage, B.: The impact of biomass burning on upper tropospheric carbon monoxide: a study using MOCAGE global model and IAGOS airborne data, *Atmospheric Chemistry and Physics*, 20, 9393–9417, <https://doi.org/10.5194/acp-20-9393-2020>, 2020.
- De Wachter, E., Barret, B., Le Flochmoën, E., Pavelin, E., Matricardi, M., Clerbaux, C., Hadji-Lazaro, J., George, M., Hurtmans, D., Coheur, P.-F., et al.: Retrieval of MetOp-A/IASI CO profiles and validation with MOZAIC data, *Atmospheric Measurement Techniques*, 5, 2843–2857, <https://doi.org/10.5194/amt-5-2843-2012>, 2012.
- 610 Deeter, M., Martínez-Alonso, S., Andreae, M. O., and Schlager, H.: Satellite-based analysis of CO seasonal and interannual variability over the Amazon Basin, *Journal of Geophysical Research: Atmospheres*, 123, 5641–5656, <https://doi.org/10.1029/2018JD028425>, 2018.
- Dufour, G., Eremenko, M., Orphal, J., and Flaud, J.-M.: IASI observations of seasonal and day-to-day variations of tropospheric ozone over three highly populated areas of China: Beijing, Shanghai, and Hong Kong, *Atmospheric Chemistry and Physics*, 10, 3787–3801, <https://doi.org/10.5194/acp-10-3787-2010>, 2010.
- 615 Duncan, B. N., Lamsal, L. N., Thompson, A. M., Yoshida, Y., Lu, Z., Streets, D. G., Hurwitz, M. M., and Pickering, K. E.: A space-based, high-resolution view of notable changes in urban NO_x pollution around the world (2005–2014), *Journal of Geophysical Research: Atmospheres*, 121, 976–996, <https://doi.org/10.1002/2015JD024121>, 2016.
- 620 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., et al.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- Eremenko, M., Dufour, G., Foret, G., Keim, C., Orphal, J., Beekmann, M., Bergametti, G., and Flaud, J.-M.: Tropospheric ozone distributions over Europe during the heat wave in July 2007 observed from infrared nadir spectra recorded by IASI, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL034803>, 2008.
- 625

- Galanter, M., Levy, H., and Carmichael, G. R.: Impacts of biomass burning on tropospheric CO, NO_x, and O₃, *Journal of Geophysical Research: Atmospheres*, 105, 6633–6653, <https://doi.org/10.1029/1999JD901113>, 2000.
- Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J., Clerbaux, C., Coheur, P.-F., Cuesta, J., Cuevas, E., et al.: Tropospheric Ozone Assessment Report: Present-day distribution and trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation, *Elementa: science of the anthropocene*, 6, <https://doi.org/10.1525/elementa.291>, 2018.
- Gaudel, A., Cooper, O. R., Chang, K.-L., Bourgeois, I., Ziemke, J. R., Strode, S. A., Oman, L. D., Sellitto, P., Nédélec, P., Blot, R., et al.: Aircraft observations since the 1990s reveal increases of tropospheric ozone at multiple locations across the Northern Hemisphere, *Science Advances*, 6, eaba8272, <https://doi.org/10.1525/elementa.291>, 2020.
- Gauss, M., Myhre, G., Pitari, G., Prather, M., Isaksen, I., Bernsten, T., Brasseur, G., Dentener, F., Derwent, R., Hauglustaine, D., et al.: Muller, JF, Plantevin, P., Pyle, JA, Rogers, HL, Stevenson, DS, Sundet, JK, van Weele, M., and Wild, O.: Radiative forcing in the 21st century due to ozone changes in the troposphere and the lower stratosphere, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JD002624>, 2003.
- George, M., Clerbaux, C., Hurtmans, D., Turquety, S., Coheur, P.-F., Pommier, M., Hadji-Lazaro, J., Edwards, D., Worden, H., Luo, M., et al.: Carbon monoxide distributions from the IASI/METOP mission: evaluation with other space-borne remote sensors, *Atmospheric Chemistry and Physics*, 9, 8317–8330, <https://doi.org/10.5194/acp-9-8317-2009>, 2009.
- Granier, C., Bessagnet, B., Bond, T., D’Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., et al.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic change*, 109, 163–190, <https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- Gressent, A., Sauvage, B., Defier, E., Pätz, H. W., Thomas, K., Holle, R., Cammas, J.-P., Nédélec, P., Boulanger, D., Thouret, V., et al.: Lightning NO_x influence on large-scale NO_y and O₃ plumes observed over the northern mid-latitudes, *Tellus B: Chemical and Physical Meteorology*, 66, 25 544, <https://doi.org/10.3402/tellusb.v66.25544>, 2014.
- Gressent, A., Sauvage, B., Cariolle, D., Evans, M., Leriche, M., Mari, C., and Thouret, V.: Modeling lightning-NO_x chemistry on a sub-grid scale in a global chemical transport model, *Atmospheric Chemistry and Physics*, 16, 5867–5889, <https://doi.org/10.5194/acp-16-5867-2016>, 2016.
- Hedelius, J. K., Toon, G. C., Buchholz, R. R., Iraci, L. T., Podolske, J. R., Roehl, C. M., Wennberg, P. O., Worden, H. M., and Wunch, D.: Regional and urban column CO trends and anomalies as observed by MOPITT over 16 years, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 967, <https://doi.org/10.1029/2020JD033967>, 2021.
- Heue, K.-P., Coldewey-Egbers, M., Delcloo, A., Lerot, C., Loyola, D., Valks, P., and Van Roozendaal, M.: Trends of tropical tropospheric ozone from 20 years of European satellite measurements and perspectives for the Sentinel-5 Precursor, *Atmospheric Measurement Techniques*, 9, 5037–5051, <https://doi.org/10.5194/amt-9-5037-2016>, 2016.
- IPCC: Climate change 2021: The physical science basis, Tech. rep., <http://www.ipcc.ch/report/ar6/wg1/>, 2021.
- Jaeglé, L., Martin, R., Chance, K., Steinberger, L., Kurosu, T., Jacob, D. J., Modi, A., Yoboué, V., Sigha-Nkamdjou, L., and Galy-Lacaux, C.: Satellite mapping of rain-induced nitric oxide emissions from soils, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004787>, 2004.
- Jerrett, M., Burnett, R. T., Pope III, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E., and Thun, M.: Long-term ozone exposure and mortality, *New England Journal of Medicine*, 360, 1085–1095, <https://doi.org/10.1056/NEJMoa0803894>, 2009.
- Jia, J., Ladstätter-Weissenmayer, A., Hou, X., Rozanov, A., and Burrows, J. P.: Tropospheric ozone maxima observed over the Arabian Sea during the pre-monsoon, *Atmospheric Chemistry and Physics*, 17, 4915–4930, <https://doi.org/10.5194/acp-17-4915-2017>, 2017.

- Kaiser, J., Heil, A., Andreae, M., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M., Suttie, M., et al.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., et al.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmospheric Chemistry and Physics*, 16, 4605–4629, <https://doi.org/10.5194/acp-16-4605-2016>, 2016.
- Kumar, R., Naja, M., Pfister, G., Barth, M., Wiedinmyer, C., and Brasseur, G.: Simulations over South Asia using the Weather Research and Forecasting model with Chemistry (WRF-Chem): chemistry evaluation and initial results, *Geoscientific Model Development*, 5, 619–648, <https://doi.org/10.5194/gmd-5-619-2012>, 2012.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., et al.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- Lannuque, V., Sauvage, B., Barret, B., Clark, H., Athier, G., Boulanger, D., Cammas, J.-P., Cousin, J.-M., Fontaine, A., Le Flochmoën, E., et al.: Origins and characterization of CO and O₃ in the African upper troposphere, *Atmospheric chemistry and physics*, 21, 14535–14555, <https://doi.org/10.5194/acp-21-14535-2021>, 2021.
- Lawrence, M. and Lelieveld, J.: Atmospheric pollutant outflow from southern Asia: a review, *Atmospheric Chemistry and Physics*, 10, 11017–11096, <https://doi.org/10.5194/acp-10-11017-2010>, 2010.
- Lelieveld, J., Hoor, P., Jöckel, P., Pozzer, A., Hadjinicolaou, P., Cammas, J.-P., and Beirle, S.: Severe ozone air pollution in the Persian Gulf region, *Atmospheric Chemistry and Physics*, 9, 1393–1406, <https://doi.org/10.5194/acp-9-1393-2009>, 2009.
- Lelieveld, J., Gromov, S., Pozzer, A., and Taraborrelli, D.: Global tropospheric hydroxyl distribution, budget and reactivity, *Atmospheric Chemistry and Physics*, 16, 12477–12493, <https://doi.org/10.5194/acp-16-12477-2016>, 2016.
- Lelieveld, J. o., Crutzen, P., Ramanathan, V., Andreae, M., Brenninkmeijer, C., Campos, T., Cass, G., Dickerson, R., Fischer, H., De Gouw, J., et al.: The Indian Ocean experiment: widespread air pollution from South and Southeast Asia, *Science*, 291, 1031–1036, <https://doi.org/10.1126/science.1057103>, 2001.
- Li, Y., Lau, A. K., Fung, J. C., Zheng, J., and Liu, S.: Importance of NO_x control for peak ozone reduction in the Pearl River Delta region, *Journal of Geophysical Research: Atmospheres*, 118, 9428–9443, <https://doi.org/10.1002/jgrd.50659>, 2013.
- Liu, J., Logan, J. A., Jones, D., Livesey, N., Megretskaia, I., Carouge, C., and Nedelec, P.: Analysis of CO in the tropical troposphere using Aura satellite data and the GEOS-Chem model: insights into transport characteristics of the GEOS meteorological products, *Atmospheric Chemistry and Physics*, 10, 12207–12232, <https://doi.org/10.5194/acp-10-12207-2010>, 2010.
- Livesey, N., Logan, J., Santee, M., Waters, J., Doherty, R., Read, W., Froidevaux, L., and Jiang, J.: Interrelated variations of O₃, CO and deep convection in the tropical/subtropical upper troposphere observed by the Aura Microwave Limb Sounder (MLS) during 2004–2011, *Atmospheric Chemistry and Physics*, 13, 579–598, <https://doi.org/10.5194/acp-13-579-2013>, 2013.
- Logan, J. A.: An analysis of ozonesonde data for the troposphere: Recommendations for testing 3-D models and development of a gridded climatology for tropospheric ozone, *Journal of Geophysical Research: Atmospheres*, 104, 16115–16149, <https://doi.org/10.1029/1998JD100096>, 1999.
- Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: A global perspective, *Journal of Geophysical Research: Oceans*, 86, 7210–7254, <https://doi.org/10.1029/JC086iC08p07210>, 1981.

- Lu, X., Zhang, L., Zhao, Y., Jacob, D. J., Hu, Y., Hu, L., Gao, M., Liu, X., Petropavlovskikh, I., McClure-Begley, A., et al.: Surface and tropospheric ozone trends in the Southern Hemisphere since 1990: possible linkages to poleward expansion of the Hadley circulation, *Science Bulletin*, 64, 400–409, <https://doi.org/10.1016/j.scib.2018.12.021>, 2019.
- 705 Marais, E. A. and Wiedinmyer, C.: Air quality impact of diffuse and inefficient combustion emissions in Africa (DICE-Africa), *Environmental science & technology*, 50, 10 739–10 745, <https://doi.org/10.1021/acs.est.6b02602>, 2016.
- Marengo, A., Thouret, V., Nédélec, P., Smit, H., Helten, M., Kley, D., Karcher, F., Simon, P., Law, K., Pyle, J., et al.: Measurement of ozone and water vapor by Airbus in-service aircraft: The MOZAIK airborne program, An overview, *Journal of Geophysical Research: Atmospheres*, 103, 25 631–25 642, <https://doi.org/10.1029/98JD00977>, 1998.
- Mari, C., Jacob, D. J., and Bechtold, P.: Transport and scavenging of soluble gases in a deep convective cloud, *Journal of Geophysical Research: Atmospheres*, 105, 22 255–22 267, <https://doi.org/10.1029/2000JD900211>, 2000.
- 710 McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector-and fuel-specific sources (1970–2017): an application of the Community Emissions Data System (CEDS), *Earth System Science Data*, 12, 3413–3442, <https://doi.org/10.5194/essd-12-3413-2020>, 2020.
- 715 Monks, P. S.: Gas-phase radical chemistry in the troposphere, *Chemical Society Reviews*, 34, 376–395, <https://doi.org/10.1039/B307982C>, 2005.
- Monks, P. S., Archibald, A., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G., et al.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmospheric Chemistry and Physics*, 15, 8889–8973, <https://doi.org/10.5194/acp-15-8889-2015>, 2015.
- 720 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., et al.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmospheric Chemistry and Physics*, 13, 1853–1877, <https://doi.org/10.5194/acp-13-1853-2013>, 2013.
- Naus, S., Domingues, L. G., Krol, M., Lujikx, I. T., Gatti, L. V., Miller, J. B., Gloor, E., Basu, S., Correia, C., Koren, G., et al.: Sixteen years of MOPITT satellite data strongly constrain Amazon CO fire emissions, *European Geoscience Union (EGU) General Assembly 2022*, pp. 1–25, <https://doi.org/10.5194/egusphere-2022-450>, 2022.
- 725 Nédélec, P., Blot, R., Boulanger, D., Athier, G., Cousin, J.-M., Gautron, B., Petzold, A., Volz-Thomas, A., and Thouret, V.: Instrumentation on commercial aircraft for monitoring the atmospheric composition on a global scale: the IAGOS system, technical overview of ozone and carbon monoxide measurements, *Tellus B: Chemical and Physical Meteorology*, 67, 27 791, <https://doi.org/10.3402/tellusb.v67.27791>, 2015.
- 730 Pandey, A., Sadavarte, P., Rao, A. B., and Venkataraman, C.: Trends in multi-pollutant emissions from a technology-linked inventory for India: II. Residential, agricultural and informal industry sectors, *Atmospheric environment*, 99, 341–352, <https://doi.org/10.1016/j.atmosenv.2014.09.080>, 2014.
- Park, M., Randel, W. J., Gettelman, A., Massie, S. T., and Jiang, J. H.: Transport above the Asian summer monsoon anticyclone inferred from Aura Microwave Limb Sounder tracers, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD008294>, 2007.
- 735 Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A., and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, *Atmospheric Chemistry and Physics*, 8, 757–764, <https://doi.org/10.5194/acp-8-757-2008>, 2008.

- Petetin, H., Thouret, V., Athier, G., Blot, R., Boulanger, D., Cousin, J.-M., Gaudel, A., Nédélec, P., and Cooper, O.: Diurnal cycle of ozone throughout the troposphere over Frankfurt as measured by MOZAIC-IAGOS commercial aircraft Diurnal cycle of ozone throughout the troposphere, *Elementa: Science of the Anthropocene*, 4, <https://doi.org/10.12952/journal.elementa.000129>, 2016.
- 740 Petetin, H., Jeoffrion, M., Sauvage, B., Athier, G., Blot, R., Boulanger, D., Clark, H., Cousin, J.-M., Gheusi, F., Nedelec, P., et al.: Representativeness of the IAGOS airborne measurements in the lower troposphere, *Elementa: Science of the Anthropocene*, 6, <https://doi.org/10.1525/elementa.280>, 2018a.
- 745 Petetin, H., Sauvage, B., Parrington, M., Clark, H., Fontaine, A., Athier, G., Blot, R., Boulanger, D., Cousin, J.-M., Nédélec, P., et al.: The role of biomass burning as derived from the tropospheric CO vertical profiles measured by IAGOS aircraft in 2002–2017, *Atmospheric Chemistry and Physics*, 18, 17 277–17 306, <https://doi.org/10.5194/acp-18-17277-2018>, 2018b.
- Petzold, A., Thouret, V., Gerbig, C., Zahn, A., Brenninkmeijer, C. A., Gallagher, M., Hermann, M., Pontaud, M., Ziereis, H., Boulanger, D., et al.: Global-scale atmosphere monitoring by in-service aircraft—current achievements and future prospects of the European Research Infrastructure IAGOS, *Tellus B: Chemical and Physical Meteorology*, 67, 28 452, <https://doi.org/10.3402/tellusb.v67.28452>, 2015.
- 750 Pickering, K. E., Thompson, A. M., Tao, W.-K., Rood, R. B., McNamara, D. P., and Molod, A. M.: Vertical transport by convective clouds: Comparisons of three modeling approaches, *Geophysical research letters*, 22, 1089–1092, <https://doi.org/10.1029/95GL00889>, 1995.
- Sadavarte, P. and Venkataraman, C.: Trends in multi-pollutant emissions from a technology-linked inventory for India: I. Industry and transport sectors, *Atmospheric environment*, 99, 353–364, <https://doi.org/10.1016/j.atmosenv.2014.09.081>, 2014.
- 755 Safieddine, S., Boynard, A., Coheur, P.-F., Hurtmans, D., Pfister, G., Quennehen, B., Thomas, J. L., Raut, J.-C., Law, K. S., Klimont, Z., et al.: Summertime tropospheric ozone assessment over the Mediterranean region using the thermal infrared IASI/MetOp sounder and the WRF-Chem model, *Atmospheric chemistry and physics*, 14, 10 119–10 131, <https://doi.org/10.5194/acp-14-10119-2014>, 2014.
- Safieddine, S., Boynard, A., Hao, N., Huang, F., Wang, L., Ji, D., Barret, B., Ghude, S. D., Coheur, P.-F., Hurtmans, D., et al.: Tropospheric ozone variability during the East Asian summer monsoon as observed by satellite (IASI), aircraft (MOZAIC) and ground stations, *Atmospheric Chemistry and Physics*, 16, 10 489–10 500, <https://doi.org/10.5194/acp-16-10489-2016>, 2016.
- 760 Sahu, L., Sheel, V., Kajino, M., Gunthe, S. S., Thouret, V., Nedelec, P., and Smit, H. G.: Characteristics of tropospheric ozone variability over an urban site in Southeast Asia: A study based on MOZAIC and MOZART vertical profiles, *Journal of Geophysical Research: Atmospheres*, 118, 8729–8747, <https://doi.org/10.1002/jgrd.50662>, 2013.
- Sahu, L., Sheel, V., Kajino, M., Deushi, M., Gunthe, S. S., Sinha, P., Sauvage, B., Thouret, V., and Smit, H. G.: Seasonal and interannual variability of tropospheric ozone over an urban site in India: A study based on MOZAIC and CCM vertical profiles over Hyderabad, *Journal of Geophysical Research: Atmospheres*, 119, 3615–3641, <https://doi.org/10.1002/2013JD021215>, 2014.
- 765 Saunio, M., Reeves, C., Mari, C., Murphy, J., Stewart, D. J., Mills, G., Oram, D., and Purvis, R.: Factors controlling the distribution of ozone in the West African lower troposphere during the AMMA (African Monsoon Multidisciplinary Analysis) wet season campaign, *Atmospheric Chemistry and Physics*, 9, 6135–6155, <https://doi.org/10.5194/acp-9-6135-2009>, 2009.
- 770 Sauvage, B., Thouret, V., Cammas, J.-P., Gheusi, F., Athier, G., and Nédélec, P.: Tropospheric ozone over Equatorial Africa: regional aspects from the MOZAIC data, *Atmospheric Chemistry and Physics*, 5, 311–335, <https://doi.org/10.5194/acp-5-311-2005>, 2005.
- Sauvage, B., Thouret, V., Thompson, A., Witte, J., Cammas, J.-P., Nédélec, P., and Athier, G.: Enhanced view of the “tropical Atlantic ozone paradox” and “zonal wave one” from the in situ MOZAIC and SHADOZ data, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006241>, 2006.

- 775 Sauvage, B., Gheusi, F., Thouret, V., Cammas, J.-P., Duron, J., Escobar, J., Mari, C., Mascart, P., and Pont, V.: Medium-range mid-tropospheric transport of ozone and precursors over Africa: two numerical case studies in dry and wet seasons, *Atmospheric Chemistry and Physics*, 7, 5357–5370, <https://doi.org/10.5194/acp-7-5357-2007>, 2007a.
- Sauvage, B., Martin, R., Van Donkelaar, A., Liu, X., Chance, K., Jaeglé, L., Palmer, P., Wu, S., and Fu, T.-M.: Remote sensed and in situ constraints on processes affecting tropical tropospheric ozone, *Atmospheric Chemistry and Physics*, 7, 815–838, <https://doi.org/10.5194/acp-7-815-2007>, 2007b.
- 780 Sauvage, B., Martin, R. V., Van Donkelaar, A., and Ziemke, J.: Quantification of the factors controlling tropical tropospheric ozone and the South Atlantic maximum, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD008008>, 2007c.
- Sauvage, B., Thouret, V., Cammas, J.-P., Brioude, J., Nedelec, P., and Mari, C.: Meridional ozone gradients in the African upper troposphere, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL028542>, 2007d.
- 785 Sauvage, B., Fontaine, A., Eckhardt, S., Auby, A., Boulanger, D., Petetin, H., Paugam, R., Athier, G., Cousin, J.-M., Darras, S., et al.: Source attribution using FLEXPART and carbon monoxide emission inventories: SOFT-IO version 1.0, *Atmospheric Chemistry and Physics*, 17, 15 271–15 292, <https://doi.org/10.5194/acp-17-15271-2017>, 2017.
- Sheel, V., Sahu, L., Kajino, M., Deushi, M., Stein, O., and Nedelec, P.: Seasonal and interannual variability of carbon monoxide based on MOZAIC observations, MACC reanalysis, and model simulations over an urban site in India, *Journal of Geophysical Research: Atmospheres*, 119, 9123–9141, <https://doi.org/10.1002/2013JD021425>, 2014.
- 790 Singh, H., Herlth, D., Kolyer, R., Chatfield, R., Viezee, W., Salas, L., Chen, Y., Bradshaw, J., Sandholm, S., Talbot, R., et al.: Impact of biomass burning emissions on the composition of the South Atlantic troposphere: Reactive nitrogen and ozone, *Journal of Geophysical Research: Atmospheres*, 101, 24 203–24 219, <https://doi.org/10.1029/96JD01018>, 1996.
- Stein, O., Schultz, M. G., Bouarar, I., Clark, H., Huijnen, V., Gaudel, A., George, M., and Clerbaux, C.: On the wintertime low bias of Northern Hemisphere carbon monoxide found in global model simulations, *Atmospheric chemistry and physics*, 14, 9295–9316, <https://doi.org/10.5194/acp-14-9295-2014>, 2014.
- 795 Stevenson, D., Young, P., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsoren, S. B., Myhre, G., Berntsen, T. K., et al.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmospheric Chemistry and Physics*, 13, 3063–3085, <https://doi.org/10.5194/acp-13-3063-2013>, 2013.
- 800 Stohl, A., Bonasoni, P., Cristofanelli, P., Collins, W., Feichter, J., Frank, A., Forster, C., Gerasopoulos, E., Gäggeler, H., James, P., et al.: Stratosphere-troposphere exchange: A review, and what we have learned from STACCATO, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002490>, 2003.
- Tang, Q., Prather, M., and Hsu, J.: Stratosphere-troposphere exchange ozone flux related to deep convection, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2010GL046039>, 2011.
- 805 Tarasick, D., Carey-Smith, T., Hocking, W., Moeini, O., He, H., Liu, J., Osman, M., Thompson, A., Johnson, B., Oltmans, S., et al.: Quantifying stratosphere-troposphere transport of ozone using balloon-borne ozonesondes, radar windprofilers and trajectory models, *Atmospheric Environment*, 198, 496–509, <https://doi.org/10.1016/j.atmosenv.2018.10.040>, 2019.
- Thompson, A. M., Witte, J. C., Hudson, R. D., Guo, H., Herman, J. R., and Fujiwara, M.: Tropical tropospheric ozone and biomass burning, *Science*, 291, 2128–2132, <https://doi.org/10.1126/science.291.5511.2128>, 2001.
- 810 Thompson, A. M., Witte, J. C., McPeters, R. D., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W., Posny, F., Coetzee, G. J., et al.: Southern hemisphere additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 1. Comparison

- with Total ozone mapping spectrometer (TOMS) and ground-based measurements, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2001JD000967>, 2003a.
- 815 Thompson, A. M., Witte, J. C., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W., Posny, F., Coetzee, G. J., Hoegger, B., et al.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 2. Tropospheric variability and the zonal wave-one, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002241>, 2003b.
- Thompson, A. M., Stauffer, R. M., Wargan, K., Witte, J. C., Kollonige, D. E., and Ziemke, J. R.: Regional and Seasonal trends in tropical ozone from SHADOZ profiles: Reference for models and satellite products, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034691, <https://doi.org/10.1029/2021JD034691>, 2021.
- 820 Thouret, V., Clark, H., Petzold, A., Nédélec, P., and Zahn, A.: IAGOS: Monitoring Atmospheric Composition for Air Quality and Climate by Passenger Aircraft, pp. 1–14, Singapore: Springer Nature Singapore, https://doi.org/10.1007/978-981-15-2527-8_57-1, 2022.
- van der A, R. J., Mijling, B., Ding, J., Koukouli, M. E., Liu, F., Li, Q., Mao, H., and Theys, N.: Cleaning up the air: effectiveness of air quality policy for SO₂ and NO_x emissions in China, *Atmospheric Chemistry and Physics*, 17, 1775–1789, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-17-1775-2017)
- 825 17-1775-2017, 2017.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr, A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmospheric Chemistry and Physics*, 6, 3423–3441, [https://doi.org/10.5194/acp-6-3423-](https://doi.org/10.5194/acp-6-3423-2006)
- 2006, 2006.
- Wang, T., Xue, L., Brimblecombe, P., Lam, Y. F., Li, L., and Zhang, L.: Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects, *Science of the Total Environment*, 575, 1582–1596, <https://doi.org/10.1016/j.scitotenv.2016.10.081>, 2017.
- 830 Wild, O., Pochanart, P., and Akimoto, H.: Trans-Eurasian transport of ozone and its precursors, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004501>, 2004.
- Yamasoe, M. A., Sauvage, B., Thouret, V., Nédélec, P., Le Flochmoen, E., and Barret, B.: Analysis of tropospheric ozone and carbon monoxide profiles over South America based on MOZAIC/IAGOS database and model simulations, *Tellus B: Chemical and Physical Meteorology*, 67, 27884, <https://doi.org/10.3402/tellusb.v67.27884>, 2015.
- 835 Zhang, Y., Cooper, O. R., Gaudel, A., Thompson, A. M., Nédélec, P., Ogino, S.-Y., and West, J. J.: Tropospheric ozone change from 1980 to 2010 dominated by equatorward redistribution of emissions, *Nature Geoscience*, 9, 875–879, <https://doi.org/10.1038/ngeo2827>, 2016.
- Zhang, Y., West, J. J., Emmons, L. K., Flemming, J., Jonson, J. E., Lund, M. T., Sekiya, T., Sudo, K., Gaudel, A., Chang, K.-L., et al.: Contributions of world regions to the global tropospheric ozone burden change from 1980 to 2010, *Geophysical Research Letters*, 48, e2020GL089184, <https://doi.org/10.1029/2020GL089184>, 2021.
- 840 Zheng, B., Chevallier, F., Ciais, P., Yin, Y., Deeter, M. N., Worden, H. M., Wang, Y., Zhang, Q., and He, K.: Rapid decline in carbon monoxide emissions and export from East Asia between years 2005 and 2016, *Environmental Research Letters*, 13, 044007, <https://doi.org/10.1088/1748-9326/aab2b3>, 2018.
- 845 Zheng, B., Chevallier, F., Yin, Y., Ciais, P., Fortems-Cheiney, A., Deeter, M. N., Parker, R. J., Wang, Y., Worden, H. M., and Zhao, Y.: Global atmospheric carbon monoxide budget 2000–2017 inferred from multi-species atmospheric inversions, *Earth System Science Data*, 11, 1411–1436, <https://doi.org/10.5194/essd-11-1411-2019>, 2019.
- Ziemke, J., Chandra, S., Duncan, B., Schoeberl, M., Torres, O., Damon, M., and Bhartia, P.: Recent biomass burning in the tropics and related changes in tropospheric ozone, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL039303>, 2009.

850 Ziemke, J. R., Oman, L. D., Strode, S. A., Douglass, A. R., Olsen, M. A., McPeters, R. D., Bhartia, P. K., Froidevaux, L., Labow, G. J., Witte, J. C., et al.: Trends in global tropospheric ozone inferred from a composite record of TOMS/OMI/MLS/OMPS satellite measurements and the MERRA-2 GMI simulation, *Atmospheric Chemistry and Physics*, 19, 3257–3269, <https://doi.org/10.5194/acp-19-3257-2019>, 2019.