

Supplement of

Tropical tropospheric ozone and carbon monoxide distributions: characteristics, origins and control factors, as seen by IAGOS and IASI

Maria Tsivlidou¹, Bastien Sauvage^{1,*}, Brice Barret^{1,*}, Pawel Wolff², Hannah Clark³,
Yasmine Bennouna¹, Romain Blot¹, Damien Boulanger², Philippe Nédélec¹, Eric Le Flochmoën¹, and
Valérie Thouret¹

¹Laboratoire d'Aérologie (LAERO), Université Toulouse III – Paul Sabatier, CNRS, Toulouse, France

²Observatoire Midi-Pyrénées (OMP-SEDOO), Université Toulouse III - Paul Sabatier, CNRS, Toulouse, France

³IAGOS-AISBL, 98 Rue du Trône, Brussels, Belgium

*These authors contributed equally to this work.

Correspondence: Maria Tsivlidou (maria.tsivlidou@aero.obs-mip.fr)

S1 CO emissions

S1.1 Biomass burning

GFAS Wildfire CO Emissions 2003-2017




Figure S1. GFAS BB CO emissions (in $\text{kg m}^{-2} \text{ s}^{-1}$) averaged from 2003 to 2017.

S1.2 Anthropogenic

CEDS v2 Anthropogenic CO emissions 2000-2019




Figure S2. CEDS AN CO emissions (in $kg m^{-2} s^{-1}$) averaged from 2000 to 2019.

Horizontal wind at 850 hPa - 2008 to 2019




Figure S3. ECMWF Reanalysis 5th Generation (ERA5) wind vectors at 850 (a–d), 550 (e–h) and 250 (i–l) hPa averaged from 2008 to 2020 at $0.25 \times 0.25^\circ$ resolution.

Horizontal wind at 550 hPa - 2008 to 2019




Figure S3. Continued.

Horizontal wind at 250 hPa - 2008 to 2019




Figure S3. Continued.