



Hydrogen peroxide in the upper tropical troposphere over the Atlantic Ocean and western Africa during the CAFE-Africa aircraft campaign

5 Zaneta Hamryszczak¹, Dirk Dienhart¹, Bettina Brendel¹, Roland Rohloff¹, Daniel Marno¹, Monica Martinez¹, Hartwig Harder¹, Andrea Pozzer^{1,4}, Birger Bohn², Martin Zöger³, Jos Lelieveld^{1,4}, and Horst Fischer¹

¹Atmospheric Chemistry Department, Max Planck Institute for Chemistry, Mainz, 55128, Germany

²Institute of Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich GmbH, Jülich, 52428, Germany

³Flight Experiments, German Aerospace Center (DLR), Oberpfaffenhofen, 82234 Germany

10 ⁴Climate and Atmosphere Research Center, The Cyprus Institute, Nicosia, 1645, Cyprus

Correspondence: Zaneta Hamryszczak (z.hamryszczak@mpic.de) and Horst Fischer (horst.fischer@mpic.de)

Abstract. This study focuses on the distribution of hydrogen peroxide (H₂O₂) in the upper tropical troposphere at altitudes between 8 and 15 km based on *in situ* observations during the Chemistry of the Atmosphere – Field Experiment in Africa (CAFE-Africa) campaign conducted in August–September 2018 over the tropical Atlantic Ocean and western Africa. The measured hydrogen peroxide mixing ratios in the upper troposphere show a nearly uniform latitudinal distribution with locally increased levels (up to 1 ppb_v) within the Intertropical Convergence Zone (ITCZ), over the African coastal area, as well as during measurements performed in proximity of the tropical storm Florence (later developing into a hurricane), indicating the influence of convective transport processes. The measurements are compared to observation-based photostationary steady-state (PSS) calculations and numerical simulations by the global EMAC model. North of the ITCZ, PSS calculations produce lower H₂O₂ mixing ratios relative to the observations. Here observed mixing ratios exceed the PSS calculations by up to a factor of 2. On the other hand, PSS calculations overestimate the H₂O₂ mixing ratios south of the ITCZ by a factor of up to 3. The significant influence of convection in the ITCZ and the enhanced presence of clouds towards the southern hemisphere indicate contributions of atmospheric transport and cloud scavenging in the probed region. Differences between H₂O₂ observations and simulations of local PSS indicate that convective transport in the ITCZ region and consequent redistribution of H₂O₂ towards the north and south impacts the spatial distribution of H₂O₂ in the upper troposphere. Simulations performed by EMAC analogously overestimate hydrogen peroxide levels particularly in the southern hemisphere, most likely due to underestimated cloud scavenging. Latitudinal distribution analysis indicates a gradient from the equator towards the subtropics both in the EMAC simulations and the PSS calculations. On the other hand, the measurements display nearly uniform mixing ratios of the species in the upper troposphere with a slight decrease from the ITCZ towards the subtropics, indicating a relatively low dependency on the solar radiation inclination and the corresponding photolytic activity. The highest deviations relative to the observations correspond with the underestimated hydrogen peroxide loss due to enhanced cloud presence, scavenging, and rainout in the ITCZ and towards the south.

15
20
25
30



1 Introduction

35 The key role of hydrogen peroxide in the oxidative chemistry of the troposphere is well acknowledged (Lelieveld and Crutzen, 1990; Crutzen et al., 1999). On the one hand, H₂O₂ serves as a reservoir of HO_x (OH + HO₂ = HO_x) species, which are the most prominent oxidants controlling the self-cleansing capacity of the atmosphere (Levy, 1971; Logan et al., 1981; Kleinman, 1991). On the other hand, hydrogen peroxide can oxidize SO₂ and to a minor extent NO₂ and convert them into H₂SO₄ and HNO₃ in clouds, rain, and fog, leading to their acidification (Hoffmann and Edwards, 1975; Penkett et al., 1979; Robbin Martin and Damschen, 1981; Damschen and Martin, 1983; Calvert et al., 1985).

40 The most prominent pathway leading to H₂O₂ production is the recombination of HO₂ radicals (R4). These precursors of H₂O₂ are mainly produced via oxidation of carbon monoxide by OH radicals, which are formed during the photolysis of ozone and the subsequent reaction of O¹D with water vapor in the troposphere (R1 – R3). The budget of H₂O₂ is thus controlled by the fate of the HO₂ radical, which is determined by a competition between its self-reaction leading to H₂O₂ formation and its conversion to OH via reaction with NO and O₃.



The global distribution of hydrogen peroxides is not only dependent on the chemical composition of the atmosphere but also on meteorological conditions. The amount of hydrogen peroxide is strongly dependent on the availability of water vapor and near-UV radiation (Jacob and Klockow, 1992; Perros, 1993; Slemr and Tremmel, 1994; Snow, 2003; Snow et al., 2007; Klippel et al., 2011). Towards the tropopause as well as towards the poles, the amount of water vapor generally decreases, resulting in a reduced primary production of HO_x radicals. Additionally, with increasing altitudes near-UV radiation and therefore H₂O₂ photolysis increase leading to a pronounced production of OH via H₂O₂ photolysis in R5 (Jaeglé et al., 1997; Jaeglé et al., 2000; Faloon et al., 2000; Faloon et al., 2004; Lee et al., 2000). On the other hand, the availability of hydrogen peroxide precursors is diminished with increasing latitude, due to the decreasing inclination of solar radiation and reduced amount of water vapor towards the poles. Physical loss of hydrogen peroxide occurs through deposition processes promoted by its high solubility (Walcek, 1987; Chang et al., 2004; Nguyen et al., 2015). The aqueous uptake and subsequent removal of H₂O₂ strongly depends upon the uptake by aerosols and clouds (O'Sullivan et al., 1999). In effect, the vertical distribution of H₂O₂ often follows an inverted c-shape with decreased mixing ratios within the boundary layer and the upper troposphere and a local maximum in the middle troposphere at altitudes between 2 and 5 km. Additionally, observations indicate a decreasing trend approximately from the equator towards the north and south (Daum et al., 1990; Heikes, 1992; O'Sullivan et al., 1996; Weinstein-Lloyd et al., 1998; Snow, 2003; Snow et al., 2007; Klippel et al., 2011).

60
65



In general, the global budget of hydrogen peroxide is significantly affected by anthropogenic as well as natural emissions of nitrogen oxides. In urban areas, the formation of hydrogen peroxide is diminished by the increased mixing ratios of NO_x ($\text{NO}_x = \text{NO}_2 + \text{NO}$) derived from anthropogenic sources, as the self-reaction of HO_2 to H_2O_2 is suppressed by the much faster reaction of HO_2 with NO (Lee et al., 2000; Reeves and Penkett, 2003). In contrast, biomass burning events lead to significant
70 injections of additional hydrogen peroxide through primary as well as secondary chemical production (Lee et al., 1997; Rinsland et al., 2007; Snow et al., 2007; Allen et al., 2022). Finally, convection processes are considered to increase the mixing ratios of hydrogen peroxide in the upper troposphere (Jaeglé et al., 1997; Jaeglé et al., 2000; Klippel et al., 2011; Bozem et al., 2017). Especially within the Intertropical Convergence Zone (ITCZ), convective processes play a key role in the transport of a large suite of trace species to higher tropospheric layers. The ITCZ is a low-pressure region, which marks the meeting zone
75 of airmasses transported from both hemispheres and constitutes the ascending branch of the Hadley circulation (Waliser and Gautier, 1993). Due to the high sea surface temperatures, strong solar radiation and increased air humidity, this band-like area near the equator is mainly characterized by highly dynamic weather phenomena, namely, large convective cumulonimbus clouds penetrating deep into the upper troposphere (Hastenrath and Lamb, 1977; Waliser and Gautier, 1993). Thus, convective transport processes might contribute to increased levels of hydrogen peroxide in the upper troposphere and promote elevated
80 HO_x levels via subsequent photochemical processes involving H_2O_2 degradation (Jaeglé et al., 1997; Jaeglé et al., 2000). Numerous airborne measurements of hydrogen peroxide were performed over the past decades over the Atlantic Ocean and in proximity to the ITCZ. The majority of these studies focused on the troposphere in the northern hemisphere, providing an overview on the vertical and latitudinal distribution of hydrogen peroxide.

In September and October 1992, as part of the NASA's Global Tropospheric Experiment (GTE) program, the Transport and
85 Atmospheric Chemistry Near the Equatorial Atlantic (TRACE A) mission took place over the Atlantic Ocean. The mean observed mixing ratio of hydrogen peroxide was approximately 0.2 ppb_v in the upper troposphere (8–12 km) (Prather and Jacob, 1997; O'Sullivan et al., 1999). During the Subsonic Assessment Ozone and Nitrogen Oxide Experiment (SONEX) campaign, which took place in autumn 1997 over the North Atlantic, mean values of 0.12 ppb_v (median: 0.08 ppb_v) specifically in the upper troposphere were observed (Snow et al., 2007). Allen et al. (2013) presented satellite-based global distribution
90 data of H_2O_2 in the mid-to-upper troposphere obtained by the Atmospheric Chemistry Experiment (ACE) mission and reported mean levels of 0.10–0.28 ppb_v in the upper tropical troposphere (>8 km), symmetrically decreasing towards the poles. During the Atmospheric Tomography Mission (ATom) performed in August 2016 (ATom-1) and October 2017 (ATom-3), mean values ranging between 0.05 ppb_v up to 0.25 ppb_v were measured over the Mid-Atlantic Ocean (–20°–20°N; Allen et al., 2022). Please note that the average values within the upper troposphere cited here are obtained from the information presented in the
95 figures in the study. Further, Hottmann et al. (2020) deduced mean ($\pm 1\sigma$) and median hydrogen peroxide mixing ratios of 0.17 (± 0.09) ppb_v and 0.15 ppb_v , respectively, during the Oxidation Mechanism Observation (OMO) mission in summer 2015, which took place over the Arabian Peninsula, the eastern Mediterranean, and northern Indian Ocean covering the marine ITCZ region east of the African continent.



100 Numerous measurements have been performed in the marine tropical troposphere. In this study, we address the budget of hydrogen peroxide specifically in the upper tropical troposphere within the equatorial Atlantic region with a main focus on the ITCZ. Our objective was to study the distribution of trace gases and radicals over the Central Atlantic and the possible impact of convection in the ITCZ on the abundance of H₂O₂ in the upper troposphere.

2 CAFE-Africa campaign

105 The distribution of hydrogen peroxide (H₂O₂) was measured in the free troposphere over the Atlantic Ocean during the Chemistry of the Atmosphere: Field Experiment in Africa (CAFE-Africa) campaign. The major objective of the mission was to investigate the large-scale distribution of trace gases, radicals, and aerosols in the tropical eastern Atlantic and along the western coast of Africa. In particular, the influence of biomass burning emissions and long-distance pollution transport on the atmosphere's oxidation capacity and the chemical processing of trace gases and aerosols in clean and polluted air masses were studied.

110 The campaign took place in August and September 2018 during the West African monsoon. During this period, 14 measurement flights were made over the Atlantic Ocean and the African coast with the High Altitude and Long-Range research aircraft (HALO) operating from the international airport on Sal, Cape Verde (16.75°N, -22.95°E). The flights focused on the upper troposphere up to an altitude of 15 km with a few vertical profiles mostly in the northern hemisphere. The probed area covered a latitudinal and longitudinal range from approximately -10°N–50°N and -50°E–15°E. An overview of the
115 corresponding flight dates and the objectives of the individual flights was presented by Tadic et al. (2021). The flight tracks color-coded by GPS flight altitudes are presented in Fig. 1.




Figure 1. Flight tracks and the sampled region during the CAFE-Africa campaign color-coded by the GPS flight altitude. The majority of flights were performed from the base of operations in Sal, Cape Verde.

120 Related to the location of the base of operations on Cape Verde, the majority of the flights were performed in close proximity to the ITCZ, which allowed the study of tropical trace gases and aerosol distributions in both hemispheres. During the campaign, the ITCZ roughly covered positions between approximately 5°N and 15°N (Tadic et al., 2021). Information on the meteorological conditions with special emphasis on the total cloud coverage and convective precipitation located mostly within the ITCZ (5°N–20°N) are presented in the Supplement of this work (Figs. S1–3).

125 3 Methods

3.1 Hydrogen peroxide measurements

Hydroperoxides were measured as the sum of organic hydroperoxides and hydrogen peroxide and were determined using a wet chemical system named the Hydrogen Peroxide and Higher Organic Peroxides monitor (HYPHOP;(Klippel et al., 2011; Bozem et al., 2017; Hottmann et al., 2020; Hamryszczak et al., 2022) based on a previous design by Lazrus et al. (Lazrus et al., 1985; Lazrus et al., 1986). The ambient air was sampled from the top of the aircraft fuselage via a trace gas inlet (TGI) with a 1/2" PFA liner that left the cabin again through a second exhaust line. From this bypass, a 1/4" PFA sampling line is
130 connected to a Teflon-coated membrane pump (Type MD 1C; Vacuubrand, Wertheim, Germany) and a pressure control unit regulating the pump speed to a line pressure of 1000 hPa (Constant Pressure Inlet; CPI). The CPI provides a constant inlet pressure covering external pressure variations between 1000 and 150 hPa. Following the CPI inlet, the ambient air passed



135 through a stripping coil with a buffered sampling solution (potassium hydrogen phthalate/NaOH; pH 6; stripping efficiency of
1 for hydrogen peroxide and between 0.6 and 1 for organic peroxides; Lee et al., 2000). The hydroperoxide solution was probed
in two individual channels in response to addition of p-hydroxyphenyl acetic acid (POPHA) and horseradish peroxidase (HRP).
The formed chemiluminescent 6,6'-dihydroxy-3,3'-biphenyldiacetic acid was detected via fluorescence spectroscopy using a
Cd pen-ray lamp at 326 nm. The hydroperoxide-specific fluorescence (Guilbault et al., 1968) was detected at 400–420 nm
140 using photomultiplier tubes for each channel separately.

Hydrogen peroxide mixing ratios are then calculated from the difference between the entirety of the measured peroxides
(Channel A) and the sum of organic ROOH hydroperoxides (Channel B), where H₂O₂ is selectively destroyed by the addition
of catalase. Further information on the mixing ratios of individual organic peroxides cannot be provided by the monitor, due
to the characteristics of the measurement technique. Please note that especially within the boundary layer and due to biomass
145 burning, a variety of organic peroxides might contribute to the total measured signal of organic peroxide (Fels and Junkermann,
1994; Slemr and Tremmel, 1994; Valverde-Canossa et al., 2005; Hua et al., 2008; Dienhart et al., 2022).

Prior to ambient measurements, both channels are simultaneously calibrated using a liquid standard (0.98 μmol L⁻¹) produced
from serial dilution of a H₂O₂ stock solution. The H₂O₂ destruction efficiency in channel B corresponding to the added catalase
was determined to be 0.95–0.98 based on liquid calibrations. In-flight background measurements were performed using
150 purified zero air, generated by a cartridge with silica gel (type IAC-502; Infiltec, Speyer, Germany) and hopcalite (type IAC-
330; Infiltec, Speyer, Germany). Using a gas-phase calibration source (LDPE permeation devices), the H₂O₂ transmission
efficiency through the inlet (± 1σ) was determined regularly by measuring the difference between the addition of the standard
before and after the CPI and was found to be 0.61 ± 0.06. Due to a positive ozone interference, the H₂O₂ data was further
corrected by subtraction of 0.056 ppb_v H₂O₂/100 ppb_v O₃ based on a scatter plot of hydrogen peroxide vs. ozone mixing ratios
155 in the lower stratosphere, assuming that ambient H₂O₂ above the tropopause is essentially zero. The total measurement
uncertainty (TMU) of the monitor was estimated as:

$$\text{TMU} = \sqrt{((P)^2 + (US)^2 + (UOI)^2 + (UTE)^2)} \quad (1)$$

by considering the instrument's precision (P), uncertainty of the standard (US), uncertainty of the H₂O₂ transmission efficiency
(UTE), and the uncertainty of the ozone interference (UOI). The determined precision with 1 sigma confidence interval was
160 determined from the reproducibility of the liquid calibrations performed during the campaign to be 1.3% at 5.46 ppb_v for
hydrogen peroxide and 0.8% at 5.64 ppb_v for the organic hydroperoxides. The uncertainty of the standard was included in the
instrument precision calculation. The uncertainty of the transmission efficiency was calculated to be 6%. The total
measurement uncertainty was determined to be 9% for hydrogen peroxide and 41% for the sum of organic hydroperoxides.
The total measurement uncertainty of organic hydroperoxides is increased by 40% due to the varying solubility of individual
165 organic hydroperoxides in aqueous solution, which ranges between 60% (e.g., MHP) and 100% (e.g., PAA). The instrumental
time resolution was determined to be 122 s based on the calibration signal rise and fall time from 10% to 90% and 90% to
10%, respectively. The detection limit with a 2σ confidence was derived from the reproducibility of the in-flight background
measurements as 15 ppt_v for hydrogen peroxide and 6 ppt_v for the sum of organic hydroperoxides, respectively. For the



170 purposes of this study, hydrogen peroxide data were filtered for stratospheric influences by removing all data points with ozone mixing ratios higher than 100 ppb_v.

3.2 Measurement of other relevant species

175 GPS, temperature, pressure, and windspeed were obtained using the Basic HALO Measurement And Sensor System, BAHAMAS. Water vapor mixing ratios and the corresponding air humidity were measured with the Sophisticated Hygrometer for Atmospheric ResearCh (SHARC) based on a tunable diode laser (TDL) setup (Krautstrunk and Giez, op. 2012). HO_x radicals were measured by laser-induced fluorescence with the HydrOxyl Radical measurement Unit (HORUS; (Marno et al., 2020). Spectrally resolved upward and downward actinic flux density was obtained with two spectroradiometers (Bohn and Lohse, 2017). A brief overview of the campaign instrumentation, measurement methods, their TMU values, and the corresponding technical references are listed in Table 1.

180 **Table 1. Overview of observed species with corresponding measurement method, total measurement uncertainty (TMU), and references regarding the instrumentation.**

Measurement	Method	TMU	References
H ₂ O ₂	Chemiluminescence	H ₂ O ₂ : 9%; OrgPer: 41%	Hamryszczak et al., 2022
HO _x	Laser-induced Fluorescence (LIF; additional chem. conversion for HO ₂)	50%	Marno et al., 2020
Actinic flux density	Spectroradiometer	7–8% (15% for j(H ₂ O ₂))	Bohn and Lohse, 2017
H ₂ O	TDLAS	5%	Krautstrunk and Giez, 2012

3.3 Atmospheric Chemistry Model ECHAM/MESSy (EMAC)

185 For the purposes of this study, the *in situ* observations are compared to numerical simulations from the global chemistry and climate 3-D model EMAC (ECHAM/MESSy for Atmospheric Chemistry, Jöckel et al., 2010). The model numerically simulates the chemistry and dynamics of the troposphere and stratosphere using a large variety of submodels describing chemical and meteorological processes and the influences arising from anthropogenic and natural emissions from continental and marine environments (Jöckel et al., 2006). The basis of the atmospheric model is the 5th generation of the European Centre HAMburg general circulation model (ECHAM5; Roeckner et al., 2003; Roeckner et al., 2006). The communication between the various submodels is achieved by the Modular Earth Submodel System (MESSy; Jöckel et al., 2005; Jöckel et al., 2010; Jöckel et al., 2016). Atmospheric chemistry is simulated by the Module for Efficiently Calculating the Chemistry of the Atmosphere (MECCA) submodel (Sander et al., 2005; Sander et al., 2011; Sander et al., 2019), using the Mainz Organic Mechanism (MOM) and photolysis rate calculations from a radiation transfer model (Sander et al., 2014; Sander et al., 2019). 190 Primary emissions and dry deposition as well as aqueous phase chemistry in clouds and cloud scavenging are simulated by the ONLEM, OFFLEM, TNUDGE, and DRYDEP submodels (Kerkweg et al., 2006a; Kerkweg et al., 2006b) and the Scavenging of Tracers submodel (SCAV; Tost et al., 2006). Anthropogenic emissions are based on the EDGARv4.3.2 inventory (European



195 Joint Center; JRC; Crippa et al., 2018) and are distributed vertically according to (Pozzer et al., 2009). Biomass burning
emissions were simulated based on the Global Fire Assimilation System (GFAS; Kaiser et al., 2012). The model has a vertical
and horizontal resolution of 47 vertical levels up to 0.01 hPa at T63 (i.e., approximately $1.8^\circ \times 1.8^\circ$), respectively, and a time
resolution of 6 min. The model was further weakly nudged towards the ECMWF ERA-Interim data (Tadic et al., 2021). This
model set-up has been extensively evaluated for different trace gases and aerosols (Pozzer et al., 2022). For comparison to
200 observations, the simulation results were interpolated on the GPS flight tracks using the S4D submodel (Jöckel et al., 2010).

3.4 Simulations based on photostationary steady-state conditions

The hydrogen peroxide mixing ratios in the upper troposphere under the assumption of photostationary steady-state conditions
were calculated based on measured precursors and photochemical loss processes of hydrogen peroxide (HO_2 , OH, $j(\text{H}_2\text{O}_2)$)
and the rate coefficient data from Atkinson et al., 2004).

205 In the upper troposphere, the production rate $P(\text{H}_2\text{O}_2)$ of hydrogen peroxide due to the self-reaction of HO_2 can be calculated
from Eq. (2). The photochemical loss rate of hydrogen peroxide, $L(\text{H}_2\text{O}_2)$ can be derived from H_2O_2 photolysis and the reaction
with OH as shown in Eq. (3). Please note that because of low mixing ratios of other trace gases producing hydrogen peroxide
in the upper troposphere, other potential photochemical sources and sinks of H_2O_2 were not considered. The equations are
derived from the most prominent reaction pathways presented in Sect. 1 (R4–R6).

$$210 \quad P(\text{H}_2\text{O}_2) = k_{\text{HO}_2+\text{HO}_2} \cdot [\text{HO}_2]^2 \quad (2)$$

$$L(\text{H}_2\text{O}_2) = \left(k_{\text{H}_2\text{O}_2+\text{OH}} \cdot [\text{OH}] + j(\text{H}_2\text{O}_2) \right) \cdot [\text{H}_2\text{O}_2] \quad (3)$$

The calculations of the rate coefficients were based on the measured parameters along the flight tracks according to Atkinson
et al., 2004; Eqs. 4–5). Due to the water dependence of the hydrogen peroxide production rate coefficient, both expressions in
Eq. (4) were further extended by the factor $1 + 1.4 \times 10^{-21} \text{ cm}^3 [\text{H}_2\text{O}] \exp(2200/T)$.

$$215 \quad k_{\text{HO}_2+\text{HO}_2} = 2.2 \cdot 10^{-13} \cdot \exp\left(\frac{600}{T}\right) \text{ cm}^3 \text{ s}^{-1} + 1.9 \cdot 10^{-33} [\text{N}_2] \cdot \exp\left(\frac{980}{T}\right) \text{ cm}^6 \text{ s}^{-1} \quad (4)$$

$$k_{\text{H}_2\text{O}_2+\text{OH}} = 2.9 \cdot 10^{-12} \cdot \exp\left(\frac{-160}{T}\right) \text{ cm}^3 \text{ s}^{-1} \quad (5)$$

Neglecting deposition and transport processes impacting the hydrogen peroxide budget, the mixing ratio of H_2O_2 was
calculated via Eq. (6).

$$[\text{H}_2\text{O}_2]^{\text{PSS}} = \frac{[\text{HO}_2]^2 \cdot k_{\text{HO}_2+\text{HO}_2}}{[\text{OH}] \cdot k_{\text{H}_2\text{O}_2+\text{OH}} + j(\text{H}_2\text{O}_2)} \quad (6)$$

220 3.5 Data processing details

For the purpose of the present study, we used spatially interpolated EMAC simulations of H_2O_2 , OH, HO_2 , water vapor,
 $j(\text{H}_2\text{O}_2)$, temperature, and pressure to supplement the measurement observations. To synchronize the time stamp of the



simulated data with the measurement output, we calculated a mean of the measurement data with a matching temporal resolution of 6 min.

225 Vertical profiles of all species under investigation were calculated as 1000 m bins (means and medians) over the entire probed atmospheric column. Profile information is restricted to 30 take-offs and landings at Sal, while other areas are not considered due to a lack of statistically significant data.


The spatially resolved data based on measurements, PSS model calculations, and EMAC simulations were binned into $1^\circ \times 1^\circ$ subsets over the full extension of the flight tracks in the upper troposphere (≥ 8 km).

230 The latitudinal distribution of the species was examined using 2.5° bins over the upper tropospheric region. Please note that due to a reduced amount of data in the lower troposphere, the analysis of spatial and latitudinal distributions was restricted to measurements performed in the upper troposphere.

4 Results

4.1 Observations of hydrogen peroxide during CAFE-Africa and previous airborne measurements

235 The observed mixing ratios of hydrogen peroxide during the CAFE-Africa campaign are presented as a latitude vs. longitude plot with mean mixing ratio values binned into a subset of $1^\circ \times 1^\circ$ bins for the entirety of the upper troposphere (≥ 8 km) along the flight tracks (Fig. 2). The color scale represents the measured mixing ratio of H_2O_2 .



240 **Figure 2.** Spatial distribution of measured hydrogen peroxide in the upper troposphere (≥ 8 km) during the CAFE-Africa campaign. Data were binned into $1^\circ \times 1^\circ$ bins over the full extension of the flight tracks.



The mean ($\pm 1\sigma$) and median mixing ratios during the campaign were $0.18 (\pm 0.13)$ ppb_v and 0.15 ppb_v, respectively, with maximum hydrogen peroxide mixing ratios reaching 1.03 ppb_v. Slightly higher H₂O₂ levels were observed in the ITCZ (approx. 5°N–20°N), where locally mixing ratios up to 0.67 ppb_v were observed. These maxima are most likely due to atmospheric transport of H₂O₂ in continental air masses transported from central and southern Africa into the upper troposphere. This is in accordance with previous reports about increased hydrogen peroxide mixing ratios due to biomass burning and convective activity, elevating the H₂O₂ mixing ratios in the upper troposphere (Lee et al., 1998; O'Sullivan et al., 1999; Allen et al., 2022). Locally enhanced H₂O₂ was further observed during the measurement flight in close proximity to the tropical storm Florence on 2nd of September 2018 (approx. 18°N, - 38.5°E). Here, the mean mixing ratios were between 0.10 ppb_v and 0.43 ppb_v with a maximum of up to 0.94 ppb_v. Based on the high convective activity during the tropical storm, as reported by Nussbaumer et al. (2021), the H₂O₂ mixing ratios were expected to rise due to the rapid transport of air masses from the MBL into the UT. Table 2 gives an overview of the estimated mean and median hydrogen peroxide mixing ratios measured during CAFE-Africa in relation to previous airborne measurements performed at a comparable latitudinal range.

Table 2. Comparison of hydrogen peroxide mean and median mixing ratios (ppb_v) in the upper troposphere during CAFE-Africa with measurements from previous campaigns (TRACE A, SONEX, OMO, ATom-1 and ATom-3; O'Sullivan et al., 1999; Snow et al., 2007; Hottmann et al., 2020; Allen et al., 2022).

	CAFE-Africa	TRACE A	SONEX	OMO	ATom-1		ATom-3	
	10°S–40°N	40°S–15°N	15°–60°N	0–50°N	20°–60°N	20°S–20°N	20°–60°N	20°S–20°N
Mean	0.18	<0.20	0.12	0.16	0.55	0.61	0.18	0.40
Median	0.15	0.15	0.08	0.15	0.29	0.27	0.12	0.20

The mean and median values during CAFE-Africa are comparable to previously reported mixing ratios during TRACE A and OMO campaigns, which covered a comparable latitude and altitude range (Tab. 2). Enhanced mixing ratios for CAFE-Africa relative to observations during the SONEX campaign are most likely due to differences in the examined regional range, since the later campaign focused on the north Atlantic. Mean and median values in the northern hemisphere (20–40°N; Tab. S1) during CAFE-Africa (0.14 ± 0.11 ppb_v and 0.12 ppb_v, respectively) are comparable to observations during SONEX. During the ATom campaigns, slightly higher mean values were observed, although median values are comparable. This could be due to differences in the probed altitudes, since ATom measurements were generally restricted to altitudes below 12 km.

Based on the comparison with previous studies, the observed mixing ratios of hydrogen peroxide during CAFE-Africa fit well into the general range of recent studies over the equatorial and subtropical Atlantic. The observed H₂O₂ distribution confirms further that mixing ratios of hydrogen peroxide in the upper troposphere seem to be far less dependent on latitude than those at lower altitudes. The latitudinal distribution of H₂O₂ during the CAFE-Africa displays a rather small symmetrical latitudinal decrease from the inner tropics into the subtropics.



270 4.2 Comparison of measured hydrogen peroxide with photostationary steady-state and EMAC calculations

In order to investigate the impact of deep convection in the ITCZ on the H_2O_2 budget in the upper troposphere, a comparison of the *in situ* data with the output of photostationary steady-state (achieved) calculations and EMAC simulations was performed. The complementary spatial distributions of the hydrogen peroxide levels were expressed as latitude versus longitude plots of mean mixing ratios aggregated over a spatial grid of $1^\circ \times 1^\circ$ in the upper troposphere (≥ 8 km) (Fig. S4 in the Supplement).

275 The calculated PSS- H_2O_2 levels range between approximately 0.01 ppb_v and 0.88 ppb_v with mean ($\pm 1\sigma$) and median mixing ratios of 0.14 (± 0.16) ppb_v and 0.07 ppb_v, respectively, which is a factor of 1.3 lower than the observations. PSS-based H_2O_2 mixing ratios tend to be higher at the southernmost coastal area (-2.5°N , -10.5°E), where the levels range between 0.40 and 0.88 ppb_v, and in the proximity to the tropical storm Florence at up to 0.40 ppb_v (approx. 18°N , -38.5°E). Hydrogen peroxide mixing ratios simulated by EMAC vary between 0.10 ppb_v and 0.75 ppb_v. The mean ($\pm 1\sigma$) and median simulated mixing ratios are 0.30 (± 0.19) ppb_v and 0.29 ppb_v, respectively, with maximum mixing ratios up to 1.04 ppb_v (-3.5°N , -9.5°E ; Fig. S4b), which is slightly higher than the observations. The spatial distributions of the point-by-point ratio between PSS calculations and EMAC simulations versus the observations in the upper troposphere above 8 km are presented in Fig. 3, which give an overview on the local differences relative to the observations varying from low ratios (yellow) to high (deep blue) values.

285 Please note that for resolution purposes the color scaling is restricted to ratios up to 4.5.




Figure 3. Spatial distribution of $\text{H}_2\text{O}_2(\text{PSS})/\text{H}_2\text{O}_2(\text{measurements})$ ratios (a) and $\text{H}_2\text{O}_2(\text{EMAC})/\text{H}_2\text{O}_2(\text{measurements})$ ratios (b) in the upper troposphere (≥ 8 km) during the CAFE-Africa campaign. Data were binned into $1^\circ \times 1^\circ$ bins over the full extension of the flight tracks.

290 Generally, the $\text{H}_2\text{O}_2(\text{EMAC})/\text{H}_2\text{O}_2(\text{measurement})$ ratios indicate a good agreement between the simulations and the measurements in the northern hemisphere ($\geq 20^\circ\text{N}$; Fig. 3b), with values between 0.95 and 1.3. With decreasing latitudes, the model tends to overestimate hydrogen peroxide, with $\text{H}_2\text{O}_2(\text{EMAC})/\text{H}_2\text{O}_2(\text{measurement})$ ratios increasing with latitude from approximately 2 to 4. Locally, most likely due to underestimated cloud scavenging, EMAC simulates highly elevated hydrogen peroxide with a factor of up to 14 higher than the measurements (4.5°N , -9.5°E).

295 Good agreement between the observation-based PSS calculations and the measurements was found in proximity to the tropical storm Florence. Here, the $\text{H}_2\text{O}_2(\text{PSS})/\text{H}_2\text{O}_2(\text{measurement})$ ratios agree between 0.83 to 1.04. Beyond the measurements taken



here, the agreement with respect to the measured hydrogen peroxide levels is generally less satisfactory. Towards the southern subtropics as well as locally in the coastal area and at the base of operations in Sal (Cape Verde), the ratios increase to 2.8, indicating an overestimation of the hydroperoxide levels relative to the observations, similar to the EMAC simulations. On the other hand, the PSS calculations tend to strongly underestimate hydrogen peroxide concentrations in the ITCZ (5–20°N) and in the northern part of the investigated region ($\geq 20^\circ\text{N}$) by factors of up to 10 and 12, respectively. This indicates that, in the upper troposphere, the local photostationary steady-state conditions based on observed radical levels do not account for all sources and sinks of hydrogen peroxide or additional injections of the H_2O_2 , which are most likely related to transport and cloud scavenging. The calculated difference between production (Eq. 2) and loss (Eq. 3) of hydrogen peroxide, $(\text{P}-\text{L})\text{H}_2\text{O}_2$ based on the observations is expressed as a latitude vs. longitude plot with mean mixing ratios binned into a subset of $1^\circ \times 1^\circ$ bins for the entirety of the upper troposphere (≥ 8 km) along the flight tracks (Fig. 4).




Figure 4. Color-coded spatial resolution of calculated deviation from PSS based on the difference between the observed hydrogen peroxide photochemical production and loss. The calculated data were binned into $1^\circ \times 1^\circ$ bins over the full extension of the flight tracks. Shaded background is the average high cloud coverage (≥ 6 km) during the measurement period based on ERA5 reanalysis results (HCC; Hersbach et al., 2019).

Generally, the majority of the probed region is loss-dominated, especially in the ITCZ (approx. 5°N – 20°N) and towards the north, where an H_2O_2 deficit of up to approximately -0.01 ppbv h^{-1} was determined. H_2O_2 production-dominated regions of up to 0.03 ppbv h^{-1} are observed towards the south and in the coastal area. The difference between the photochemical production and loss paths is directly linked to the deviations from the photostationary steady state and thus to the additional sources and



sinks of hydrogen peroxide on a local scale. These are associated with the local meteorological conditions and transport processes. Based on ERA5 reanalysis results, especially towards the south and in the coastal area, enhanced presence of clouds at altitudes above 6 km (gray shading in Fig. 4) and convective precipitation was observed during the measurement period (Hersbach, H. et al., 2019). At a mean ($\pm 1\sigma$) horizontal windspeed of 14.3 (± 7.3) m/s measured along the flight tracks and a species lifetime of 3–4 days, transport towards the subtropics impacts H_2O_2 levels and further justifies the difference between the local PSS calculations and the observations.

Based on the coincidence with the latitudinal range of the ITCZ, the enhanced loss in the H_2O_2 budget relative to the PSS in the upper troposphere is most likely due to convective injection of H_2O_2 from lower layers into the upper troposphere and the subsequent redistribution of the species towards the north and south. Below, we show that, based on the comparison with EMAC simulation output, convective transport is important for the budget of hydrogen peroxide in the upper troposphere not only in the ITCZ but also in the subtropics.

As discussed above, the comparison between observations and both PSS calculations and EMAC simulations indicate large deviations at the most southern latitudes that were visited by HALO. This is clearly demonstrated in Fig. 5, which shows observations, PSS calculations, and EMAC simulations of H_2O_2 as functions of latitude in the upper troposphere (above 8 km). The mean values of each dataset with 6 min time resolution are binned into subsets of 2.5° of latitude for the investigated region from 6°S to 40°N . The lines and the complementary shading represent mean values and the supplementary standard deviations. The area of the ITCZ between 5°N and 20°N is highlighted by gray shading.




Figure 5. Latitudinal dependence of hydrogen peroxide mixing ratios (mean $\pm 1\sigma$) compared to EMAC simulations and calculations based on PSS (red: CAFE-Africa; black: PSS CAFE-Africa; blue: EMAC; a) and calculated ratios between the simulations and the



observations and PSS-modelled calculations, and the observations, respectively (b). The data with 6 min time resolution and mean values was binned for 2.5° of latitude for altitudes ≥ 8 km. The corresponding numbers indicate the total amount of data points per bin. The shaded pattern marks the ITCZ region.

340 Contrary to the calculations based on local photostationary steady-state conditions and the simulations by EMAC, the observations show very little latitudinal variation with mean values of approximately 0.1–0.2 ppb_v. A tendency towards slightly enhanced values is observed in the ITCZ, where the mixing ratios increase up to 0.22 ppb_v. However, considering the standard deviation range (up to 0.19 ppb_v), the rather flat distribution in the upper troposphere indicates nearly constant hydrogen peroxide levels throughout the whole investigated tropospheric region (see also Tab. S1).

345 Both the EMAC-simulated and PSS-calculated latitudinal hydrogen peroxide distributions display decreasing trends with increasing latitudes towards the north. The highest mean values of 0.49 (± 0.29) ppb_v for PSS and 0.583 (± 0.40) ppb_v for EMAC are found in the southernmost part of the probed region. The elevated levels of H₂O₂ in the EMAC simulations starts already in the ITCZ, while the PSS calculations only increase south of 5°N.


350 Overall H₂O₂ mixing ratios from the PSS calculations decrease from the equator towards the subtropics with at a rate of -0.004 (± 0.002) ppb_v per degree of latitude. At approximately 5°N, the PSS-based mixing ratios decrease to a range of 0.09 ppb_v to 0.25 ppb_v and tend to fall below the measured levels at the northern edge of the examined region by a factor of up to 5 (at approx. 32°N). The mixing ratios of hydrogen peroxide simulated by EMAC remain elevated at approximately 0.30 ppb_v to 0.35 ppb_v from 6°S to 15°N, yielding H₂O₂(EMAC)/H₂O₂(measurements) ratios of 2.2 to 2.5. Further north of 15°N, the H₂O₂ levels in EMAC decrease by almost half to 0.15 ppb_v, resulting in H₂O₂(EMAC)/H₂O₂(measurements) ratios between 1.1 and 1.2 and a rather good agreement between simulations and observations. The linear fit applied to the latitudinal distribution of
355 simulated hydrogen peroxide levels displays a mean ($\pm 1\sigma$) decrease of -0.008 (± 0.001) ppb_v per degree of latitude. An overview of numerical values for measured means ($\pm 1\sigma$) and medians, PSS-calculated and EMAC-simulated hydrogen peroxide levels subdivided into three hemispheric regions, the northern (approximately 20°N–40°N), ITCZ (approximately 5°N < 20°N), and southern hemisphere (approx.-10°N < 5°N) is given in Tab. S1 in the Supplement of this work. Please note that the steady-state calculations, which are based on observed levels of OH, HO₂, and H₂O₂ photolysis rates, only account for photochemical production and loss of hydrogen peroxide. EMAC simulations additionally account for vertical and horizontal transport
360 processes, as well as local losses due to cloud scavenging (Hamryszczak et al., 2022). Thus, deviations between local photostationary steady-state budget calculations and EMAC simulations can be used to identify and quantify the impact of convective processes in the ITCZ on the budget of H₂O₂ in the upper troposphere. This requires that the EMAC model correctly simulates precursors (HO₂) and sinks (OH, H₂O₂ photolysis) of photochemical H₂O₂ formation.

365 A comparison between observations and EMAC simulations for the basic species that affect photochemical production (HO₂) and loss (OH and H₂O₂ photolysis rates) of H₂O₂ reveals that while HO₂ is adequately reproduced by EMAC, the simulations tend to underestimate OH and the photolysis frequencies, in particular, south of 15°N, where EMAC underestimates OH by up to 0.500 ppt_v (Fig. S5b). At the same time, model results partly tend to overestimate HO₂ by approximately 3.5 ppt_v, indicating issues with HO₂/OH partitioning. Since the production of OH in the UT depends to an extent on the reaction of HO₂



370 with NO, these might be associated with the underestimation of NO by EMAC in the southern hemisphere as shown by Tadic
et al. (2021). Additionally, the measured H₂O₂ photolysis frequencies show minor discrepancies with those from the model
(Fig. S5c). Due to the overestimated HO₂ mixing ratios, EMAC simulates higher levels of H₂O₂. At the same time, the lower
OH mixing ratios and the underestimated H₂O₂ photolysis frequencies might cause decreased loss rates, thus leading to
underestimation of the H₂O₂ loss paths. Therefore, overestimation of the photochemical source and underestimation of the
375 photochemical H₂O₂ sinks by EMAC explain the differences between PSS calculations and simulations at southern latitudes;
however, these observations fail to explain the observed differences between PSS calculations and EMAC simulations versus
the observations south of the ITCZ. Also, it is highly unlikely that a measurement error is responsible for the discrepancies, as
this would have to be local and restricted to the most southerly latitudes. Potential causes leading to the observed discrepancy
between the measurement and the simulations might be an underestimation of cloud scavenging and removal of hydrogen
380 peroxide by wet and dry deposition in the ITCZ and also south of this region. A number of flights south of 15°N were performed
in close proximity to the western shores of Africa or even over land (Fig. 2), close to enhanced convective precipitation (Fig.
S3). Based on the ERA5 reanalysis results on cloud coverage during the measurement period (Hersbach, H. et al., 2019; Fig.
S2), we hypothesize that the air masses probed in this area were affected by cloud processing especially in the UT, causing the
deviations towards the model calculations.

385 Since the amount of the H₂O₂ in the upper atmosphere depends on convective transport from the lower troposphere as well as
on losses caused by cloud droplets and the subsequent rainout, it is important that EMAC simulations and PSS calculations
reproduce the levels of hydrogen peroxide in the middle troposphere and the boundary layer. In Fig. 6, averages of hydrogen
peroxide mixing ratios based on *in situ* observations, PSS calculations, and EMAC simulations and the corresponding
H₂O₂(EMAC)/H₂O₂(measurements) and H₂O₂(PSS)/H₂O₂(measurements) ratios are shown. The data are binned into subsets
390 of 1 km of altitude with respect to the center of the bin width based on take-off and landings in proximity to the base of
operations in Sal, Cape Verde. The lines and the shadings represent mean values and the 1σ-standard deviations. Dashed lines
represent median values.



395

Figure 6. Vertical profiles of observed (red), simulated (blue), and calculated based on the PSS assumption (black) hydrogen peroxide means and medians (top panel; a-b) and vertical profiles of mean and median H₂O₂(EMAC)/H₂O₂(measurements) and H₂O₂(PSS)/H₂O₂(measurements) ratios (bottom panel; c–d). Vertical profile estimations were calculated within 1 km means and medians over the atmospheric column based on the data obtained in the region in proximity to the base of operations in Sal, Cape Verde (approx.16°35′–16°51′N; -22°52′–23°E).




400 In general, the observed, PSS-calculated and EMAC-simulated vertical profiles of hydrogen peroxide follow the expected trend throughout the troposphere (Fig. 6a–b). The lowest hydrogen peroxide mixing ratios of 0.141 ppb_v were measured in the upper troposphere, where the availability of the H₂O₂ precursor HO₂ is limited due to low water vapor concentrations. The highest mean values ($\pm 1\sigma$) of 2.44 (± 0.78) ppb_v were measured directly above the boundary layer (2–5 km) in the free troposphere. Below 2 km, the levels of observed H₂O₂ decrease to 1.7 (± 1.1) ppb_v, reflecting the impact of deposition processes on H₂O₂ in the boundary layer in proximity to Cape Verde. Based on the good agreement of the observed vertical distribution with the expected trend as well as log book reports, the presence of clouds and their subsequent scavenging is assumed to have a minor impact on the local budget of the species.

405 A good agreement between the measured and EMAC-simulated datasets with a H₂O₂(EMAC)/H₂O₂(measurement) ratio of approximately 1 was found in the lower troposphere (2–6 km; Fig. 6c). Model results tend to overestimate hydrogen peroxide in the boundary layer, which might be due to the model resolution (1.8° x 1.8°) and the corresponding restrictions in resolving small-scale variations in hydrogen peroxide deposition processes in proximity to the base of operations on the island. The vertical profiles of the observations and the model show that the differences arise mainly in the upper troposphere, with increased ratios of simulated vs. measured H₂O₂ of up to a factor 5.6 at 12 km altitude, which might indicate convective outflow in the model. Vertical profiles of observed and simulated HO₂, OH, and H₂O₂ photolysis rates are in excellent agreement (Fig. 415 S6), indicating that the model accurately simulates photochemical processes throughout the troposphere, so that the remaining differences for H₂O₂ are most likely caused by physical processes (e.g., deposition and transport).

The comparison of measured and PSS-calculated H₂O₂ vertical profiles indicates a missing source or an overestimated sink below altitudes of 5 km (Fig. 6b). Here, the PSS-calculated hydrogen peroxide levels fall short by about 1 ppb_v at altitudes below 4 km, yielding H₂O₂(PSS)/H₂O₂(measured) ratios smaller than unity at these altitudes (Fig. 6d). The absolute difference 420 between the measured and calculated mixing ratios seems to be very prominent in and right above the boundary layer (< 5 km) and can be associated with air masses affected by Saharan dust, which was often probed during take-off and landings at Sal. Heterogeneous loss of HO₂ on desert dust particles, while modest, is expected to lower the production of H₂O₂ (Reus et al., 2005). Thus, local mixing ratios of HO₂ could be affected by heterogeneous loss, while PSS is not yet achieved (Fig. S7).

425 In order to investigate the extent of the potential hydrogen peroxide injection into the upper troposphere, we calculate excess hydrogen peroxide mixing ratios as the difference between the observed and the corresponding H₂O₂ based on PSS. Analogously, potential excess of H₂O₂ using model-simulated data was determined. The spatial distribution of the calculated transport rates in the upper troposphere is presented in Fig. 7 as latitude vs. longitude plots of mean hydrogen peroxide levels aggregated over a spatial 1° x 1° grid at altitudes above 8 km. The color scale represents the average excess mixing ratios determined for the species in ppt_v.



430

Figure 7. Color-coded spatial resolution of calculated H_2O_2 excess based on the difference between the observations and PSS-based calculations (a) and EMAC simulations and calculations at steady state based on the EMAC data output (b). Data were binned into $1^\circ \times 1^\circ$ bins over the full extension of the flight tracks.

435

The H_2O_2 observations exceed the calculated values based on PSS mostly in the range of 70 ppt_v to 110 ppt_v with the exception of the values derived in the southern hemisphere. Highest deviations were derived in the ITCZ and reach up to 310 ppt_v at their maximum. The excess mixing ratios show a clear trend with the most impactful injection events in the ITCZ region (5°N–15°N), where convective transport is expected (Waliser and Gautier, 1993; Fontaine et al., 2011). From there, a subsequent redistribution of hydrogen peroxide towards the northern and southern hemispheres occurs, which agrees well with the decreasing gradient towards the north and south. EMAC simulates exceeding H_2O_2 mixing ratios mostly in the range of 20 ppt_v to 150 ppt_v, with maximum excess up to 240 ppt_v in the ITCZ (12.5°N, -25.5°E; Fig. 7b). EMAC reproduces the transport rates to a lower extent in the ITCZ, but local convective events such as those occurring within tropical storm Florence and above the African coast are not simulated by the model. In contrast, no significant excess of H_2O_2 was determined between 1.5°N, -4.5°E.

440

445

The calculated absolute difference between the measured and the PSS-calculated H_2O_2 in the upper troposphere displays an average excess of 44 (± 120) ppt_v hydrogen peroxide over the entire region relative to the PSS-modelled conditions. In comparison, injections based on EMAC simulations show about 60% lower values of 18 (± 120) ppt_v. The mixing ratios of the H_2O_2 measurements below 4 km at Sal were in the range of 1.7–2.4 ppb_v, which would contribute with up to 1.8–2.6% of H_2O_2 in the outflow, assuming potential inflow below 4 km. A similar calculation based solely on EMAC data indicates a contribution of 0.6–0.8 % within the model (based on EMAC mixing ratios of 2.1–3.0 ppb_v). Thus, although EMAC reproduces potential transport processes from the lower troposphere, the discrepancy in the transmission efficiencies indicates a smaller contribution to the simulated hydrogen peroxide levels. Consequently, the enhanced hydrogen peroxide mixing ratios cannot be justified exclusively by photochemical reactions within the upper troposphere. Additional injections from the lower troposphere via convective transport and the subsequent redistribution towards the subtropics have to be considered.

450



5 Conclusions

455 Hydrogen peroxide was measured during the CAFE-Africa campaign over the tropical Atlantic and West Africa in the upper
troposphere (above 8 km). Generally, the measured levels of hydrogen peroxide in the upper troposphere fit well to the
previously observed H_2O_2 at latitudes 10°S – 40°N . At high altitudes, a minor symmetrical decrease from the ITCZ towards
northern and southern latitudes was observed, which deviates from previously reported observations in the upper troposphere.
According to previous reports, the H_2O_2 mixing ratios are expected to be elevated in the equatorial upper troposphere due to
460 biomass burning and atmospheric transport. However, the H_2O_2 mixing ratios measured during the CAFE-Africa campaign
show only very little latitudinal variation relative to previous measurements over the Atlantic with a shift of the maximum
mixing ratios towards the ITCZ. The measured hydrogen peroxide mixing ratios show a rather uniform distribution with peak
events in the ITCZ and over the African coast, indicating the influence of convective transport processes on the distribution of
hydrogen peroxide in the upper troposphere.

465 Whilst the observations of hydrogen peroxide are in good general agreement with the range of previous observations performed
in the upper troposphere, the measured H_2O_2 mixing ratios deviate from the PSS calculations based on OH and HO_2
measurements and the simulations performed by EMAC. The local PSS calculations significantly underestimate the H_2O_2
mixing ratios in the north of the probed region. There, the comparison of the H_2O_2 measurements with PSS calculations reveals
a large impact of vertical transport within the ITCZ and the associated redistribution in the upper troposphere on the spatial
470 distribution of hydrogen peroxide. Further, the enhanced presence of clouds in the ITCZ and towards the southern hemisphere
indicates significant cloud scavenging in the probed region, justifying the deviations to the local photostationary steady-state
calculations, which only account for photochemical sources and sinks of H_2O_2 . The EMAC simulations of H_2O_2 , HO_2 , and OH
agree with the observations in the lower tropospheric layers. An overestimation of model results compared with the
observations of hydrogen peroxide mixing ratios due to inaccuracies in cloud scavenging was observed in the upper
475 troposphere towards the southern hemisphere. Based on our calculations, the model simulates only partially the impacts of
atmospheric transport on the H_2O_2 budget. In fact, the calculated excess hydrogen peroxide mixing ratios based on EMAC are
lower compared to those based on the PSS calculations by approximately 60%. The comparison between the EMAC- and PSS-
calculated versus measured hydrogen peroxide confirms that convective transport and consequent redistribution most likely
by northerly and southerly winds towards the subtropics has a significant impact on H_2O_2 . This redistribution alters the spatial
480 distribution of H_2O_2 towards more uniform mixing ratios in the marine tropical upper troposphere than would be expected
based exclusively on photochemical production and loss processes in the UT.

Data availability. All CAFE-Africa data sets used in this study are permanently stored in an archive on the KEEPER service
of the Max Planck Digital Library (<https://keeper.mpdl.mpg.de>; last access: 17 October 2022) and are available to all scientists,
485 who agree to the CAFE-Africa data protocol.



Author contributions. JL and HF planned the campaign; DD, BBr, RR, DM, MM, HH, and BB performed the measurements; ZH and HF designed the study; ZH, DD, RR, BB, and MZ processed and analyzed the data; AP developed the model code and performed the simulation; ZH wrote the manuscript draft with contributions of all co-authors.

490

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments. The authors are very grateful to the CAFE-Africa team, Forschungszentrum Jülich, Karlsruhe Institute of Technology and Deutsches Zentrum für Luft- und Raumfahrt (DLR) in Oberpfaffenhofen for their great support. Their work

495

References

- Allen, H. M., Crouse, J. D., Kim, M. J., Teng, A. P., Ray, E. A., McKain, K., Sweeney, C., and Wennberg, P. O.: H₂O₂ and CH₃OOH (MHP) in the Remote Atmosphere: 1. Global Distribution and Regional Influences, *JGR Atmospheres*, 127, <https://doi.org/10.1029/2021JD035701>, 2022.
- Allen, N. D.C., González Abad, G., Bernath, P. F., and Boone, C. D.: Satellite observations of the global distribution of hydrogen peroxide (H₂O₂) from ACE, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 115, 66–77, <https://doi.org/10.1016/j.jqsrt.2012.09.008>, 2013.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, <https://doi.org/10.5194/acp-4-1461-2004>, 2004.
- Bohn, B. and Lohse, I.: Calibration and evaluation of CCD spectroradiometers for ground-based and airborne measurements of spectral actinic flux densities, *Atmos. Meas. Tech.*, 10, 3151–3174, <https://doi.org/10.5194/amt-10-3151-2017>, 2017.
- Bozem, H., Pozzer, A., Harder, H., Martinez, M., Williams, J., Lelieveld, J., and Fischer, H.: The influence of deep convection on HCHO and H₂O₂ in the upper troposphere over Europe, *Atmos. Chem. Phys.*, 17, 11835–11848, <https://doi.org/10.5194/acp-17-11835-2017>, 2017.
- Calvert, J. G., Lazrus, A., Kok, G. L., Heikes, B. G., Walega, J. G., Lind, J., and Cantrell, C. A.: Chemical mechanisms of acid generation in the troposphere, *Nature*, 317, 27–35, <https://doi.org/10.1038/317027a0>, 1985.
- Chang, W., Lee, M., and Heikes, B. G.: One-dimensional photochemical study of H₂O₂, CH₃OOH, and HCHO in the marine boundary layer during Pacific Exploratory Mission in the Tropics (PEM-Tropics) B, *J. Geophys. Res.*, 109, n/a–n/a, <https://doi.org/10.1029/2003jd004256>, 2004.
- Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded Emissions of Air Pollutants for the period 1970–2012 within EDGAR v4.3.2, 2018.

500

510

515



- Crutzen, P. J., Lawrence, M. G., and Pöschl, U.: On the background photochemistry of tropospheric ozone, *Tellus B: Chemical and Physical Meteorology*, 51, 123–146, <https://doi.org/10.3402/tellusb.v51i1.16264>, 1999.
- Damschen, D. E. and Martin, L.R.: Aqueous aerosol oxidation of nitrous acid by O₂, O₃ AND H₂O₂, *Atmospheric Environment* (1967), 17, 2005–2011, [https://doi.org/10.1016/0004-6981\(83\)90357-8](https://doi.org/10.1016/0004-6981(83)90357-8), 1983.
- Daum, P. H., Kleinman, L. I., Hills, A. J., Lazrus, A. L., Leslie, A. C. D., Busness, K., and Boatman, J.: Measurement and interpretation of concentrations of H₂O₂ and related species in the upper midwest during summer, *J. Geophys. Res.*, 95, 9857–9871, <https://doi.org/10.1029/JD095iD07p09857>, 1990.
- Dienhart, D., Brendel, B., Crowley, J. N., Eger, P. G., Harder, H., Martinez, M., Pozzer, A., Rohloff, R., Schuladen, J., Tauer, S., Lelieveld, J., and Fischer, H.: Formaldehyde and hydroperoxide distribution around the Arabian Peninsula – evaluation of EMAC model results with ship-based measurements, 2022.
- Faloona, I., Tan, D., Brune, W. H., Jaeglé, L., Jacob, D. J., Kondo, Y., Koike, M., Chatfield, R., Pueschel, R., Ferry, G., Sachse, G., Vay, S., Anderson, B., Hannon, J., and Fuelberg, H.: Observations of HO_x and its relationship with NO_x in the upper troposphere during SONEX, *J. Geophys. Res.*, 105, 3771–3783, <https://doi.org/10.1029/1999JD900914>, 2000.
- Faloona, I. C., Tan, D., Leshner, R. L., Hazen, N. L., Frame, C. L., Simpas, J. B., Harder, H., Martinez, M., Di Carlo, P., Ren, X., and Brune, W. H.: A Laser-induced Fluorescence Instrument for Detecting Tropospheric OH and HO₂ Characteristics and Calibration, *J Atmos Chem*, 47, 139–167, <https://doi.org/10.1023/B:JOCH.0000021036.53185.0e>, 2004.
- Fels, M. and Junkermann, W.: The occurrence of organic peroxides in air at a mountain site, *Geophys. Res. Lett.*, 21, 341–344, <https://doi.org/10.1029/93GL01892>, 1994.
- Fontaine, B., Roucou, P., Gaetani, M., and Marteau, R.: Recent changes in precipitation, ITCZ convection and northern tropical circulation over North Africa (1979-2007), *Int. J. Climatol.*, 31, 633–648, <https://doi.org/10.1002/joc.2108>, 2011.
- Guilbault, G. G., Brignac, P. J., and Juneau, M.: New substrates for the fluorometric determination of oxidative enzymes, *Anal. Chem.*, 40, 1256–1263, <https://doi.org/10.1021/ac60264a027>, 1968.
- Hamryszczak, Z. T., Pozzer, A., Obersteiner, F., Bohn, B., Steil, B., Lelieveld, J., and Fischer, H.: Distribution of hydrogen peroxide over Europe during the BLUESKY aircraft campaign, 2022.
- Hastenrath, S. and Lamb, P.: Some Aspects of Circulation and Climate over the Eastern Equatorial Atlantic, *Mon. Wea. Rev.*, 105, 1019–1023, [https://doi.org/10.1175/1520-0493\(1977\)105<1019:SAOCAC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1977)105<1019:SAOCAC>2.0.CO;2), 1977.
- Heikes, B. G.: Formaldehyde and hydroperoxides at Mauna Loa Observatory, *J. Geophys. Res.*, 97, 18001, <https://doi.org/10.1029/92JD00268>, 1992.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N.: ERA5 monthly averaged data on single levels from 1979 to present CDS [data set] <https://doi.org/10.24381/CDS.F17050D7>, last access: 22-07-22, 2019.



- Hoffmann, M. R. and Edwards, J. O.: Kinetics of the oxidation of sulfite by hydrogen peroxide in acidic solution, *J. Phys. Chem.*, 79, 2096–2098, <https://doi.org/10.1021/j100587a005>, 1975.
- 555 Hottmann, B., Hafermann, S., Tomsche, L., Marno, D., Martinez, M., Harder, H., Pozzer, A., Neumaier, M., Zahn, A., Bohn, B., Stratmann, G., Ziereis, H., Lelieveld, J., and Fischer, H.: Impact of the South Asian monsoon outflow on atmospheric hydroperoxides in the upper troposphere, *Atmos. Chem. Phys.*, 20, 12655–12673, <https://doi.org/10.5194/acp-20-12655-2020>, 2020.
- Hua, W., Chen, Z. M., Jie, C. Y., Kondo, Y., Hofzumahaus, A., Takegawa, N., Chang, C. C., Lu, K. D., Miyazaki, Y., Kita, K., Wang, H. L., Zhang, Y. H., and Hu, M.: Atmospheric hydrogen peroxide and organic hydroperoxides during PRIDE-
560 PRD'06, China: their concentration, formation mechanism and contribution to secondary aerosols, *Atmos. Chem. Phys.*, 8, 6755–6773, <https://doi.org/10.5194/acp-8-6755-2008>, 2008.
- Jacob, P. and Klockow, D.: Hydrogen peroxide measurements in the marine atmosphere, *J Atmos Chem*, 15, 353–360, <https://doi.org/10.1007/BF00115404>, 1992.
- 565 Jaeglé, L., Jacob, D. J., Wennberg, P. O., Spivakovsky, C. M., Hanisco, T. F., Lanzendorf, E. J., Hintsä, E. J., Fahey, D. W., Keim, E. R., Proffitt, M. H., Atlas, E. L., Flocke, F., Schauffler, S., McElroy, C. T., Midwinter, C., Pfister, L., and Wilson, J. C.: Observed OH and HO₂ in the upper troposphere suggest a major source from convective injection of peroxides, *Geophys. Res. Lett.*, 24, 3181–3184, <https://doi.org/10.1029/97GL03004>, 1997.
- 570 Jaeglé, L., Jacob, D. J., Brune, W. H., Faloon, I., Tan, D., Heikes, B. G., Kondo, Y., Sachse, G. W., Anderson, B., Gregory, G. L., Singh, H. B., Poeschel, R., Ferry, G., Blake, D. R., and Shetter, R. E.: Photochemistry of HO_x in the upper troposphere at northern midlatitudes, *J. Geophys. Res.*, 105, 3877–3892, <https://doi.org/10.1029/1999JD901016>, 2000.
- Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) - a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433–444, <https://doi.org/10.5194/acp-5-433-2005>, 2005.
- 575 Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmos. Chem. Phys.*, 6, 5067–5104, <https://doi.org/10.5194/acp-6-5067-2006>, 2006.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.:
580 Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717–752, <https://doi.org/10.5194/gmd-3-717-2010>, 2010.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy)
585 version 2.51, *Geosci. Model Dev.*, 9, 1153–1200, <https://doi.org/10.5194/gmd-9-1153-2016>, 2016.



- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- 590 Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal processes DRY DEPosition and SEDImentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617–4632, <https://doi.org/10.5194/acp-6-4617-2006>, 2006a.
- Kerkweg, A., Sander, R., Tost, H., and Jöckel, P.: Technical note: Implementation of prescribed (OFFLEM), calculated (ONLEM), and pseudo-emissions (TNUDGE) of chemical species in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 3603–3609, <https://doi.org/10.5194/acp-6-3603-2006>, 2006b.
- 595 Kleinman, L. I.: Seasonal dependence of boundary layer peroxide concentration: The low and high NO_x regimes, *J. Geophys. Res.*, 96, 20721, <https://doi.org/10.1029/91JD02040>, 1991.
- Klippel, T., Fischer, H., Bozem, H., Lawrence, M. G., Butler, T., Jöckel, P., Tost, H., Martinez, M., Harder, H., Regelin, E., Sander, R., Schiller, C. L., Stickler, A., and Lelieveld, J.: Distribution of hydrogen peroxide and formaldehyde over Central Europe during the HOOVER project, *Atmos. Chem. Phys.*, 11, 4391–4410, [https://doi.org/10.5194/acp-11-4391-](https://doi.org/10.5194/acp-11-4391-2011)
600 2011, 2011.
- Krautstrunk, M. and Giez, A.: The Transition From FALCON to HALO Era Airborne Atmospheric Research, in: *Atmospheric physics: Background-methods-trends*, edited by: Schumann, U., Springer-Verlag, Berlin, Heidelberg, 609–624, https://doi.org/10.1007/978-3-642-30183-4_37, op. 2012.
- Lazrus, A. L., Kok, G. L., Gitlin, S. N., Lind, J. A., and McLaren, S. E.: Automated fluorimetric method for hydrogen peroxide in atmospheric precipitation, *Anal. Chem.*, 57, 917–922, <https://doi.org/10.1021/ac00281a031>, 1985.
- 605 Lazrus, A. L., Kok, G. L., Lind, J. A., Gitlin, S. N., Heikes, B. G., and Shetter, R. E.: Automated fluorometric method for hydrogen peroxide in air, *Anal. Chem.*, 58, 594–597, <https://doi.org/10.1021/ac00294a024>, 1986.
- Lee, M., Heikes, B. G., Jacob, D. J., Sachse, G., and Anderson, B.: Hydrogen peroxide, organic hydroperoxide, and formaldehyde as primary pollutants from biomass burning, *J. Geophys. Res.*, 102, 1301–1309,
610 <https://doi.org/10.1029/96jd01709>, 1997.
- Lee, M., Heikes, B. G., and Jacob, D. J.: Enhancements of hydroperoxides and formaldehyde in biomass burning impacted air and their effect on atmospheric oxidant cycles, *J. Geophys. Res.*, 103, 13201–13212,
<https://doi.org/10.1029/98jd00578>, 1998.
- Lee, M., Heikes, B. G., and O'Sullivan, D. W.: Hydrogen peroxide and organic hydroperoxide in the troposphere: a review, *Atmospheric Environment*, 34, 3475–3494, [https://doi.org/10.1016/S1352-2310\(99\)00432-X](https://doi.org/10.1016/S1352-2310(99)00432-X), 2000.
- 615 Lelieveld, J. and Crutzen, P. J.: Influences of cloud photochemical processes on tropospheric ozone, *Nature*, 343, 227–233, <https://doi.org/10.1038/343227a0>, 1990.
- Levy, H.: Normal atmosphere: large radical and formaldehyde concentrations predicted, *Science (New York, N.Y.)*, 173, 141–143, <https://doi.org/10.1126/science.173.3992.141>, 1971.



- 620 Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: A global perspective, *J. Geophys. Res.*, 86, 7210, <https://doi.org/10.1029/JC086iC08p07210>, 1981.
- Marno, D., Ernest, C., Hens, K., Javed, U., Klimach, T., Martinez, M., Rudolf, M., Lelieveld, J., and Harder, H.: Calibration of an airborne HO_x instrument using the All Pressure Altitude-based Calibrator for HO_x Experimentation (APACHE), *Atmos. Meas. Tech.*, 13, 2711–2731, <https://doi.org/10.5194/amt-13-2711-2020>, 2020.
- 625 Nguyen, T. B., Crounse, J. D., Teng, A. P., St Clair, J. M., Paulot, F., Wolfe, G. M., and Wennberg, P. O.: Rapid deposition of oxidized biogenic compounds to a temperate forest, *Proceedings of the National Academy of Sciences of the United States of America*, 112, E392–401, <https://doi.org/10.1073/pnas.1418702112>, 2015.
- Nussbaumer, C. M., Tadic, I., Dienhart, D., Wang, N., Edtbauer, A., Ernle, L., Williams, J., Obersteiner, F., Gutiérrez-Álvarez, I., Harder, H., Lelieveld, J., and Fischer, H.: Measurement report: In situ observations of deep convection without lightning during the tropical cyclone Florence 2018, *Atmos. Chem. Phys.*, 21, 7933–7945, <https://doi.org/10.5194/acp-21-7933-2021>, 2021.
- 630 O'Sullivan, D. W., Lee, M., Noone, B. C., and Heikes, B. G.: Henry's Law Constant Determinations for Hydrogen Peroxide, Methyl Hydroperoxide, Hydroxymethyl Hydroperoxide, Ethyl Hydroperoxide, and Peroxyacetic Acid, *J. Phys. Chem.*, 100, 3241–3247, <https://doi.org/10.1021/jp951168n>, 1996.
- 635 O'Sullivan, D. W., Heikes, B. G., Lee, M., Chang, W., Gregory, G. L., Blake, D. R., and Sachse, G. W.: Distribution of hydrogen peroxide and methylhydroperoxide over the Pacific and South Atlantic Oceans, *J. Geophys. Res.*, 104, 5635–5646, <https://doi.org/10.1029/98JD01250>, 1999.
- Penkett, S. A., Jones, B.M.R., Brich, K. A., and Eggleton, A.E.J.: The importance of atmospheric ozone and hydrogen peroxide in oxidising sulphur dioxide in cloud and rainwater, *Atmospheric Environment (1967)*, 13, 123–137, [https://doi.org/10.1016/0004-6981\(79\)90251-8](https://doi.org/10.1016/0004-6981(79)90251-8), 1979.
- 640 Perros, P.E.: Large-scale distribution of hydrogen peroxide from aircraft measurements during the TROPOZ II experiment, *Atmospheric Environment. Part A. General Topics*, 27, 1695–1708, [https://doi.org/10.1016/0960-1686\(93\)90232-N](https://doi.org/10.1016/0960-1686(93)90232-N), 1993.
- Pozzer, A., Jöckel, P., and van Aardenne, J.: The influence of the vertical distribution of emissions on tropospheric chemistry, *Atmos. Chem. Phys.*, 9, 9417–9432, <https://doi.org/10.5194/acp-9-9417-2009>, 2009.
- 645 Pozzer, A., Reifenberg, S. F., Kumar, V., Franco, B., Kohl, M., Taraborrelli, D., Gromov, S., Ehrhart, S., Jöckel, P., Sander, R., Fall, V., Rosanka, S., Karydis, V., Akritidis, D., Emmerichs, T., Crippa, M., Guizzardi, D., Kaiser, J. W., Clarisse, L., Kiendler-Scharr, A., Tost, H., and Tsimpidi, A.: Simulation of organics in the atmosphere: evaluation of EMACv2.54 with the Mainz Organic Mechanism (MOM) coupled to the ORACLE (v1.0) submodel, *Geosci. Model Dev.*, 15, 2673–2710, <https://doi.org/10.5194/gmd-15-2673-2022>, 2022.
- 650 Prather, M. J. and Jacob, D. J.: A persistent imbalance in HO_x and NO_x photochemistry of the upper troposphere driven by deep tropical convection, *Geophys. Res. Lett.*, 24, 3189–3192, <https://doi.org/10.1029/97GL03027>, 1997.



- Reeves, C. E. and Penkett, S. A.: Measurements of peroxides and what they tell us, *Chemical reviews*, 103, 5199–5218, <https://doi.org/10.1021/cr0205053>, 2003.
- 655 Reus, M. de, Fischer, H., Sander, R., Gros, V., Kormann, R., Salisbury, G., van Dingenen, R., Williams, J., Zöllner, M., and Lelieveld, J.: Observations and model calculations of trace gas scavenging in a dense Saharan dust plume during MINATROC, *Atmos. Chem. Phys.*, 5, 1787–1803, <https://doi.org/10.5194/acp-5-1787-2005>, 2005.
- Rinsland, C. P., Coheur, P. F., Herbin, H., Clerbaux, C., Boone, C., Bernath, P., and Chiou, L. S.: Detection of elevated tropospheric hydrogen peroxide (H₂O₂) mixing ratios in atmospheric chemistry experiment (ACE) subtropical infrared solar occultation spectra, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 107, 340–348, <https://doi.org/10.1016/j.jqsrt.2007.02.009>, 2007.
- 660 Robbin Martin, L. and Damschen, D. E.: Aqueous oxidation of sulfur dioxide by hydrogen peroxide at low pH, *Atmospheric Environment* (1967), 15, 1615–1621, [https://doi.org/10.1016/0004-6981\(81\)90146-3](https://doi.org/10.1016/0004-6981(81)90146-3), 1981.
- Sander, R., Jöckel, P., Kirner, O., Kunert, A. T., Landgraf, J., and Pozzer, A.: The photolysis module JVAL-14, compatible with the MESSy standard, and the JVal PreProcessor (JVPP), *Geosci. Model Dev.*, 7, 2653–2662, <https://doi.org/10.5194/gmd-7-2653-2014>, 2014.
- 665 Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Groß, J.-U., Harder, H., Huijnen, V., Jöckel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Riede, H., Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geosci. Model Dev.*, 12, 1365–1385, <https://doi.org/10.5194/gmd-12-1365-2019>, 2019.
- 670 Slemr, F. and Tremmel, H. G.: Hydroperoxides in the marine troposphere over the Atlantic Ocean, *J Atmos Chem*, 19, 371–404, <https://doi.org/10.1007/BF00694493>, 1994.
- Snow, J. A.: Winter-spring evolution and variability of HO_x reservoir species, hydrogen peroxide, and methyl hydroperoxide, in the northern middle to high latitudes, *J. Geophys. Res.*, 108, 1890, <https://doi.org/10.1029/2002JD002172>, 2003.
- 675 Snow, J. A., Heikes, B. G., Shen, H., O'Sullivan, D. W., Fried, A., and Walega, J.: Hydrogen peroxide, methyl hydroperoxide, and formaldehyde over North America and the North Atlantic, *J. Geophys. Res.*, 112, 8353, <https://doi.org/10.1029/2006JD007746>, 2007.
- Tadic, I., Nussbaumer, C. M., Bohn, B., Harder, H., Marno, D., Martinez, M., Obersteiner, F., Parchatka, U., Pozzer, A., Rohloff, R., Zöger, M., Lelieveld, J., and Fischer, H.: Central role of nitric oxide in ozone production in the upper tropical troposphere over the Atlantic Ocean and western Africa, *Atmos. Chem. Phys.*, 21, 8195–8211, <https://doi.org/10.5194/acp-21-8195-2021>, 2021.
- 680 Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVenging submodel for global atmospheric chemistry modelling, *Atmos. Chem. Phys.*, 6, 565–574, <https://doi.org/10.5194/acp-6-565-2006>, 2006.
- 685



- Valverde-Canossa, J., Wieprecht, W., Acker, K., and Moortgat, G. K.: H₂O₂ and organic peroxide measurements in an orographic cloud: The FEBUKO experiment, *Atmospheric Environment*, 39, 4279–4290, <https://doi.org/10.1016/j.atmosenv.2005.02.040>, 2005.
- 690 Walcek, C. J.: A theoretical estimate of O₃ and H₂O₂ dry deposition over the northeast United States, *Atmospheric Environment* (1967), 21, 2649–2659, [https://doi.org/10.1016/0004-6981\(87\)90196-x](https://doi.org/10.1016/0004-6981(87)90196-x), 1987.
- Waliser, D. E. and Gautier, C.: A Satellite-derived Climatology of the ITCZ, *Journal of Climate*, 6, 2162–2174, [https://doi.org/10.1175/1520-0442\(1993\)006<2162:ASDCOT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<2162:ASDCOT>2.0.CO;2), 1993.
- Weinstein-Lloyd, J. B., Lee, J. H., Daum, P. H., Kleinman, L. I., Nunnermacker, L. J., Springston, S. R., and Newman, L.: Measurements of peroxides and related species during the 1995 summer intensive of the Southern Oxidants Study in
695 Nashville, Tennessee, *J. Geophys. Res.*, 103, 22361–22373, <https://doi.org/10.1029/98JD01636>, 1998.