

We thank the reviewer for their supportive and constructive comments on the manuscript. We feel that the paper has been improved by the review process. Below, we address each of the reviewer's specific and technical line-by-line comments. The reviewer comments are in black text, while the responses are in *blue italics* and new text added to the manuscript is in *blue*.

GENERAL COMMENTS

The authors present new shipborne measurements during winter and spring of the stable isotopes of nitrogen (d15N) and oxygen (d18O, D17O) isotopes in the coarse mode of atmospheric nitrate collected in the marine boundary layer (MBL) between South Africa and the marginal sea ice zone in Antarctica. d15N values are used to attribute primary sources of atmospheric nitrate: during spring/summer lightning, ocean (alkyl nitrates) and snowpack NO_x emissions dominated at low, mid and high latitudes, respectively. During winter transport of NO_x precursors such as PAN from lower latitudes as well as potentially stratospheric nitrate contribute mostly to the atmospheric nitrate background. Using D17O and d18O values in an isotope end member mixing analysis the authors confirm the current understanding that oxidation during daytime is dominated by OH and during night time/ winter by O₃. They speculate that a third end member emerging at sunrise in spring may be attributed to the onset of halogen chemistry and contribution to oxidation via peroxy radicals.


These are important new atmospheric data from the Southern Ocean MBL covering seasons which are notoriously under-sampled, and therefore should be published. However there are some weaknesses in data interpretation, some gaps in the cited literature as well as presentation of results can be improved.

Major points:

- the introduction should expand on the nitrogen chemistry relevant for the oxygen and nitrogen isotope transfer, i.e. spell out key reactions of the relevant pathways: Step1) NO,NO₂ interconversion (fast) and Step2) NO₂ oxidation to form nitrate (slower). This will help the reader to follow the arguments presented and assess key uncertainties and missing variables for future studies aiming at a quantitative isotope budget.

Based on the reviewers suggestion, key reactions and relevant pathways have been included in the introduction as follows, with original text in black and new text in blue:

"In addition to there being multiple NO_x sources across the Southern Ocean MBL, several different oxidation pathways can be responsible for NO_x to NO₃⁻ conversion, varying with chemistry and time of day (Savarino et al., 2007). Once emitted, NO is rapidly oxidised by ozone (O₃) (R1), peroxy radicals (RO₂ or HO₂) (R2), and/or halogen oxides (XO; where X = Br, Cl, or I) (R3), to NO₂.



Under sunlit conditions, NO_2 is readily photolyzed to regenerate NO and O_3 (R4). The recycling of NO_x between NO and NO_2 happens much faster than NO_x oxidation to NO_3^- during the day (Michalski et al., 2003). On a global scale, NO is primarily oxidised to NO_2 by O_3 , followed by HO_2 and RO_2 , while NO to NO_2 oxidation via XO is relatively minor (Alexander et al., 2020).

During summer in the Southern Ocean MBL, NO_2 is subsequently oxidised primarily by hydroxyl radicals (OH) to form HNO_3 (R5).





In winter, under dark conditions, when the photolytic production of OH stops, NO_2 is oxidised primarily by O_3 to form nitrate radicals (NO_3) (R6). NO_3 can then react with NO_2 to form dinitrogen pentoxide (N_2O_5) followed by hydrolysis on a wet particle surface to form HNO_3 (R7-R8).



Alternatively, HNO_3 can be formed by the reaction of NO_3 with hydrocarbons (HC) (e.g., dimethylsulphide (DMS)) (R9).



Lastly, halogen chemistry may result in NO_3^- formation via the production and subsequent hydrolysis of halogen nitrates (R10-R11), as has been suggested for coastal Antarctica in summer (Bauguitte et al., 2012).



- halogen chemistry in step1) NO , NO_2 interconversion and step2) NO_2 oxidation to form nitrate with respective implications for the oxygen isotope transfer is currently not considered (Section 3.3) and not included in the oxygen isotope mixing model. However, halogens are important in the MBL particularly near/above sea ice or polar ice caps. There is evidence that halogen chemistry acts as a major NO_x sink and source of nitrate via the production and subsequent hydrolysis of XNO_3 species as observed in coastal Antarctica in summer (e.g. Bauguitte et al., 2012). Thus increases in D17O (or d18O) in nitrate may reflect increased oxidation by XO during step1 and step2 during daytime (mostly spring), possibly closely linked to local NO_x emissions of NO_x (e.g, Morin et al., 2012). This is because reaction of halogen radicals X ($=\text{Cl}, \text{Br}, \text{I}$) with ozone lead to the formation of XO

1) $\text{X} + \text{O}_3 \rightarrow \text{XO} + \text{O}_2$ followed by

2) $\text{XO} + \text{NO} \rightarrow \text{X} + \text{NO}_2$ (e.g. at 2-3 pptv BrO small impact on D17O in NO_2 and NO_3^- ; Savarino, 2016)

3) $\text{XO} + \text{NO}_2 + \text{M} \rightarrow \text{XNO}_3 + \text{M}$, $\text{XNO}_3 + \text{H}_2\text{O} \rightarrow \text{HNO}_3 + \text{HOX}$ (efficient transfer of D17O of XO and NO_2) at halogen levels of only a few pptv there is considerable impact on NO/NO_2 ratios (e.g. Savarino et al.,

2016), NO_x lifetime (Bauguitte et al., 2012; Frey et al., 2015) and impact on D17O/d18O in atmospheric nitrate. This needs to be mentioned and included in the discussion on latitudinal gradients of d18O/D17O(NO₃⁻).

A section pertaining to the potential influence of halogen chemistry during spring and its impact on the oxygen isotopic composition of atmospheric nitrate has now been included in the discussion as follows:

“Higher $\delta^{18}\text{O}\text{-NO}_3^-$ values in spring compared to summer may originate from NO_x oxidation by XO. In the Antarctic boundary layer, enhanced levels of BrO occur in spring, over sea ice covered areas (Theys et al., 2011). The production of inorganic bromine has been proposed to be related to frost flowers on thin sea ice (Kaleschke et al., 2004) and blowing of saline snow on sea ice (Yang et al., 2010). Significant interaction with sea ice cover was experienced in spring, particularly at the ice edge transect, which could have promoted NO₃⁻ formation via the BrO pathway, resulting in increased values of $\delta^{18}\text{O}\text{-NO}_3^-$.”

Additionally the relevant halogen chemistry reactions have been included in the introduction as discussed above.

- related to the above: negative correlation between d15N and D17O observed in atmospheric nitrate during Arctic spring (Morin et al., 2012) and in inner Antarctica (e.g. Savarino et al., 2016) indicate that snowpack emissions result in enhanced D17O transfer to nitrate. Possible processes include reactions with XO near halogen sources (sea ice, open ocean) or HONO co-emitted with NO_x from the snow pack contributing to the local OH budget (e.g. Legrand et al., 2014; Bond et al., 2023). Correlations between the reported D17O(d18O) and d15N especially during spring need to be analysed to discuss the impact of snow emissions and halogens on the isotope transfer. It seems to me that by overlaying Fig. 3 & 5 there is a notable anti-correlation between d18O and d15N in the spring ice edge measurements.

We thank the reviewer for this suggestion, and we do indeed see a strong anti-correlation between $\Delta^{17}\text{O}\text{-NO}_3^-$ and $\delta^{15}\text{N}\text{-NO}_3^-$ for samples collected in spring that experienced large sea ice influence (> 75%) as opposed to open ocean influence, determined using the AMBTs of the samples. As per the reviewer's suggestion, we now include a short discussion about the $\Delta^{17}\text{O}\text{-NO}_3^-$, $\delta^{15}\text{N}\text{-NO}_3^-$ relationship in spring, and invoke HONO and halogen chemistry as potential pathways of enhanced $\Delta^{17}\text{O}\text{-NO}_3^-$ production:

“Additionally, a strong anti-correlation ($r = -0.86$) is observed between $\delta^{15}\text{N}\text{-NO}_3^-$ and $\Delta^{17}\text{O}\text{-NO}_3^-$ for samples collected in spring which experience a greater than 75% sea ice influence, determined based on air mass history. A similar relationship was observed at Dome C during summer (Erbland et al., 2013; Savarino et al., 2016). Previous studies found that the production of enhanced $\Delta^{17}\text{O}\text{-NO}_3^-$ in polar regions is linked to the intensity of NO_x emissions from the snowpack (Moring et al., 2012; Savarino et al., 2016). The correlation between $\Delta^{17}\text{O}\text{-NO}_3^-$ and $\delta^{15}\text{N}\text{-NO}_3^-$ could arise from an increased contribution of HONO photolysis to total OH production, which is co-emitted with NO_x from the snowpack (Grannas et al., 2007), and induces a greater ¹⁷O excess in OH compared to the OH production pathway: O(¹D) + H₂O (Savarino et al., 2016). It could also arise from the coupling of snowpack emissions with reactive halogen chemistry as suggested by Morin et al. (2012). The $\Delta^{17}\text{O}/\delta^{15}\text{N}$ relationship presented here for the spring samples with air mass histories that indicate extensive

influence from snow covered sea ice, suggests that snowpack emissions may lead to enhanced $\Delta^{17}\text{O}$ transfer to NO_3^- .”




Figure X. The relationship between $\Delta^{17}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ in spring (square symbols). In both panels, samples collected along the ice edge are denoted by the red edge colour, with all other samples collected on the north and southbound legs of the voyage denoted by the green edge colour. The colour bar (blues) indicates the percentage sea ice influence experienced by each filter sample as determined using AMBTs. In panel a, all spring samples are included. In panel b, only samples that experienced a sea ice influence > 75% are included. A straight line (grey) is fitted to the data in panel b. Note the difference in x axis scale between panels.

SPECIFIC COMMENTS

L18 I think you mean "the dominating primary NO_x sources"

This is correct, I have updated the text to indicate the main NO_x sources.

L22-24 is the threshold for when you think O₃ oxidation dominates 60 or 70 permil? It does not make sense to have two threshold values or you have to explain why they are different in spring vs summer.

To clarify, the intention here is not to define a threshold value for $\delta^{18}\text{O}$ above which we think O₃ oxidation dominates. Instead, we are assuming that given that most of summertime $\delta^{18}\text{O}-\text{NO}_3^-$ values are relatively low (below 70‰), OH oxidation is the dominant NO₃⁻ formation pathway. Likewise, in winter and spring, given that the majority of $\delta^{18}\text{O}-\text{NO}_3^-$ values are relatively higher (greater than 60‰), additional oxidation pathways must be contributing to NO₃⁻ formation that involve O₃.

To avoid any confusion, we have removed these values from the abstract, and simply state that: "Greater values of $\delta^{18}\text{O}-\text{NO}_3^-$ in spring and winter compared to summer, suggest an increased influence of oxidation pathways that incorporate oxygen atoms from O₃ into the end product NO₃⁻ (i.e., N₂O₅, DMS and XO)."

L26-27 not only HO₂/RO₂ but also oxidation by XO (see related comments)

The influence of XO in spring has now been included in the abstract as follows:

“Significant linear relationships between $\delta^{18}\text{O}$ and $\Delta^{17}\text{O}$ suggest isotopic mixing between $\text{H}_2\text{O}_{(\text{v})}$ and O_3 in winter, and isotopic mixing between $\text{H}_2\text{O}_{(\text{v})}$ and O_3/XO in spring with the addition of a third endmember (atmospheric O_2) becoming relevant in spring.”

L48 NO_x emissions from snow are not considered a primary NO_3^- source, as this is recycled nitrate from atmospheric deposition (oceanic and lower latitude sources) and NO_3^- produced in snow or coming from the sea ice surface/ ocean. Please clarify.



We agree with the reviewer that NO_x emissions from snow are not a primary source for NO_3^- . The use of the word primary in this context is used to describe the main/dominant contributors to nitrate that are natural as opposed to anthropogenic. It is not referring to the nature/phase of the source. We have edited the sentence for clarity: “However, regional budgets of NO_x sources can have a variety of anthropogenic and natural contributors. In the summertime Southern Ocean MBL, natural NO_x sources are the main contributors to atmospheric NO_3^- formation (Morin et al., 2009; Burger et al., 2022).”

L61-62 or by halogens (see comment above)

This has been included in the text: “In addition to there being multiple NO_x sources across the Southern Ocean MBL, several different oxidation pathways can be responsible for NO_x to NO_3^- conversion, varying with chemistry and time of day (Savarino et al., 2007). Once emitted, NO is rapidly oxidised by ozone (O_3) (R1), peroxy radicals (RO_2 or HO_2) (R2), and halogen oxides (XO ; where $\text{X} = \text{Br}, \text{Cl}, \text{or I}$) (R3), to NO_2 ”

L67-68 a few pptv of BrO are sufficient. Please expand following above comment.

We have expanded on this based on the above comment as follow: “Lastly, halogen chemistry may result in NO_3^- formation via the production and subsequent hydrolysis of halogen nitrates (R10-R11), as has been suggested for coastal Antarctica in summer (Baugitte et al., 2012).



L83 Note that using d_{15}N in nitrate as a source tracer works only if any of the processes involved does not induce any significant isotopic fractionation. Please clarify.

We assume that isotope fractionation is negligible in our system. Based on a similar comment from Reviewer 1, we outline the basis for this assumption in section 3.2 as follows: “In remote environments where O_3 concentrations largely exceed NO_x concentrations, as is the case for the remote Southern Ocean, NO_x isotopic exchange occurs at a much slower rate than the Leighton Cycle reactions. Therefore, little to no equilibrium isotope fractionation is expressed, and the $\delta^{15}\text{N}$ of NO_3^- is assumed to reflect the $\delta^{15}\text{N}$ of the NO_x source (Walters et al., 2016).”

L84-85 It is misleading especially for the modellers amongst the readers to cite only a single number for d_{15}N in atmospheric nitrate originating from snow nitrate photolysis. In particular, d_{15}N in the atmospheric nitrate above snow is not constant but changes after polar sunrise as photolytic recycling and isotope fractionation between snow and atmosphere proceed into summer, going from very negative values to near zero. Thus cite

here a range of values observed in spring (when they are still strongly negative) at relevant polar locations were year-round observations are available (e.g. Wagenbach et al., 1998: Neumayer coastal Antarctica, 1986-92; Winton et al., 2020: Dome C East Antarctic Plateau 2009-15)

We now cite a range of negative values (-50 to -20‰) as per the reviewers suggestion and cite the literature Wagenbach et al., 1998 and Winton et al., 2020. The text has been amended to read: “This is distinct from the snowpack NO_x source, which typically has a very low δ¹⁵N signature (Berhanu et al., 2014; Berhanu et al., 2015) on the order of -50 to -20‰ (Wagenbach et al., 1998; Winton et al., 2020), depending on the degree of snowpack NO₃⁻ ¹⁵N enrichment (Shi et al., 2018).”

L88 what is the uncertainty (standard deviation) of this value?

The uncertainty of the value (7.6‰) has been included.

L97 can serve as a proxy (see comment above on halogen chemistry)

This sentence has been updated to: ‘Δ¹⁷O-NO₃⁻ therefore can serve as a proxy for the influence of O₃ and/or XO during NO₃⁻ formation (Berhanu et al., 2012)’.

L101-102 I strongly recommend to summarise in a table assumed isotope ratios for both

d18 and D17O in the discussed end members (O₃, OH, RO₂/HO₂, H₂O etc), including

respective uncertainties and references. Place it either here or later in section 3.3 when

prevalent oxidation pathways are discussed.

A table has now been included in section 3.3, which outlines the assumed oxygen isotope ratios of all the relevant oxidants, or sources of oxygen molecules such as H₂O, utilised for the interpretation of our results.

Table X: A summary of the oxygen isotope ratios (δ¹⁸O and Δ¹⁷O) for the end member oxidants and/or oxidant sources (O₃, OH, HO₂/RO₂ and H₂O) utilised in Sect. 3.3.

Oxidant/source	δ ¹⁸ O (‰)	References	Δ ¹⁷ O (‰)	References
Terminal O ₃	126.3 ± 11.9	Vicars & Savarino (2014)	39.3 ± 2	Vicars & Savarino (2014)
OH	-52.7 ± 2.8 ^a	Walters & Michalski (2016)	~0	Michalski et al. (2011)
HO ₂ /RO ₂	23.88 ± 0.03	Barkan & Luz (2005)	~0	Michalski et al. (2011)
H ₂ O	-13.9 ± 1.4	Dar et al. (2020)	~0	Michalski et al. (2011)

^aThe average δ¹⁸O-OH was calculated from the equilibrium fractionation between OH and H₂O_(v) (Walters & Michalski, 2016) using the observed atmospheric temperature range for winter and spring and the average δ¹⁸O-H₂O (Dar et al., 2020).

L103-04 or XO; of course the O in XO originates from O₃

This has been corrected as follows: “As such, a higher $\delta^{18}\text{O}$ or $\Delta^{17}\text{O}$ for atmospheric NO_3^- reflects the increased influence of O_3 and/or XO on NO_3^- formation, while a lower $\delta^{18}\text{O}$ or $\Delta^{17}\text{O}$ occurs when there is an increased contribution from other oxidants (Hastings et al., 2003; Fang et al., 2011; Altieri et al., 2013).”

L157-70 were the steps prior to freezing carried out on the ship right after filter exchange? please clarify.

The filter samples were immediately stored at -20°C on ship once removed from the cascade impactor. Once back in the laboratory at UCT, filter samples were extracted as outlined in the methods. To clarify this, we now write: “Once back on land, filters were extracted using ultra-clean deionised water (DI; 18 M Ω) under a laminar flow cabinet (Air Science).”

L166 the blank values are large compared to the ambient values. What are the N and O isotope ratios of the blanks? Were reported sample isotope ratios also corrected for the blank contribution? This may actually have quite an impact on the reported values if the blank comes from an isotopically very different pool.


A similar comment was made by reviewer 1 and our response is as follows:

The amount of nitrate on each stage of the field blank filters is roughly similar (e.g., average of 214 and standard deviation of 41 nmols per filter across stages 1 through 4). The spring and winter sample concentrations are much lower than the summer concentrations, therefore the percentage contribution of the blank to the total signal is larger in spring and winter. To facilitate a seasonal comparison, it was important to not increase the number of sampling hours too much from summer to winter to spring. The blank extract concentrations were all less than 1.5 μM , therefore we did not have enough volume to measure the isotopic composition directly.

In evaluating the potential sources of the blank, we concluded that it was unlikely to have a vastly different $\delta^{15}\text{N}$ than the sample nitrate for several reasons. First, and importantly, in the figures below the percent contribution of the blank vs. $\delta^{15}\text{N}$ - and $\delta^{18}\text{O}$ - NO_3^- for spring and winter show no significant relationship indicating that the measured signal is not being driven by a blank. Second, the sodium and chloride values are not unusually high in the blank filters, which lead us to conclude that there was no contamination with seawater. There is also not an unusually high value for sulfate, which makes us confident that ship stack emissions are not the source of the blank.

Finally, the coarse mode $\delta^{15}\text{N}$ is a mass weighted average of stages 1 through 4 for each filter deployment. As a result, samples where the blank is a high proportion of the total signal result in low sample nitrate concentrations, and that stage will then have a relatively low influence on the resulting mass weighted average $\delta^{15}\text{N}$. If the blank was greater than the sample concentration for a given stage that value was not used in the mass weighted average $\delta^{15}\text{N}$.

This is now noted in section 2.2.2: “It is important to note that given the low $[\text{NO}_3^-]$ of the field blanks ($< 1.5 \mu\text{M}$), no isotopic analysis could be performed on the blank filters and therefore the blank was not subtracted from the isotope results. However, we note that there was no relationship found between the blank percent contribution and $\delta^{15}\text{N}$ - or $\delta^{18}\text{O}$ - NO_3^- for spring and winter. This indicates that the measured signal is not driven by the blank contribution.”









The blank percentage of sampled (Blank %), versus $\delta^{15}\text{N-NO}_3^-$ and $\delta^{18}\text{O-NO}_3^-$ in spring (a and b, respectively) and winter (c and d, respectively).

L192-95 Considering the stability of NO_2^- in solution - When was NO_2^- measured? Were samples frozen and kept in the dark? please clarify.

Seawater sampled collected for NO_2^- determination where immediately frozen at -20°C and stored in the dark until analysis. Sample analysis was mostly conducted while on board the vessel. This is now included in the text as follows: "Seawater samples for NO_2^- determination where immediately frozen at -20°C and stored in dark conditions until analysis. $[\text{NO}_2^-]$ was analysed using the colorimetric method of Grasshof et al. (1983) using a Thermo Scientific Genesys 30 visible spectrophotometer (detection limit of $0.05 \mu\text{mol L}^{-1}$). The majority of seawater $[\text{NO}_2^-]$ analysis was conducted while at sea."

L197-204 Please provide also vertical information on the calculated back trajectories (this is output produced by default in your HYSPLIT runs), e.g. in the figures. Further below you discuss interaction with ocean/ sea ice/ snow surfaces, this applies only when the air mass arriving at the ship location spent time in the boundary layer.

Figure 1 has been amended to include the vertical information of calculated air mass back trajectories. This has been done by adding a colour bar to each subplot.






Figure 1. 72-hour AMBTs computed for each hour of every filter deployment made in winter on both the southbound (Winter S) and northbound (Winter N) voyages, in spring on the southbound voyage (Spring S), northbound voyage (Spring N) and ice edge transect (Spring ice edge) and in summer on the southbound (Summer S) and northbound (Summer N) voyages. Red triangles indicate the ships cruise track during each filter deployment. The AMBTs are coloured by height (m) (blue to green colour bar).

L210, 213 interaction with sea ice. See previous comment.

The relatively low height (< 100 m) of air mass back trajectories confirms the potential for sea ice interaction. This is included in the discussion with reference to figure 1 as follows: “The potential for sea ice influence is supported by the relatively low height (< 100 m) of AMBTs (Fig. 1).”

L241 cite also other atmospheric nitrate observation in the relevant sector of coastal Antarctica: Halley 2004-05 (Wolff et al., 2008); Neumayer 1986-92 (Wagenbach et al., 1998)

As per the reviewer’s recommendation we have cited the atmospheric nitrate observations in the relevant sector of coastal Antarctica as follows: “Atmospheric $[\text{NO}_3^-]$ ranging from tens of ng m^{-3} to approximately 100 ng m^{-3} have been observed for the Southern Ocean MBL during late spring (Morin et al., 2009; Shi et al., 2021) and observations from coastal Antarctic sites in the Atlantic sector showed elevated $[\text{NO}_3^-]$ (~ 20 to 70 ng m^{-3}) in late spring and early summer (Wagenbach et al., 1998; Wolff et al., 2008).”

L252 highest in early summer - supposedly due to the spring time depletion in stratospheric ozone. Please clarify.

Yes, increased UV radiation in spring and early summer is attributed to stratospheric ozone depletion and low noon solar zenith angle, this is now included in the text as follows: “Ground-based studies in Antarctica

demonstrate that UV radiation is highest in spring and early summer, when stratospheric O₃ concentrations are at a minimum and the noon solar zenith angle is low (Aun et al., 2020; Lakkala et al., 2020).”

L265-66 Please check vertical information of the corresponding trajectory to support this.

We have isolated the initial filter deployment in winter and plotted the air mass history of the sample, colour coded by AMBT vertical height. This has now been included as an additional supplementary figure as seen below. This new supplementary figure shows that near the sea ice edge, some AMBTs originated from 300 to 400 m and descend towards the sampling location where most air masses are at < 100 m.




Figure S1. 72-hour AMBTs computed for each hour of the first filter deployment made in winter on both the southbound (Winter S). Red triangles indicate the ships cruise track during the filter deployment. The AMBTs are coloured by height (m) (blue to green colour bar).

L272-73 Having a combined figure of all isotopes would make it easier to show this (see comment below).

We have now combined figures 2,3,5 and S3, as seen below.




Figure 2. The average coarse mode ($> 1 \mu\text{m}$) atmospheric nitrate concentration $[\text{NO}_3^-]$ (ng m^{-3}) (a), weighted average $\delta^{15}\text{N}$ of atmospheric nitrate ($\delta^{15}\text{N}-\text{NO}_3^-$ (‰ vs. N_2)) (b), $\delta^{18}\text{O}$ of atmospheric nitrate ($\delta^{18}\text{O}-\text{NO}_3^-$ (‰ vs. VSMOW)) (c) and $\Delta^{17}\text{O}$ of atmospheric nitrate ($\Delta^{17}\text{O}-\text{NO}_3^-$ (‰)) (d) as a function of latitude (° S). Winter, spring and summer are denoted by blue diamonds, green squares, and orange circles, respectively. For the summer data, where error bars (± 1 SD) are not visible, the standard deviation is smaller than the size of the marker. Spring data are separated into northbound (NB), southbound (SB) and ice edge legs by clear, light grey

and dark grey fills, respectively for panels b-d. Vertical lines indicate the approximate location of the sea ice edge in summer (orange), winter (blue) and spring (green), identified visually using satellite derived sea ice concentration obtained from passive microwave sensors AMSR2 (Advanced Microwave Scanning Radiometer 2; Spreen et al., 2008).

L283 I am surprised, NO_x atmospheric lifetimes are considerably shorter than for instance those of PAN (which in turn is admittedly stable at winter temperatures). How can NO_x reach Antarctica from lower latitudes? Can you clarify?

Previous modelling studies suggest that tropospheric transport of NO_x emitted in the mid to low latitudes (i.e., soil emissions, lightning thermal decomposition of peroxyacetylnitrate (PAN) and fossil fuel combustion), contributes to the Antarctic NO₃⁻ budget in winter (Lee et al., 2014). PAN decomposition has previously been suggested as a NO_x source to coastal Antarctica during winter and early spring (Savarino et al., 2007; Jones et al., 2011). This has now been included in the discussion for clarification as follows: “Previous modelling studies suggest that tropospheric transport of NO_x emitted in the mid to low latitudes (i.e., soil, lightning, thermal decomposition of peroxyacetylnitrate (PAN) and fossil fuel combustion), contributes to the Antarctic NO₃⁻ budget in winter (Lee et al., 2014). PAN decomposition has previously been suggested as a NO_x source to coastal Antarctica during winter and early spring (Savarino et al., 2007; Jones et al., 2011).”

L316 the triple stable isotopic composition ...

This has been corrected in the text: “The initial winter sample had a low concentration indicative of the background conditions; however, the triple stable isotopic composition of the sample confirms that it originated from the stratosphere (see sect. 3.2.1).”

L320-25 Cite also relevant Antarctic observations e.g. Wagenbach et al., 1998: Neumayer coastal Antarctica, 1986-92; Winton et al., 2020: Dome C East Antarctic Plateau 2009-15

The above Antarctic observations have now been cited: “Springtime $\delta^{15}\text{N-NO}_3^-$ is also consistent with long-term records of $\delta^{15}\text{N-NO}_3^-$ measured at coastal Antarctica (Wagenbach et al., 1998) and on the east Antarctic Plateau (Winton et al., 2020), for the same season.”

L328 refrence here also Frey et al., 2009; Erbland et al., 2013.

These references have been added. “There is a large isotope effect associated with snow NO₃⁻ photolysis during summer in the Antarctic (Berhanu et al., 2014, 2015; Frey et al., 2009; Erbland et al., 2013), resulting in the emission of low $\delta^{15}\text{N-NO}_x$ (~ -48‰) to the overlying atmosphere (Savarino et al., 2007; Morin et al., 2009; Shi et al., 2018; Walters et al., 2019).”

L332 I suppose there are no measurements of $\delta^{15}\text{N}$ in nitrate of the snowpack source? this is an important measurement gap to be addressed in the future.

Currently there are no measurements of $\delta^{15}\text{N-NO}_3^-$ in snowpack on sea ice in Antarctica that we are aware of.

This measurement gap is now addressed in the conclusions as follows: “Furthermore, the large sea ice extent characteristic of spring highlights the importance of snow-covered sea ice as a NO_x source, in addition to the

well documented summer source from snow covered continental ice (Jones et al., 2001; Walters et al., 2019; Winton et al., 2020). Currently no measurements of $\delta^{15}\text{N-NO}_3^-$ from snowpack on sea ice exist, which is an important measurement gap that should be addressed in future studies.”

L332-34 Please rephrase in light of the non-stationarity of the $\delta^{15}\text{N}$ in atmospheric nitrate from snow emissions (see comment above)

In order to acknowledge the fact that snow NO_3^- photolysis does not only lead to very low values of $\delta^{15}\text{N-NO}_x$ but rather a range of values (-50 to -20‰), depending on the degree of NO_3^- loss from the snowpack and subsequent enrichment in the snow and atmosphere, we amended the text as follows:

“This suggests that the low $\delta^{15}\text{N-NO}_x$ likely comes from snow nitrate photolysis from the snow on sea ice, before a net loss of NO_3^- from the snowpack leads to any large ^{15}N enrichment in the snow and subsequently the atmosphere (Shi et al., 2018). We conclude that NO_x as a result of photolysis of snow nitrate on sea ice can explain the relatively low $\delta^{15}\text{N-NO}_3^-$ observed in samples collected at the high latitudes on the spring southbound leg and during the ice edge transect (Fig. 3 grey filled squares).”

L343-44 limited influence from ... - the caveat is that this depends on the $\delta^{15}\text{N}$ in nitrate of the local snow source. higher $\delta^{15}\text{N}$ in atmospheric nitrate later in spring/ summer can also originate from snowpack emissions, when the source has become increasingly enriched (see comment above). Please balance your conclusion here.

We have included a caveat to our conclusion here, explaining that higher atmospheric $\delta^{15}\text{N-NO}_3^-$ values in spring and summer can also originate from snowpack emissions if the local NO_x source becomes enriched in ^{15}N , as per the reviewer’s suggestion. We then go on to explain that a lack of interaction between sampled air masses and sea ice, as indicated by the AMBTs, suggests that a snowpack emission source was unlikely influencing the samples.

The text has been edited as follows: “Higher $\delta^{15}\text{N-NO}_3^-$ values (-22.7 to -1.0‰) were observed during spring for the northbound leg (Fig. 3 open squares; Fig. 4b). The $\delta^{15}\text{N}$ of atmospheric NO_3^- that originates from snowpack emissions, depends on the $\delta^{15}\text{N}$ of the local snowpack NO_x source. ^{15}N enrichment in the snow due to NO_3^- loss, can lead to increased $\delta^{15}\text{N-NO}_x$ via photolysis, and ultimately higher values of atmospheric $\delta^{15}\text{N-NO}_3^-$ (Shi et al., 2018). However, the air mass histories of the samples indicate no contact with surrounding sea ice (i.e., the northbound leg; Fig. 3 open squares; Fig. 4b), suggesting that any influence from snowpack NO_x emissions was limited.”

L345-47 I don't understand this sentence. Please rephrase.

This sentence has been rephrased to: “The NO which originates from nitrite in seawater is thought to limit sea surface RONO_2 production. As a result, elevated nitrite concentrations are required for RONO_2 production to occur in seawater (Dahl & Saltzman 2008; Dahl et al., 2012).”

L347 oceanic RONO_2 has been long proposed as an important net primary nitrate source to the Antarctic. This should be mentioned.

We have adjusted the discussion to state that: “Oceanic RONO_2 has been long proposed as an important primary NO_3^- source to the Antarctic (Jones et al., 1991). Recent studies have used modelling and isotopic

approaches to investigate the potential importance of oceanic RONO_2 compared to other sources of NO_3^- in the Southern Ocean MBL, particularly in summer (Fisher et al., 2018; Burger et al., 2022). ”

L374 reference here some of the earlier literature (Frey et al., 2009; Berhanu 2014, 2015)

The earlier references suggested have now been added. “We know that NO_3^- photolysis in snow is associated with a large fractionation, leading to the emission of isotopically light NO_x while the remaining NO_3^- pool becomes enriched in ^{15}N (eg., Frey et al., 2009; Berhanu et al., 2014;2015; Shi et al., 2018). ”

L379-80 What is the expected time scale (or lifetime) of aerosol nitrate photolysis? If similar to snow nitrate (on the order of weeks), then it may not be relevant compared to the time scales of transport and deposition.

The formation of HNO_3 was thought to be a permanent NO_x sink in the boundary layer, due to slow photolysis rate of gaseous HNO_3 , in comparison to deposition. However, this view has been challenged by lab and field studies that show that HNO_3 adsorbed on particle surfaces is photolyzed at much higher rates than gaseous HNO_3 (Ye et al., 2016). Therefore, we refrain from using this argument to discount the potential of aerosol nitrate photolysis as a NO_x source. Given that the time scale of aerosol nitrate photolysis may be relevant against deposition, we have softened the language around our concluding statement as follows:

“Since neither of the above scenarios matches the observations, the potential influence of aerosol NO_3^- photolysis as a significant NO_x source to the region during our study is unlikely. ”

L390 qualitatively - this study is not a quantitative isotope budget

This has been changed to qualitatively.

L406 higher $\delta^{18}\text{O}$ in spring possibly also due to oxidation by XO (see above)

Additional text has been included in the discussion, which refers to the potential for XO oxidation in spring: “Higher $\delta^{18}\text{O}\text{-NO}_3^-$ values in spring compared to summer may originate from NO_x oxidation by XO. In the Antarctic boundary layer, enhanced levels of BrO occur in spring, over sea ice covered areas (Theys et al., 2011). The production of inorganic bromine has been proposed to be related to frost flowers on thin sea ice (Kaleschke et al., 2004) and blowing of saline snow on sea ice (Yang et al., 2010). Significant interaction with sea ice cover was experienced in spring, particularly at the ice edge transect, which could have promoted NO_3^- formation via the BrO pathway, resulting in increased values of $\delta^{18}\text{O}\text{-NO}_3^-$. ”

L417 Water vapour is not an oxidant. I am a bit confused here - oxygen isotope transfer from oxidants: O_3 , OH (O source atmospheric H_2O and O(1D) from O_3 photolysis), HO_2/RO_2 (O source atmospheric O_2) please clarify, also how is $\text{H}_2\text{O}(\text{v})$ mixing here.

We thank the reviewer for this comments, and agree that we need to clarify this point in the discussion. H_2O does not act as an oxidant during NO_x to NO_3^- conversion, but rather serves as an oxygen source during the oxidation process (Michalski et al., 2011). Gas phase H_2O can be incorporated into NO_3^- , during NO_2 reaction with OH (RX). OH exchanges with $\text{H}_2\text{O}(\text{g})$, such that $\delta^{18}\text{O}\text{-OH}$ depends on the $\delta^{18}\text{O}$ of the H_2O vapor it exchanged with.

Previous studies that assess the correlation between the oxygen isotopes of NO_3^- in the marine boundary layer during summer (Shi et al., 2021), similarly found that a mixing line between H_2O vapor and O_3 was the best fit to the observations. They go on to explain that if an equilibrium isotope fractionation of ^{18}O between OH and H_2O vapor is excluded (Michalski et al., 2011), such that the $\delta^{18}\text{O}$ of OH is close to that of H_2O vapor, then the lower end-member of the mixing line likely results from OH oxidation. In our case it is also likely that oxygen atoms from O_3 and $\text{H}_2\text{O}(\text{v})$ control the oxygen isotopes of NO_3^- in our study, with OH and N_2O_5 oxidation being the dominant pathways. In spring, XO oxidation may also lead to high $\delta^{18}\text{O}\text{-NO}_3^-$ and $\Delta^{17}\text{O}\text{-NO}_3^-$, similar to the N_2O_5 oxidation pathway. The figure has been updated and labelled appropriately to reflect this. Our data also suggest that a large equilibrium fractionation between OH and H_2O cannot account for the winter and spring observations, which are best explained by $\delta^{18}\text{O}\text{-OH} \sim \delta^{18}\text{O}\text{-H}_2\text{O}$. This is now included in the discussion. In addition, given that not all the spring samples are collected at the high latitudes (60°S to 70°S), for which an additional $\text{H}_2\text{O}(\text{v})$ end member equivalent to the minimum observed by Dar et al., 2020 (-27.5‰) was included, we now also include average $\delta^{18}\text{O}\text{-H}_2\text{O}(\text{v})$ determined between $\sim 33^\circ\text{S}$ and $\sim 60^\circ\text{S}$ ($-13.9 \pm 1.4\text{‰}$). Therefore the figure now includes two $\text{H}_2\text{O}(\text{v})/\text{O}_3$ mixing lines, both shown in orange. The updated figure is shown below.




Figure 4. Winter and spring $\delta^{18}\text{O}\text{-NO}_3^-$ vs. $\Delta^{17}\text{O}\text{-NO}_3^-$ are plotted in panels (a) and (b), respectively. A straight line (black) is fitted to the data in each panel. In both panels the grey line represents the OH/ O_3 mixing line, the orange line represents the $\text{H}_2\text{O}(\text{v})/\text{O}_3$ mixing line and the red line represents the O_2/O_3 mixing line. In panel (b), an additional $\text{H}_2\text{O}(\text{v})/\text{O}_3$ mixing line is included (also in orange) to account for potentially lower values of $\delta^{18}\text{O}\text{-H}_2\text{O}(\text{v})$ ($\sim -27.5\text{‰}$) at 60° to 70°S .

L414-31 This paragraph will greatly benefit from a table (see comment above on L101-102) to better follow your argument.

A table has now been included as discussed above.

L459 increase control of O_3 or XO ...

This has been corrected.

L492 potentially powerful, but complex (see previous comments); N & O stable isotope measurements of the regional sources (snow, sea ice) are required to achieve a more quantitative budget analysis. consider rephrasing.

This paragraph was re-phrased as follows:

“Our observations highlight the potential power of N and O isotopes of nitrate in distinguishing between the various natural NO_x sources that result in NO₃⁻ formation, and constraining formation pathways of aerosol NO₃⁻. In order to improve the utility of the N and O isotopes in the polar atmosphere, more measurements of the isotopic composition of the regional sources, e.g., snow on sea ice, and regional processes, e.g., OH from HONO and sea ice oxidant emissions, is needed. Even though it is complex, the utility of the N isotopes in distinguishing between the various natural NO_x sources that result in NO₃⁻ formation in the MBL of the Atlantic Southern Ocean, especially in the less frequently sampled seasons of winter and spring is evident. Furthermore, the O isotopes were able to help constrain formation pathways of aerosol NO₃⁻ seasonally. This is especially important in the Atlantic Southern Ocean where oxidation chemistry is poorly constrained (Beygi et al., 2011).”

TECHNICAL CORRECTIONS

L19 emissions ... originated from ...

The sentence was modified to indicate where the snowpack emissions come from. “Based on δ¹⁵N-NO₃⁻, the main NO_x sources were likely a combination of lightning, biomass burning and/or soil emissions at the low latitudes, as well as oceanic alkyl nitrates and snowpack emissions from continental Antarctica or the sea ice at the mid and high latitudes, respectively.”

L68 typo: from

Corrected

L106 typo: atmospheric

Corrected

L107 Antarctic tropospheric oxidation chemistry ...

Corrected

L225 In Fig2 I cannot see the second highest winter value of 22 ng/m³, is it covered by other symbols?

Yes, this is hidden by the orange circle at the same location. I have re ordered the symbols in the updated figure to make this value easier to see.

L721 typo: atmospheric

Corrected

Figures

Fig1: Label each subplot to help the reader navigate more easily, e.g. 1a. Winter-S 1d. Spring-N ... and include dates in the caption. There is a typo in the caption: ice edge transect should be (e) and N voyage (d)

The typo in the caption has been corrected, dates of each transect will be included in the caption and each subplot will be given a more descriptive label, as per the reviewer's suggestion.

Fig2,3 and 5: I strongly recommend to combine these figures including also Fig. S3. This will help to detect a lot more easily common features in [NO₃-] and N & O isotope ratios. After all they are related.

Figures 2, 3, 5 and S3 will be combined into one figure as separate panels.

To aid interpretation I also suggest to add a panel (or as a separate figure) showing air temperature, radiation (or solar elevation angel) and wind speed at the ship location.

While solar radiation data is unfortunately unavailable for these cruises we have included a figure of atmospheric temperature and wind speed in the supplementary material as seen below.




Figure S5. Daily averaged wind speed (a) and air temperature (b) for summer (orange circles), winter (blue diamonds) and spring (green squares), respectively.

Fig.4: Add labels to subplots, e.g. Spring-N ...; 4c: I suspect only the trajectories in bluish colours were within the atmospheric boundary layer above sea ice, whereas the ones with higher d15N (reddish colours) were likely higher up in the free troposphere. This is a point easily supported by including vertical AMBT info (see above).

More descriptive labels have been added to all subplots.

REFERENCES

- Bauguitte et al., Summertime NO_x measurements during the CHABLIS campaign: can source and sink estimates unravel observed diurnal cycles?, *Atmos. Chem. Phys.*, 12(2), pp 989--1002, doi:10.5194/acp-12-989-2012, 2012.
- Berhanu et al., Laboratory study of nitrate photolysis in Antarctic snow. II. Isotopic effects and wavelength dependence, *J. Chem. Phys.*, 140(24), doi:10.1063/1.4882899, 2014.
- Berhanu et al., Isotopic effects of nitrate photochemistry in snow: a field study at Dome C, Antarctica, *Atmos. Chem. Phys.*, 15(19), pp 11243--11256, doi:10.5194/acp-15-11243-2015, 2015.
- Bond et al., 2023, Snowpack nitrate photolysis drives the summertime atmospheric nitrous acid (HONO) budget in coastal Antarctica, *Atmos. Chem. Phys. Disc.*, doi:10.5194/acp-2022-845, 2023.
- Erbland et al., Air--snow transfer of nitrate on the East Antarctic Plateau -- Part 1: Isotopic evidence for a photolytically driven dynamic equilibrium in summer, *Atmos. Chem. Phys.*, 13, pp 6403-6419, doi:10.5194/acp-13-6403-2013, 2013.
- Frey et al., 2009, Photolysis imprint in the nitrate stable isotope signal in snow and atmosphere of East Antarctica and implications for reactive nitrogen cycling, *Atmos. Chem. Phys.*, doi:10.5194/acp-9-8681-2009, 2009.
- Frey et al., Atmospheric nitrogen oxides (NO and NO₂) at Dome C, East Antarctica, during the OPALE campaign, *Atmos. Chem. Phys.*, 15(14), pp 7859--7875, doi:10.5194/acp-15-7859-2015, 2015.
- Legrand et al., 2014, Large mixing ratios of atmospheric nitrous acid (HONO) at Concordia (East Antarctic Plateau) in summer: a strong source from surface snow?, *Atmos. Chem. Phys.*, 14(18), pp 9963--9976, doi:10.5194/acp-14-9963-2014, 2014.
- Morin et al., An isotopic view on the connection between photolytic emissions of NO_x from the Arctic snowpack and its oxidation by reactive halogens, *J. Geophys. Res.*, 117, doi:10.1029/2011JD016618, 2012.
- Savarino et al., Oxygen isotope mass balance of atmospheric nitrate at Dome C, East Antarctica, during the OPALE campaign, *Atmos. Chem. Phys.*, 16(4), pp 2659--2673, doi:10.5194/acp-16-2659-2016, 2016.
- Wagenbach et al., Atmospheric near-surface nitrate at coastal Antarctic sites, *J. Geophys. Res.*, 103(D9), pp 11007--11020, doi:10.1029/97JD03364, 1998.
- Winton et al., Deposition, recycling, and archival of nitrate stable isotopes between the air--snow interface: comparison between Dronning Maud Land and Dome C, Antarctica, *Atmos. Chem. Phys.*, 20(9), 5861--5885, doi:10.5194/acp-20-5861-2020, 2020.

Wolff et al., The interpretation of spikes and trends in concentration of nitrate in polar ice cores, based on evidence from snow and atmospheric measurements, Atmos. Chem. Phys., 8(18), pp 5627--5634, 2008.

Additional references utilised have now been cited.