

The authors investigate collocated in situ and lidar observations for two classes of high and low particulate linear depolarization ratio (PLDR) measured by the WALES lidar during the ML-CIRRUS field campaign as identified by Urbanek et al. (2018). The in situ measurements of interest are ice effective diameter and ice number concentration, and measurements in high PLDR (formed in air traffic regions) and low PLDR (formed in “pristine” regions) cirrus clouds are compared. After a quick presentation of the ML-CIRRUS campaign, of the WALES lidar system, and of the in situ instrumentation, comparisons are shown for two contrail cirrus clouds, two warm conveyor belt cirrus clouds, and finally for all flights, for which the comparisons are shown at 10 temperatures between 208 and 217 K. The authors conclude that in the 210-215 K temperature range, chosen to have a “sufficient contribution of both cloud types”, high PLDR mode clouds have larger effective diameter and lower number concentration, which is “an indication for more heterogeneous freezing due to aviation induces emissions”.

Even though not clarified in the abstract and in the introduction, this paper is an extension of the work presented by Urbanek et al. (2018), who found 2 classes of cirrus clouds with higher and lower PLDR, and who showed that they formed in busy air traffic regions and in regions with low aviation emissions, respectively. Urbanek et al. (2018) stated that heterogeneous freezing on emitted exhaust particles could explain the lower super saturations and higher PLDR that they found.

The lidar results derived from Urbanek et al. (2018) are well presented, which is convenient for the reader, but I was expecting to see more solid material about the in situ measurements. Section 2.3 about the in situ instrumentation does not discuss the expected performances of the instruments. It is not clear if NIXE-CAPS alone is sufficient for this study. I was expecting detailed presentations and discussions of the “combined” and “coordinated” lidar and in situ measurements, with discussions regarding the spatial and temporal collocations of the in situ and lidar legs. In the two case studies, the authors compare cirrus clouds of the same type and discuss the relevance of the comparisons. No conclusion could be drawn for the second case study (sect. 3.2) because temperatures differed by about 10 K (223 K and 232 K). In section 3.3 where all flights are combined for the comparisons, the relevance of the comparisons is not discussed. The authors present comparisons vs. temperature (Fig. 8), but only between 208 and 217 K (10 temperature values). As a matter of fact, not all in situ data were used. The authors need to detail which flights were selected for Fig. 8, why, how many PSDs per temperature range in the high and low PLDR modes, etc... I am not convinced that different number of samples justifies ignoring the cases which do not match the expectations. This might indicate that other phenomena come into play. I understand that such comparisons are challenging and that the campaign was not designed for this type of analysis.

In my opinion, the analyses presented in the manuscript are incomplete, and this manuscript does not represent a sufficient contribution to scientific progress to be accepted for publication in ACP. However, the scientific question is important, and perhaps the following suggestions and questions will help the authors pursuing this effort.

We thank this reviewer for the careful reading and the helpful suggestions to improve the manuscript! The answers to the reviewer’s comments are given in italic and bold.

Specific comments

Abstract:

1. Lines 20-22: this is misleading. These findings were published by Urbanek et al. (2018).

We now clearly indicate already in the Abstract that this study is a follow-on study of the work started by Urbanek et al. (2018).

2. Lines 22-23: in my opinion, this is an overstatement.

We skipped that sentence.

Introduction:

3. Line 83: I strongly suggest to clarify that the cirrus clouds formed in air traffic regions or in pristine regions are identified according to the classification established by Urbanek et al. (2018) which is based on lidar measurements of PLDR. Lines 55 to 59 could be moved here.

We now mention clearly that this is a work that extends the study by Urbanek et al. (2018) and that the same clouds as in this former study are investigated.

Section 2 method

Section 2.1:

4. Only 8 flights are listed in Table 1 (which should be introduced in the text), whereas 16 flights are shown in Fig. 1, with combined remote sensing and in situ observations for all these flights if I understand the text correctly. Please confirm that only the 8 flights listed in Table 1 are relevant for this study and explain why. For clarity, only the 8 flights listed in Table 1 should be shown in Fig. 1.

This is right, we did not explain why only those eight flights are uses. We corrected this in the revised manuscript and also explain better the measurement strategy for the collocated lidar and in-situ measurements by adding the following paragraph:

'However, only eight of the 16 flights were designed in a way, that they provide coordinated lidar and in-situ measurements. The sampling strategy during these flights were as follows: First, the HALO aircraft flew at higher altitudes for sounding the cirrus clouds with lidar. Subsequently, the cirrus clouds were probed by in-situ measurements at several flight altitudes within the cirrus clouds. For our study only these eight flights with coordinated lidar and in-situ measurements are relevant. Information on the flights (including their Mission ID to make it comparable to Urbanek et al. (2018)) are given in Table 1.'

5. Later in the paper, the authors refer to Table 1 when discussing number of observations in various conditions. I suggest providing for each flight information such as the number of PSDs, temperature and altitude range, PLDR range, etc...A suggestion is to add a dedicated table at the beginning of Section 3.

We like this suggestion and modified Table 1; we included information on the median PLDR (as given in Urbanek et al., 2018), of the altitude range of the cloud and the temperature of the in-situ measurements within the cloud (that was observed also by lidar). However, we did not include information on the PSDs as we did not perform the comparisons on a cloud by cloud base but a temperature base.

6. Please define the PLDR: ratio of which quantities?

Done

Section 2.3

7. How many instruments were involved for this study? How do NIXE-CAPS, CAS-DLR and CIPg-UniM compare in terms of sensitivity range? Which instrument(s) was/were not available when only data from the NIXE-CAPS instrument were available and what are the possible consequences for this study? Please specify when you state that “comparison of the data sets for all other days showed a good agreement”.

The study was conducted using the data from the combination of the CAS, CIPg-UniM and the PIP. The probe combination offers a measurement size range between 3 and 6400 μm . Deff never exceeded a value of 200 μm , therefore only the CAS and CIPg-UniM were the main instruments feeding into the analysis. This combination has been used in various other publications (Righi et al., 2020, Wang et al., 2023). We added the NIXE CAPS data (with the same size range) for one flight where no data from the main-probe-combination was available due to a failure of the CIPg-UniM. The data sets agreed well when compared for selected flights. We clarified this in the manuscript.

8. Please define the effective diameter Deff and explain how it is computed from the PSDs. I anticipate that assumptions are necessary. I could not find the Schumann et al. (2010) reference.

The reference was change to Schumann et al. 2011. Please excuse the confusion.

Schumann, U.; Mayer, B.; Gierens, K.; Unterstrasser, S.; Jessberger, P.; Petzold, A.; Voigt, C. & Gayet, J.-F. Effective Radius of Ice Particles in Cirrus and Contrails, Journal of the Atmospheric Sciences, American Meteorological Society, 2011, 68, 300 - 321

9. Line 139: the data are averaged over 5 s intervals. Can you comment on the number of PSDs for each flight? This piece of information should be provided.

The number of data points n for N_{part} and $Deff$ and each temperature interval are mentioned in Fig. 8, next to the median. Each data point n refers to a 5 s interval, where one averaged PSD in that time segment was used to derive N and $Deff$.

Section 3 results

10. It might be worth clarifying or reminding that the goal is to compare in situ measurements in two classes of cirrus clouds exhibiting large and low PLDR as established by Urbanek et al. (2018). The authors actually investigate the microphysical properties.

We have now added the following sentences to clarify that we extend the former study by comparing in-situ measurements:

'In a previous study (Urbanek et al., 2018) we found, that cirrus clouds evolved in regions with enhanced air traffic show larger mean values of the PLDR than cirrus clouds forming in rather pristine regions. In this study we extend the investigation of an impact of aviation on the microphysical properties of the cirrus clouds by comparing in-situ measurements performed within these two cloud classes (of high and low PLDR).'

Section 3.1

11. Can you explain why the cirrus with embedded fresh contrails observed on April 7th, 2014 is not somewhat affected by aviation exhaust. I could not have access to Stettler et al. (2013), but nevertheless, I think that this deserves an explanation in this paper.

We included this information in Section 2.1.

12. Line 171: why care about a missing instrument on 7 March? Unless you meant 7 April? Why no impact on the results if an instrument is missing?

This was a typo. We meant 7 April.

13. Is it possible to point to the regions with embedded fresh contrails in the lidar plots? Are these regions identified from observations or from the model?

The regions with embedded contrails were identified using the CoCIP Model (Schumann et al., 2012).

14. The authors should present in details the spatial and temporal collocations of the lidar measurements and in situ measurements presented in this paper. I note that in situ measurements are shown in Wang et al. (2022) for the 26 March, 2014 flight. The authors should clarify which in situ legs are used in this work, and how they were chosen. The number of PSDs for each flight, altitudes and temperatures could be added in Figure 3 for clarity.

We added information on the temperature of the in-situ flight legs, that was used for our comparison, in Table 1. For most of the comparison we did not restrict ourselves to only one flight leg (at one temperature) but used all the available co-located information of lidar and in-situ measurements. For the comparison of the contrail cirrus, we are in the good position, that the in-situ flight tracks were performed at the same temperature levels.

15. I see large $Deff > 100$ um on April 7th even though median $Deff$ is smaller than on March 26th. This should be acknowledged. Authors should discuss in this section

the various possible reasons for the larger N on April 7th compared to March 26th. For instance, it seems that the in situ measurements were higher in the cloud on April 7th, and perhaps closer to an embedded fresh contrail? Plots showing N vs. Deff for each case could be useful for this discussion.

The large values were due to the sampling on the base of the cloud; partly inside and partly outside. A closer look at this in-situ leg, showed, that it is not reasonable to use this flight leg as in in-situ in-cloud measurement. Thus, we better filtered our data and included a new version of the analysis. We also did a better filtering for the second case study with respect to collocated measurements.

Section 3.2

16. Lines 183-186: are you describing liquid origin cirrus clouds (e.g. Luebke et al., ACP, 2016)?

Yes, warm conveyor belts are a type of liquid origin cirrus, but at their top they can be connected to in-situ formed ice clouds.

17. The authors should present in details the spatial and temporal collocations of the lidar and in situ measurements. Same comments as for the previous case study.

We added information on the co-location of the lidar and in-situ measurements in the general description of our study (Section 2).

'However, only eight of the 16 flights were designed in a way, that they provide coordinated lidar and in-situ measurements. The sampling strategy during these flights were as follows: First, the HALO aircraft flew at higher altitudes for sounding the cirrus clouds with lidar (lidar leg). Subsequently, the cirrus clouds were probed by in-situ measurements at several flight altitudes within the cirrus clouds (in-situ leg). Typical lidar legs took about 30 min to 50 min; with a typical aircraft speed of 200 m/s that result in an observed cloud dimension of about 360 km to 600 km. The in-situ legs took a minimum of 10 min per constant flight altitude. For our study only these eight flights with coordinated lidar and in-situ measurements are relevant.'

Section 3.3

18. I believe that Fig. 7 is not really useful here and that Fig. 8 is the most interesting. That being said, the warmest temperature is 217 K, which indicates that the comparisons presented in section 3.2 (223 and 232 K) are not included. I really do not understand. I see that Fig. 8 uses data from CAS-DLR/CIPg-UniM and NIXE-CAPS, but having only NIXE-CAPS did not seem to be an issue for the case study presented in section 3.1. Please justify this choice and detail which flights were used to create Fig. 8 and which cirrus types. I suggest to give the number of PSDs at each temperature for the high and low PLDR range to avoid vague discussions.

The reviewer is right, the left part of Figure 7 is not of any value, so we removed it. However, we believe that the right part provides useful information. As we now also included the temperature in which the in-situ measurements were performed in Table 1 it gets more obvious why we restricted our comparison in Figure 8 only to the temperature range between 208 K and

217 K. For the warmer temperatures we have either only measurements in a 'high PLDR cloud' or in a 'low PLDR cloud'.

19. I see $Deff$ larger in the high PLDR mode (dark blue) than in the low PLDR mode (light blue) only at 210 K and between 212 and 214 K. N is smaller in the high PLDR mode except at 215 K and 217 K. I am not convinced that the different number of samples justifies ignoring the cases which do not match the expectations. This might indicate that other phenomena come into play.

This might have been misleading in the text. We did not omit the cases which did not match our expectations, we omitted the temperature ranges, where no comparison was possible due to a highly imbalanced number of available datapoints. We now include the number of datapoints in Figure 8. To our understanding an imbalanced number of datapoints for both cloud types provides no significant comparison and are thus omitted in the general view of Figure 7.

20. Line 247: The “tendency towards larger $Deff$ with temperatures” is consistent with numerous publications found in the literature, which should be cited.

Done

Discussion and conclusions (should be section 4?):

21. Lines 265-266: respectfully, I think that this is an overstatement. I see these findings only at 210 K and between 212 and 214 K.

We removed this sentence and tried to show our indications more carefully.