

Montreal Protocol's impact on the ozone layer and climate

Tatiana Egorova^{1,*}, Jan Sedlacek^{1,*}, Timofei Sukhodolov^{1,4,2,*}, Arseniy Karagodin-Doyennel^{2,1,3}, Franziska Zilker³, and Eugene Rozanov^{1,*4,3}

¹ Physikalisch-Meteorologisches Observatorium Davos and World Radiation Center, Davos, Switzerland

² ~~Department of Earth Sciences, Vrije Universiteit Amsterdam, Amsterdam, Netherlands~~^{Institute of Meteorology and Climatology, University of Natural Resources and Life Sciences, Vienna, Austria}

³ Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

⁴~~Saint Petersburg State University, Saint Petersburg, Russia~~

*These authors contributed equally to this work.

Correspondence to: Tatiana Egorova (tegorova@pmo-dwrc.ch)

Abstract. It is now recognized and confirmed that the ozone layer shields the biosphere from dangerous solar UV radiation and is also important for the global atmosphere and climate. The observed massive ozone depletion forced the introduction of limitations on the production of halogen-containing ozone-depleting substances (hODS) by the Montreal Protocol and its Amendments and adjustments (MPA). ~~Previous research has demonstrated the success of~~ ~~Further research was aimed at analyzing the role played by~~ the Montreal Protocol ~~to~~ ~~and~~ increase public awareness of its necessity. In this study, we evaluate the benefits of the Montreal Protocol on climate and ozone evolution using the Earth system model (ESM) SOCOLv4.0 which includes dynamic ocean, sea ice, interactive ozone, and stratospheric aerosol modules. Here, we analyze the results of the numerical experiments performed with and without limitations on the ~~ODS~~~~ozone-depleting substances~~ emissions. In the experiments, we have used CMIP6 SSP2-4.5 and SSP5-8.5 scenarios for future forcing behavior. We confirm previous results regarding catastrophic ozone layer depletion ~~and substantial climate warming~~ in the case without MPA limitations. ~~We show that~~~~The-~~ ~~the~~ climate effects of MPA consist of additional global mean warming by up to 2.5 K in 2100 caused by the direct radiative effect of the hODS, ~~which is comparable to large climate warming obtained with SSP5-8.5 scenario.~~ ~~We also~~ ~~For the first time we reveal~~ ~~obtained~~ dramatic effects of MPA on chemical species and cloud cover. ~~The response of surface temperature, precipitation and sea-ice fields~~ was demonstrated for the first time with the model that has interactive tropospheric and stratospheric chemistry. ~~We have found some differences in the climate response compared to the model with prescribed ozone, which should be further addressed.~~ ~~changes in several essential climate variables such as regional surface air temperature, sea ice cover, and precipitation fields.~~ Our research updates and complements previous modeling studies on the quantifying of MPA benefits for the terrestrial atmosphere and climate.

Short description: for submission purposes

This paper describes the climate and atmosphere benefits of the Montreal Protocol, simulated with the ~~state-of-the-art~~ Earth system model SOCOLv4.0. We have added to and confirmed the previous studies by showing that without the Montreal Protocol by the end of the 21st century there would be a dramatic reduction of the ozone layer as well as substantial

Formatted: Superscript

35 perturbation of the essential climate variables in the troposphere caused by the warming from increasing ozone-depleting
substances.

1 Introduction

40 The ~~evolution of~~ ozone layer ~~evolution~~ remains a major problem in contemporary science because of its significance for the
sustainable development of human civilization. The ozone not only shields the biosphere from dangerous solar UV radiation
but is also essential for the global atmosphere and climate (e.g., Bais et al., 2018, Barnes et al., 2019, Neale et al., 2021).
Ozone depletion was registered first using the ground station data (Farman et al., 1985). Afterward, it was confirmed by
space observations, which showed that this ozone depletion coined as the Antarctic ozone “hole” can cover the entire
Antarctic (Stolarski et al., 1986). Physical explanations for the massive southern polar ozone loss (see e.g., review by
45 Solomon, 1999), established the role of heterogeneous chemistry and previously suggested impact of anthropogenic halogen-
containing ozone-depleting substances (hODS) (Molina and Rowland, 1974). These discoveries led to limitations on hODS
production by the Montreal Protocol and its Amendments and adjustments (MPA) (e.g., Rowland, 2006). Further research
was aimed at analyzing the benefits of the MPA to increase public acceptance of its necessity, projection, and estimation of
the time at which ozone layer recovery will be reached, and careful analysis of observational and model data to characterize
50 the ozone layer evolution. According to the last Executive summary of Scientific Assessment of Ozone Depletion: 2022
(WMO, 2022), Montreal Protocol is continuing to decrease atmospheric abundances of regulated hODS and contribute to
ozone recovery. However, the widely expected future ozone recovery ~~and the efficiency of the MPA~~ are now being
questioned due to disagreement between models and observations and inspire a search for missing processes or unaccounted
driving factors in models (Ball et al., 2018, 2019, Dietmüller et al. 2021).

55 Several attempts have been made to estimate the ~~efficacy~~~~efficiency~~ of the MPA by comparing the future state of the ozone
layer simulated under two hypothetical scenarios: with and without MPA limitations. ~~The so called “world avoided” case is
interesting because it allowed estimating advantages of Montreal Protocol implementation, which was associated with
certain difficulties and motivating reasons to perform extreme case without MPA restrictions are described in Newman et al.
(2009).~~ The first estimates were made with relatively simple models (e.g., Prather et al., 1996). Later, Egorova et al. (2001),
60 and recently Chipperfield et al. (2015), performed more accurate estimates using 3-D chemistry-transport models driven by
observed meteorological fields. An improved understanding of the influence and importance of ozone on the climate led to
the application of chemistry-climate models that can account for stratospheric changes related to the ozone evolution
(Morgenstern et al., 2008; Newman et al., 2009; Egorova et al., 2013). The results of ~~these studies confirmed that the
absence of MPA limitations would have led to dramatic ozone loss and an increase in the level of dangerous UV radiation~~
65 near the surface. However, these models have certain limitations for modeling the effect of the Montreal Protocol restrictive
measures on ozone evolution and climate. For example, in Newman et al. (2009) and Egorova et al. (2013), the use of

prescribed sea-surface surface temperature did not allow considering tropospheric temperature response to halogen forcing or stratospheric changes. And the lack of troposphere chemistry (Morgenstern et al., 2008, Newman et al., 2009, Egorova et al., 2013) led to incorrect modeling of the tropospheric response with possible implications for the stratosphere. In addition, ~~halogen-containing ozone-depleting substances (hODS)~~ are also very potent greenhouse gases (GHG). They can directly affect radiation balance (e.g., Polvani et al., 2020) in the present-day atmosphere, but in the case of hODS uncontrolled increase their radiative forcing can be comparable to or even exceed the other GHG (Liang et al., 2022). Therefore, proper evaluation of the MPA influence on climate is not possible without the application of dynamically coupled ocean-atmospheric models. First, Garcia et al. (2012) applied a coupled chemistry-climate-ocean model and showed that, in addition to the collapse of the ozone layer, the uncontrolled growth of hODS would have caused enhanced global mean warming from 2010 to 2070 between 2 K, in the tropics, and 6 K, in the Arctic, ~~by the end of 2070~~; this is comparable to the effects of carbon dioxide increase from 325 to 560 ppmv (IPCC RCP 4.5 scenario) and would have had substantial implications for sea ice coverage. However, the robustness of these conclusions cannot be established because this experiment had only one run and was not performed with other models. Further, Goyal et al. (2019) tried to increase awareness for a wider community about the MPA's success in climate warming reduction. They used ACCESS1.0 coupled atmosphere-ocean-land-sea-ice but without interactive atmospheric chemistry to estimate the impact of MPA on surface temperature warming. In this work, the stratospheric ozone field was prescribed from GEOS CCM (Pawson et al., 2008) for no MPA and reference scenarios, while the tropospheric ozone concentration was fixed for both experiments. Goyal et al. (2019) showed that for the year 2019 the MPA already prevented additional 0.5-1.0°C warming over land in middle latitudes (Africa, North America, Eurasia), and in the Arctic. For 2050, they obtained larger MPA effects, reaching 1.5-2.0°C warming in the extra-polar areas and 3-4°C in the Arctic with the greenhouse forcing from the RCP8.5 scenario. However, there are various shortcomings in this study. First, the fixed tropospheric ozone approach is difficult to justify for the no MPA case due to the expected dramatic increase of UV radiation in the troposphere caused by stratospheric ozone depletion and strong tropospheric ozone radiative forcing. Second, the provided results were not supported by statistical significance analysis, and therefore the conclusions about MPA effects in Goyal et al. (2019) require further verification. Morgenstern et al. (2020) obtained larger cooling by ozone depletion than Goyal et al. (2019) and Polvani et al. (2020) and pointed out that the effect of MPA on surface climate would be smaller, therefore there is a need to reduce the uncertainties in the estimation of the MPA regulation effect on surface temperature (Neale et al., 2021). Because of the uncertainties in MPA climate effects estimations, more studies are necessary to clarify and obtain robust results of MPA benefits for the Earth's climate. In this paper, we applied the Earth system model (ESM) SOCOLv4.0 (Sukhodolov et al., 2021) to simulate the dramatic global scale depletion of the ozone layer and its consequences on Earth's climate. In Section 2 we describe the model, experiment design, and forcing. The results of this study are provided in Section 3 followed by the conclusions summarized in Section 4.

2 Method

100 2.1 Model description

SOCOLv4.0 used in this study has the standard configuration with horizontal resolutions (T63, approximately 1.9x1.9 degrees) and 47 layers in the vertical direction and has been validated against observations and other models using comprehensive statistical tools (Sukhodolov et al., 2021). The model consists of the Earth system model (MPI-ESM, Hamburg, Germany) (Mauritsen et al., 2019), the chemistry (MEZON) (Egorova et al., 2003), and the size resolving sulfate aerosol microphysics (AER) (Sheng et al., 2015) modules. The chemical module treats more than 100 gas species linked by 216 gas-phase, 72 photolysis, and 16 heterogeneous reactions in/on aqueous sulfuric acid aerosols and polar stratospheric clouds. The photolysis rates are calculated online using the look-up-table scheme. All modules are interactively coupled by the 3-dimensional meteorological fields of wind and temperature as well as by the radiative forcing of sulfate aerosol and radiatively active gases including chlorofluorocarbons (CFCs). The model directly simulates many important forcings and 110 feedbacks in the climate system, including oceanic processes and dynamic vegetation. Thus, the model accounts for most of the known atmospheric processes involved in the ozone net chemical production and transport as well as chemistry-dynamics feedbacks. Extensive model validation showed that the climatology and variability of the temperature/circulation fields, as well as the distribution of several key atmospheric tracers, are successfully reproduced (see Sukhodolov et al., 2021 for further details). Despite some differences related to interactive chemistry and aerosol the representation of several climate parameters such as clouds, precipitation and sea-ice coverage is very similar to the core MPI-ESM model discussed in Mauritsen et al. (2019).

115 2.2 Experimental design and forcing

To assess the benefits of the Montreal Protocol and its Amendments and adjustments for atmospheric composition and climate, we performed two main 3-member 120-year long (1980-2100) ensemble simulations with SOCOLv4.0 switching on 120 (MPA) and off (noMPA) limitations on the emissions of hODS. For the model spin-up time we used December 1949 restart from MPI-ESM 100 years historical experiment (Maher et al., 2019). For the MPA and noMPA cases, the model boundary conditions mostly follow the recommendations of CMIP6 provided by the input4MIPs database, (<https://esgf-node.llnl.gov/search/input4mips/>) as described in Sukhodolov et al. (2021). All anthropogenic forcing except hODS is historical before 2015 and then switched to the SSP2-4.5 scenarios till 2100, accordingly. The experiment with MPA 125 restrictions under the SSP2-4.5 scenario further will be named MPA_ssp245. For the long- and short-lived halogenated source gases, we used the baseline mixing ratio scenario from WMO (2018), which is a combination of the observation record up to the year 2017 (Engel et al., 2018) and CMIP6 data (Eyring et al., 2016). Several newly discovered and unregulated hODSs (CFC-112, CFC-112a, CFC-113a, CFC114a, and HCFC-133a) have been introduced to the model chemistry scheme together with some additional chlorine-containing very short-lived substances (VSLs: CHCl₃, CH₂Cl₂, 130 C₂Cl₄, C₂HCl₃, C₂H₄Cl₂) that are not controlled by the MPA (Hossaini et al., 2017). For the noMPA experiment, we

applied the same boundary conditions except for the hODS, whose surface mixing ratio was increased by 3% per year since 1987 (Velders et al., 2007) for regulated species. For unregulated species we follow the recommendations of WMO (2018): from 2016 CFC112a grow 1.5%/yr, CFC113a grow 6.5%/yr, HCFC-133a grow 5.4%/yr. HCFC-141b, CH3Cl, CHBr3, CH2Br2, and H-2402 are assumed to be equal to the reference case value. In addition to the main two experiments, we performed a run with the MPA limitations using the SSP5-8.5 scenario (MPA_ssp585) to compare climate effects with the MPA_ssp245 experiment. To distinguish between the ozone and direct greenhouse effects of CFCs, we performed a model run, where increasing CFCs were active only chemically but not radiatively under the SSP2-4.5 scenario (noMPA_noRadCFC). The statistical significance of all results shown in the following sections has been calculated using t-test t-test with a two sided 90 % -level.

Table 1. Model experiments

<u>Experiment name</u>	<u>SSP used</u>	<u>MPA limitations</u>
<u>MPA_ssp245</u>	<u>2-45</u>	<u>yes</u>
<u>MPA_ssp585</u>	<u>5-85</u>	<u>yes</u>
<u>noMPA</u>	<u>2-45</u>	<u>no</u>
<u>noMPA_noRadCFC</u>	<u>2-45</u>	<u>no for chemistry</u> <u>yes or radiation</u>

3 Results and discussion

3.1 MPA benefits for the total column ozone

Figure 1 shows the global, annual and ensemble mean total ozone from the MPA_ssp245, MPA_ssp585, and noMPA experiments simulated with SOCOLv4 and the amount of saved total ozone by MPA, calculated as the absolute difference between the MPA_ssp245 and noMPA experiments. Without MPA, ozone could be almost entirely depleted by the end of the 21st century and this result agrees very well with previous publications (e.g., Newman et al., 2009, Garcia et al., 2012, Egorova et al., 2013). The model results demonstrate that the continuous increase of stratospheric halogens destroys about 80% of the total ozone at the end of the 21st century, and it is difficult to overemphasize the importance of the ~~Montreal Protocol and its Amendments and adjustments~~MPA for preserving the ozone layer. Besides the direct consequences for human health, the increased surface UV radiation would also strongly affect the terrestrial and aquatic ecosystems (Neale et al., 2021) and biogeochemical cycles (Young et al., 2021).





Figure 1: Time evolution of the global mean annual mean total ozone (DU) for the MPA_ssp245 (blue line), MPA_ssp585 (green line), and noMPA (orange line) simulations, the red line illustrates saved total ozone by the MPA limitations ($MPA_ssp245 - noMPA$).

Figure 2 shows the geographical distribution of annual, and ensemble mean total ozone changes due to the absence of MPA regulations for the time periods 2021-2040 and 2080-2099. The most pronounced changes take place over the northern high latitudes and mid-latitudes of both hemispheres, where halogen chemistry is the most active. It is interesting to note that the southern polar area is less sensitive. It is explained by already large ozone depletion caused by the present-day hODS level, leading to slight saturation of the effects. As expected, smaller, but still considerable changes occur over the tropical latitudes where the ozone layer is still maintained to some extent by the in-situ production through photolysis of molecular oxygen. All obtained results are statistically significant even for the near present time 2021-2040, which means that in no MPA world some dramatic consequences could already be observable.



165 **Figure 2:** Geographical distribution of the annual mean total ozone difference (in D.U.) between noMPA-ssp245 and noMPA-ssp245 simulations, averaged over the periods 2021-2040 (left upper part) and 2080-2099 (right lower part). All differences are statistically significantly different from zero at the 90% level! The colored areas mark statistically significant results at the better than 95% level.

3.2 Chemical composition and temperature changes

170 It is of interest to analyze the impact of hODS on chemical composition and temperature because it helps to understand the response of the ozone and climate. Moreover, the extreme conditions that the atmosphere would be put in the luckily avoided noMPA scenario presents an interesting exercise that allows us to provide new insights learn more and verify our knowledge about the chemical and dynamic links in the atmosphere/climate system. Figure 3 shows the response of several essential chemical species and temperature to the absence of MPA limitations for the 2080-2099 period. The period is chosen because the response pattern is quite close to the earlier periods, while the magnitude of the signal is larger. An uncontrolled increase of the hODS emissions dramatically (around 2000%) enhances Cl_y concentration. This primary forcing leads to ozone depletion in most of the stratosphere. The ozone decrease maximizes in the upper and lower stratosphere. In the upper stratosphere, the efficiency of gas-phase chlorine-based catalytic ozone destruction cycles is the highest (e.g., Revell et al., 2012). In the lower stratosphere, ozone depletion is related to the enhanced liquid aerosol concentration in the Junge layer (Junge and Mandson, 1961), a colder environment, and the presence of polar stratospheric clouds. In the lower stratosphere, almost complete ozone depletion similar to the finding of Newman et al. (2009) is explained by the acceleration of heterogeneous chlorine activation, which is caused by the cooling in this area. The cooling accelerates heterogeneous reactions and enhances surface area density (SAD) due to the additional generation of the liquid sulfate aerosol in the Junge layer (Junge and Mandson, 1961) and the appearance of the polar stratospheric clouds (PSCs). The obtained increase of annual mean SAD in the noMPA run relative to the reference exceeds 100% in the entire lower stratosphere, except southern high latitudes (not shown). Lower SAD in this area is explained by a substantial decrease of PSC1 due to HNO_3 depletion caused by enhanced chlorine loading. In the tropical lower stratosphere (10-20 km) the model shows about a 20% ozone increase caused by the increase in the oxygen photolysis rates in the lower stratosphere related to strong ozone depletion

Formatted: Subscript

higher up leading to a downward shift of the net ozone production. The obtained pattern of the ozone response is very close to the other model results (e.g., Morgenstern et al., 200; Newman et al., 2009).

190 The ozone abundance is tightly related to atmospheric chemistry due to the modulation of the temperature structure and photolysis rates. Well-pronounced cooling in the stratosphere is mostly related to ozone depletion and reaches maxima in the upper stratosphere, where the ozone absorption of solar UV radiation is the main energy source. The secondary maximum cooling in the lower stratosphere is formed by both ozone depletion and intensification of the Brewer-Dobson circulation (e.g., Zubov et al., 2013) caused by the tropospheric warming owed to the direct radiative forcing of hODS as well as by the modulation of the temperature gradients in the stratosphere. The tropical pattern of the temperature response is similar to the Garcia et al. (2012) and Newman et al. (2009) results. More details of climate warming will be discussed in section 3.3.

195 Tropospheric warming is responsible for the increase of tropospheric water vapor mostly because warmer air can hold more water vapor according to Clausius–Clapeyron law and partially due to enhanced evaporation from the warmer surface.

~~Tropospheric warming is responsible for the increase of tropospheric water vapor due to enhanced evaporation in a warmer climate.~~ Together with enhanced ozone photolysis followed by more intensive $O(^1D)$ production, it leads to substantially higher (up to 100%) hydroxyl concentration, with a negative impact on tropospheric ozone that is also lowered because of the reduced transport from the stratosphere. The stratospheric water vapor is slightly decreasing in the upper stratosphere, where the methane oxidation cycle is still operational. However, in the lower stratosphere, where the influx from the troposphere dominates, the water vapor concentration drops by up to 70% due to a cooler environment and a more efficient

200 cold trap at the entry level. This process and lower ozone concentration (less $O(^1D)$) suppresses hydroxyl radical production and decreases the stratospheric oxidation capacity. A substantial amount of HO_x is also removed via $HO_2 + ClO = HOCl + O_2$ due to very high chlorine loading. The effect of a lower OH mixing ratio is more visible in the weaker ozone depletion in the lower mesosphere and partially in the tropical lower stratosphere. The major source gases such as methane and N_2O demonstrate rather similar behaviour, with very weak changes in the troposphere and substantial depletion above approximately 30 km. The methane decrease in the troposphere is caused by enhanced hydroxyl concentration. Above 30 km, methane is being destroyed by atomic chlorine via the reaction $CH_4 + Cl = HCl + CH_3$. Tropospheric N_2O is not extensively sensitive to other species and photolysis rates, so it only very weakly reacts to the introduced forcing. Two spots of positive N_2O changes over the extratropics are probably formed due to the deficit of the $O(^1D)$ related to ozone depletion.

205 N_2O decrease above 25 km is most likely related to the enhanced photolysis ($N_2O + h\nu = N_2 + O$), which does not enhance NO_y production. As a result, NO_y concentration drops in the entire stratosphere because of enhanced NO photolysis followed by the cannibalistic reaction $NO + N = N_2 + O$ in combination with a substantial N_2O decrease. Reactive NO_x radical depletion is even more intensive than NO_y . It can be explained by the conversion to $ClONO_2$ (via $NO_2 + ClO = ClONO_2$) in the high chlorine loading atmosphere. NO_y increase in the upper tropical troposphere is partly related to the enhanced lightning flashes frequency and convective activity in the warmer climate (e.g., Mareev and Volodin, 2014; Revell et al., 2015). The


220 obtained results show that an uncontrolled increase in halogen loading affects chemical processes and the temperature distribution in the entire atmosphere from the ground to the mesosphere.

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Font: Italic






Figure 3: Zonal, annual, and ensemble mean of total inorganic chlorine (Cl_y), O_3 , $\text{NO}_x(\text{N}+\text{NO}+\text{NO}_2)$, OH, H_2O , total reactive nitrogen (NO_y), CH_4 , H_2O , N_2O (in %), and temperature (in K) response to the absence of the MPA regulations (noMPA relative to - MPA-ssp245) averaged over 2080–2099. All differences are statistically significantly different from zero at the 90% level except areas with black dots. The colored areas mark statistically significant results at the better than 95% level.


3.3 MPA implications for the surface air temperature

As it was mentioned above, the troposphere would warm by up to 3 K without MPA restrictions at the end of the 21st century (see figure 3). It means that the MPA not only protects the ozone layer but also helps to limit greenhouse warming (Velders et al., 2007) described in IPCC (2013). Here, we analyze surface air temperature in more detail to illustrate the benefits of the MPA limitations for the surface climate. Figure 4 shows the time evolution of global mean temperature response for all considered cases in comparison with the other model results published by Garcia et al. (2012), Goyal et al. (2019), and Young et al. (2021). Garcia et al. (2012) exploited Whole Atmosphere Community Climate Model with interactive chemistry, coupled to a deep ocean model and IPCC RCP4.5 scenario for greenhouse gas (GHG) emissions. Australian Community Climate and Earth-System Simulator v1.0 (ACCESS1.0) which is a coupled atmosphere ocean-land-

240 sea-ice model without interactive chemistry was applied in Goyal et al. (2019) to simulate the atmospheric state using IPCC RCP8.5 scenario for GHG with and without MPA. Young et al. (2021) simulated climate using NIWA–UKCA chemistry–ocean-climate model in combination with land surface model driven by IPCC RCP 6.0 scenario for GHG and also with and without MPA limitations. Figure 4 demonstrates the difference between noMPA and reference runs, which does not dramatically depends on the basic scenario. Before approximately 2030, all models show a slightly (within 0.25 K) warmer climate. Then the warming started to grow at a higher rate for Garcia et al. (2012), Goyal et al. (2019), and Young et al. (2021). Our model demonstrates weaker warming until approximately 2060 and a sharp temperature increases during the second half of the century. In 2100 the warming from our model is even higher than in Young et al. (2021). This disagreement looks statistically significant, but the reason is not easy to understand. The model’s peculiarities could play a role. It is known (e.g., IPCC, 2013) that climate sensitivities are not in perfect agreement among the models. The ocean inertia could also strongly affect the magnitude of the response. Another aspect is related to the treatment of the hODS distribution. In climate models without interactive chemistry, hODS are usually prescribed as a single value for the entire atmosphere. In the chemistry-climate models, there is the possibility to do the same or utilize 3-D hODS concentration from the chemical module for calculating radiation fluxes and heating rates. Due to intensive destruction by solar UV radiation, hODS almost disappear from the stratosphere, which diminishes their radiative forcing. This process is insignificant for the hODS concentration close to the present-day values but becomes more important in the future for the noMPA case. It can explain a smaller temperature response in our model in comparison with the prescribed chemistry model of Goyal et al. (2019). Still, the reason for the difference with results from Garcia et al. (2012) remains unclear because we do not know how hODS were treated in their model.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt






Figure 4: Global mean surface air temperature deviation for the noMPA (redorange line), MPA-ssp585 (brownmagenta line), and noMPA_noRadCFC (violet line) from the reference MPA-ssp245 run. Light red shadow illustrate internal variability in noMAP run. For comparisons, the same quantity is shown using digitized data from Garcia et al. (2012, blue line), Goyal et al. (2019, yellowred line), and Young et al. (2021, light greenblue line).

An interesting fact to note is that the noMPA case gives virtually the same global warming rate as the MPA-ssp885, which emphasizes the importance of MPA for global climate. The influence of the ozone change is also shown in Figure 4. For the noMPA_noRadCFC run, uncontrolled hODS increase was applied only for the chemical module, while infrared radiative forcing from hODS was kept as in the reference (MPA_ssp245) case. The ozone depletion discussed earlier in section 3.2 caused significant global cooling by up to 0.8 K. This can be explained mainly by the decrease of ozone in the troposphere and stratosphere (see Figure 3), where it has a strong radiative forcing potential. The separation of these two physical processes requires several additional experiments and cannot be done in the present paper framework.

Formatted: Centered

Formatted: English (United States)





Figure 5: Geographical distribution of annual mean difference of the surface air temperature between noMPA and MPA_ssp245 simulations, averaged over the periods 2040-2059, and 2080-2099 (in K). Areas, where the statistical significance of the signal is less than 90%, are marked by dots.

Figure 5 shows the regional effects of MPA regulations on surface air temperature for annual mean values for two periods. Uncontrolled hODS increase leads by the end of the 21st century to robust warming all around the globe, ~~except marginally significant cooling in equatorial Africa.~~ The pattern of the warming is typical for the warmer climate (IPCC, 2013) with up to 8K warming in the northern high latitude, reflecting well-known Arctic amplification (e.g. Prevedi et al., 2021). In general, the warming maximizes over the land masses and is only about 1 K over the oceans. The smaller warming as in the case of GHG increase (e.g., IPCC, 2013) is visible over the northern Atlantic. The temperature response is insignificant during the 2020-2040 period (not shown) and started to appear over North America, the Arctic, Asia, and Africa only after 2040. Small and even not robust warming just after 2020 does not agree with the results of Goyal et al. (2019), who demonstrate substantial warming already during the early 21st century. It looks reasonable due to their larger global mean response (see Figure 4). However, their results are also difficult to interpret because Goyal et al. (2019) did not provide statistical significance of the results and internal variability of the climate system can provide quite substantial noise even with 5 ensemble members.


3.4 MPA impact on cloud cover, precipitation, and sea ice

Among essential climate variables, cloud amount and precipitation intensity play an important role because they demonstrate energy and water flux changes in the climate system, which are dramatically important for the environment (e.g., IPCC, 2013). Figure 6 (left panel) illustrates the evolution of the global and annual mean cloud cover for the MPA (reference) and noMPA cases relative to the 1980-1990 period. The cloud amount response is ~~very~~ small (within 1%) until approximately

2060 for both cases, and the difference between cases is not statistically significant. After 2060 global mean cloud amount decrease due to the absence of the MPA limitations is observable, reaching about 2% at the end of the century and more pronounced than in the reference case. The small global mean response could be related to the redistribution of the cloud field caused by the circulation changes in the warmer climate. The geographical distribution of the cloud amount supports this explanation, showing a very inhomogeneous pattern. The cloud amount increase over the high and tropical latitudes is competing with a pronounced decrease over the extratropics leading to very small changes (up to 2% at the end of the 21st century) in the global mean. The obtained pattern resembles the results of climate models obtained for the RCP8.5 scenario (see IPCC, 2013, Figure 12.17), but there are some deviations in the maritime continent, South America, Australia, and Southern Africa, where the cloud response to hODS forcing has a different sign. It should be noted, however, that these differences could be related rather to our model peculiarities.




Formatted: Centered



305 **Figure 6:** Time evolution of global and annual mean cloud cover change (left panel) for MPA-ssp245 (blue) and for noMPA (orange) cases relative to 1980-1990 mean and geographical distribution (right panel) of the annual mean difference in cloud cover (%) between noMPA and MPA_ssp245 simulations, averaged over the period 2090–2100. Areas, where the statistical significance of the signal is less than 90%, are marked by dots.

310 Figure 7 shows substantial changes (up to 9% at the end of the 21st century) in global mean precipitation and its spatial distribution. In a warmer climate, we expect more water vapor in the atmosphere, and consequently substantial changes in precipitation intensity. It is known (IPCC, 2013, 2022) that the global mean precipitation increases in a warming climate. The rate is not perfectly constrained and can be within 1-4 % per 1 K warming. In our case, we have about 2 K warming for the noMPA case (see Figure 4) and a consistent extra 7% precipitation amount. As expected, the geographical distribution of the precipitation changes resembles the cloud cover response.




315

Figure 7: Time evolution of global and annual mean precipitation change (left panel) for reference MPA-ssp245 (blue) and for noMPA (orange) cases relative to 1980-1990 mean and geographical distribution (right panel) of the annual mean difference in precipitation between noMPA and MPA-ssp245 simulations averaged over the period 2090–2100 (in %). Areas, where the statistical significance of the signal is less than 90.5%, are marked by dots.

320 High latitude areas, the tropical belt, Australia, and the maritime continent are experiencing substantial (more than 20%), statistically significant precipitation increase. Greater changes over the high latitudes are explained by increased water vapor transport from the tropical latitudes (IPCC, 2013). The Mediterranean region becomes substantially (by up to 20%) drier, responding to the absence of MPA limitations. Over the Pacific Ocean, the pattern is not homogeneous. The obtained precipitation changes resemble the results presented in IPCC (2022, Figure 4.24) but are not identical. Our model produces
 325 very different precipitation changes over Australia and South Africa. However, the magnitude of the changes due to the absence of MPA limitations is comparable with the precipitation response obtained for the SSP3-7.0 scenario (IPCC, 2022).

The status of the Northern Sea ice cover is essential for the Arctic biosphere as well as for the economic issues related to transportation, tourism, population life, and fossil fuel extraction (e.g., Notz et al., 2016). All these factors make sea ice one of the main climate indicators. As shown above, in Figure 5, the Arctic region experiences large warming if the MPA had not been implemented. The implications on the September sea-ice cover are visible in Figure 8 (left panel). The Arctic would be ice-free (cover less than 1 Mio. km²) in September around 40 years earlier, i.e., around 2050 without MPA compared to 2090 with MPA. A similar early timing of an ice-free Arctic is also seen for simulations with the SSP5-8.5 scenario (see for example Lee et al., 2021: IPCC AR6 Figure 4.2c). The minimum sea-ice cover is decreased on the whole surface quite uniformly (Figure 8 right panel). In spring, however, which is the time of the maximum sea-ice extension, the reduction is
 330 confined to the edge of the sea-ice cover. Hudson Bay and Baffin Bay would be largely ice-free at the end of the century without the MPA. Additionally, the sea-ice edge between Svalbard and Siberia will move to the North and Fram Strait has a reduced cover, leading to the conclusion that there is less transport of sea ice out of the Arctic Basin.



340 **Figure 8:** Left panel: September sea-ice extent time evolution for reference MPA-ssp245 (blue) and for no MPA (orange) cases; Right panel: Sea-ice loss due to hODSs for 2080-2099, the red line represents sea ice extent of the reference MPA-ssp245 run.

4 Summary and conclusions

This paper describes the climate and atmosphere benefits of the ~~Montreal Protocol and its Amendments and adjustments (MPA)~~ simulated with the ~~state-of-the-art~~ Earth system model SOCOLv4.0 using the CMIP6 boundary conditions and socioeconomic pathway scenarios. We have added to and confirmed the previous studies by showing that without the MPA
345 by the end of the 21st century there would be a dramatic reduction of the ozone layer as well as a huge perturbation of the essential climate variables in the troposphere caused by the warming from increasing hODSs.

With performed simulations, we have confirmed the value and benefits of MPA limitations for the ozone layer and climate, considering multiple feedbacks within the Earth system. The uncontrolled increase in hODS influences the chemical processes and temperature distribution in the entire atmosphere from the earth to the mesosphere. Ozone could be almost
350 entirely depleted, and surface temperature could be warmer all around the globe and up to 8K in the northern high latitude, reflecting well-known Arctic amplification. Also, cloud cover and precipitation would be affected. Our simulations with ESM SOCOLv4 showed very small changes in global mean cloud amount, by about a 2% decrease at the end of the century due to the competing between an increase over the high and tropical latitudes and a pronounced decrease over the extratropics, and the geographical distribution of the precipitation changes resemble the cloud cover response. The modeling
355 results for the sea ice show that the Arctic would be ice-free around 40 years earlier, i.e., around 2050 without MPA compared to 2090 with MPA.

The warming rate of the noMPA scenario under the SSP2-4.5 conditions would be very similar to that from the SSP5-8.5 conditions with MPA, i.e., the harsh future without MPA would combine both the negative health effects from the increased UV radiation at the surface and most of the negative consequences of the high-end future climate pathway described in detail
360 by IPCC (2021). Such a combination of fast and strong negative environmental changes would trigger further socioeconomic problems like shortage of food and water security (Barnes et al., 2019), unprecedented migration (Ionesco et al., 2016), civil and international conflicts (e.g., Hsiang et al., 2011), with many of the negative effects starting to emerge already in the current decade.

Our paper reminds us once more how timely and extremely important was the MPA. The Montreal Protocol is one of the
365 few cases in human history when all countries urgently agreed on measures benefiting the survival of humanity. The ozone layer made life on terrestrial land possible (e.g., Ratner and Walker, 1972), and we were very close to severely damaging it. The Montreal Protocol science should continue to be conveyed in the future to remind society how vulnerable and close to a total disaster we could be, and that international dialogue and consensual decisions on environmental questions are possible.

370 *Data availability.* The data were downloaded to a general-purpose open repository Zenodo at
https://doi.org/10.5281/zenodo.7234665 (Egorova et al., 2022) and can also be provided by the corresponding authors upon
request.

375 *Author contributions.* TE wrote the manuscript draft, prepared CFC and VLSL scenarios for the experiments, and
participated in the experiments' performance. TS and JS run the SOCOLv4 experiments. ADK took part in the discussion of
the results. TE, ER, TS, JS, and FZ participated in the visualizing and analysis of the results. All authors participated in
manuscript editing and discussions about the results.

Competing interests. The authors declare that they have no conflict of interest.

380 *Acknowledgments.* We (TE, TS, JS, ER, and AKD) thank the Swiss National Science Foundation for supporting this study
through the № 200020-182239 project POLE (Polar Ozone Layer Evolution). Calculations were supported by a grant from
the Swiss National Supercomputing Center (SCSC) under projects S-901 (ID 154), S-1029 (ID 249), and S-903. Part of the
model development was performed on the ETH Zürich cluster EULER. The work of ER and TS has been partly performed
in the SPbSU "Ozone Layer and Upper Atmosphere Research" laboratory, supported by the Ministry of Science and Higher
Education of the Russian Federation under agreement 075-15-2021-583.-

References

385 Bais, A. F., Lucas, R. M., Bornman, J. F., Williamson, C. E., Sulzberger, B., Austin, A. T., Wilson, S. R., Andraday, A. L.,
Bernhard, G., McKenzie, R. L., Aucamp, P. J., Madronich, S., Neale, R. E., Yazar, S., Young, A. R., de Grijl, F. R., Norval,
M., Takizawa, Y., Barnes, P. W. and Robson, T. M. and Robinson, S. A. and Ballaré, C. L. and Flint, S. D. and Neale, P. J.
and Hylander, S. and Rose, K. C., Wängberg, S.-Å., Häder, D.-P., Worrest, R. C., Zepp, R. G., Paul, N. D., Cory, R. M.,
Solomon, K. R., Longstreth, J., Pandey, K. K., Redhwi, H. H., Torikai, A., and Heikkilä, A. M.: Environmental effects of
ozone depletion, UV radiation and interactions with climate change: UNEP Environmental Effects Assessment Panel, update
390 2017, Photochem. Photobiol. Sci., 17, 127–79, <http://dx.doi.org/10.1039/C7PP90043K>, 2018.

395 Barnes, P. W., Williamson, C. E., Lucas, R. M., Robinson, S. A., Madronich, S., Paul, N. D., Bornman, J. F., Bais, A. F.,
Sulzberger, B., Wilson, S. R., Andraday, A. L., McKenzie, R. L., Neale, P. J., Austin, A. T., Bernhard, G. H., Solomon, K. R.,
Neale, R. E., Young, P. J., Norval, M., Rhodes, L. E., Hylander, S., Rose, K. C., Longstreth, J., Aucamp, P. J., Ballaré, C. L.,
Cory, R. M., Flint, S. D., de Grijl, F. R., Hader, D. -P., Heikkilä, A. M., Jansen, M. A.K., Pandey, K. K., Robson, T.
Matthew., Sinclair, C. A., Wangberg, S., Worrest, R. C., Yazar, S., Young, A. R., and Zepp, R. G.: Ozone depletion,
ultraviolet radiation, climate change and prospects for a sustainable future. Nature Sustainability,
<http://dx.doi.org/10.1038/s41893-019-0314-2>, 2019.

400 Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J.,
Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P.,
Ziemke, J. R., and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer
recovery, Atmos. Chem. Phys., 18, 1379–1394, <https://doi.org/10.5194/acp-18-1379-2018>, 2018.

405 Ball, W. T., Alsing, J., Staehelin, J., Davis, S. M., Froidevaux, L., and Peter, T.: Stratospheric ozone trends for 1985–2018:
sensitivity to recent large variability, Atmos. Chem. Phys., 19, 12731–12748, <https://doi.org/10.5194/acp-19-12731-2019>,
2019.

- 410 Chipperfield, M.P., Dhomse, S.S., Feng, W., McKenzie, R.L., Velders, G.J.M., and Pyle, J.A.: Quantifying the ozone and
ultraviolet benefits already achieved by the Montreal Protocol, *Nat. Commun.*, 6,
7233, <https://doi.org/10.1038/ncomms8233>, 2015.
- 415 Dietmüller, S., Garny, H., Eichinger, R., and Ball, W. T.: Analysis of recent lower-stratospheric ozone trends in chemistry
climate models, *Atmos. Chem. Phys.*, 21, 6811–6837, <https://doi.org/10.5194/acp-21-6811-2021>, 2021.
- Egorova, T. A., Rozanov, E. V., Schlesinger, M. E., Andronova, N. G., Malyshev, S. L., Karol, I. L., and Zubov, V. A.:
Assessment of the effect of the Montreal Protocol on atmospheric ozone, *Geophys. Res. Lett.*, 28, 2389–2392,
<https://doi.org/10.1029/2000GL012523>, 2001.
- 420 Egorova, T., Rozanov, E., Zubov, V., and Karol, I.: Model for investigating ozone trends (MEZON), *Izvestiya Atmos.
Ocean. Phys.*, 39, 277–292, 2003.
- Egorova, T., Rozanov, E., Gröbner, J., Hauser, M., and Schmutz, W.: Montreal Protocol Benefits simulated with CCM
SOCOL, *Atmos. Chem. Phys.*, 13, 3811–3823, <https://doi.org/10.5194/acp-13-3811-2013>, 2013.
- 425 Egorova, T., Sedlacek, J., Sukhodolov, T., Karagodin-Doyennel, A., Zilker, F., and Rozanov, E.: Montreal Protocol's impact
on the ozone layer and climate [data set], <https://doi.org/10.5281/zenodo.7234665>, 2022.
- 430 Engel, A. and Rigby, M. (Lead Authors), Burkholder, J.B., Fernandez, R.P., Froidevaux, L., Hall, B.D., Hossaini, R., Saito,
T., Vollmer, M.K., and Yao, B.: Update on Ozone-Depleting Substances (ODSs) and Other Gases of Interest to the Montreal
Protocol, Chapter 1 in *Scientific Assessment of Ozone Depletion: 2018, Global Ozone Research and Monitoring Project–
Report No. 58*, World Meteorological Organization, Geneva, Switzerland, 2018.
- 435 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled
Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958,
<https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 440 Farman, J. C., Gardiner, B. G., and Schaklin, J. D.: Large losses of the total ozone in Antarctica reveal seasonal ClO_x/NO_x
interaction, *Nature*, 315, 207–210, <http://dx.doi.org/10.1038/315207a0>, 1985.
- Garcia, R. R., Kinnison, D. E., and Marsh, D. R.: “World avoided” simulations with the Whole Atmosphere Community
Climate Model, *J. Geophys. Res.*, 117, D23303, <https://doi.org/10.1029/2012JD018430>, 2012.
- 445 Goyal R., Matthew H England M.H., Gupta A.S., and Jucker M.: Reduction in surface climate change achieved by the 1987
Montreal Protocol, *Environ. Res. Lett.*, 14, 124041, <https://doi.org/10.1088/1748-9326/ab4874>, 2019.
- Hsiang, S., Meng, K., and Cane, M.: Civil conflicts are associated with the global climate, *Nature*, 476, 438–441,
<https://doi.org/10.1038/nature10311>, 2011.
- 450 Hossaini, R., Chipperfield, M. P., Montzka, S. A., Leeson, A. A., Dhomse, S., and Pyle, J. A.: The increasing threat to
stratospheric ozone from dichloromethane, *Nat. Commun.*, 8, 15962, <https://doi.org/10.1038/ncomms15962>, 2017.

Ionesco, D., Mokhnacheva, D., and Gemenne, F.: The Atlas of Environmental Migration (1st ed.). Routledge. <https://doi.org/10.4324/9781315777313>, 2016.

455

Intergovernmental Panel on Climate Change (IPCC), 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.

460

Intergovernmental Panel on Climate Change (IPCC), 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2391 pp., 2021.

465

Intergovernmental Panel on Climate Change (IPCC), 2022: Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Lösschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp., 2022.

470

Junge C.E. and Manson J.E.: Stratospheric aerosol studie., J. Geophys. Res., 66, 2163-2182, <https://doi.org/10.1029/JZ066i007p02163>, 1961.

475

Lee, J.-Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J.P., Engelbrecht, F., Fischer, E., Fyfe, J.C., Jones, C., Maycock, A., Mutemi, J., Ndiaye, O., Panickal, S., and Zhou, T.: Future Global Climate: Scenario-Based Projections and Near-Term Information. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 553–672, <https://doi.org/10.1029/JZ066i007p02163>, 2021.

480

Ionesco, D., Mokhnacheva, D., and Gemenne, F.: The Atlas of Environmental Migration (1st ed.). Routledge. <https://doi.org/10.4324/9781315777313>, 2016.

485

Liang, Y.-Ch., Polvani, L. M., Previdi, M., Smith, K. L., England, M. R., and Chiodo, G.: Stronger Arctic amplification from ozone-depleting substances than from carbon dioxide. Environ. Res. Lett., 17, 024010. <https://dx.doi.org/10.1088/1748-9326/ac4a31>, 2022.

490

Mareev, E. A., and Volodin, E. M.: Variation of the global electric circuit and Ionospheric potential in a general circulation model. Geophys. Res. Lett., 41, 9009–9016, <https://doi.org/10.1002/2014GL062352>, 2014.

495

Maher, N., Milinski, S., Suarez-Gutierrez, L., Botzet, M., Dobrynin, M., Kornbluh, L., Kröger, J., Takano, Y., Ghosh, R., Hedemann, Ch., Li, Chao., Li, H., Manzini, E., Notz, D., Putrasahan, D., Boysen, L., Claussen, M., Ilvina, T., Olonscheck, D., Raddatz, Th., Stevens, B., Marotzke, J.: The Max Planck Institute Grand Ensemble: Enabling the exploration of climate system variability. Journal of Advances in Modeling Earth Systems, 11, 2050–2069. <https://doi.org/10.1029/2019MS001639>, 2019.

Formatted: Justified, Space Before: 0 pt, After: 0 pt

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United States)

- 500 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M.,
Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C.,
Hohenegger, C., Ilyina, T., Jahn, Th., Jimenez-de-la-Cuesta, D., Jungclaus, J., Kleinen, Th., Kloster, S., Kracher, D., Kinne,
S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B.,
Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann,
505 H., Pongratz, J., Popp, M., Raddatz, Th. J., Rast, S., Redler, R., Reick, Ch. H., Rohrschneider, T., Schemann, V., Schmidt,
H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese,
Ph., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model
version 1.2 (MPI-ESM1.2) and its response to increasing CO₂, *J. Adv. Model. Earth Sys.*, 11, 998–1038,
<https://doi.org/10.1029/2018MS001400>, 2019.
- 510 Molina, M. J. and Rowland, F. S.: Stratospheric sink of chlorofluoromethanes: Chlorine atom catalyzed sink of ozone,
Nature, 249, 810-814, <https://doi.org/10.1038/249810a0>, 1974.
- Morgenstern, O., Braesicke, P., Hurwitz, M. M., O'Connor, F. M., Bushell, A. C., Johnson, C. E., and Pyle, J. A.: The world
515 avoided by the Montreal Protocol, *Geophys. Res. Lett.*, 35, L16811, doi:10.1029/2008GL034590, 2008.
- Morgenstern, O., O'Connor, F. M., Johnson, B. T., Zeng, G., Mulcahy, J. P., Williams, J., Teixeira, J., Michou, M., Nabat, P.,
Horowitz, L. W., Naik, V., Sentman, L. T., Deushi, M., Bauer, S. E., Tsigaridis, K., Shindell, D. T., and Kinnison, D. E.:
Reappraisal of the climate impacts of ozone-depleting substances, *Geophys. Res. Lett.*, 47, e2020GL088295,
<https://doi.org/10.1029/2020GL088295>, 2020.
- 520 Neale R.E., Barnes P.W., Robson T.M., Neale, P. J., Williamson, C. E., Zepp, R. G., Wilson, S. R., Madronich, S., Andraday,
A. L.,
Heikkilä, A. M., Bernhard, G. H., Bais, A. F., Aucamp, P. J., Banaszak, A. T., Bornman, J. F., Bruckman, L. S., Byrne, S.
N., Foereid, B., Häder, D.-P., Hollestein, L. M., Hou, W.-C., Hylander, S., Jansen, M. A. K., Klekociuk, A. R., Liley, J. B.:
525 Longstreth, J. Lucas, R. M., Martinez-Abaigar, J., McNeill, K., Olsen, C. M., Pandey, K. K., Rhodes, L. E., Robinson, S. A.,
Rose, K. C., Schikowski, T., Solomon, K. R., Sulzberger, B., Ukpebor, J. E., Wang, Q.-W., Wängberg, S.-Å., White, C. C.,
Yazar, S., Young, A. R.,
Young, P. J., Zhu, L., and Zhu, M.: Environmental effects of stratospheric ozone depletion, UV radiation, and interactions
with climate change: UNEP Environmental Effects Assessment Panel, Update 2020, *Photochem. Photobiol. Sci.*, 20, 1-67,
530 <https://doi.org/10.1007/s43630-020-00001-x> 2021.
- Newman, P. A., Oman, L. D., Douglass, A. R., Fleming, E. L., Frith, S. M., Hurwitz, M. M., Kawa, S. R., Jackman, C. H.,
Krotkov, N. A., Nash, E. R., Nielsen, J. E., Pawson, S., Stolarski, R. S., and Velders, G. J. M.: What would have happened to
535 the ozone layer if chlorofluorocarbons (CFCs) had not been regulated?, *Atmos. Chem. Phys.*, 9, 2113–2128,
<https://doi.org/10.5194/acp-9-2113-2009>, 2009.
- Notz, D., Jahn, A., Holland, M., Hunke, E., Massonnet, F., Stroeve, J., Tremblay, B., and Vancoppenolle, M.: The CMP6
Sea-Ice Model Intercomparison Project (SIMIP): understanding sea ice through climate-model simulations, *Geosci. Model
540 Dev.*, 9, 3427–3446, <https://doi.org/10.5194/gmd-9-3427-2016>, 2016.

Pawson, S., Stolarski, R. S., Douglass, A. R., Newman, P. A., Nielsen, J. E., Frith, S. M., and Gupta, M. L.: Goddard earth observing system chemistry-climate model simulations of stratospheric ozone-temperature coupling between 1950 and 2005, *J. Geophys. Res. Atmos.*, 113, 1–16, <https://doi.org/10.1029/2007JD009511>, 2008.

545 Polvani, L.M., Previdi, M., England, M.R., Chiodo, G., and Smith, K. L.: Substantial twentieth-century Arctic warming caused by ozone-depleting substances, *Nat. Clim. Chang.*, 10, 130–133, <https://doi.org/10.1038/s41558-019-0677-4>, 2020.

[Previdi, M., Smith, K.L., and Polvani, L.M.: Arctic amplification of climate change: a review of underlying mechanisms, *Env.Res.Lett.*, 16 093003, <https://doi.org/10.1088/1748-9326/ac1c29>, 2021.](#)

Formatted: Left, Space Before: Auto, After: Auto

550 Prather, M., Midgley, P., Sherwood R., F., and Stolarski, R.: The ozone layer: The road not taken, *Nature*, 381, 551–554, <https://doi.org/10.1038/381551a0>, 1996.

555 Ratner, M. I. and Walker, J. C. G.: Atmospheric Ozone and the History of Life, *J. Atmos. Sci.*, 29, 803–808, [https://doi.org/10.1175/1520-0469\(1972\)029%3C0803:AOATHO%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1972)029%3C0803:AOATHO%3E2.0.CO;2), 1972.

Revell, L. E., Bodeker, G. E., Huck, P. E., Williamson, B. E., and Rozanov, E.: The sensitivity of stratospheric ozone changes through the 21st century to N₂O and CH₄, *Atmos. Chem. Phys.*, 12, 11309–11317, <https://doi.org/10.5194/acp-12-11309-2012>, 2012.

560 Revell, L. E., Tummon, F., Stenke, A., Sukhodolov, T., Coulon, A., Rozanov, E., Garny, H., Grewe, V., and Peter, T.: Drivers of the tropospheric ozone budget throughout the 21st century under the medium-high climate scenario RCP 6.0, *Atmos. Chem. Phys.*, 15, 5887–5902, <https://doi.org/10.5194/acp-15-5887-2015>, 2015.

565 Rowland, F.S. Stratospheric ozone depletion. *Phil.Trans.R.Soc. B*, 361,769-790, <https://doi.org/10.1098/rstb.2005.1783>, 2006.

570 Sheng, J.-X., Weisenstein, D. K., Luo, B.-P., Rozanov, E., Stenke, A., Anet, J., Bingemer, H., and Peter T.: Global atmospheric sulfur budget under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *J. Geophys. Res.-Atmos.*, 120, 256–276, <https://doi.org/10.1002/2014JD021985>, 2015.

575 Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, <https://doi.org/10.1029/1999RG900008>, 1999.

Stolarski, R., Krueger, A., Schoeberl, McPeters, R. D., Newman, P. A., and Alpert, J. C.: Nimbus 7 satellite measurements of the springtime Antarctic ozone decrease, *Nature* 322, 808–811, <https://doi.org/10.1038/322808a0>, 1986

580 Sukhodolov, T., Egorova, T., Stenke, A., Ball, W. T., Brodowsky, C., Chiodo, G., Feinberg, A., Friedel, M., Karagodin-Doyennel, A., Peter, T., Sedlacek, J., Vattioni, S., and Rozanov, E.: Atmosphere–ocean–aerosol–chemistry–climate model SOCOLv4.0: description and evaluation, *Geosci. Model Dev.*, 14, 5525–5560, <https://doi.org/10.5194/gmd-14-5525-2021>, 2021.

Velders, G. J. M., Andersen, S. O., Daniel, J. S., Fahey, D. W., and McFarland, M.: The importance of the Montreal Protocol in protecting climate, *Proc. Natl. Acad. Sci.*, 104, 4814–9, <https://doi.org/10.1073/pnas.0610328104>, 2007.

585

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2018, Global Ozone Research and Monitoring Project–Report No. 58, 588 pp., Geneva, Switzerland, 2018.

WMO (World Meteorological Organization). Executive Summary. Scientific Assessment of Ozone Depletion: 2022, GAW Report No. 278, 56 pp.; WMO: Geneva, 2022.

590

Zubov, V., Rozanov, E., Egorova, T., Karol, I., and Schmutz, W.: Role of external factors in the evolution of the ozone layer and stratospheric circulation in 21st century, *Atmos. Chem. Phys.*, 13, 4697–4706, <https://doi.org/10.5194/acp-13-4697-2013>, 2013.

595 Young, P.J., Harper, A.B., Huntingford, C., Paul, N.D., Morgenstern, O., Newman, P.A., Oman, L.D., Madronich, S., and Garcia, R.R.: The Montreal Protocol protects the terrestrial carbon sink, *Nature*, 596, 384–388, <https://doi.org/10.1038/s41586-021-03737-3>, 2021.