

Supplement

Journal: Atmospheric Chemistry and Physics

Title: High-resolution regional emission inventory contributes to the evaluation of policy effectiveness: A case study in Jiangsu province, China

Authors: Chen Gu, Lei Zhang, Zidie Xu, Sijia Xia, Yutong Wang, Li Li, Zeren Wang, Qiuyue Zhao, Hanying Wang, Yu Zhao

Table List

Table S1 Source categories of the emission inventory for Jiangsu Province, China.

Table S2 Comparisons between observed and simulated meteorological parameters in D3.

Table S3 Annual air pollutant emissions in Jiangsu from 2015 to 2019 ($\text{Gg}\cdot\text{yr}^{-1}$).

Table S4 The emission data of SO_2 , NO_x , AVOCs, $\text{PM}_{2.5}$, and NH_3 by year and city for 2015 and 2019 ($\text{Gg}\cdot\text{yr}^{-1}$).

Table S5 Comparison of observed and simulated hourly $\text{PM}_{2.5}$ concentrations by month in Jiangsu. In total, 110 state-operated observation sites were included in the comparison.

Table S6 The same as Table S5 but for MDA8 O_3 .

Figure List

Figure S1 The modeling domain (a) and the locations of Jiangsu province (with its 13 cities) and state-operating ground stations of air quality observation (b). The map data provided by Resource and Environment Data Cloud Platform are freely available for academic use (<http://www.resdc.cn/data.aspx?DATAID=201>), © Institute of

Geographic Sciences & Natural Resources Research, Chinese Academy of Sciences.

Figure S2 Summary of major air pollution control measures taken between 2015 and 2019. De-S, desulfurization; De-N, NO_x control; NG, natural gas.

Figure S3 The locations of key industries in Jiangsu. The blue line indicates the Yangtze River.

Figure S4 Spatial distribution of air pollutant emissions in 2019 at a horizontal resolution of 3km. The species include (a) SO₂, (b) NO_x, (c) CO, (d) AVOCs, (e) NH₃, (f) PM₁₀, (g) PM_{2.5}, (h) BC, and (i) OC.

Figure S5 The spatial distribution of BVOCs emissions from 2015 to 2019 at a horizontal resolution of 3 km.

Figure S6 Annual emissions reductions of SO₂, NO_x, AVOCs, NH₃, and PM_{2.5} during 2015-2017 and 2017-2019.

Figure S7 Average pollutant removal efficiencies in the key industries including power, steel and cement production.

Figure S8 Spatial distribution of MDA8 O₃ from CMAQ simulation in January, April, July and October from 2015 to 2019 at a horizontal resolution of 3km (μg·m⁻³).

Figure S9 The same as Figure S8 but for PM_{2.5}.

Table S1 Source categories of the emission inventory for Jiangsu Province, China.

Sector (1st level)	Subsector(2nd level)	Fuel/product/process(3rd level)
Power sector	Power plant	Coal (8 categories)/ Fuel oil/ Natural gas/other 8 categories
Industry	Boilers	Coal (8 categories)/Fuel oil/Diesel/Natural gas/Biomass
	Mining	Coal mining Oil and gas extraction Metal mining Non-metallic mining
	Petroleum refining	Process devices Equipment leak Storage tanks Loading and unloading Flares Wastewater treatment Cooling Tower Petrochemical furnace
	Coking	Coke production
	Chemical manufacturing	Organic chemical manufacturing (18 categories) Inorganic chemical manufacturing (10 categories) Fertilizer (7 categories) Pesticide (8 categories) Coating production (16 categories) Ink production Synthetic resin/ Synthetic rubber (6 categories) Synthetic fiber monomer

	Specialty chemicals
	Household chemicals (11 categories)
	Chemical fiber (6 categories)
Rubber and plastic manufacturing	Rubber products/ Plastic products (30 categories)
Non-metallic mineral manufacturing	Cement manufacturing/ Lime manufacturing/ Plaster manufacturing
	Commodity concrete/ Brick and tile products/ Crushed Stone processing
	Abrasive processing/ Asphalt products /
	Glass products/ Glass fiber/ Gypsum board
	Ceramic products/ Refractory products
Ferrous metal manufacturing	Raw material yard
	Sintering
	Pellet
	Blast furnace
	Converter
	Electric furnace
	Casing steel
	Rolling steel
	Ferroalloy production
Non-ferrous metal manufacturing	Primary copper/ Primary aluminum/ Primary lead
	Primary zinc/ Secondary copper/ Secondary aluminum
Food manufacturing	
Tobacco manufacturing	
Wood processing	
Papermaking	Pulp manufacturing
Agricultural products processing	
Beverage manufacturing	Alcohol/ Liquor/ Red wine/ Beer

	Textile	
	Leather tanning	
	Furniture manufacturing	
	Package and printing	
	Pharmaceutical manufacturing	
	Shoemaking	
	Metal products	
	Machinery equipment manufacturing	
	Railway equipment	
	Auto manufacturing	
	Shipbuilding	
	Appliance manufacturing	
	Electronic equipment	
	Other industry	
Transportation	Motor vehicles	Light-duty gasoline vehicles
		Light-duty diesel vehicles
		Heavy-duty gasoline vehicles
		Heavy-duty diesel vehicles
		Heavy-duty gasoline truck
		Heavy-duty diesel truck
		Taxi/Bus/ Motorcycles
	Non-road machinery	Agricultural machinery
		Construction machinery
		Garden equipment
		Ground handling equipment
		Factory machinery

		Port machinery
Residential	Marine	
	Aviation aircraft	
	Residential combustion	
	Architectural coating	
	Housing retrofitting emissions	
	Ship fittings and repairs	
	Consumer products	
	Solvent degreasing	
	Auto repair	
	Dry cleaning	
	Hospital solvent	
	Catering	
	Sewage treatment	
Agriculture	Waste treatment	Landfill/Composting
	Oil depot	Crude oil/Gasoline/Diesel/Heavy oil/ Chemicals
	Gas station	
	Livestock	Cow/Horse/Donkey/Mule/Pig/Sheep/Rabbit/Poultry
	Nitrogen fertilizer application	Fertilizer-N/Fertilizer-P/Fertilizer-K
	Soil background	
	Nitrogen fixing plants	
	Biofuel	Rice straw/Wheat/Corn/Bean/Tuber/Cotton
	Human excrement	
	Biogenic source	Biogenic volatile organic compounds emissions

Table S2 Comparisons between observed and simulated meteorological parameters in D3 (OBS and SIM represent observation and simulation, respectively).

Meteorological parameters	Evaluation Indicators	2015	2016	2017	2018	2019	Recommended range
WS10	OBS (m/s)	2.91	2.63	2.88	2.37	2.55	
	SIM (m/s)	2.92	2.67	2.89	2.63	2.81	
	Bias (m/s)	0.01	0.04	0.04	0.27	0.26	$\leq \pm 0.5$
	RMSE (m/s)	0.27	0.29	0.33	0.27	0.3	≤ 2.0
	IOA	0.82	0.72	0.94	0.87	0.85	≥ 0.6
	R	0.89	0.82	0.82	0.89	0.75	
WD10	OBS (°)	184.29	163.29	169.28	174.34	151.75	
	SIM (°)	165.37	147.09	159.80	155.66	143.8	
	Bias	-18.92	-16.2	-9.48	-18.68	-7.95	$\leq \pm 10$
	R	0.69	0.74	0.8	0.71	0.66	
T2	OBS (°C)	15.96	16.02	16.54	16.85	15.78	
	SIM (°C)	16.19	15.73	16.33	16.35	15.45	
	Bias (°C)	0.23	-0.29	-0.21	-0.5	-0.33	$\leq \pm 0.5$
	RMSE (°C)	1.76	0.95	1.27	0.69	1.36	≤ 2.0
	IOA	0.95	0.98	0.95	0.99	0.91	≥ 0.8
	R	0.94	0.91	0.89	0.96	0.88	
RH2	OBS (%)	65.16	75.69	79.81	71.96	68.35	
	SIM (%)	61.87	76.21	82.24	70.80	63.91	
	Bias (%)	-3.29	0.52	2.43	-1.16	-4.44	
	RMSE (%)	6.61	3.86	4.24	4.07	5.18	
	IOA	0.97	0.98	0.95	0.98	0.83	≥ 0.7
	R	0.92	0.85	0.89	0.94	0.87	

Table S3 Annual air pollutant emissions in Jiangsu from 2015 to 2019 (Gg·yr⁻¹).

Year	SO ₂	NO _x	NMVOCs	NH ₃	CO	PM ₁₀	PM _{2.5}	BC	OC
Power	222	351	29	0	607	71	46	0	0
Industry	325	465	854	14	4974	481	300	15	7
Transportation	35	546	195	6	1132	31	29	12	4
Residential	45	49	270	15	1022	128	116	7	33
Agriculture	0	0	0	433	0	0	0	0	0
Biogenic source	0	0	150	0	0	0	0	0	0
2015 Total	627	1411	1498	468	7735	711	491	34	44
Power	191	282	28	0	549	51	35	0	0
Industry	305	450	838	14	4623	477	295	14	10
Transportation	39	613	208	7	1207	33	32	13	4
Residential	45	46	272	16	1018	126	113	7	29
Agriculture	0	0	0	415	0	0	0	0	0
Biogenic source	0	0	188	0	0	0	0	0	0
2016 Total	580	1391	1535	452	7397	687	475	34	43
Power	123	263	26	0	493	54	36	0	0
Industry	207	359	837	14	4527	465	288	12	9
Transportation	41	664	209	8	1272	35	33	14	5
Residential	45	45	271	20	1013	122	111	7	27
Agriculture	0	0	0	392	0	0	0	0	0
Biogenic source	0	0	213	0	0	0	0	0	0
2017 Total	416	1331	1556	434	7305	676	468	33	41
Power	128	251	26	0	490	51	32	0	0
Industry	158	184	848	14	4452	455	282	11	9
Transportation	43	717	163	7	1323	44	42	15	5
Residential	45	46	269	19	987	120	106	7	26
Agriculture	0	0	0	390	0	0	0	0	0
Biogenic source	0	0	204	0	0	0	0	0	0
2018 Total	374	1198	1510	430	7252	670	462	33	40
Power	80	186	26	0	484	43	28	0	0
Industry	127	137	839	14	4349	374	240	10	7
Transportation	44	754	151	8	1356	45	43	15	5
Residential	45	45	255	17	974	103	100	7	24
Agriculture	0	0	0	383	0	0	0	0	0
Biogenic source	0	0	193	0	0	0	0	0	0
2019 Total	296	1122	1464	422	7163	565	411	32	36

Table S4 The emission data of SO₂, NO_x, AVOCs, PM_{2.5}, and NH₃ by year and city for 2015 and 2019 (Gg·yr⁻¹).

	SO ₂	NO _x	PM _{2.5}	AVOCs	NH ₃
Southern cities	334.5	684.1	229.1	602.8	81.4
Nanjing	58.5	192.7	63.7	94.4	17.2
Suzhou	150.8	205.5	55.0	237.2	25.1
Wuxi	62.9	107.2	36.4	108.6	13.9
Changzhou	39.0	111.9	48.4	92.0	13.7
Zhenjiang	23.3	66.9	25.6	70.7	11.5
Northern cities	186.4	463.5	167.1	462.8	279.9
Xuzhou	73.1	131.4	50.6	92.3	74.3
Lianyungang	24.9	80.9	29.9	86.4	39.3
Suqian	18.4	52.8	18.8	80.1	39.3
Huaian	28.5	100.9	28.2	103.4	39.7
Yancheng	41.4	97.5	39.7	100.7	87.3
Central cities	106.7	264.5	94.5	282.3	107.2
Yangzhou	24.9	68.1	23.2	66.1	21.8
Taizhou	32.5	69.1	37.3	99.7	27.2
Nantong	49.3	127.3	34.0	116.4	58.3
2015 Total	627.6	1412.1	490.7	1347.9	468.5
Southern cities	136.3	524.5	174.9	665.0	73.2
Nanjing	28.6	135.3	47.7	119.6	18.3
Suzhou	43.7	145.4	39.4	284.3	19.8
Wuxi	32.0	109.1	31.0	128.0	13.9
Changzhou	23.7	70.9	35.8	83.8	12.2
Zhenjiang	9.4	63.7	21.0	49.4	9.0
Northern cities	87.0	379.7	154.0	335.9	253.9
Xuzhou	22.4	85.3	47.1	78.0	65.0
Lianyungang	22.1	87.1	24.5	77.7	34.9
Suqian	11.5	56.9	18.0	49.4	37.4
Huaian	18.2	78.6	24.5	62.2	37.9
Yancheng	12.7	71.8	40.0	68.7	78.7
Central cities	71.5	217.9	82.4	270.2	95.3
Yangzhou	21.4	52.0	16.1	74.5	22.5
Taizhou	20.4	48.8	36.3	92.0	25.9
Nantong	29.7	117.1	30.0	103.7	46.9
2019 Total	295.8	1122.2	411.4	1271.1	422.5

Table S5 Comparison of observed and simulated hourly PM_{2.5} concentrations by month in Jiangsu. In total, 110 state-operated observation sites were included in the comparison.

	Observation Simulation ($\mu\text{g}\cdot\text{m}^{-3}$)		R		NMB (%)		NME (%)		
	($\mu\text{g}\cdot\text{m}^{-3}$)	This study ^a	MEIC ^b	This study	MEIC	This study	MEIC	This study	MEIC
2015/1	97.06	80.29	72.11	0.40	0.25	-10.47	-25.69	48.07	53.10
2015/4	54.31	42.42	41.14	0.26	0.24	-17.28	-19.79	49.83	48.84
2015/7	39.36	32.98	30.09	0.37	0.34	-8.58	-17.23	59.96	58.15
2015/10	56.55	42.33	39.66	0.44	0.29	-14.53	-30.27	50.15	52.58
2016/1	81.87	71.80	64.32	0.41	0.30	-12.29	-17.25	43.64	43.80
2016/4	50.42	49.03	41.34	0.24	0.20	-2.77	-15.33	56.48	49.15
2016/7	30.60	34.23	34.80	0.24	0.19	11.86	16.81	49.25	57.36
2016/10	28.93	26.44	24.33	0.31	0.30	-8.62	-14.65	36.79	51.98
2017/1	66.51	64.65	50.68	0.33	0.35	-19.71	-21.43	49.46	58.15
2017/4	49.09	46.59	40.81	0.27	0.15	-17.32	-15.39	54.50	58.69
2017/7	29.87	33.80	33.45	0.25	0.20	-9.13	11.57	28.82	35.89
2017/10	34.48	36.77	29.40	0.45	0.42	-7.29	-15.09	37.44	51.90
2018/1	83.95	61.24	53.66	0.57	0.40	-27.05	-33.49	49.50	58.57
2018/4	47.93	37.86	41.70	0.28	0.10	-23.67	-12.99	54.85	56.16
2018/7	24.42	23.03	28.63	0.37	0.24	-5.71	17.50	56.55	28.66
2018/10	38.97	20.01	11.16	0.20	0.17	-28.67	-43.36	41.21	43.34
2019/1	72.29	77.79	64.55	0.29	0.10	7.61	-11.53	38.72	52.74
2019/4	40.32	49.62	43.96	0.27	0.20	23.07	7.87	45.58	54.51
2019/7	25.20	26.86	27.35	0.19	0.10	6.58	21.45	42.08	53.40
2019/10	38.40	20.63	16.54	0.20	0.12	-29.32	-38.15	49.43	52.46

This study^a: simulation with a horizontal resolution of 3 km (D3).

MEIC^b: simulation with a horizontal resolution of 9 km (D2).

Table S6 The same as Table S5 but for MDA8 O₃.

	Observation Simulation ($\mu\text{g}\cdot\text{m}^{-3}$)			R		NMB (%)		NME (%)	
	($\mu\text{g}\cdot\text{m}^{-3}$)	This study ^a	MEIC ^b	This study	MEIC	This study	MEIC	This study	MEIC
2015/1	50.93	35.83	39.54	0.25	0.17	-29.64	-25.20	47.13	42.58
2015/4	108.70	82.61	79.20	0.37	0.26	-24.01	-27.14	36.43	41.89
2015/7	107.18	85.58	86.53	0.48	0.41	-20.15	-19.85	36.74	41.82
2015/10	114.98	78.27	78.27	0.37	0.37	-31.93	-31.93	37.34	37.34
2016/1	60.19	40.62	38.52	0.29	0.32	-32.52	-36.00	41.43	42.69
2016/4	117.37	85.40	85.10	0.49	0.47	-27.24	-27.49	37.23	39.03
2016/7	118.38	95.91	110.76	0.58	0.50	-11.10	-6.44	31.68	31.61
2016/10	74.36	67.71	64.41	0.51	0.45	-8.94	-13.50	30.37	31.39
2017/1	66.37	44.2	42.70	0.43	0.44	-29.39	-35.67	38.71	41.95
2017/4	134.06	103.52	98.58	0.57	0.60	-17.48	-26.47	26.68	31.23
2017/7	134.59	116.08	145.43	0.44	0.37	-11.86	8.05	31.58	35.81
2017/10	91.28	85.92	81.97	0.57	0.37	-3.69	-10.20	24.93	26.75
2018/1	57.95	52.61	52.32	0.12	0.26	-9.21	42.04	38.74	66.28
2018/4	134.87	106.68	85.74	0.67	0.29	-20.90	-36.00	27.36	42.66
2018/7	118.01	117.53	114.78	0.45	0.43	-0.41	3.26	32.50	35.30
2018/10	108.75	79.42	70.85	0.33	0.31	-26.97	-34.66	34.99	39.03
2019/1	50.48	48.43	43.41	0.28	0.20	-9.99	-13.99	44.02	57.07
2019/4	112.93	101.95	94.46	0.50	0.41	-21.24	-16.35	31.71	39.57
2019/7	138.71	124.22	118.45	0.33	0.31	-12.65	-16.70	43.33	34.17
2019/10	109.55	88.26	80.72	0.28	0.20	-21.57	-24.10	43.93	35.89

This study^a: simulation with a horizontal resolution of 3 km (D3).

MEIC^b: simulation with a horizontal resolution of 9 km (D2).

Figure S1




Figure S2

		2015	2016	2017	2018	2019
Ultra-low emission of coal-fired power plants		33% meet "ultralow emission" standard	84% meet "ultralow emission" standard	100% meet "ultralow emission" standard		
Extensive management of coal-fired boilers		100 MW of coal-fired power generation capacity were phased out Eliminated non-heating coal-fired power units (< 300 MW)		Remediation of coal-fired boilers (< 10 steam tons/hour)		
Upgrade and renovation of non-electrical industry	Iron and steel	80% with high efficiency de-S devices		60% meet "ultralow emission" standard		
	Cement	89% with high efficiency de-N devices		Ultra-low emission transformation of 22 cement clinker enterprises		
	Flat glass	40% & 17% equipped with de-S and de-N devices		30% equipped with de-N devices		
Phase out outdated industrial capacities	Iron and steel	Eliminated 8 million ton outdated capacity		Eliminated 18 million ton outdated capacity Eliminated 50% of outdated coking capacity		
	Cement	Eliminated 9 million ton outdated capacity		Eliminated 8 million ton outdated capacity		
	Flat glass	Eliminated 3 million weight boxes outdated capacity		Eliminated 8 million weight boxes outdated capacity		
Phase out small polluting factories		1,400 petrochemical industry enterprises were phased out or rectified		1750 enterprises were phased out or rectified		
Construction of port shore power		15% equipped with shore power in port Reach 40% electric coverage (Oil to Electric)		30% equipped with shore power in port Reach 60% electric coverage		
Comprehensive treatment of mobile source pollution	Emission standard	"China V" standard (on road)		"China VI" standard (on road) > "China II" standard (non-on road)		
	Old cars	Eliminated 0.7 million old vehicles		Eliminated 30,000 old trucks (< China III standard)		
	New Energy cars	Promote 50,000 new energy vehicles		Promote 150,000 new energy vehicles Replacement 80% of buses with new energy		
VOCs emission control in key sectors		17% of the industry enterprises treat the waste gas in the process	32% of the industry enterprises treat the waste gas in the process; Average end-of-pipe treatment efficiency of 30%			
Oil and gas recovery		90% of gas stations, tanker trucks, and tank farms have achieved oil and gas recovery treatment		Average oil and gas recovery efficiency reached 95%		
Replacement with low-VOC paints		Switch to water-based paints with low VOCs content in 7 key industries		20% reduction in the use of highly reactive solvents and auxiliary products		
Application of leak detection and repair (LDAR)		Implement LDAR in all chemical parks				
Control of non-point pollution		Comprehensive utilization rate of straw reached 95% Fertilizer utilization rate reached 40% Fume management for restaurants in key areas				
Promote clean energy		Applied washed clean coal Coal substituted by NG and electricity in 30,000 households		NG has covered 70% of the province		

List of Policies and Clean Air Actions

Air Pollution Remediation Work Plan for Coal-fired Boilers in Jiangsu, (Department of ecological environment of Jiangsu, 2013)
 VOCs Pollution Control Special Action Implementation Plan, (Department of ecological environment of Jiangsu, 2015)
 Action Plan for Energy Saving, Emission Reduction and Renovation of Coal Power Units in Jiangsu (2014-2020), (The People's Government of Jiangsu, 2014)
 The Implementation plan for Ultra-low Emission Transformation of Iron and Steel Enterprises in Jiangsu (2018), (Department of ecological environment of Jiangsu, 2018)

Figure S3




Figure S4






Figure S5


2015




2016




2017



tons·grid⁻¹



2018



2019




Figure S6




Figure S7




Figure S8





Figure S9



References

Department of ecological environment of Jiangsu, Notice of the Jiangsu Provincial Government on issuing the air pollution prevention and control action plan.

http://www.jiangsu.gov.cn/art/2014/1/6/art_46690_2586827.html.

Department of ecological environment of Jiangsu, Air Pollution Remediation Work Plan for Coal-fired Boilers in Jiangsu. <http://sthjt.jiangsu.gov.cn>.

Department of ecological environment of Jiangsu, VOCs Pollution Control Special Action Implementation Plan in Jiangsu.

http://www.jiangsu.gov.cn/art/2018/1/25/art_46143_7431312.html.

Department of ecological environment of Jiangsu, The Implementation plan for Ultra-low Emission Transformation of Iron and Steel Enterprises in Jiangsu (2018).

http://sthjt.jiangsu.gov.cn/art/2019/5/22/art_83589_10046297.html.

Ministry of Ecology and Environment of the People's Republic of China, 2020 Report on the state of the ecology and environment in China.

<http://english.mee.gov.cn/Resources/Reports/soe/SOEE2019/202204/P020220407417638702591.pdf>.

The People's Government of Jiangsu, Notice of the Jiangsu Provincial Government on issuing the Three-year Action Plan to fight Air Pollution.

http://www.jiangsu.gov.cn/art/2018/10/17/art_64797_7842355.html.

The People's Government of Jiangsu, Action Plan for Energy Saving, Emission Reduction and Renovation of Coal Power Units in Jiangsu (2014-2020).

http://www.jiangsu.gov.cn/art/2014/11/10/art_46144_2545297.html.