

Answers to the review for our manuscript “Ice Nucleating Particles in Northern Greenland: annual cycles, biological contribution and parameterizations”

We very much thank the reviewer for helping to advance our manuscript by doing this review. Below, find the original comments in black, [our answers in blue](#) and new text added to the manuscript in “*blue, italic and quotation marks*”. All line numbers refer to the new version with tracked changes.

---

## Major comments

- Parameterization: Which fraction of your sample is within a certain factor of the developed parameterization? E.g., are you able to predict the INP concentration accurately within a factor of 10, as deviations larger than this value can impact cloud microphysical and radiative properties (Phillips et al., 2003)?

Variations in INP concentrations may indeed influence cloud glaciation, particularly when ice multiplication processes are assumed, as in the simulations in the study you refer to here. However, both, ice multiplication processes and with this the relevance of a change in INP concentration (relative or absolute) are still a matter of scientific research today and not well constrained. As such, the influence of e.g., a change of a factor of 10 on cloud glaciation is uncertain. Still, we added some text and a related table to the study, and added a plot to the SI, Fig. S12. (We replaced the appendix by an SI, please check for the figure there.) The text is as follows (line 383 ff) (for the table, please refer to the revised version of the manuscript):

*“We furthermore examined which fraction of the corresponding samples falls within a range covering one order of magnitude above and below our parameterizations. This fraction is generally well above 90%, as shown at three selected temperatures of  $-12$ ,  $-16$  and  $-20^{\circ}\text{C}$  in Table 3 (for more see SI, Fig. S12), emphasizing the representativeness of our parameterizations.”*

- Case study: The linkage of an observed higher cloud ice fraction to an increased INP concentration in April 2020 as compared to the previous year is insufficient. First, to my knowledge, there is no evidence that ground-based INP concentrations can impact cloud properties observed on top of the cloud. Second, other parameters differ between April 2019 and 2020, for example, mean wind speed (potential impact from blowing snow) or a lower surface temperature in April 2019 (e.g., a greater impact of pre-activation of INPs; Conen et al., 2015). Moreover, it would be interesting to investigate if there was an enhanced impact from glacial dust sources or a higher biological activity in the ocean. In addition, in the late winter months, the Arctic haze phenomena (e.g., Shaw, 1995) can impact aerosol populations, which is not discussed here.

The impression that this part may be insufficient may arise from the fact that we were not able to pin down INP sources. Still, we did examine a broad range of meteorological data as well as

back-trajectories. It is also valuable information to the research community to see what did not work out. Also, past studies show a connection from ground based INP measurements to glaciation of clouds under coupling conditions (Griesche et al., 2021) and to the boundary layer in general (Gong et al., 2022), connecting ground based and lidar derived data. Further examinations are for sure needed, but this is not the topic of our study.

As for parameters differing between April 2019 and April 2020, none of the parameters differs throughout the whole month (the parameters you mentioned differ for the second half of the month, but not for the first half), while INP concentrations are higher throughout the month. So no conclusions can be drawn from that.

We agree with the reviewer that an examination of additional factors such as glacial dust, blowing snow or oceanic activity would be interesting. But this is far beyond the scope of this study. We did examine the effect of biological activity in the ocean, and did not see an effect on a first glance, which lead us to drop this and not include it. Arctic haze can be excluded as a source for INP in the here examined temperature range, as we have described previously (Hartmann et al., 2019; Wex et al., 2019). On that, we added in the introduction (line 89/90):

*“in line with the observation that Arctic haze (Wex et al., 2019; Hartmann et al., 2019), or anthropogenic pollution in general (Chen et al., 2018; Tarn et al., 2018; Welti et al., 2020), do not contribute INP that are ice active in the temperature range down to  $\approx -30^{\circ}\text{C}$ .”*

Blowing snow is now mentioned in the text, as we elaborate on in our answer to your second minor comment.

As our data will be available on Pangaea, maybe you or others can pick up the thread and add additional examinations to this two year long data-set.

## **Minor comments**

- What is the relevance of mixed-phase clouds in the Arctic winter regarding the radiative forcing?

We do not see what this has to do with the topic of our study and hence did not elaborate on this in order to not lengthen the text unnecessarily.

- Is there an impact of blowing snow on the aerosol filters?

For the atmosphere, INP may be resuspended through blowing snow. However, due to the use of a PM10 inlet for the low volume aerosol sampler, snow will not have been collected. Still, we added a comment on blowing snow (line 456 ff):

*“Blowing snow occurring in the Arctic (Yang et al., 2019) may contribute to atmospheric INP. However, in our case, due to the use of a PM<sub>10</sub> inlet during filter sampling, we do*

*not expect that snow was collected in considerable amounts and hence we do not expect an influence of blowing snow on our measured INP concentrations.”*

- In some cases, the mentioned publications are examples and do not represent all existing literature. Please check and make use of "e.g." in such cases or complete the cited literature

You are right: certainly not all publications we could have cited on all the mentioned topics are included. But your request would mean to add an “e.g.” to quasi all citations, or at least to those where we added citations concerning more general statements. This would flood particularly the introduction with “e.g.” which reduces the readability.

We would want to leave it up to the editor to decide if we should really add this. For the time being, nothing was changed.

- You mention that blank filters were collected weekly, but do not present them here. How high were the INP concentrations from those filters and did you consider using them for a background correction of the INP concentration?

Thanks for pointing this out. We now added respective figures and some explanatory text in the supplemental information (SI, Fig. S4 and corresponding text). (As said before: We replaced the appendix by an SI.) We refer to them in the main text at the end of section 2.2.5: *“Regarding the blank filters, measured  $f_{ice}$  were generally clearly below the respective values of the atmospheric samples and a background subtraction was not done (for details see SI, section S5).”*

Text in the SI:

Figure caption:

*“Measured  $f_{ice}$  for blank filters and respective atmospheric samples for LINA (left panels) and INDA (right panels) for the month of January (upper panels) and July (lower panels). Blank filter data are always shown in black, atmospheric sample data in gray, except for INDA data for July (panel C, lower left side), for which data for the blank filters with the highest values are given in blue and the corresponding filters in cyan.”*

Main text:

*“Fig. S4 exemplarily shows measured frozen fractions ( $f_{ice}$ ) for blank filters and respective atmospheric samples. Data are given for both measurement devices LINA and INDA (see main text for more details), and for all samples collected in the months of January and July. Data are shown separately for the two months, as blank filter values are generally higher in summer than in winter, an observation which was made before (Wex et al., 2019). Even within July, blank filters with the highest values correspond to atmospheric samples with higher concentrations.*

*The blank filter values were clearly below the values from the atmospheric samples. It also can be seen that the temperature range covered by the data from the blank filters is at lower temperatures than that of the measurements. Therefore, correcting the measurements with background data would only be partially possible. Also, a correction is done based on concentrations (Vali et al., 2019), i.e., using the logarithms of  $f_{ice}$ , such that corrections for the data used in this study are small and no influence on the outcome of our overall results is observed. Therefore, a background subtraction was not done.”*

- Lines 35 – 36: Reference for this statement missing.

The sentence was adjusted slightly as given below, and the following references have been added.

*“Also, exposure of non snow-covered surfaces to the atmosphere likely will enhance emission of soil dust, and therewith could largely contribute to ice-nucleating particles (INPs) (Šantl-Temkiv et al., 2019; Tobo et al., 2019; Sanches-Marroquin et al., 2020).”*

- Lines 94 – 97: It might be worth mentioning that also the impact of glacial dust is increasing due to retreating glaciers (e.g., AMAP, 2021).

We adjusted the text and added the citation as follows:

*“... where snow and sea ice cover are expected to decline in the upcoming years, glacial dust can also increase due to the retreating glaciers (AMAP, 2021), which will ... “*

- Lines 125: What is the pore size of the polycarbonate filters?

The pore size was 200 nm. Information was added to the text.

- Line 152: What is the uncertainty in temperature regarding the 6 second time resolution of the camera and using a 1 °C/min cooling rate?

A picture is taken every 0.1 K, but that is independent of the temperature uncertainty. Both LINA and INDA are regularly calibrated, and this is accounted for in the data. The temperature uncertainty per se is 0.5 K, determined from these regularly executed calibration measurements, for both LINA and INDA. Information was added to the text.

- Line 253: Can you quantify the variability in INP concentration better, e.g., using the standard deviation?

We already quantified the variability of  $N_{INP}$  by giving the span between the 10% and the 90% percentile (based on values given in Table 1). Therefore, we do not think it is necessary to add also standard deviation. However, in the respective text we made it clearer that the "1 to 2 orders of magnitude" mentioned in the respective sentence refer to these percentile values. Additionally, a better quantification is now also given with respect to our answer to your first major comment.

- Line 277: Statement „Their respective INP parameterizations are often used in atmospheric models“ needs a reference.

We added DeMott et al. (2010) and Curry & K Khvorostyanov (2012).

- Lines 333 – 334: Are there publications that can strengthen this statement („A common background of mineral dust particles throughout the year may exist“)?

Mineral dust as a contribution to the atmosphere in the Arctic has been described. We carefully expressed this statement by using “may”, and now added the following citations: “(e.g., Si et al., 2019; Tobo et al., 2019; Sanches-Marroquin et al., 2020)”

- Lines 354 – 355: It might be worth explaining on which measurements the parameterizations from Cooper (1986) and Fletcher (1962) are based to understand the difference of three orders of magnitudes as compared to your parameterization.

Concerning the data by Cooper (1986), we wrote before (line 292 ff): “In Cooper (1986), a selection of previously made measurements from literature was examined. However, it was not well described based on what criteria certain data was included or rejected. Still, when sighting this literature, it can be seen that data at higher temperatures up to  $-5^{\circ}\text{C}$  was included in Cooper (1986), ...”. Indeed, after repeated careful reading of Cooper (1986), we are still unable to give more detail. Should you be aware of more, please let us know and we will add it happily.

As for Fletcher (1962), he collected literature data from a number of different cloud chambers used for atmospheric measurements. We already wrote (line 291 ff), “that Fletcher (1962) reported the value of  $-0.6$  as the usual value, but commented that values between  $-0.4$  and  $-0.8$  were still common.” Similarly, concerning the value given for A (the y-intercept), he said that it is “more variable, typically being about  $10^{-5} \text{ L}^{-1}$ , with variations of several orders of magnitude sometimes occurring.” We added this information to our text at the location you indicated (see line 368 ff). Overall, this shows that the parameterization often cited as originating from Fletcher (1962) is merely a small fraction of what was really reported in this older study, which we hope to clarify.

- Line 426: In April 2019 there is a correlation coefficient of 0.86 between INP concentrations at  $-18^{\circ}\text{C}$  and surface temperature.

We had to update some meteorological data due to a new quality control check. This changed values slightly. The value you are referring to is now 0.73.

### Technical comments

- Line 5: The abbreviation „VRS“ is not used in the abstract, thus should not be introduced here.

Done.

- Line 473: „biolgocial“ should be „biological“.

Done.

- Check the use of hyphens: I believe it should be „ice-active“, „temperature-dependent“, etc.

Done. (Also, language check by ACP will take care of those we did not see.)

## References:

AMAP, 2021. Arctic Climate Change Update 2021: Key Trends and Impacts. Summary for Policy-makers. Arctic Monitoring and Assessment Programme (AMAP), Tromsø, Norway. 16 pp

Chen, J., Z. Wu, S. Augustin-Bauditz, S. Grawe, M. Hartmann, X. Pei, Z. Liu, D. Ji, and H. Wex (2018), Ice nucleating particle concentrations unaffected by urban air pollution in Beijing, China, *Atmos. Chem. Phys.*, 18, 3523–3539, doi:10.5194/acp-18-3523-2018.

Curry, J. A., and V. I. Khvorostyanov (2012), Assessment of some parameterizations of heterogeneous ice nucleation in cloud and climate models, *Atmos. Chem. Phys.*, 12(2), 1151-1172, doi:10.5194/acp-12-1151-2012.

DeMott, P. J., A. J. Prenni, X. Liu, S. M. Kreidenweis, M. D. Petters, C. H. Twohy, M. S. Richardson, T. Eidhammer, and D. C. Rogers (2010), Predicting global atmospheric ice nuclei distributions and their impact on climate, *Proc. Natl. Acad. Sci. USA*, 107(25), 11217-11222, doi:10.1073/pnas.0910818107.

Gong, X., M. Radenz, H. Wex, P. Seifert, F. Ataei, S. Henning, H. Baars, B. Barja, A. Ansmann, and F. Stratmann (2022), Significant continental source of ice-nucleating particles at the tip of Chile's southernmost Patagonia region, *Atmos. Chem. Phys.*, 22, 10505-10525, doi:10.5194/acp-22-10505-2022.

Griesche, H. J., K. Ohneiser, P. Seifert, M. Radenz, R. Engelmann, and A. Ansmann (2021), Contrasting ice formation in Arctic clouds: surface-coupled vs. surface-decoupled clouds, *Atmos. Chem. Phys.*, 21(13), 10357-10374, doi:10.5194/acp-21-10357-2021.

Hartmann, M., T. Blunier, S. O. Brügger, J. Schmale, M. Schwikowski, A. Vogel, H. Wex, and F. Stratmann (2019), Variation of ice nucleating particles in the European Arctic over the last centuries, *Geophys. Res. Lett.*, 46, doi:10.1029/2019GL082311.

Sanchez-Marroquin, A., O. Arnalds, K. J. Baustian-Dorsi, J. Browse, P. Dagsson-Waldhauserova, A. D. Harrison, E. C. Maters, K. J. Pringle, J. Vergara-Temprado, I. T. Burke, J. B. McQuaid, K. S. Carslaw, and B. J. Murray (2020), Iceland is an episodic

source of atmospheric ice-nucleating particles relevant for mixed-phase clouds, *Science Advances*, 6(26), doi:10.1126/sciadv.aba8137.

Šantl-Temkiv, T., R. Lange, D. Beddows, U. Rauter, S. Pilgaard, M. Dall'Osto, N. Gunde-Cimerman, A. Massling, and H. Wex (2019), Biogenic sources of Ice Nucleation Particles at the high Arctic site Villum Research Station, *Environ. Sci. Technol.*, 53(18), 10580-10590, doi:10.1021/acs.est.9b00991.

Si, M., E. Evoy, J. Yun, Y. Xi, S. J. Hanna, A. Chivulescu, K. Rawlings, D. Veber, A. Platt, D. Kunkel, P. Hoor, S. Sharma, W. R. Leitch, and A. K. Bertram (2019), Concentrations, composition, and sources of ice-nucleating particles in the Canadian High Arctic during spring 2016, *Atmos. Chem. Phys.*, 19, 3007–3024, doi:10.5194/acp-19-3007-2019.

Tarn, M. D., S. N. F. Sikora, G. C. E. Porter, D. O'Sullivan, M. Adams, T. F. Whale, A. D. Harrison, J. Vergara-Temprado, T. W. Wilson, J. U. Shim, and B. J. Murray (2018), The study of atmospheric ice-nucleating particles via microfluidically generated droplets, *Microfluidics and Nanofluidics*, 22(5), doi:10.1007/s10404-018-2069-x.

Tobo, Y., K. Adachi, P. J. DeMott, T. C. J. Hill, D. S. Hamilton, N. M. Mahowald, N. Nagatsuka, S. Ohata, J. Uetake, Y. Kondo, and M. Koike (2019), Glacially sourced dust as a potentially significant source of ice nucleating particles, *Nat. Geosci.*, 12(4), 253+, doi:10.1038/s41561-019-0314-x.

Vali, G. (2019), Revisiting the differential freezing nucleus spectra derived from drop-freezing experiments: methods of calculation, applications, and confidence limits, *Atmos. Meas. Tech.*, 19, 1219–1231, doi:10.5194/amt-12-1219-2019.

Welti, A., E. K. Bigg, P. J. DeMott, X. Gong, M. Hartmann, M. Harvey, S. Henning, P. Herenz, T. C. J. Hill, B. Hornblow, C. Leck, M. Löffler, C. S. McCluskey, A. M. Rauker, J. Schmale, C. Tatzelt, M. van Pinxteren, and F. Stratmann (2020), Ship-based measurements of ice nuclei concentrations over the Arctic, Atlantic, Pacific and Southern Ocean, *Atmos. Chem. Phys.*, 20, 15191-15206, doi:10.5194/acp-20-15191-2020.

Wex, H., L. Huang, W. Zhang, H. Hung, R. Traversi, S. Becagli, R. J. Sheesley, C. E. Moffett, T. E. Barrett, R. Bossi, H. Skov, A. Hünerbein, J. Lubitz, M. Löffler, O. Linke, M. Hartmann, P. Herenz, and F. Stratmann (2019), Annual variability of ice nucleating particle concentrations at different Arctic locations, *Atmos. Chem. Phys.*, 19, 5293–5311, doi:10.5194/acp-19-5293-2019.

Yang, X., M. M. Frey, R. H. Rhodes, S. J. Norris, I. M. Brooks, P. S. Anderson, K. Nishimura, A. E. Jones, and E. W. Wolff (2019), Sea salt aerosol production via sublimating wind-blown saline snow particles over sea ice: parameterizations and relevant microphysical mechanisms, *Atmos. Chem. Phys.*, 19(13), 8407-8424, doi:10.5194/acp-19-8407-2019.