

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

**Changes in surface ozone in South Korea on diurnal to decadal time scale
for the period of 2001-2021**

Si-Wan Kim¹, Kyoung-Min Kim², Yujoo Jeong^{1,2}, Seunghwan Seo², Yeonsu Park², and
Jeongyeon Kim²

¹Irreversible Climate Change Research Center, Yonsei University, Seoul, South Korea
²Department of Atmospheric Sciences, Yonsei University, Seoul, South Korea

*Corresponding author. E-mail: siwan.kim@yonsei.ac.kr

Short title: Ozone changes in South Korea

Abstract

Several studies have reported an increasing trend of surface ozone in South Korea over the past few decades, using different measurement metrics. In this study, we examined the surface ozone trends in South Korea by analyzing the hourly or daily maximum 8-hour average ozone concentrations (MDA8) measured at the surface from 2001 to 2021. We studied the diurnal, seasonal, and multi-decadal variations of this parameter at city, province, and background sites.

We found that the 4th highest MDA8 values exhibited positive trends in 7 cities and 9 provinces from 2001 to 2021, with an approximate annual increase of 1-2 ppb. After early 2010, all sites consistently recorded MDA8 values exceeding 70 ppb, despite reductions in precursor pollutants such as NO₂ and CO. The diurnal and seasonal characteristics of ozone exceedances, defined as the percentage of data points with hourly ozone concentrations exceeding 70 ppb, differed between the Seoul Metropolitan Area (SMA) and the background sites.

In the SMA, the exceedances were more prevalent during summer compared to spring, whereas the background sites experienced higher exceedances in spring than in summer. This indicates the efficient local production of ozone in the SMA during summer and the strong influence of long-range transport during spring. The rest of the sites showed similar exceedance patterns during both spring and summer. The peak exceedances occurred

1 around 4-5 PM in the SMA and most locations, while the background sites primarily
2 recorded exceedances between 7-8 PM and throughout the night.

3 During the spring of the COVID-19 pandemic (2020-2021), ozone exceedances decreased
4 at most locations due to significant reductions in NO_x emissions in South Korea and China
5 compared to the period of 2010-2019. The largest decreases in exceedances were observed
6 at the background sites during spring. For instance, in Gosung, Gangwondo (approximately
7 600 m above sea level), the exceedances dropped from 30% to around 5% during the
8 COVID-19 pandemic.

9 Regional model simulations confirmed the concept of decreased ozone levels in the
10 boundary layer in Seoul and Gangwon-do in response to emission reductions. However,
11 these reductions in ozone exceedances were not observed in major cities and provinces
12 during the summer of the COVID-19 pandemic, as the decreases in NO_x emissions in South
13 Korea and China were much smaller compared to spring. This study highlights the
14 distinctions between spring and summer in the formation and transport of surface ozone in
15 South Korea, emphasizing the importance of monitoring and modeling specific processes
16 for each season or finer time scales.

17

1 **1. Introduction**

2 Ozone, a greenhouse gas and harmful air pollutant, can accumulate in the lower atmosphere
3 through photochemical reactions involving nitrogen oxides and volatile organic
4 compounds from both human activities and natural sources (National Research Council,
5 1991; Monks et al., 2015). The increasing concentrations of ozone near the surface and in
6 the troposphere are concerning. Gaudel et al. (2018) reported a significant increase in ozone
7 levels in South Korea from 2000 to 2014, while North America and Europe experienced
8 decreasing trends, using data from surface monitors, ozonesondes, and aircraft
9 observations. Other studies have also observed rising ozone trends in South Korea between
10 2001 and 2018 in their analysis of the long-term variations of multiple pollutants over Seoul
11 (Kim and Lee, 2018) and South Korea (Kim et al., 2018) or in the reviews of current status
12 and future directions of tropospheric ozone studies in South Korea (Lee et al., 2020) or in
13 the trend estimates of the surface ozone observations (Yeo and Kim, 2021). Ozone in South
14 Korea can be influenced by ozone and its precursor in China (Oh et al., 2010; Lee and Park,
15 2022; Colombi et al., 2023). However, Gaudel et al. (2018) did not include Chinese data
16 due to a lack of reported information. Recent studies have highlighted a rapid increase in
17 ozone levels in China from 2004 to 2020, especially after 2013 (Li et al., 2019; Wang et
18 al., 2020; Wang et al., 2022). Gaudel et al. (2020) also found that tropospheric ozone in

1 China and South Korea increased between 1996 and 2016. Considering the proximity of
2 the two countries and their potential for ozone and precursor exchange, it is essential to
3 study the ozone trends in South Korea in relation to those in China. Additionally, as spring
4 and summer have distinct transport patterns and source-receptor relationships relevant to
5 surface and tropospheric ozone (e.g., Cooper et al., 2010), it would be valuable to
6 investigate ozone trends separately for these seasons.

7 The COVID-19 pandemic brought about significant changes in atmospheric
8 composition (Bauwens et al., 2020; Koo et al., 2020; Seo et al., 2021). Analyzing deviations
9 from long-term trends during the pandemic can provide valuable insights for future
10 environmental policies aimed at mitigating ozone pollution. In this study, we examine
11 ozone trends and exceedances in South Korea from 2001 to 2021, focusing on the warm
12 seasons of spring and summer, including the COVID-19 period. In this study, we analyzed
13 the 4th highest daily maximum 8 hours-average ozone concentrations (MDA8 O₃) at
14 various locations in South Korea for a global comparison because this is a metric used for
15 the US Environmental Protection Agency National Ambient Air Quality Standard and the
16 recent study by Wang et al. (2022) utilized the same metric for their study of Chinese ozone
17 pollution. We also introduced a new metric of ozone exceedance, defined as the percentage
18 of data points with hourly ozone concentrations exceeding 70 ppb. Previous published

1 works about surface ozone in South Korea have not focused on the two metrics used in our
2 study. We analyze diurnal, seasonal, and decadal variations at 7 cities, 9 provinces, and 2
3 background sites. Furthermore, we discuss the factors contributing to the observed
4 temporal changes based on regional model results.

5 The manuscript is organized as follows. In section 2, the surface and satellite data,
6 global and regional modeling, and other methods to utilize the data are explained. In section
7 3, the results are summarized as long-term trends of ozone and its precursors,
8 characteristics of diurnal variations, and spatiotemporal variations during the pandemic.
9 The regional model results based on various emission scenarios are also shown to identify
10 the source-receptor relationship. Finally, the results are summarized and future research
11 directions are suggested in the conclusions.

12

13 **2. Data and Method**

14 **2.1. Long-term surface observational data**

15 The hourly surface air quality monitoring data are obtained from the Airkorea website
16 (<https://www.airkorea.or.kr>), including ozone (O₃), NO₂, SO₂, CO, PM₁₀, and PM_{2.5} (PM_{2.5}
17 data are provided since 2015). As of March 2020, there are about 500 monitoring stations
18 over South Korea. These routine monitor data are available for many decades and can serve

1 as a main data set to examine long-term trends. We utilized hourly and daily maximum 8
2 hour-average O₃ concentrations. The surface monitoring sites used in this study and the
3 data availability are summarized in the Supporting Information 1 (SI 1, Table S1) and
4 Supporting Information 2 (SI 2). O₃, NO₂ and CO data are also averaged for spring and
5 summer months. These surface monitoring data were used to investigate the impact of the
6 COVID-19 pandemic in the Seoul Metropolitan Area.

7

8 **2.2. Highway toll number and mobile phone usage data**

9 To examine changes in mobility pattern during the COVID-19 pandemic, traffic counts
10 from the Korea Expressway Corporation daily transit data were used
11 (<http://data.ex.co.kr/portal/>). The expressway transit data covering 3 years (2019-2021) of
12 traffic passing toll gates were quantified from Hi-Pass (electronic toll collection system)
13 and cash toll collection. Vehicles passing toll gates were not classified in details.

14 To examine changes in mobility pattern during the COVID-19 pandemic, daily
15 mobile phone movement provided by Android (Google COVID-19 Community Mobility
16 Reports, 2020) and Apple (Apple COVID-19 Mobility Trends Report, 2020) are used.
17 Android mobility data tracked movements of people using cell phones at the same spot,
18 while Apple's mobility report collects personal vehicle routing requests from Apple Maps.

1 For Google and Apple mobility report, we used the Transit station Mobility metrics and
2 driving mobility index in Seoul Metropolitan Area, respectively. The reports must be
3 carefully used as it does not directly quantify on-road traffic.

4

5 **2.3. Satellite data: tropospheric NO₂ columns**

6 The TROPOspheric Monitoring Instrument (TROPOMI) on board of a low Earth polar
7 orbiting satellite, European Space Agency (ESA) Sentinel-5 Precursor (S-5P) satellite with
8 equator passing time 13:30 local time. The instrument provides measurements at
9 unprecedentedly high spatial, temporal, and spectral resolutions (Veefkind et al., 2012). In
10 this study we utilized two available tropospheric NO₂ datasets from TROPOMI, NASA's
11 standard product (SP) version 4.0 (Lamsal et al., 2021) and KNMI's (Royal Netherlands
12 Meteorological Institute) product obtained from DOMINO v2.0 and QA4ECV v1.1
13 (Derivation of TROPOMI tropospheric NO₂) processing systems (Boersma et al., 2018).
14 The spatial resolution of KNMI's tropospheric NO₂ retrieval product is 3.5 km x 7 km (3.5
15 km x 5.5 km since 6 August 2019) and that of NASA's product is 3.5 km x 5.5 km. Level
16 2 data with pixels passing quality assurance > 0.75 and the cloud fraction < 0.5 were
17 selected for analysis following recommendations provided by Sentinel-5 precursor
18 TROPOMI Level 2 product User Manual for nitrogen dioxide (Eskes et al., 2019).

1 TROPOMI data are regridded to a standard grid with a horizontal resolution of 0.1° latitude
2 $\times 0.1^\circ$ longitude (11×11 km) and monthly averaged values were derived. As the random
3 error in the TROPOMI single-pixel uncertainties influence 40 to 60% of the tropospheric
4 column abundance, temporal and spatial averaging may remove the random errors
5 (Bauwens et al., 2020).

6 We conducted the sensitivity test by applying different sampling conditions and
7 found consistent results irrespective of quality control parameters: larger tropospheric NO_2
8 column reduction during spring than during summer between 2019 and 2020-2021
9 (COVID-19 periods). Differences between KNMI and NASA retrievals are large when the
10 the filtering condition of quality assurance > 0.5 and cloud radiance fraction < 0.4 is applied.
11 When the stricter filter is applied, differences between KNMI and NASA retrievals are
12 small. Therefore, the stricter filter (quality assurance > 0.75 and cloud radiance fraction $<$
13 0.5) is selected. Since the NASA product released in November, 2022 were generated in a
14 consistent manner for May 2018-December 2021, we mainly present the NASA MINDS
15 product. We summarized the sensitivity tests in the Supporting Information 3 (SI3). The
16 distribution of absolute tropospheric NO_2 columns for different years are also shown in the
17 SI3.

18

1 **2.4. CAM-Chem model simulations**

2 The atmospheric component of Community Earth System model (CESMv2.2), Community
3 Atmosphere Model with Chemistry version 6 (CAM6-chem) is developed by National
4 Center for Atmospheric Research (NCAR) (<https://www2.acom.ucar.edu/gcm/cam-chem>).

5 The CAM-chem adapted MOZART-T1 as the tropospheric chemistry mechanism
6 (Emmons et al., 2020). The simulation used in this study was configured with 1° horizontal
7 resolution. The sea surface temperature was prescribed, and meteorological fields were
8 nudged to Modern-Era Retrospective analysis for Research and Applications version 2
9 (MERRA-2) instead of using self-produced meteorological field
10 (<https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/>) (refer to SI1 Figure S1 for
11 performance of the model wind). The simulation was performed from 2000 to 2020 and
12 applied CMIP6 emission inventory (2000-2014) and SSP5-8-5 emission inventory (2015-
13 2020). The first 3 years were regarded as a spin-up. In this study, we utilized the CAM-
14 Chem results to estimate the impact of stratospheric ozone on the surface in each season.

15 CAM-Chem calculates the contribution of stratospheric ozone to tropospheric ozone, O_{3S}
16 as a three-dimensional variable in space. Originally, O_{3S} is ozone value above tropopause.
17 Then O_{3S} is transported below tropopause and undergoes chemical losses in the model.

18 Evaluations of the CAM-Chem results against the data from the ozonesondes that were

1 launched in Pohang, South Korea are shown in the Supporting Information (SI1, Figure S2;
2 Jeong et al., 2023). The model results and observations reasonably agree in terms of
3 seasonal variability and absolute values. Especially, the CAM-Chem results agree with the
4 observations at the 200 hPa level, close to tropopause.

5

6 **2.5. WRF-Chem model simulations**

7 The Weather Research and Forecasting (WRF) model coupled with Chemistry (WRF-
8 Chem) is developed by National Oceanic and Atmospheric Administration (NOAA) and
9 National Center for Atmospheric Research (NCAR) and collaborating institutes (Grell et
10 al., 2005). We utilized WRF-Chem v4.4 to simulate regional meteorological fields and
11 chemical compositions.

12 Our WRF-Chem set up utilizes the horizontal resolution of 28 x 28 km² and 60
13 vertical levels. The simulation period is from 24th April 12 UTC to 11th June 12 UTC in
14 2016. We restart the simulation at 12 UTC every day to reduce computing errors. The first
15 7 days of model simulation is regarded as spin-up period. The analysis period is selected
16 as 1st May to 10th June based on local time. The Global Forecast System (GFS) Final (FNL)
17 analysis data are used for meteorological input and boundary conditions
18 (<https://rda.ucar.edu/datasets/ds083.2/>). We used The Community Atmosphere Model with

1 Chemistry (CAM-Chem) output to the chemical boundary and first initial conditions
2 (<https://www.acom.ucar.edu/cam-chem/cam-chem.shtml>) (Buchholz et al., 2019). The
3 Model of Emissions of Gases and Aerosols from Nature (MEGAN) is used for biogenic
4 emissions (Guenther et al., 2006).

5 There are 7 model sensitivity runs that adopt different emission scenarios. The
6 control run is based on the standard EDGAR-HTAPv3 emission inventory representing
7 2016 (Crippa et al., 2023). Park et al (2021) informed that biomass burning was not an
8 important factor affecting air quality in South Korea during KORUS-AQ. Therefore,
9 biomass burning emissions are omitted in this study. “No China” case removes all
10 anthropogenic emissions in China. “No Seoul” case eliminates all anthropogenic emissions
11 in Seoul. There is one case that decreased Chinese VOC emissions by 50%. There are two
12 cases that reduced Chinese NO_x emissions by 50%: the one case has the same VOC
13 emissions as in the control case while the other case has the 50% reductions of VOC
14 emissions as well. Lastly, there is one case that reduced Chinese NO_x emissions by 75%.
15 The WRF-Chem sensitivity runs are summarized and are discussed in Section 3. The
16 extensive evaluations of the model results against the surface and airborne data from the
17 KORUS-AQ field campaign in 2016 are shown in the Supporting Information (SI1 Table
18 S2-S4 and Figure S3-S8) and Kim K.-M. et al (2023).

1

2 **3. Results**

3 **3.1. Surface ozone trends**

4 In this study, ozone and its precursor concentrations in 7 cities, 9 provinces, and 2
5 background sites in South Korea (Figure 1) are analyzed at diurnal, seasonal, and decadal
6 time scales. Figure 2 and 3 shows the 4th highest daily maximum 8 hours-average ozone
7 concentrations (MDA8 O₃) for the cities and provinces for ozone season (May-September)
8 from 2001 to 2021. The results from statistical analysis (slope, standard-deviation, p-value,
9 signal-to-noise ratio) are summarized in Table 1. P-values were presented as suggested by
10 Chang et al. (2021) and Wasserstein et al. (2019) for the purpose of estimating uncertainties
11 in trends. Because of discontinuity of data records, the background sites are omitted in the
12 trend analysis in Figure 2 and 3 and Table 1. The 4th highest MDA8 O₃ increases by 1.0-
13 1.5 ppb yr⁻¹ with very high certainty for most of cities and provinces across South Korea
14 in this period. In nearly all cities and provinces, the 4th highest MDA8 O₃ has been higher
15 than 70 ppb since 2010 or earlier (see gray dashed line in Figures 2 and 3). The trend in
16 Jeollanam-do (JLN) is small with very low certainty (p=0.67) partly because the MDA8
17 O₃ was high before 2010. The monitoring sites in Jeollanam-do include the Yeosu-
18 Kwangyang region in which many petrochemical industries and iron steel complexes are
19 located. This region experienced severe ozone problems in the 1990's to early 2000's
20 (Ghim, Y. S. 2000). Widely increasing long-term ozone trends in South Korea indicate a

1 regional nature of this pollutant, potentially influenced by Asian emissions, chemical
2 transformations, and long-range transport (Colombi et al., 2023; Lee and Park, 2022).
3 Therefore, it is imperative to understand the local and regional processes that enhance
4 surface ozone. Ozone originated from Asia is known to be efficiently transported to North
5 America during springtime (Jacob et al., 1999; Jaffe et al., 1999; Jaffe et al., 2003; Cooper
6 et al., 2010; Lin et al., 2012; Langford et al., 2017; Jaffe et al., 2018) and summertime
7 (Fiore et al., 2002; Liang et al., 2007) as well. Investigating seasonal differences in ozone
8 in South Korea may provide insights on the relative importance of local and regional
9 processes.

10

11 **3.2. Difference between spring and summer: background value, exceedance,** 12 **stratospheric influence, and precursor concentrations**

13 3.2.1. Background values at the base and peak times

14 Table 2 summarizes the abundances and differences between spring and summer ozone
15 concentrations averaged for the peak time (10-20 Local Time (LT)) and the base time (01-
16 06 LT). For the base time, the ozone concentration in spring is always higher than that in
17 summer: differences between the two seasons range from 3.1 to 14.5 ppb. This clearly
18 indicates the importance of large-scale influences in spring. The results are the same for
19 the peak time except for Seoul and Gyeonggi-do: the mean ozone concentrations in Seoul
20 and Gyeonggi-do in summer are slightly higher than those in spring. The differences at the

1 peak time are small for Incheon, Daegu, and Chungcheongbuk-do, suggesting the
2 importance of local chemistry in the areas during summer.

3 The surface ozone data from the base time (01-06 LT) over polluted regions are
4 often omitted in the analysis because ozone loss reacting with NO is an important process
5 to control ozone levels at nighttime. In this study, we utilized the ozone data at this time to
6 find information about background ozone because ozone is transported throughout a day
7 and this process is essential in the studied region. WRF-Chem sensitivity runs
8 demonstrated increase of ozone from upwind sources at this time (refer to SII Figure S9).

9

10 3.2.2. Ozone exceedances

11 Figure 4 illustrates the ratio of summer ozone exceedances to spring ozone exceedances
12 for the cities, provinces, and background sites. In Seoul, Incheon, and Gyeonggi-do, there
13 are more exceedances in summer than in spring, indicating the significance of local ozone
14 production during the summer season in these areas. Conversely, at the background sites
15 such as Gosung and Ulleung Island, springtime exceedances dominate, highlighting the
16 importance of high springtime ozone levels and their transport within and beyond Asia. For
17 the remaining regions, springtime and summertime exceedances are comparable, or
18 springtime exceedances are slightly higher than those in summer. Note that meteorological
19 conditions in Seoul and Gyeonggi-do (differences between the two seasons) are similar to
20 other cities and provinces (see SII Table S5). Therefore, the meteorological factors are not

1 main drivers of high summertime exceedances in Seoul and Gyeonggi-do region.

2 The diurnal variations of exceedances, as shown in Figure 5, confirm these
3 findings. The summertime ozone exceedances are notably enhanced during the daytime,
4 from 13 to 20 local time (LT), suggesting efficient photochemical ozone production during
5 this season. The peak exceedances occur at 17 LT in Seoul and Gyeonggi-do, and one hour
6 earlier at 16 LT in Incheon. Incheon, being situated adjacent to the West Sea (as depicted
7 in Figure 1), experiences airflow from Incheon to Seoul under typical westerly or seabreeze
8 conditions. The late-afternoon peaks (4-5 PM) in the region and the one-hour delay in peak
9 exceedances in Seoul compared to the time of exceedances in Incheon imply that local
10 circulation plays a significant role in the buildup and distribution of ozone within the
11 Incheon, Seoul, and Gyeonggi-do region.

12 Springtime and summertime ozone exceedances predominantly occur during the
13 daytime, with some extent of exceedances at night, in Daejeon, Busan, and Daegu (Figure
14 5). Notably, the peaks in spring occur approximately two hours later than those in summer
15 for the three cities, indicating a potential influence of transport during the spring season.
16 Negligible exceedances are observed from midnight to 10 LT in the three cities due to high
17 NO_x pollution and the depletion of ozone associated with NO_x during this time period.

18 At the background sites, springtime exceedances are much higher compared to
19 summer, and nighttime exceedances are as frequent as daytime exceedances. In Gosung,
20 springtime exceedances account for approximately 20% of the observations throughout the

1 day, whereas summertime exceedances are less than or equal to 10% (Figure 5). The
2 observation site in Gosung is located at an altitude of approximately 600 meters above sea
3 level, providing a unique opportunity to examine long-range transported plumes and
4 background information at higher altitudes (refer to SI1 Table S6 and Figure S10). Diurnal
5 variations of exceedances during spring and summer for all individual sites are illustrated
6 in SI1 (Figure S11-S13).

7

8 3.2.3. Influence of stratospheric ozone

9 Stratospheric ozone can deeply intrude into the lower troposphere, leading to elevated
10 surface ozone levels, particularly during the spring season (Lin et al., 2012; Lin et al., 2015).
11 It is important to assess the contribution of stratospheric ozone to surface ozone in South
12 Korea and understand its potential impact on surface ozone trends in the region using
13 results from the CAM-Chem model. The derivation of the contribution of stratospheric
14 ozone in the CAM-Chem is explained in the Supporting Information. Figure 6 presents the
15 contribution of stratospheric ozone to surface ozone in South Korea for each season.
16 According to our global chemistry-climate model simulations, stratospheric ozone has the
17 greatest influence on surface ozone during winter and spring, increasing levels by 17-23
18 ppb. The model suggests that approximately 37% and 76% of surface ozone in spring and
19 winter, respectively, can be attributed to stratospheric ozone (refer to SI1 Table S7 for
20 summary of the CAM-Chem results for all seasons at surface and 1 km above ground level).

1 However, during the summer season, the impact of stratospheric ozone on surface ozone is
2 minimal, accounting for only around 4% of the surface ozone concentration. Therefore, it
3 would be valuable to analyze ozone trends and exceedances separately for spring and
4 summer. It is worth noting that the contribution of stratospheric ozone to surface ozone
5 does not exhibit clear trends during the period from 2001 to 2021 (not shown). Note that
6 the contribution of stratospheric ozone to tropospheric ozone at each altitude and time
7 shown in this study should be a qualitative measure since the representation of this process
8 has uncertainties and needs further assessment.

9

10 3.2.4. Long-term trends of surface NO₂ and CO concentrations

11 In contrast to the trends of ozone, NO₂ and CO that are ozone precursors decreased both in
12 spring and summer from 2001 to 2021 (Table 3 and 4). There are no systematic differences
13 in the trends of NO₂ and CO between the two seasons. NO₂ has declined in Seoul, Busan,
14 Daegu, Gwangju, Incheon, Gyeongsangbuk-do, and Gyeonggi-do with very high certainty.
15 For the rest of sites, the declining NO₂ trends were found with medium-to-high certainty
16 (refer to Chang et al., 2021 for assessment of uncertainty in the trend analysis). Seo et al.
17 (2021) investigated the trend of NO₂ in the Seoul area utilizing satellite tropospheric NO₂
18 columns and surface NO₂ observations from 2005 to 2019 and found decrease of NO₂ only
19 between 2015 and 2019. They did not find significant trends between 2005 and 2015.
20 Therefore, the trends in our study are strongly influenced by recent NO₂ reductions prior
21 to and during the COVID-19 pandemic. CO reductions are evident for a wider region with
22 very high certainty. Only the CO trend in Jeollanam-do was estimated with high certainty

1 (instead of very high certainty). The decreasing trends of NO₂ and CO were estimated with
2 very low certainty in Ulsan throughout this period. Overall, signs of slopes agree between
3 emission inventory and ambient concentrations at least for the cities, but site-to-site
4 variations do not agree even for the cities. There are disagreements of signs of slopes
5 between emission inventory and ambient concentrations for the provinces (refer to SII
6 Table S8 and S9). This can be attributed to the uncertainties in the bottom-up emission
7 inventories of NO_x and CO in South Korea.

8 Ozone increases in South Korea despite reduction of main precursors at local scale
9 can be attributed to the increase of long-range transport of ozone or potentially “VOC-
10 limited” (or “NO_x-saturated”) local photochemical regime of South Korea. “VOC-limited”
11 regime is the condition in which NO_x (sum of NO and NO₂) concentration is high and VOC
12 is a limiting factor to form ozone. In this case, VOC reduction would decrease ozone, while
13 NO_x reduction would nonlinearly increase ozone. Since long-range transport from China
14 is frequent during spring, it is useful to identify characteristics of ozone exceedance in
15 spring separate from summer.

16

17 **3.3. Changes detected during the COVID-19 pandemic (2020-2021) compared to** 18 **2002-2019**

19 Nationwide social distancing protocol enforced by Korean government started February 25
20 of 2020 and lasted until April 18 of 2022, although levels of protocol differ. During spring
21 in 2020 (until May 6, 2020), facilities for public use (libraries, swimming pools, museums,

1 and national parks) and religious, indoor sports, entertainment facilities were forced to
2 close, and people were refrain from going out except for buying necessities, visiting a
3 doctor, and commuting to/from work. Since May 6 of 2020, as number of new confirmed
4 COVID-19 cases remain relatively steady, the guidelines have shifted from social
5 distancing to distancing in daily life, no restrictions on people going out. Because a cluster
6 of new COVID-19 cases emerged in mid-August, social distancing protocol (since August
7 16 until early October) was again forced by the government, people were strongly
8 recommended to stay indoors. After August 16 of 2020, there were well-defined
9 government protocols as Level 1, 2, and 3: Level 1 is no restricted personal gathering and
10 daily life, Level 2 allows personal gathering up to 8 people and discourage unnecessary
11 and unurgent travel, and Level 3 allows personal gathering up to 3 people, requires remote
12 work and online classes, and discourage travels. Most days in spring and summer in 2021
13 were the period under the Level 2 protocol. In summary, most distinct changes in social-
14 distancing protocols and traffic/mobile activities occurred between spring and summer in
15 2020 in South Korea (refer to SI1 Figure S14-15).

16

17 3.3.1. Changes in ozone exceedances and local precursors during springtime

18 The frequency of springtime ozone exceedances increases from period P1 (2002-2010) to
19 period P2 (2011-2019) across all observation sites in South Korea (Figure 7). However,
20 during the COVID-19 period (P3: 2020-2021), the frequency of exceedances significantly

1 decreases at most sites. Notable reductions are observed in Daejeon, Daegu,
2 Chungcheongbuk-do, Gyeongsangnam-do, Gyeongsangbuk-do, Gangwon-do, as well as
3 the background sites Gosung (Gangwon-do) and Ulleung Island. In Gosung, the percentage
4 of ozone exceedances drops from 30% during P2 to 5% during P3 in spring. Although
5 Gosung is located close to the East Sea and is the region farthest from China within a
6 similar latitude range, it is still susceptible to long-range transported ozone due to its high
7 elevation (see S11 Figure 10 for the elevation map and diagram of a possible ozone
8 transport path).

9 Across all sites, the concentration of NO₂ shows little change from P1 to P2, with
10 an average decrease of 5%. However, during the COVID-19 period (P2 to P3), there was
11 an average reduction of 25% in NO₂ concentrations. CO concentrations also experienced a
12 decrease of 22% from P1 to P2 and a further decrease of 14% from P2 to P3. However, the
13 reductions in CO are relatively minor compared to the changes in NO₂ observed during the
14 COVID-19 period. The decrease in ozone exceedances during COVID-19 may be
15 associated with the reductions in NO₂ concentrations during this time.

16 A notable finding is the significant reduction in ozone levels at the background
17 sites, such as Gosung and Ulleung Island, between P2 and P3. This suggests a cleaner
18 background influenced by changes in emissions from sources in Asia and long-range
19 transport. It is important to note that there were no significant changes in NO₂ and CO
20 concentrations observed at the background sites from P2 to P3. There are several studies

1 reporting the increase of near-surface ozone after COVID lockdowns in the urban areas
2 (e.g., Shi & Brasseur, 2020) because of expected non-linear relationship between ozone
3 and NO_x in the highly polluted regions. However, there are also studies reporting reductions
4 of ozone concentrations from 1 to 8 km altitude in the northern extra-tropics during COVID
5 (Steinbrecht et al., 2021). Parrish et al. (2020) reported zonal similarity of tropospheric
6 ozone changes at northern mid-latitudes. Therefore, ozone reductions from P2 to P3 across
7 the sites in South Korea may be associated with decreased background ozone at northern
8 mid-latitudes to some extents. On top of this, local and regional emission changes during
9 COVID may also play a role in reducing ozone exceedances in South Korea in this season.

10

11 3.3.2. Changes in ozone exceedances and local precursors during summertime

12 During summer, ozone exceedance frequencies also increase from P1 to P2 for all sites:
13 Chungcheongnam-do has the largest increase from 3.2% to 11.3% and Gyeonggi-do,
14 Daejeon, Jeollabuk-do, Gyeongsangnam-do and Gyeongsangbuk-do have similar increases
15 (Figure 8). The ozone exceedances in the background sites Gosung, and Ulleung Island
16 also increase in this period. NO₂ and CO concentrations decreased marginally from P1 to
17 P2. During COVID-19, the ozone exceedance frequencies in summer increase in Seoul,
18 Incheon, Gyeonggi-do, and Chungcheongnam-do, substantially decrease in Gangwon-do
19 and the background sites, and does not show changes from P2 for the rest of sites. Because
20 NO₂ concentrations decrease from P2 to P3 for Seoul, Incheon, Gyeonggi-do, and

1 Chungcheongnam-do contrasting with increases of ozone exceedance, chemical regime for
2 these regions during summer is likely to be VOC-limited (NO_x -saturated) as mentioned
3 above and as in previous studies (e.g., Kim et al., 2020). Ozone exceedance substantially
4 decreases in the background sites from P2 to P3 during summer, indicating cleaner air at
5 large-scale as shown in Steinbrecht et al. (2021).

6

7 3.3.3. Changes in precursor concentrations at a regional scale during spring and summer:

8 TROPOMI tropospheric NO_2 columns

9 Figure 9 presents the spatial distributions of NASA TROPOMI tropospheric NO_2 columns
10 (Lamsal et al., 2022) in spring (MAM) and summer (JJA) across East Asia, along with their
11 changes from 2019 to 2020 and from 2019 to 2021. The plot illustrates significant
12 reductions in NO_2 columns during the spring of COVID-19 in most areas of China, South
13 Korea, and the surrounding seas. Changes in traffic activities in the Seoul Metropolitan
14 Area were also detected between 2019 and 2020 (refer to SI1 Figure S14 and 15). The
15 number of cars counted at the highway tolls in this region decreased by 6% in March, April,
16 and May in 2020 compared to 2019, but this trend was reversed in June (SI1 Figure S14).
17 Furthermore, observed concentrations of NO_2 , SO_2 , CO, PM_{10} , and $\text{PM}_{2.5}$ during the spring
18 of 2020 showed reduction of 15-30% (SI1 Figure S15). Changes in traffic counts in the
19 Seoul Metropolitan Area between 2019 and 2021 were small (SI1 Figure S14 and S15).
20 But observed concentrations of NO_2 , SO_2 , CO, PM_{10} , and $\text{PM}_{2.5}$ were also reduced during

1 spring in 2021 compared to 2019 by 10-30% except for PM₁₀ that was enhanced due to
2 Asian dust events in spring 2021 (SI1 Figure S15).

3 As depicted in Figure 9, TROPOMI tropospheric NO₂ columns also decreased
4 during the summer in the same region, although in fewer locations and to a lesser extent
5 compared to the spring, during the COVID-19 period. The observed NO₂, SO₂, CO, PM₁₀,
6 and PM_{2.5} concentrations in the Seoul Metropolitan Area were also reduced during summer
7 in 2020 or 2021 compared to 2019 by 2-20%. Surface NO₂ concentrations reduced by ~10%
8 during summer, which is smaller than the reductions during spring (~20%; see SI1 Figure
9 S14 and S15). Overall, the reductions in 2020/2021 from 2019 during summer are smaller
10 than those during spring at the surface in the Seoul Metropolitan Area, which is similar to
11 the seasonal changes detected from space.

12 The substantial decrease in NO₂ in China during spring, observed by satellite, is
13 likely to contribute to significant reductions in ozone levels in South Korea due to long-
14 range transport. Additionally, local reductions in NO_x emissions in South Korea can lead
15 to ozone decreases if the reductions are significant enough, especially in the "VOC-limited"
16 chemical regime prevalent in this area. However, further investigation is required to
17 understand the detailed source-receptor mechanism of ozone and its precursors in each
18 season, which warrants long-term air quality model simulations in future studies. The next
19 section of this study discusses the sensitivity of ozone concentrations in Seoul and
20 Gangwon-do to various emission scenarios in China and South Korea, albeit within a

1 limited time period.

2

3 **3.4. Impacts of changes in East Asian emissions on surface/boundary layer ozone in**
4 **South Korea: a modeling analysis**

5 3.4.1. Changes in surface/boundary layer ozone due to emission reductions: East Asian
6 region

7 In this section, we will discuss WRF-Chem model simulations conducted during the
8 KORUS-AQ 2016 field campaign (primarily in May; refer to Crawford et al., 2021 for
9 detailed information) to gain insights into the impacts of emission changes on ozone
10 concentrations in East Asia, including South Korea. We have extensively evaluated our
11 model results with the airborne and surface observations acquired during the KORUS-AQ
12 campaign and the routine surface monitors in China and South Korea. The model decently
13 simulated boundary-layer ozone over South Korea (3% difference) for the cases that were
14 strongly influenced by long-range transport. For local emission dominating cases, the
15 model underestimated boundary-layer ozone over South Korea by 20%. The results are
16 summarized in SI1 (Table S3 and Figure S8) and Kim K.-M. et al. (2023). This study
17 considers two extreme cases: the "No China" case, where all anthropogenic emissions in
18 China are removed, and the "No Seoul" case, where all anthropogenic emissions in Seoul
19 are removed. Additionally, several other scenarios are examined, including a 50% reduction
20 in Chinese NO_x emissions only, a 50% reduction in Chinese VOC emissions only, a 50%

1 reduction in both Chinese NO_x and VOC emissions, and a 75% reduction in Chinese NO_x
2 emissions only.

3 Our study reveals both increases and decreases in ozone concentrations due to
4 emission changes resembling those during the COVID-19 period. Specifically, near-
5 surface ozone concentrations in polluted regions increase, while ozone concentrations in
6 the elevated layer show reductions (refer to Figures 10 and 11). A novel finding is the
7 decrease in downwind ozone, from the near surface to the upper layer, resulting from
8 reductions in NO_x/VOC emissions in upwind pollution hotspots (refer to Figures 10 and
9 11 for several sensitivity runs). For instance, a 50%-75% reduction in Chinese NO_x
10 emissions leads to decreased ozone concentrations in Korea, surrounding seas, and the
11 Pacific Ocean, from the surface to the upper layers. However, near-surface ozone in
12 Northeast China increases due to these emission changes.

13 Reductions in Chinese VOC emissions result in decreased ozone concentrations
14 from the surface to the upper layer and from hotspots to downwind areas. Our study
15 suggests potential changes in photochemical regimes with altitude over pollution hotspots,
16 indicating NO_x-saturated conditions near the surface and NO_x-limited conditions in the
17 elevated layer. Thus, the combined effects of vertical and horizontal ozone transport, as
18 well as local production dependent on altitude, would determine the ultimate changes in
19 ozone concentrations at specific locations and altitudes. It is important to note that the

1 accuracy of VOC emission estimates also influences the assessment, but this aspect is
2 highly uncertain and requires further study.

3

4 3.4.2. Vertical sensitivity of ozone changes in South Korea to East Asian emission 5 reductions

6 Figure 11 presents the vertical profiles of simulated ozone concentrations for different
7 emission scenarios. In Seoul, the 50% reduction in Chinese NO_x emissions only slightly
8 decreases ozone concentrations near the surface but decreases them above 500 m AGL
9 (above ground level) to a larger extent. The 50% reduction in Chinese VOC emissions
10 causes a decrease in ozone concentrations from the surface to 2000 m AGL. In the elevated
11 layer (> 1500 m AGL) in Seoul, the reduction in Chinese NO_x emissions leads to a greater
12 decrease in ozone concentrations compared to the reduction in Chinese VOC emissions.
13 The scenario with a 50% reduction in both Chinese NO_x and VOC emissions efficiently
14 decreases ozone concentrations from the surface to 2000 m AGL, particularly above 1000
15 m AGL. The scenario with a 75% reduction in NO_x emissions decreases ozone
16 concentrations near the surface similarly to the scenarios with a 50% reduction in NO_x and
17 VOC emissions, but it causes the largest ozone reductions above 1000 m AGL, except for
18 the "No China" emission scenario. The "No China" emission scenario results in ozone
19 concentrations 10-15 ppb lower than the control case at all altitudes. On the other hand, the
20 "No Seoul" emission scenario leads to ozone concentrations about 20 ppb higher than the

1 control case near the surface, partly due to significantly reduced ozone depletion reactions
2 with NO. The sensitivity test results for Seoul and Gosung, Gangwon-do are similar, except
3 that all emission scenarios (including "No Seoul" and 50% reduction in Chinese NO_x
4 scenarios) cause a decrease in ozone concentrations in Gangwon-do. Both NO_x and VOC
5 emission reductions in China contribute to cleaner air in Gangwon-do, with the largest
6 cleaning effect observed above 500 m AGL. This may explain the sharp decline in ozone
7 exceedances observed in Gosung, located at an elevation of approximately 600 m AGL,
8 during the COVID-19 pandemic (Figure 7). Refer to S11 (Table S6 and Figure S10) about
9 altitudes of monitoring sites in Gangwon-do including Gosung. The sensitivity runs clearly
10 demonstrate the long-range transport of Chinese ozone or the influence of Chinese
11 emissions on the eastern part of the Korean Peninsula, such as Gangwon-do, from May to
12 the beginning of June 2016. Both reductions in Chinese VOC emissions and NO_x
13 emissions contribute to improving ozone pollution in the boundary layer (1-3 km) in South
14 Korea.

15

16 3.4.3. Comparisons with recent modeling research

17 Lee and Park (2022) investigated seasonal differences in ozone utilizing a chemical
18 transport model. They reported the April mean ozone concentration of 39.3 ppb, which is
19 slightly higher than the July counterpart (38.3 ppb) from their model simulations for the
20 year 2016 and the selected surface monitor sites for 4 main regions (Seoul,

1 Chungcheongbuk-do, Gwangju, and Busan). Our study summarizes the differences
2 between spring (March, April, May) and summer (June, July, August) for 21 years
3 including 192 monitoring sites covering the whole of South Korea focusing on the analysis
4 of long-term surface ozone observations. On average, the observed spring mean ozone is
5 34.3 ppb and the summer mean ozone is 29.0 ppb over South Korea in our study. Lee and
6 Park (2022) indicated that ozone air quality in South Korea is determined mainly by year-
7 round regional background contributions (peak in spring). With some differences in details,
8 the results from the two studies are qualitatively similar arguing high springtime
9 background ozone value. One unique aspect of our modeling study is demonstrations of
10 the impact of emissions in Seoul on Gangwon-do, causing slight ozone decrease in
11 Gangwon-do with zero-Seoul emissions from surface to 2 km in May 2016. Our study
12 highlights the diverse impacts of surface emission changes (over China or Seoul) on
13 downwind ozone at different altitudes (Figure 11).

14 Colombi et al. (2023) performed an analysis on the effect of precursor changes on
15 observed surface ozone increases in South Korea. A main difference between Colombi et
16 al. (2023) and our study is the period of the study and whether it focuses on the surface
17 ozone or vertical sensitivity explaining ozone variability at different locations in South
18 Korea. Our study investigated surface ozone and ozone at various altitudes to consider the
19 transport within and above the boundary layer between China and South Korea. Colombi
20 et al. (2023) analyzed the surface ozone and NO₂ concentrations mainly over the Seoul

1 Metropolitan Area from 2015 to 2019. The increase of ozone was mostly attributed to
2 decrease in NO₂ for the studied period.

3 Both Lee and Park (2022) and Colombi et al. (2023) indicated high background
4 ozone concentration external to East Asia (or South Korea), suggesting difficulty of
5 achieving ozone standards. Our study agrees to this point. Probably one different message
6 is that reducing emissions of NO_x and VOC here and there all together have positive
7 impacts on reducing ozone downwind. For example, emission reductions associated with
8 the COVID-19 would lead to decrease of ozone at most sites over South Korea in spring.

9

10 **4. Conclusions**

11 We conducted a study on the spatiotemporal variability of surface ozone in 7 cities, 9
12 provinces, and 2 background sites in South Korea from 2001 to 2021. The 4th highest
13 maximum daily 8-hour average (MDA8) ozone concentrations showed an increasing trend
14 in all cities, most provinces, and background sites during this period, with a yearly increase
15 of 1-2 ppb. After 2010, these concentrations reached approximately 70 ppb or higher. If the
16 US EPA National Ambient Air Quality Standards were applied, most of the monitoring sites
17 in South Korea would have been considered nonattainment areas for the past decade.

18 Ozone exceedances in this study were defined as the ratio of data with
19 concentrations exceeding 70 ppb to the total data, which aligns with the US EPA standard.
20 In Seoul, Incheon, and Gyeonggi-do, ozone exceedances were more frequent in summer

1 than in spring. However, the opposite trend was observed in Daejeon, Gwangju, Jeollanam-
2 do, Gyeongsangbuk-do, Gangwon-do, Jeju Island and the background site Gosung and
3 Ulleung Island. In other areas, the frequencies of exceedances were similar between spring
4 and summer. The majority of ozone exceedances occurred between 16-19 LT (4-7 PM).
5 Interestingly, exceedances also occurred frequently at night in background sites such as
6 Gosung and Ulleung Island, indicating a strong influence of long-range transport on surface
7 ozone levels in these locations.

8 Ozone exceedances increased from period P1 (2002-2010) to period P2 (2011-2019)
9 across all observation sites in South Korea during spring and summer. Overall, NO₂
10 concentrations showed declining trends from 2001 to 2021, but significant and relatively
11 large decreases were only evident after the mid 2010s. NO₂ concentrations for P1 and P2
12 were similar and increase of CO/VOC concentrations between the two periods were not
13 detected or reported. Therefore, it is not clear what drove increase of ozone exceedances
14 over South Korea from P1 to P2. We observed significant reductions in ozone exceedances
15 across all monitoring sites in South Korea during the spring of the COVID-19 pandemic
16 (period P3, 2020-2021), which was attributed to decreased anthropogenic activities and
17 subsequent lower emissions in both China and South Korea. We conducted sensitivity tests
18 using a regional chemical model to investigate the impact of emission changes on ozone
19 pollution in South Korea for a limited period in spring. The results suggest that reductions
20 in Chinese NO_x emissions as well as VOC emissions can contribute to the improvement of

1 ozone pollution in South Korea. These findings provide valuable insights for future efforts
2 to address ozone pollution in South Korea and emphasize the need for further research to
3 project air quality and prioritize actions for the next decade or so.

4 In the future, employing multidecadal mathematical modeling on a local to global
5 scale in both hindcast and forecast modes would be beneficial for better understanding
6 ozone trends in South Korea. Additionally, reliable VOC observations and conducting
7 intensive field campaigns, similar to the KORUS-AQ 2016, would provide crucial
8 information to unravel the complexities of ozone chemistry in this region and facilitate the
9 careful monitoring of changes in atmospheric composition relevant to ozone.

10

11 **Code/Data availability**

- 12 • The surface monitor data for South Korea can be downloaded from
13 <https://www.airkorea.or.kr/web/>.
- 14 • Korea Expressway Corporation transit data: Daily traffic counts using highway,
15 available at: <http://data.ex.co.kr/portal/>, last access: 31 December 2022.
- 16 • KORUS-AQ data: NASA/LARC/SD/ASDC. (2022). KORUS-AQ Aircraft Merge
17 Data Files [Data set]. NASA Langley Atmospheric Science Data Center DAAC.
18 Retrieved from
19 https://doi.org/10.5067/ASDC/SUBORBITAL/KORUSAQ_Merge_Data_1
- 20 • NASA TROPOMI NO₂ columns are available at

1 [https://disc.gsfc.nasa.gov/datasets/TROPOMI_MINDS_NO2_1.1/summary?keyw](https://disc.gsfc.nasa.gov/datasets/TROPOMI_MINDS_NO2_1.1/summary?keywords=tropomi%20no2)
2 [ords=tropomi%20no2.](https://disc.gsfc.nasa.gov/datasets/TROPOMI_MINDS_NO2_1.1/summary?keywords=tropomi%20no2)

3 • KNMI TROPOMI NO₂ columns are available at

4 [https://disc.gsfc.nasa.gov/datasets/S5P_L2_NO2_HiR_2/summary?keywords](https://disc.gsfc.nasa.gov/datasets/S5P_L2_NO2_HiR_2/summary?keywords=tropomi%20no2)
5 [=tropomi%20no2.](https://disc.gsfc.nasa.gov/datasets/S5P_L2_NO2_HiR_2/summary?keywords=tropomi%20no2)

6 • CAM-Chem (CESM) code is available at

7 [https://www.cesm.ucar.edu/models/cesm2/release_download.html.](https://www.cesm.ucar.edu/models/cesm2/release_download.html)

8 • WRF-Chem model can be downloaded from

9 [https://www2.mmm.ucar.edu/wrf/users/download/get_sources.html.](https://www2.mmm.ucar.edu/wrf/users/download/get_sources.html)

10

11 **Author contribution**

12 SWK initiates, designs, analyzes surface monitor data, and writes the manuscript, KMK,
13 SHS, and SWK design and conduct WRF-Chem model runs, JYJ, JYJ, and SWK design
14 and conduct CAM-Chem model runs, SHS processes the airkorea data, YSP and SHS
15 process, analyze, and visualize TROPOMI data, and YSP and JYJ collect and analyze the
16 highway traffic data. All authors edit the manuscript.

17

18 **Competing interests**

19 Authors declare no competing interests.

20

1 **Acknowledgements**

2 This subject is also supported by the National Research Foundation of Korea (NRF) grant
3 funded by the Korea government (MSIT) (No. 2020R1A2C2014131). The first author also
4 acknowledges support from NRF-2018R1A5A1024958. KMA and KISTI supercomputing
5 center support computing resources used (KSC-2021-RND-0040). We thank Louisa
6 Emmons for helping with setting CAM-Chem. The National Center for Atmospheric
7 Research (NCAR) is sponsored by the National Science Foundation (NSF)
8 (NNX16AD96G). We also thanks to the KORUS-AQ science team for producing the
9 extensive number of observations from ground to aircraft.

1 **References**

2 Apple 2020 COVID 19 mobility trends reports—Apple available
3 at: <https://covid19.apple.com/mobility>, last accessed April 2022.

4 Bauwens, M., Compernelle, S., Stavrakou, T., Müller, J.-F., van Gent, J., Eskers, H., Levelt,
5 P. F., van der A, R., Veefkind, J. P., Vlietinck, J., Yu, H., and Zehner, C.: Impact of
6 Coronavirus outbreak on NO₂ pollution assessed using TROPOMI and OMI
7 observations, *Geophys. Res. Lett.*, 47, e2020GL087978,
8 <https://doi.org/10.1029/2020GL087978>, 2020.

9 Boersma, K. F., Eskes, H. J., Richter, A., De Smedt, I., Lorente, A., Beirle, S., van Geffen,
10 J. H. G. M., Zara, M., Peters, E., Van Roozendaal, M., Wagner, T., Maasakkers, J. D.,
11 van der A, R. J., Nightingale, J., De Rudder, A., Irie, H., Pinardi, G., Lambert, J.-C.,
12 and Compernelle, S. C.: Improving algorithms and uncertainty estimates for satellite
13 NO₂ retrievals: results from the quality assurance for the essential climate variables
14 (QA4ECV) project, *Atmos. Meas. Tech.*, 11, 6651–6678, [https://doi.org/10.5194/amt-](https://doi.org/10.5194/amt-11-6651-2018)
15 [11-6651-2018](https://doi.org/10.5194/amt-11-6651-2018), 2018.

16 Buchholz, R. R., Emmons, L. K., Tilmes, S., & The CESM2 Development Team (2019),
17 CESM2.1/CAM-chem Instantaneous Output for Boundary Conditions. UCAR/NCAR
18 - Atmospheric Chemistry Observations and Modeling Laboratory. Lat: -5 to 45, Lon:
19 75 to 145, April 2016 – June 2016, Accessed 1 Oct 2019,
20 <https://doi.org/10.5065/NMP7-EP60>.

21 Chang, K.-L., Schultz, M. G., Lan, X. McClure-Begley, A., Petropavlovskikh, I., Xu, X.,
22 and Ziemke, J. R.: Trend detection of atmospheric time series: Incorporating
23 appropriate uncertainty estimates and handling extreme events, *Elem. Sci.*
24 *Anthropocene*, 9, 00035, <https://doi.org/10.1525/elementa.2021.00035>, 2021.

1 Colombi, N. K., Jacob, D. J., Yang, L. H., Zhai, S., Shah, V., Grange, S. K., Yantosca, R.
2 M., Kim, S., and Liao, H.: Why is ozone in South Korea and the Seoul metropolitan
3 area so high and increasing?, *Atmos. Chem. Phys.*, 23, 4031–4044,
4 <https://doi.org/10.5194/acp-23-4031-2023>, 2023.

5 Cooper, O., Parrish, D., Stohl, A., Trainer, M., Nédélec, P., Thouret, V., Cammas, J. P.,
6 Oltmans, S. J., Johnson, B. J., Tarasick, D., Leblanc, T., McDermid, I. I. S., Jaffe, D.,
7 Gao, R., Stith, J., Ryerson, T., Aikin, K., Campos, T., Weinheimer, A., and Avery, M.
8 A.: Increasing springtime ozone mixing ratios in the free troposphere over western
9 North America, *Nature*, 463, 344–348, <https://doi.org/10.1038/nature08708>, 2010.

10 Crawford, J. H., Ahn, J.-Y., Al-Saadi, J., Chang, L., Emmons, L. K., Kim, J., Lee, G., Park,
11 J.-H., Park, R. J., Woo, J. H., Song, C.-K., Hong, J.-H., Hong, Y.-D., Lefter, B. L., Lee,
12 M., Lee, T., Kim, S., Min, K.-E., Yum, S. S., Shin, H. J., Kim, Y.-W., Choi, J.-S., Park,
13 J.-S., Szykman, J. J., Long, R. W., Jordan, C. E., Simpson, I. J., Fried, A., Dibb, J. E.,
14 Cho, S., and Kim, Y. P.: The Korea–United States Air Quality (KORUS-AQ) field study,
15 *Elem. Sci. Anthropocene*, 9, 1–27, <https://doi.org/10.1525/elementa.2020.00163>, 2021.

16 Crippa, M., Guizzardi, D., Butler, T., Keating, T., Wu, R., Kaminski, J., Kuenen, J.,
17 Kurokawa, J., Chatani, S., Morikawa, T., Pouliot, G., Racine, J., Moran, M. D., Klimont,
18 Z., Manseau, P. M., Mashayekhi, R., Henderson, B. H., Smith, S. J., Suchyta, H.,
19 Muntean, M., Solazzo, E., Banja, M., Schaaf, E., Pagani, F., Woo, J.-H., Kim, J.,
20 Monforti-Ferrario, F., Pisoni, E., Zhang, J., Niemi, D., Sassi, M., Ansari, T., and Foley,
21 K.: HTAP_v3 emission mosaic: a global effort to tackle air quality issues by quantifying
22 global anthropogenic air pollutant sources, *Earth Syst. Sci. Data Discuss.* [preprint],
23 <https://doi.org/10.5194/essd-2022-442>, in review, 2023.

1 Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J.-
2 F., Marsh, D., Mills, M. J., Tilmes, S., Bardeen, Ch., Buchholz, R. R., Conley, A.,
3 Gettelman, A., Garcia, R., Simpson, I., Blake, D. R., Meinardi, S., and Pétron, G.: The
4 Chemistry Mechanism in the Community Earth System Model version 2 (CESM2), *J.*
5 *Adv. Model. Earth Sy.*, 12, e2019MS001882, <https://doi.org/10.1029/2019MS001882>,
6 2020.

7 Eskes, H., van Geffen J., Boersma, F., Eichmann, K.-U., Apituley, A., Pedernana, M.,
8 Sneep, M., Veeffkind, J. P., and Loyola, D.: Sentinel-5 precursor/TROPOMI level 2
9 product user manual nitrogen dioxide, Royal Netherlands Meteorological Institute, #
10 S5P-KNMI-L2-0021-MA, issue 3.0.0, 27, 2019.

11 Fiore, A. M., Jacob, D. J., Bey. I., Yantosca, R. M., Field, B. D., Fusco, A. C., and
12 Wilkinson, J. G.: Background ozone over the United States in summer: Origin, trend,
13 and contribution to pollution episodes, *J. Geophys. Res.*, 107, D15, 4275,
14 <https://doi.org/10.1029/2001JD000982>, 2002.

15 Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux,
16 C., Coheur, P. F., Cuesta, J., Cuevas, E., Doniki, S., Dufour, G., Ebojje, F., Foret, G.,
17 Garcia, O., Granados-Muñoz, M. J., Hannigan, J. W., Hase, F., Hassler, B., Huang, G.,
18 Hurtmans, D., Jaffe, D., Jones, N., Kalabokas, P., Kerridge, B., Kulawik, S., Latter, B.,
19 Leblanc, T., Le Flochmoën, E., Lin, W., Liu, J., Liu, X., Mahieu, E., McClure-Begley,
20 A., Neu, J. L., Osman, M., Palm, M., Petetin, H., Petropavlovskikh, I., Querel, R.,
21 Rahpoe, N., Rozanov, A., Schultz, M. G., Schwab, J., Siddans, R., Smale, D.,
22 Steinbacher, M., Tanimoto, H., Tarasick, D. W., Thouret, V., Thompson, A. M., Trickl,
23 T., Weatherhead, E., Wespes, C., Worden, H. M., Vigouroux, C., Xu, X., Zeng, G., and
24 Ziemke, J.: Tropospheric Ozone Assessment Report: Present-day distribution and

1 trends of tropospheric ozone relevant to climate and global atmospheric chemistry
2 model evaluation *Elem. Sci. Anthropocene*, 6, 39, <https://doi.org/10.1525/elementa.291>,
3 2018.

4 Gaudel, A., Cooper, O. R., Chang, K.-L., Bourgeois, I., Ziemke, J. R., Strode, S. A., Oman,
5 L. D., Sellitto, P., Nédélec P., Blot, R., Thouret, V., and Granier, C.: Aircraft
6 observations since the 1990s reveal increases of tropospheric ozone at multiple
7 locations across the Northern Hemisphere, *Sci. Adv.*, 6, eaba8272,
8 [doi:10.1126/sciadv.aba8272](https://doi.org/10.1126/sciadv.aba8272), 2020.

9 Ghim, Y. S.: Trends and factors of ozone concentration variations in Korea, *J. Korean Soc.*
10 *Atmos. Environ*, 16, 607-623, <https://pubs.kist.re.kr/handle/201004/10289>, 2000.

11 Google 2020 COVID-19 community mobility report available
12 at: www.google.com/covid19/mobility (Last accessed November 2022).

13 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and
14 Eder, B.: Fully coupled “online” chemistry within WRF model, *Atmos. Environ.*, 39,
15 6957-6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.

16 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates
17 of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases
18 and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210,
19 <https://doi.org/10.5194/acp-6-3181-2006>, 2006.

20 Jacob, D. J., Logan, J. A., and Murti, P. P.: Effect of rising Asian emissions on surface
21 ozone in the United States, *Geophys. Res. Lett.*, 26, 2175-2178,
22 [doi:10.1029/1999gl900450](https://doi.org/10.1029/1999gl900450), 1999.

1 Jaffe, D., Anderson, T., Covert, D., Kotchenruther, R., Trost, B., Danielson, J., Simpson,
2 W., Berntsen, T., Karlsdottir, S., Blake, D., Harris, J., Carmichael, G., Uno, I.:
3 Transport of Asian Air Pollution to North America, *Geophys. Res. Lett.*, 26, 711-714,
4 doi:10.1029/1999GL900100, 1999.

5 Jaffe, D., Price, H., Parrish, D., Glodstein, A., and Harris, J.: Increasing background ozone
6 during spring on the west coast of North America, *Geophys. Res. Lett.*, 30, 1613,
7 <https://doi.org/10.1029/2003GL017024>, 2003.

8 Jaffe, D. A., Cooper, O. R., Fiore, A. M., Henderson, B. H., Tonnesen, G. S., Russell, A.
9 G., Henze, D. K., Langford, A. O., Lin, M., and Moore, T.: Scientific assessment of
10 background ozone over the U.S.: Implications for air quality management, *Elem. Sci.*
11 *Anthropocene*, 6, 56, <https://doi.org/10.1525/elementa.309>, 2018.

12 Jeong, Y., et al.: Influence of ENSO on tropospheric ozone variability in Asia, 2023.
13 submitted.

14 Kim, H., Gil, J., Lee, M., Jung, J., Whitehill, A., Szykman, J., Lee, G., Kim, D.-S., Cho,
15 S., Ahn, J.-Y., Hong, J., and Park, M.-S.: Factors controlling surface ozone in the Seoul
16 Metropolitan Area during the KORUS-AQ Campaign, *Elem. Sci. Anthropocene*, 8, 46,
17 <https://doi.org/10.1525/elementa.444>, 2020.

18 Kim, J., Ghim, Y. S., Han, J.-S., Park, S.-M., Shina, H.-J., Lee, S.-B., Kim, J., Lee, G.:
19 Long-term trend analysis of Korean air quality and its implication to current air quality
20 policy on ozone and PM₁₀, *J. Korean Soc. Atmos. Environ.*, 34, 1-15,
21 <https://doi.org/10.5572/KOSAE.2018.34.1.001>, 2018.

1 Kim, K.-M., et al.: Sensitivity of the WRF-Chem v4.4 ozone, formaldehyde, and their
2 precursor simulations to multiple bottom-up emission inventories over East Asia during
3 the KORUS-AQ 2016 field campaign, 2023. *in preparation*.

4 Kim, Y. P., and Lee G. W.: Trend of Air Quality in Seoul: Policy and Science, *Aerosol Air*
5 *Qual. Res.*, 18, 2141-2156, <https://doi.org/10.4209/aaqr.2018.03.0081>, 2018.

6 Koo, J. H., Kim, J., Lee, Y. G., Park, S. S., Lee, S., Chong, H., Cho, Y., Kim, J., Choi, K.,
7 and Lee, T.: The implication of the air quality pattern in South Korea after the COVID-
8 19 outbreak, *Scientific Reports*, 10, 22462, [https://doi.org/10.1038/s41598-020-80429-](https://doi.org/10.1038/s41598-020-80429-4)
9 4, 2020.

10 Korea Expressway Corporation transit data: Daily traffic counts using highway, available
11 at: <http://data.ex.co.kr/portal/>, last access: 31 December 2020.

12 Lamsal, L. N., Krotkov, N. A., Vasilkov, A., Marchenko, S., Qin, W., Yang, E.-S., Fasnacht,
13 Z., Joiner, J., Choi, S., Haffner, D., Swartz, W. H., Fisher, B., and Bucsela, E.: Ozone
14 Monitoring Instrument (OMI) Aura nitrogen dioxide standard product version 4.0 with
15 improved surface and cloud treatments, *Atmos. Meas. Tech.*, 14, 455–479,
16 <https://doi.org/10.5194/amt-14-455-2021>, 2021.

17 Lamsal, Lok N., Nickolay A. Krotkov, Sergey V. Marchenko, Joanna Joiner, Luke Oman,
18 Alexander Vasilkov, Bradford Fisher, Wenhan Qin, Eun-Su Yang, Zachary Fasnacht,
19 Sungeon Choi, Peter Leonard, and David Haffner (2022), TROPOMI/S5P NO2
20 Tropospheric, Stratospheric and Total Columns MINDS 1-Orbit L2 Swath 5.5 km x 3.5
21 km, NASA Goddard Space Flight Center, Goddard Earth Sciences Data and
22 Information Services Center (GES DISC), Accessed: [2023-03-20],
23 10.5067/MEASURES/MINDS/DATA203

24 Langford, A. O., Alvarez II, R. J., Brioude, J., Fine, R., Gustin, M. S., Lin, M. Y.,

1 Marchbanks, R. D., Pierce, R. B., Sandberg, S. P., Senff, C. J., Weickmann, A. M., and
2 Williams, E. J.: Entrainment of stratospheric air and Asian pollution by the convective
3 boundary layer in the southwestern U.S., *J. Geophys. Res.*, 122, 1312-1337,
4 doi.org/10.1002/2016JD025987, 2017.

5 Lee, G. W., Park, J. H., Kim D. G., Koh, M. S., Lee, M., Han, J. S., and Kim, J. C.: Current
6 status and future directions of tropospheric photochemical ozone studies in Korea, *J.*
7 *Korean Soc. Atmos. Environ.*, 36, 419-441, <https://doi:10.5572/KOSAE.2020.36.4.419>,
8 2020.

9 Lee, H.-M., and Park R. J.: Factors determining the seasonal variation of ozone air quality
10 in South Korea: Regional Background versus Domestic Emission Contributions,
11 *Environmental Pollution*, 308, 119645, <https://doi.org/10.1016/j.envpo.2022.119645>,
12 2022.

13 Li, K., Jacob, D. J., Liao, H., and Bates, K. H.: Anthropogenic drivers of 2013-2017 trends
14 in summer surface ozone in China, *P. Natl. Acad. Sci. USA*, 116, 422-427,
15 <https://doi.org/10.1073/pnas.1812168116>, 2019.

16 Liang, Q., Jaeglé, L., Hudman, R. C., Turquety, S., Jacob, D. J., Avery, M. A., Browell, E.
17 V., Sachse, G. W., Blake, D. R., Brune, W., Ren, X., Cohen, R. C., Dibb, J. E., Fried,
18 A., Fuelberg, H., Porter, M., Heikes, B. G., Huey, G., Singh, H. B., and Wennberg, P.
19 O.: Summertime influence of Asian pollution in the free troposphere over North
20 America, *J. Geophys. Res.*, 112, D12D11, <https://doi:10.1029/2006JD007919>, 2007.

21 Lin, M., Fiore, A. M., Horowitz, L. W., Cooper, O. R., Naik, V., Holloway, J., Johnson, B.
22 J., Middlebrook, A. M., Oltmans, S. J., Pollack, I. B., Ryerson, T. B., Warner, J. X.,
23 Wiedinmyer, C., Wilson, J., and Wyman, B.: Transport of Asian ozone pollution into

1 surface air over the western United States in spring, *J. Geophys. Res.*, 117, D00V07,
2 <https://doi:10.1029/2011JD016961>, 2012.

3 Lin, M., Fiore, A. M., Horowitz, L. W., Langford, A. O., Oltmans, S. J., Tarasick, D., and
4 Rieder, H. E.: Climate variability modulates western US ozone air quality in spring via
5 deep stratospheric intrusions, *Nature Communications*, 6, 7105,
6 <https://doi.org/10.1038/ncomms8105>, 2015.

7 National Research Council (1991), Rethinking the ozone problem in urban and regional air
8 pollution, *National Academic Press*, 500pp.

9 Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D.,
10 Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., von
11 Schneidemesser, E., Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone
12 and its precursors from the urban to the global scale from air quality to short-lived
13 climate forcer, *Atmos. Chem. Phys.*, 15, 8889–8973, [https://doi.org/10.5194/acp-15-](https://doi.org/10.5194/acp-15-8889-2015)
14 [8889-2015](https://doi.org/10.5194/acp-15-8889-2015), 2015.

15 Oh, I.-B., Kim, Y.-K., Hwang, M.-K., Kim, C.-H., Kim, S., and Song, S.-K.: Elevated ozone
16 layers over the Seoul Metropolitan Region in Korea: Evidence for long-range ozone
17 transport from Eastern China and its contribution to surface concentrations, *Journal of*
18 *Applied Meteorology and Climatology*, 49, 203-220,
19 <https://doi.org/10.1175/2009JAMC2213.1>, 2010.

20 Park, R. J., Oak, Y. J., Emmons, L. K., Kim, C.-H., Pfister, G. G., Carmichael, G. R., Saide,
21 P. E., Cho, S.-Y., Kim, S., Woo, J.-H., Crawford, J. H., Gaubert, B., Lee, H.-J., Park,
22 S.-Y., Jo, Y.-J., Gao, M., Tang, B., Stanier, C. O., Shin, S. S., Park, H. Y., Bae, C., and
23 Kim, E.: Multi-model intercomparisons of air quality simulations for the KORUS-AQ

1 campaign, *Elementa*, 9, 00139, <https://doi.org/10.1525/elementa.2021.00139>, 2021.

2 Parrish, D. D., Derwent, R. G., Steinbrecht, W., Stübi, R., VanMalderen, R., Steinbacher,
3 M., Trickl, T., Ries, L., and Xu, X.: Zonal similarity of long-term changes and seasonal
4 cycles of baseline ozone at northern midlatitudes, *J. Geophys. Res.*, 125,
5 e2019JD031908, <https://doi.org/10.1029/2019JD031908>, 2020.

6 Seo, S., Kim, S.-W., Kim, K.-M., Lamsal, L. N., and Jin, H.: Reductions in NO₂
7 concentrations in Seoul, South Korea detected from space and ground-based monitors
8 prior to and during the COVID-19 pandemic, *Environ. Res. Commun.*, 3, 051005,
9 <https://doi.org/10.1088/2515-7620/abed92>, 2021.

10 Shi, X., and Brasseur, G. P.: The response in air quality to the reduction of Chinese
11 economic activities during the COVID-19 outbreak. *Geophys. Res. Lett.*, 47,
12 e2020GL088070, <https://doi.org/10.1029/2020GL088070>, 2020.

13 Steinbrecht, W., Kubistin, D., Plass-Dülmer, C., Davies, J., Tarasick, D. W., von der
14 Gathen, P., Deckelmann, H., Jepsen, N., Kivi, R., Lyall, N., Palm, M., Notholt, J., Kois,
15 B., Oelsner, P., Allaart, M., Pitters, A., Gill, M., van Malderen, R., Delcloo, A. W.,
16 Sussmann, R., Mahieu, E., Servais, C., Romanens, G., Stübi, R., Ancellet, G., Godin-
17 Beekmann, S., Yamanouchi, S., Strong, K., Johnson, B., Cullis, P., Petropavlovskikh,
18 P., Hannigan, J. W., Hernandez, J.-L., Rodriguez, A. D., Nakano, T., Chouza, F.,
19 Leblanc, T., Torres, C., Garcia, O., Röhling, A. N., Schneider, M., Blumenstock, T.,
20 Tully, M., Paton-Walsh, C., Jones, N., Querel, R., Strahan, S., Stauffer, R. M.,
21 Thompson, A. M., Inness, A., Engelen, R., Chang, K.L., and Cooper, O. R.: COVID-
22 19 crisis reduces free tropospheric ozone across the Northern Hemisphere, *Geophys.*
23 *Res. Lett.*, 48, e2020GL091987, <https://doi.org/10.1029/2020GL091987>, 2021.

1 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes,
2 H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R.,
3 Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser,
4 H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission
5 for global observations of the atmospheric composition for climate, air quality and
6 ozone layer applications, *Remote Sens. Environ.*, 120, 70–
7 83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.

8 Wang, W., Parrish, D. D., Li, X., Shan, M., Liu, Y., Mo, Z., Lu, S., Hu, M., Fang, X., Wu,
9 Y., Zeng, L., and Zhang, Y.: Exploring the drivers of the increased ozone production in
10 Beijing in summertime during 2005-2016, *Atmos. Chem. Phys.*, 20, 15617-15633,
11 <https://doi.org/10.5194/acp-20-15617-2020>, 2020.

12 Wang, W., Parrish, D. D., Wang, S., Bao, F., Ni, R., Li, X., Yang, S., Wang, H., Cheng, Y.,
13 and Su, H.: Long-term trend of ozone pollution in China during 2014-2020: distinct
14 seasonal and spatial characteristics and ozone sensitivity, *Atmos. Chem. Phys.*, 22,
15 8935-8949, <https://doi.org/10.5194/acp-22-8935-2022>, 2022.

16 Wasserstein, R.L., Schirm, A. L., and Lazar, N. A.: Moving to a world beyond “ $p < 0.05$ ”,
17 *The American Statistician*, 73, 1-19, <https://doi.org/10.1080/00031305.2019.1583913>,
18 2019.

19 Yeo, M. J., and Kim Y. P.: Long-term trends of surface ozone in Korea, *Journal of Cleaner*
20 *Production*, 294, 125352, <https://doi.org/10.1016/j.jclepro.2020.125352>, 2021.

21
22

1 **List of Tables**

2

3 Table 1. Trend estimates based on the 4th highest MDA8 O₃ values. The data were acquired
4 from the surface monitoring network (www.airkorea.or.kr). Unit of slope and limit (2 sigma
5 = 2 standard deviation) is ppb yr⁻¹. SNR denotes signal-to-noise ratio defined as the ratio
6 of absolute value of slope to standard deviation. For the use of P-value and SNR, refer to
7 Chang et al. (2021).

8

9 Table 2. Spring and summer ozone concentrations in Korean metropolitan cities and
10 provinces. Both peak time (10-20 LT) and base time (01-06 LT) averages are shown.
11 Differences in concentrations between spring and summer (O₃ spring - O₃ summer) are in the
12 parenthesis. The cities and provinces listed in the table are in counterclockwise order in
13 regards to the South Korean map.

14

15 Table 3. The observed trends of NO₂ concentrations in spring and summer from linear fits
16 of the data covering 2001-2021. The data were acquired from the surface monitoring
17 network (www.airkorea.or.kr). Unit of slope and limit (2 sigma = 2 standard deviation) is
18 ppb yr⁻¹. SNR denotes signal-to-noise ratio defined as the ratio of absolute value of slope
19 to standard deviation. For the use of P-value and SNR, refer to Chang et al. (2021).

20

21 Table 4. The observed trends of CO concentrations in spring and summer from linear fits
22 of the data covering 2001-2021. The data were acquired from the surface monitoring
23 network (www.airkorea.or.kr). Unit of slope and limit (2 sigma = 2 standard deviation) is
24 ppb yr⁻¹. SNR denotes signal-to-noise ratio defined as the ratio of absolute value of slope
25 to standard deviation. For the use of P-value and SNR, refer to Chang et al. (2021).

1 **Figure captions**

2

3 Figure 1. The locations of cities, provinces, and background sites in South Korea. The red,
4 black, and blue color denote city, province, and background site, respectively: Cities – SUL
5 (Seoul), INC (Incheon), DJN (Daejeon), GWJ (Gwangju), BSN (Busan), ULS (Ulsan),
6 DGU (Daegu); Provinces - GGI (Gyeonggi-do), CCB (Chungcheongbuk-do), CCN
7 (Chungcheongnam-do), JLB (Jeollabuk-do), JLN (Jeollanam-do), JEJ (Jeju Island), GSN
8 (Gyeongsangnam-do), GSB (Gyeongsangbuk-do), GWO (Gangwon-do); Background
9 sites - ULL (Ulleung Island), and GSU (Gosung, Gangwon-do).

10

11

12 Figure 2. The trend of the 4th highest daily maximum 8 hours average (MDA8) O₃
13 concentrations in the South Korean metropolitan cities from 2001 to 2021. Only the data
14 for May-September (ozone season) are used. Bars denotes standard deviations among the
15 sites within the city. The slopes (S) and correlation coefficients (r) from linear fits are
16 shown in parentheses. Grey dashed line indicates 70 ppb that is the air quality standard
17 defined by the US Environmental Protection Agency.

18

19 Figure 3. The same as in Figure 2 except for South Korean provinces.

20

21 Figure 4. Ratio of O₃ exceedances in summer to exceedances in spring. The red line
22 indicates an one to one line. X-axis denotes names of cities, provinces, and background
23 sites. Cities – SUL (Seoul), INC (Incheon), DJN (Daejeon), GWJ (Gwangju), BSN (Busan),
24 ULS (Ulsan), DGU (Daegu); Provinces - GGI (Gyeonggi-do), CCB (Chungcheongbuk-do),

1 CCN (Chungcheongnam-do), JLB (Jeollabuk-do), JLN (Jeollanam-do), JEJ (Jeju Island),
2 GSN (Gyeongsangnam-do), GSB (Gyeongsangbuk-do), GWO (Gangwon-do);
3 Background sites - ULL (Ulleung Island), and GSU (Gosung, Gangwon-do). The data for
4 2001-2019 are utilized.

5

6 Figure 5. Diurnal O₃ exceedances. (Top) Seoul area, (middle) secondary cities, (bottom)
7 remote sites. The data for 2001-2019 are utilized.

8

9 Figure 6. The contribution of stratospheric O₃ (O_{3s}) to the O₃ concentrations in each season
10 at surface and 1 km above ground level in South Korea. The plotted values are extracted
11 from the CESMv2.2 results for the entire country.

12

13 Figure 7. (Top) O₃ exceedances (%), (middle) NO₂, and (bottom) CO concentrations in
14 South Korean cities, provinces, and background sites during spring for 2002-2010, 2011-
15 2019, and 2020-2021 (COVID-19). X-axis denotes names of cities, provinces, and
16 background sites. Cities - SUL (Seoul), INC (Incheon), DJN (Daejeon), GWJ (Gwangju),
17 BSN (Busan), ULS (Ulsan), DGU (Daegu); Provinces - GGI (Gyeonggi-do), CCB
18 (Chungcheongbuk-do), CCN (Chungcheongnam-do), JLB (Jeollabuk-do), JLN
19 (Jeollanam-do), JEJ (Jeju Island), GSN (Gyeongsangnam-do), GSB (Gyeongsangbuk-do),
20 GWO (Gangwon-do); Background sites - ULL (Ulleung Island), and GSU (Gosung,
21 Gangwon-do).

22

23 Figure 8. The same as Figure 7 except for summer.

24

1 Figure 9. Differences in TROPOMI tropospheric NO₂ columns between 2019 and 2020 or
2 between 2019 and 2021 (Difference = NO₂ 2020 or 2021 - NO₂ 2019). Unit: molecules cm⁻²

3

4 Figure 10. Differences in the WRF-Chem simulated ozone concentrations ($\Delta O_3 =$
5 O₃_emission reduction case-O₃_control case) at (top) surface and (bottom) 1000 m above
6 ground level. Green to blue colors (yellow to red colors) denotes reduced (increased) ozone
7 concentration due to the emission changes.

8

9 Figure 11. Vertical profiles of ozone from the WRF-Chem model simulations based on
10 various emission scenarios: (top) Seoul, and (bottom) Gosung, Gangwon-do.

11

12

13

14

15

16

17

18

19

20

21

22

23

24

1 Table 1. Trend estimates based on the 4th highest MDA8 O₃ values. The data were acquired
 2 from the surface monitoring network (www.airkorea.or.kr). Unit of slope and limit (2 sigma
 3 = 2 standard deviation) is ppb yr⁻¹. SNR denotes signal-to-noise ratio defined as the ratio
 4 of absolute value of slope to standard deviation. For the use of P-value and SNR, refer to
 5 Chang et al. (2021).

Location		Slope (ppb yr ⁻¹)	2-Sigma (ppb yr ⁻¹)	P value	SNR
City	Seoul (SUL)	1.19	0.38	< 0.01	6.23
	Incheon (INC)	1.07	0.37	< 0.01	5.72
	Daejeon (DJN)	1.22	0.49	< 0.01	4.96
	Gwangju (GWJ)	0.98	0.46	< 0.01	4.30
	Busan (BSN)	0.98	0.36	< 0.01	5.47
	Ulsan (ULS)	1.40	0.34	< 0.01	8.14
	Daegu (DGU)	1.12	0.46	< 0.01	4.89
Province	Gyeonggi-do (GGI)	1.26	0.27	< 0.01	9.33
	Chungcheongbuk-do (CCB)	0.79	0.51	< 0.01	3.09
	Chungcheongnam-do (CCN)	1.45	0.47	< 0.01	6.12
	Jeollabuk-do (JLB)	1.83	0.32	< 0.01	11.30
	Jeollanam-do (JLN)	0.08	0.39	0.67	0.41
	Jeju Island (JEJ)	0.66	0.46	< 0.01	2.89
	Gyeongsangnam-do (GSN)	0.83	0.52	< 0.01	3.18
	Gyeongsangbuk-do (GSB)	1.10	0.35	< 0.01	6.32
	Gangwon-do (GWO)	0.67	0.48	< 0.01	2.79

6

1 Table 2. Spring and summer ozone concentrations in Korean metropolitan cities and
 2 provinces. Both peak time (10-20 LT) and base time (01-06 LT) averages are shown.
 3 Differences in concentrations between spring and summer ($O_3_{\text{spring}} - O_3_{\text{summer}}$) are in the
 4 parenthesis. The cities and provinces listed in the table are in counterclockwise order in
 5 regards to the South Korean map.

Location		Peak time	Base time
		Spring / Summer (difference)	Spring / Summer (difference)
City	Seoul (SUL)	34.4 / 35.6 (-1.2)	20.6 / 17.5 (3.1)
	Incheon (INC)	34.6 / 33.1 (1.5)	25.1 / 20.2 (4.9)
	Daejeon (DJN)	41.2 / 37.0 (4.2)	22.8 / 19.1 (3.7)
	Gwangju (GWJ)	39.9 / 35.4 (4.5)	28.5 / 24.0 (4.5)
	Busan (BSN)	40.3 / 34.2 (6.1)	30.3 / 22.4 (7.9)
	Ulsan (ULS)	38.7 / 33.4 (5.3)	25.8 / 18.7 (7.1)
	Daegu (DGU)	39.6 / 37.6 (2.0)	24.0 / 19.6 (4.4)
Province	Gyeonggi-do (GGI)	37.5 / 38.5 (-1.0)	20.8 / 18.0 (2.8)
	Chungcheongbuk-do (CCB)	42.1 / 39.4 (2.7)	24.8 / 20.6 (4.2)
	Chungcheongnam-do (CCN)	41.3 / 37.7 (3.6)	29.6 / 23.1 (6.5)
	Jeollabuk-do (JLB)	38.3 / 35.0 (3.3)	26.7 / 23.6 (3.1)
	Jeollanam-do (JLN)	42.5 / 35.1 (7.4)	33.0 / 24.1 (9.4)
	Jeju Island (JEJ)	49.0 / 35.0 (14.0)	43.7 / 29.2 (14.5)
	Gyeongsangnam-do (GSN)	44.3 / 40.0 (4.3)	28.9 / 21.9 (7.0)
	Gyeongsangbuk-do (GSB)	45.1 / 38.0 (7.1)	28.5 / 20.6 (7.9)
	Gangwon-do (GWO)	45.6 / 39.5 (6.1)	31.5 / 24.0 (7.5)

6
7
8
9
10
11
12
13

1 Table 3. The observed trends of NO₂ concentrations in spring and summer from linear fits
 2 of the data covering 2001-2021. The data were acquired from the surface monitoring
 3 network (www.airkorea.or.kr). Unit of slope and limit (2 sigma = 2 standard deviation) is
 4 ppb yr⁻¹. SNR denotes signal-to-noise ratio defined as the ratio of absolute value of slope
 5 to standard deviation. For the use of P-value and SNR, refer to Chang et al. (2021).


Stations		NO ₂ Spring (Summer)			
		Slope (ppb yr ⁻¹)	2 Sigma (ppb yr ⁻¹)	P-value	SNR
City	Seoul (SUL)	-0.77 (-0.72)	0.22 (0.15)	< 0.01 (< 0.01)	6.94 (9.57)
	Incheon (INC)	-0.37 (-0.50)	0.22 (0.17)	< 0.01 (< 0.01)	3.36 (5.88)
	Daejeon (DJN)	-0.10 (-0.12)	0.14 (0.09)	0.21 (0.02)	1.43 (2.53)
	Gwangju (GWJ)	-0.51 (-0.35)	0.15 (0.09)	< 0.01 (< 0.01)	6.94 (7.74)
	Busan (BSN)	-0.64 (-0.49)	0.16 (0.11)	< 0.01 (< 0.01)	8.12 (8.93)
	Ulsan (ULS)	-0.04 (-0.06)	0.23 (0.19)	0.73 (0.51)	0.34 (0.63)
	Daegu (DGU)	-0.65 (-0.51)	0.18 (0.13)	< 0.01 (< 0.01)	7.21 (8.15)
Province	Gyeonggi (GGI)	-0.41(-0.44)	0.22 (0.16)	< 0.01 (< 0.01)	3.80 (5.58)
	Chungcheongbuk (CCB)	-0.18(-0.16)	0.20 (0.15)	0.09 (0.05)	1.82 (2.15)
	Chungcheongnam (CCN)	-0.10(-0.12)	0.15 (0.12)	0.21 (0.08)	1.38 (1.97)
	Jeollabuk (JLB)	-0.17(-0.25)	0.18 (0.14)	0.08 (< 0.01)	1.90 (3.61)
	Jeollanam (JLN)	-0.21(-0.21)	0.16 (0.14)	0.02 (< 0.01)	2.56 (2.95)
	Jeju Island (JEJ)	-0.18(-0.16)	0.20 (0.15)	0.10 (0.04)	1.76 (2.20)
	Gyeongsangnam (GSN)	-0.12(-0.10)	0.17 (0.11)	0.18 (0.08)	1.42 (1.88)
	Gyeongsangbuk (GSB)	-0.76(-0.49)	0.18 (0.13)	< 0.01 (< 0.01)	8.47 (7.74)
	Gangwon (GWO)	-0.16(-0.20)	0.14 (0.10)	0.03 (< 0.01)	2.37 (4.18)

6
7
8
9
10
11
12
13
14
15
16

1 Table 4. The observed trends of CO concentrations in spring and summer from linear fits
 2 of the data covering 2001-2021. The data were acquired from the surface monitoring
 3 network (www.airkorea.or.kr). Unit of slope and limit (2 sigma = 2 standard deviation) is
 4 ppb yr⁻¹. SNR denotes signal-to-noise ratio defined as the ratio of absolute value of slope
 5 to standard deviation. For the use of P-value and SNR, refer to Chang et al. (2021).

Stations		CO Spring (Summer)			
		Slope (ppb yr ⁻¹)	2 Sigma (ppb yr ⁻¹)	P-value	SNR
City	Seoul (SUL)	-7.56 (-5.34)	2.94 (1.66)	< 0.01 (< 0.01)	5.15 (6.44)
	Incheon (INC)	-7.65 (-4.64)	3.62 (2.46)	< 0.01 (< 0.01)	4.23 (3.77)
	Daejeon (DJN)	-15.53 (-9.71)	5.68 (5.56)	< 0.01 (< 0.01)	5.47 (3.49)
	Gwangju (GWJ)	-10.64 (-8.00)	3.60 (3.94)	< 0.01 (< 0.01)	5.91 (4.06)
	Busan (BSN)	-12.32 (-11.05)	3.90 (3.80)	< 0.01 (< 0.01)	6.32 (5.82)
	Ulsan (ULS)	-4.80 (0.75)	5.54 (5.28)	0.10 (0.78)	1.73 (0.28)
	Daegu (DGU)	-23.49 (-19.87)	5.50 (5.30)	< 0.01 (< 0.01)	8.54 (7.50)
Province	Gyeonggi (GGI)	-14.50 (-8.82)	2.18 (1.54)	< 0.01 (< 0.01)	13.30 (11.42)
	Chungcheongbuk (CCB)	-17.68 (-6.49)	6.70 (3.92)	< 0.01 (< 0.01)	5.28 (3.31)
	Chungcheongnam (CCN)	-20.95 (-9.33)	8.32 (4.62)	< 0.01 (< 0.01)	5.04 (4.04)
	Jeollabuk (JLB)	-21.33 (-15.07)	5.88 (4.34)	< 0.01 (< 0.01)	7.26 (6.95)
	Jeollanam (JLN)	-5.86 (-5.32)	4.40 (4.60)	0.02 (0.03)	2.66 (2.31)
	Jeju Island (JEJ)	-10.74 (-6.95)	5.00 (5.64)	< 0.01 (0.02)	4.30 (2.46)
	Gyeongsangnam (GSN)	-6.76 (-3.92)	4.44 (3.58)	< 0.01 (0.04)	3.04 (2.19)
	Gyeongsangbuk (GSB)	-27.54 (-17.48)	9.00 (6.64)	< 0.01 (< 0.01)	6.12 (5.27)
	Gangwon (GWO)	-15.31 (-9.03)	4.34 (4.16)	< 0.01 (< 0.01)	7.05 (4.34)

6
 7
 8




1

2 Figure 1. The locations of cities, provinces, and background sites in South Korea. The red,
 3 black, and blue color denote city, province, and background site, respectively: Cities – SUL
 4 (Seoul), INC (Incheon), DJN (Daejeon), GWJ (Gwangju), BSN (Busan), ULS (Ulsan),
 5 DGU (Daegu); Provinces - GGI (Gyeonggi-do), CCB (Chungcheongbuk-do), CCN
 6 (Chungcheongnam-do), JLB (Jeollabuk-do), JLN (Jeollanam-do), JEJ (Jeju Island), GSN
 7 (Gyeongsangnam-do), GSB (Gyeongsangbuk-do), GWO (Gangwon-do); Background
 8 sites - ULL (Ulleung Island), and GSU (Gosung, Gangwon-do).

9

10

11



1

2 Figure 2. The trend of the 4th highest daily maximum 8 hours average (MDA8) O₃
 3 concentrations in the South Korean metropolitan cities from 2001 to 2021. Only the data
 4 for May-September (ozone season) are used. Bars denote standard deviations among the
 5 sites within the city. The slopes (S) and correlation coefficients (r) from linear fits are
 6 shown in parentheses. Grey dashed line indicates 70 ppb that is the air quality standard
 7 defined by the US Environmental Protection Agency.

8





Figure 3. The same as in Figure 2 except for South Korean provinces.

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18



1

2 Figure 4. Ratio of O₃ exceedances in summer to exceedances in spring. The red line
 3 indicates an one to one line. X-axis denotes names of cities, provinces, and background
 4 sites. Cities – SUL (Seoul), INC (Incheon), DJN (Daejeon), GWJ (Gwangju), BSN (Busan),
 5 ULS (Ulsan), DGU (Daegu); Provinces - GGI (Gyeonggi-do), CCB (Chungcheongbuk-do),
 6 CCN (Chungcheongnam-do), JLB (Jeollabuk-do), JLN (Jeollanam-do), JEJ (Jeju Island),
 7 GSN (Gyeongsangnam-do), GSB (Gyeongsangbuk-do), GWO (Gangwon-do);
 8 Background sites - ULL (Ulleung Island), and GSU (Gosung, Gangwon-do). The data for
 9 2001-2019 are utilized.

10

11

12

13

14

15

16

17

18

19


20

21


22

23


24



1



2



3

4 Figure 5. Diurnal O₃ exceedances. (Top) Seoul area, (middle) secondary cities, (bottom)
 5 remote sites. The data for 2001-2019 are utilized.


6

7

8

9

10



1

2 Figure 6. The contribution of stratospheric O₃ (O_{3s}) to the O₃ concentrations in each season
 3 at surface and 1 km above ground level in South Korea. The plotted values are extracted
 4 from the CESMv2.2 results for the entire country.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19


20

21

22

23

24



1
2 Figure 7. (Top) O₃ exceedances (%), (middle) NO₂, and (bottom) CO concentrations in
3 South Korean cities, provinces, and background sites during spring for 2002-2010, 2011-
4 2019, and 2020-2021 (COVID-19). X-axis denotes names of cities, provinces, and
5 background sites. Cities - SUL (Seoul), INC (Incheon), DJN (Daejeon), GWJ (Gwangju),
6 BSN (Busan), ULS (Ulsan), DGU (Daegu); Provinces - GGI (Gyeonggi-do), CCB
7 (Chungcheongbuk-do), CCN (Chungcheongnam-do), JLB (Jeollabuk-do), JLN
8 (Jeollanam-do), JEJ (Jeju Island), GSN (Gyeongsangnam-do), GSB (Gyeongsangbuk-do),
9 GWO (Gangwon-do); Background sites - ULL (Ulleung Island), and GSU (Gosung,
10 Gangwon-do).






Figure 8. The same as Figure 7 except for summer.

- 1
- 2
- 3
- 4
- 5




1
2
3
4
5

Figure 9. Differences in TROPOMI tropospheric NO₂ columns between 2019 and 2020 or between 2019 and 2021 (Difference = NO₂ 2020 or 2021 - NO₂ 2019). Unit: molecules cm⁻²





1
2



3

4 Figure 10. Differences in the WRF-Chem simulated ozone concentrations ($\Delta O_3 =$
 5 $O_3_{\text{emission reduction case}} - O_3_{\text{control case}}$) at (top) surface and (bottom) 1000 m above
 6 ground level. Green to blue colors (yellow to red colors) denotes reduced (increased) ozone
 7 concentration due to the emission changes.



1
 2 Figure 11. Vertical profiles of ozone from the WRF-Chem model simulations based on
 3 various emission scenarios: (top) Seoul, and (bottom) Gosung, Gangwon-do.
 4