

Production of aerosol containing ice nucleating particles (INPs) by fast growing phytoplankton

Daniel C. O. Thornton¹, Sarah D. Brooks², Elise K. Wilbourn¹, Jessica Mirrielees², Alyssa N. Alsante¹, Gerardo Gold-Bouchot¹, Andrew Whitesell^{1,3}, Kiana McFadden^{2,4}

¹Department of Oceanography, Texas A & M University, O & M Building, College Station, 77843, Texas, United States

²Department of Atmospheric Sciences, Texas A & M University, O & M Building, College Station, 77843, Texas, United States

³North Carolina State University, Raleigh, 27695, North Carolina, United States

⁴Jackson State University, Jackson, 39217, Mississippi, United States

Correspondence to: Daniel C. O. Thornton (dthornton@tamu.edu) and Sarah D. Brooks (sbrooks@tamu.edu)

Abstract. Sea spray aerosol contains ice nucleating particles (INPs), which affect the formation and properties of clouds. Here, we show that aerosols emitted from fast growing marine phytoplankton produce effective immersion INPs, which nucleate at temperatures significantly warmer than the atmospheric homogeneous freezing (-38.0 °C) of pure water. Aerosol sampled over phytoplankton cultures grown in a marine aerosol reference tank (MART) induced nucleation and freezing at temperatures as high as -15.0 °C during exponential phytoplankton growth. This was observed in monospecific cultures representative of two major groups of phytoplankton: a cyanobacterium (*Synechococcus elongatus*) and a diatom (*Thalassiosira weissflogii*). Ice nucleation occurred at colder temperatures (-28.5 °C and below), which were not different from the freezing temperatures of procedural blanks, when the cultures were in the stationary or death phases of growth. Ice nucleation at warmer temperatures was associated with relatively high values of the maximum quantum yield of photosystem II (Φ_{PSII}), an indicator of the physiological status of phytoplankton. High values of Φ_{PSII} indicate the presence of cells with efficient photochemistry and greater potential for photosynthesis. For comparison, field measurements in the North Atlantic Ocean showed that high net growth rates of natural phytoplankton assemblages were associated with marine aerosol that acted as effective immersion INPs at relatively warm temperatures. Data were collected over 4 days at a sampling station maintained in the same water mass as the water column stabilized after deep mixing by a storm. Phytoplankton biomass and net phytoplankton growth rate (0.56 day⁻¹) were greatest over the 24 hours preceding the warmest mean ice nucleation temperature (-25.5 °C). Collectively, our laboratory and field observations indicate that phytoplankton physiological status is a useful predictor of effective INPs, and more reliable than biomass or taxonomic affiliation. Ocean regions associated with fast phytoplankton growth, such as the North Atlantic during the annual spring bloom, may be significant sources of atmospheric INPs.

1 Introduction

Atmospheric aerosols significantly influence the Earth's radiative budget and climate through direct (scattering and absorption of light) and indirect effects (modifying cloud formation, properties and lifetimes) (Seinfeld et al., 2016). Clouds form through interactions between aerosol and water vapor, with cloud condensation nuclei (CCN) catalysing the formation of water droplets and ice nucleating particles (INP) catalysing the formation of ice crystals (Andrea and Rosenfeld, 2008; Hoose and Mohler et al., 2012; Hudson and Noble, 2021). Ice crystals are present in clouds at all latitudes; over the equatorial and high-latitude ocean, the

ice water path in clouds is greater than the liquid water path (Boucher et al., 2013). In the absence of INPs, homogenous freezing of water in the atmosphere occurs at temperatures below -38 °C (Kanji et al., 2017; Vali et al., 2015). INPs catalyse the freezing of water at warmer temperatures through several heterogeneous freezing modes: immersion mode in which the supercooled liquid water droplet surrounds the INP, contact mode in which freezing is initiated by an INP which has collided with the surface of a supercooled liquid water droplet, or deposition mode in which ice grows on an INP from water vapor (Kanji et al., 2017; Vali, 1985; Vali et al., 2015).

A major limitation in our ability to understand and make accurate predictions about the climate system are uncertainties associated with aerosol and cloud properties (Boucher et al., 2013). The latest generation of climate models, particularly through the Coupled Model Intercomparison project (CMIP6), indicate that the climate system is even more sensitive to cloud dynamics than previously thought (Palmer, 2020; Zelinka et al., 2020). Warming decreases cloud water content and coverage more than previously predicted, resulting in a positive feedback and more warming (Palmer, 2020; Schneider et al., 2019; Zelinka et al., 2020). These examples emphasize the role of clouds in climate uncertainty and our need to better understand cloud microphysics, to enable clouds to be better parameterized in global climate models (GCMs).

The ocean covers 71 % of the Earth's surface and is a major source of atmospheric aerosol through the production of sea spray aerosol (SSA) (Lohmann et al., 2016). Containing a significant fraction of organic matter (O'Dowd et al., 2004; Gant and Meskhidze, 2013; Quinn et al., 2014), SSA is a chemical link between atmosphere and oceans, and plays a role in climate (Cochran et al., 2017). There is a need to not only characterize SSA in the atmosphere, but also the biogeochemical processes in the ocean that affect the precursors of SSA (Brooks and Thornton, 2018; Mansour et al., 2020; Prather et al., 2013; Wilbourn et al., 2020; Quinn et al., 2019; Saliba et al., 2019, 2020; Sanchez et al., 2021; Twohy et al., 2021). Number concentrations of INP from marine sources are poorly constrained, but generally considered to be orders of magnitude lower than number concentrations of INP from terrestrial sources, particularly mineral dust (Beall et al. 2022; DeMott et al., 2016). Observations indicate that number concentrations of INP over the remote ocean are low (0.38 to 4.6 INP m⁻³ of air over the Southern Ocean nucleating at -20 °C) (McCuskey et al. 2018). Nevertheless, marine sources of INP are important over the remote oceanic areas, such as the Southern Ocean, due to the absence of terrestrial INP and large surface area of the Earth covered by the ocean (Zhao et al., 2021).

The properties of SSA have been coupled to phytoplankton blooms, indicating potential relationships between ecological processes in the epipelagic ocean, the formation of SSA, and the properties of the clouds that form on marine aerosol (O'Dowd et al., 2004; Quinn et al., 2019; Saliba et al., 2019; Mansour et al., 2020; Croft et al., 2021). Phytoplankton assemblage composition, physiological status, and interactions with other organisms are all potentially important in determining whether the organic matter produced by phytoplankton produces SSA containing effective INPs (Wilbourn et al., 2020). O'Dowd et al. (2015) found that SSA was more enriched with organic matter during bloom collapse, which was hypothesized to be due to the release of large amounts of organic matter into the water by dying cells. Exudates from phytoplankton have been proposed as a source of INPs in the sea surface microlayer (SML) (Wilson et al., 2015; Irish et al., 2017). Of the carbon fixed by phytoplankton photosynthesis, 2 to 50 % is released into the surrounding water as dissolved organic matter (DOM) (Thornton, 2014). Potentially, this DOM contains representatives from all the major groups of biological compounds (carbohydrates, proteins, lipids and nucleic acids) (Thornton, 2014). The amount of DOM released by phytoplankton increases when cells are stressed by environmental factors such as nutrient limitation, stressors that are associated with bloom collapse (Thornton, 2002; 2014).

Diatoms (Alpert et al., 2011a; Knopf et al., 2010; Wilson et al., 2015), coccolithophores (Alpert et al., 2011b), cyanobacteria (Wilbourn et al., 2020; Wolf et al., 2019), and chlorophytes (Alpert et al., 2011b) are all known sources of effective INP. There is insufficient data to determine if taxonomic affiliation and associated functional traits (e.g. chemical composition) determine whether phytoplankton act as INPs in greater quantities or warmer temperatures. It is known that changes in

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0.5"

Formatted: Superscript

Formatted: Not Highlight

80 physiological status affect resource allocation and therefore the chemical composition of phytoplankton (Klausmeier et al., 2004; Van Mooy et al., 2009; Dyhrman et al., 2012).

The objective of this work was to evaluate the link between growth and the physiology of representative phytoplankton with the properties of primary marine aerosols, focusing on factors that determine whether the aerosol are effective INPs. Growth experiments under controlled conditions were conducted with representative phytoplankton taxa in a marine aerosol reference tank (MART). Aerosol were collected onto aluminium foil substrates and their ability to act as INPs in the immersion mode was measured using our well-established ice microscope technique (Fornea et al., 2006). These data were compared with ice nucleation measurements on *in situ* aerosol collected in the North Atlantic during the annual spring bloom.

2 Methods

2.1 Growth of phytoplankton in a marine aerosol reference tank (MART)

Identifying the method which produces the most realistic primary aerosols in a laboratory is a challenge previously addressed by a number of research groups (Fuentes et al., 2010a, 2010b; Prather et al., 2013; Sellegri et al., 2006; Stokes et al., 2013). Since using a bubble plume from a plunging jet of water (Fuentes et al., 2010a; Sellegri et al., 2006) has been shown to be the best method, we used a Marine Aerosol Reference Tank (MART) featuring a plunging jet. The MART provided a controlled environment for growing phytoplankton in batch culture and a closed headspace where atmospheric conditions above the ocean were simulated. It is a 227 liter acrylic aquarium tank (52 cm wide x 122 cm long x 33 cm deep), closed with a neoprene gasket and sheet of 5 mm acrylic. The water within the tank was stirred using three magnetic stir bars (6 cm long, at a rate of approximately 180 rpm) to maintain the phytoplankton in suspension (Fig. 1). Illumination was provided by four 112 cm LED lights (Fluence Biengineering) simulating daylight. Photon flux density (PFD) was varied via a 1-10V input dimmer. PFD within the tank was measured using a Photosynthetically Active Radiation (PAR) sensor (LiCOR, Inc.). Cultures were grown at a PFD of 170 $\mu\text{mol m}^{-2} \text{s}^{-1}$ on a 14 h light: 10 hour dark cycle. Temperature on the outer surface of the tank and room air were monitored continuously using calibrated thermocouples in Labview (National Instruments), and maintained at 27 to 28 °C. Prior to introduction to the MART, phytoplankton were grown in artificial seawater (ASW) (Harrison et al., 1980; Berges et al., 2001) supplemented with L1 trace metals and vitamins (Guillard and Hargraves, 1993). The ASW was made with high purity analytical grade salts. Nevertheless, the large mass of salts in artificial seawater represents a source of potential contamination. To reduce this possibility, sodium chloride was combusted for 6 hours at 500 °C to remove organic contamination. This precaution was taken given the large amount of sodium chloride in the medium (21.19 g L⁻¹). For other salts, notably the hydrated salts, combustion to remove potential organic contamination was not an option as it would have changed the composition and solubility of the salts. The tank was inoculated with 0.63 L of exponentially growing culture (1 % of total culture volume), which was added to the 63 L of medium in the tank. *Thalassiosira weissflogii* (CCMP 1051) was grown as a representative diatom, and in separate experiment, *Synechococcus elongatus* (CCMP 1379) was grown as a representative cyanobacterium. Macronutrient concentrations were 60 μM NaNO_3 , 20 μM NaH_2PO_4 , and 60 μM Na_2SiO_3 in the *T. weissflogii* culture and 100 μM NaNO_3 , 50 μM NaH_2PO_4 , and 60 μM Na_2SiO_3 in the culture of *S. elongatus*. Nutrients (nitrate, phosphate, silicate, trace metals, and vitamins) were added on the morning of the phytoplankton addition to the tank. The nutrients were mixed into the seawater using the stir bars in the tank (Fig. 1) for approximately an hour before the phytoplankton were added. The maximum potential biomass in the tank was determined by nutrient availability. A low N:P ratio compared with the Redfield Ratio of 16:1 (Redfield, 1958) meant that the cultures were likely to become N-limited rather than P-limited. Silicate was added to both cultures for consistency, although it is not required by cyanobacteria. In both experiments, trace elements and vitamins were added as in Berges et al. (2001).

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: 1.5 lines

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), 10 pt

Formatted: Font: (Default) +Body (Times New Roman)

2.1.1 Sample collection and aerosol generation

Sample collection from the water occurred via a 2 cm port cut into the acrylic lid on the top of the tank. The port was sealed with a silicone rubber bung between water sampling events, which occurred in the morning (1.5 - 2 hours after the lights were turned on to simulate daylight). Water was removed from the tank with a 60-mL syringe that was acid-washed in 10% HCl daily and rinsed three times with ultra-high purity (UHP) prior to sampling.

Stirring of the water was switched off during aerosol generation. Immediately prior to aerosol generation, the MART was flushed with HEPA filtered air at 5 Lpm until the number concentration of particles was less than $5 \times 10^{-3} \text{ L}^{-1}$. The air was turned off, and a peristaltic pump (Watson-Marlow 550Dz) was switched on, providing water flow to a diffuser located $3.56 \times 10^{-13} \text{ m}$ above the surface of the water. The diffuser was slot shaped, creating an approximately laminar waterfall that impacted the surface of the water. Upon impaction, the waterfall entrains air, generating bubbles which rise to the surface and burst, generating aerosol. The peristaltic pump sent 70 L of water through the waterfall at the top of the tank, which took 30 minutes. Aerosol measurements were taken after a 5 minute equilibration period. Aerosol sampling occurred from a PVC pipe attached to 1.5 cm diameter tubing.

The air was dried and passed through a 46 cm long glass mixing tube, with a 0.64 cm I.D. entrance ports on one end, and 0.64 cm I.D. exit points on the opposite end. Total aerosol concentration (in the approximate range of ~ 0.01 to ~1 μm diameter) was measured with a water-based Condensation Particle Counter (CPC, TSI, Inc. Model 3786). In addition, a Portable Aerosol Spectrometer (GRIMM 1.108) was used to measure the size-resolved number concentration of 0.3 to 20 μm diameter aerosol. At the beginning of each sampling period, a Gilibrator flowmeter was used to confirm a positive flow outward through the HEPA exhaust filter.

For offline ice nucleation measurements, size-sorted aerosol samples were collected on combusted aluminium foil substrates inside a PIXE cascade impactor with the following stages: L1, 3, and 6, corresponding to 8, 1 and 0.06 μm diameter, respectively (Fig. 1). Samples collected on L1 impaction stage (0.06-1 μm aerodynamic diameter) were analysed. Aerosol were collected for 2 hours at an air flow of 1 L min^{-1} through the sampler. Samples were stored at $-80 \text{ }^\circ\text{C}$.

Formatted: Indent: First line: 0", Space After: 14 pt

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: English (US)




Figure 1. Marine aerosol reference tank (MART) and associated aerosol sampling. The MART contained a phytoplankton culture (green) and overlying headspace (white). The MART had sampling ports for water sampling (orange) and sampling the headspace above the culture (blue). A peristaltic pump was used to generate a laminar waterfall back into the culture, generating sea spray aerosol (SSA). Light to support phytoplankton growth was provided by LED lamps ($170 \mu\text{mol m}^{-2} \text{s}^{-1}$ on a 14 h light: 10 hour dark cycle) and the culture was stirred using 4x 6 cm long stir bars. Air from the headspace of the MART was dried and relative humidity was monitored (RH meter) before being mixed (46 cm long mixing tube). Total aerosol concentration was measured with a water-based Condensation Particle Counter (CPC). A Portable Aerosol Spectrometer (PAS) was used to measure the size-resolved number concentration of aerosol. Size-sorted aerosol samples were collected on combusted aluminium foil substrates inside a PIXE cascade impactor. Filters (HEPA) and mass-flow controllers (MFC) were positioned before the air pump. Arrows indicate the direction of flow.

2.2 Measurement of ice nucleation

The ice nucleation measurements in this study were conducted through off-line ice nucleation experiments using our custom-built ice microscope apparatus (Collier and Brooks, 2016; Fornea et al., 2009; Wilbourn et al. 2020) which allows us to independently determine the freezing temperature for the same sample multiple times. The ice microscope consists of an Olympus optical microscope (Model BX51M), a digital camera (Q-Imaging Micropublisher 5.0 RTV), and Linkam cooling stage (LTS 350). Temperature control was maintained to within $\pm 0.1 \text{ }^\circ\text{C}$ in the sealed Linkam stage throughout freeze-thaw cycles from $+5$ to $-40 \text{ }^\circ\text{C}$. As described in detail in Fornea et al. (2009) the stage temperature is routinely calibrated by measuring the melting points of *n*-dodecane, *n*-undecane, and *n*-decane (Collier and Brooks, 2016).

An aerosol sample collected on aluminum foil from the L1 stage of the PIXE cascade impactor was placed on a hydrophobic (silanized by immersion in a 1.0% Aquasil solution (Pierce Chemical Company)) glass microscope slide for support. To simulate immersion mode nucleation, a $2.0 \mu\text{L}$ drop of HPLC-grade water (VWR) was micro-pipetted onto the sample containing potential INPs, which were visible as a small circular spot on the aluminum foil. The Linkam stage was sealed and cooled at a rate of $1 \text{ }^\circ\text{C min}^{-1}$ from $+5 \text{ }^\circ\text{C}$ to $-40 \text{ }^\circ\text{C}$. Maintaining a dewpoint of approximately $-39 \text{ }^\circ\text{C}$ prevents droplet evaporation while ensuring that condensation does not form inside the Linkam stage. To do so, a humidified gas flow was generated by combining a flow of water-saturated nitrogen from a glass bubbler containing ultra-high purity (UHP) water (0.01 Lpm) and a flow of dry nitrogen (0.6 Lpm). Dew point was measured with a hygrometer (EdgeTech DewPrime II, Model 2000). Photographs of the

droplet were taken at 5x magnification every 6 seconds (corresponding to a 0.1 °C temperature change). After the stage reached -40 °C it was heated at a rate of 5 °C min⁻¹ until it reached 5 °C, where the temperature was held for 1 minute to ensure complete melting of the droplet. Mean ice nucleation temperature and fraction frozen over the temperature range (0 to -40 °C) were determined from multiple ice nucleation events were observed on the same sample by repeating the cooling and warming cycle 25 times. In practice, some runs resulted in a data set with fewer than 25 freezing points due to droplet evaporation. Images were analyzed on a frame-by-frame basis to visibly determine the freezing temperature. Data from the measurement of a single sample in the Linkam stage were only included in further analysis if the following criteria were met: 1) a discrete freezing event was observed in a minimum of three freeze-thaw cycles, 2) freezing was clearly observed at a discrete temperature for each observation (i.e. in a single CCD image), 3) the droplet did not visibly change size, and 4) no condensation was observed adjacent to the droplet on the slide. In procedural blank runs, homogeneous droplets of HPLC-grade water (VWR Scientific) were observed at -33.5 ± 2.0 °C on combusted aluminium foil (used as substrates in the PIXE sampler) and at -33.1 ± 0.6 °C on silanized glass slides).

The cumulative number concentration of INPs per L of air in the headspace of the MART (N_{INP}) was calculated using the following equation (DeMott et al., 2016; Vali et al., 1971):

$$N_{INP}(T) = -\ln \frac{N_u(T)}{N_0} * \frac{V_w}{V_a * V_s} \quad (1)$$

where $N_u(T)$ is the cumulative number of unfrozen droplets at a given temperature and N_0 is the total number of freezing events for a sample compiled from all replicates. V_a is the volume of a single aliquot (2 µL for this study), V_w is the total volume of all water droplets (calculated by multiplying V_a by the number of total freezing events from all replicates), and V_s is the total volume of air sampled on the impactor.

2.3 Phytoplankton biomass, growth and physiology

2.3.1 Phytoplankton biomass

Chlorophyll *a* concentrations were determined from water samples filtered onto GF/F filters (Whatman, GE Healthcare Bio-Sciences) and stored at -20°C until analysis. Chlorophyll *a* was determined using a spectrophotometric method (Parsons et al., 1984) for the experiment with *Thalassiosira weissflogii* (Jeffrey and Humphrey, 1975), and a fluorometric method was used in the experiment with *Synechococcus elongatus* (Arar and Collins, 1997). For both protocols, pigments were extracted with 90% acetone and sonication was used to break up the cells. Samples were sonicated (Qsonica, Q125 [125 Watts, 20 kHz]) for 10 min with the amplitude set at 40% in 5 s pulses. Heat build-up was prevented by keeping the extractions on ice with 5 s pauses between pulses. Pigment extraction was continued in the dark overnight at 4°C and the extractions were centrifuged at 1,000 g for 20 min at 4°C prior to measurement of chlorophyll *a* concentration in the supernatant.

Samples of water (1ml) were collected from the MART tank in triplicate and placed in small glass vials for cell counts from the culture of *T. weissflogii*. The samples were fixed with a drop of acidic Lugol's iodine (Parsons et al. 1984) and counted by light microscopy using a hemacytometer (Fuchs-Rosenthal ruling, Hauser Scientific) to count 400 cells (Guillard and Sieracki, 2005). Total (including both cellular and extracellular) carbohydrate concentrations in the medium were measured daily during the experiment with *T. weissflogii*. Carbohydrates were measured using the phenol-sulfuric acid assay (Dubois et al., 1956) calibrated with D-glucose. Culture (0.8 ml) was placed in a boiling tube mixed with 0.5 mL of 5% (w/v) phenol. Concentrated sulfuric acid (2 ml) was added rapidly using a dispenser bottle and absorbance was measured at 485 nm after 20 minutes.

2.3.2 Physiological status

Variable chlorophyll fluorescence was used as an indicator of physiological status of the phytoplankton in the MART. [The maximum quantum yield of photosystem II \(\$\phi_{PSII}\$ \) was measured using the saturating pulse method \(Genty et al., 1989; Maxwell and Johnson, 2000\). This parameter measures the proportion of light absorbed by the photosynthetic pigment chlorophyll in photosystem II that is used to drive photochemistry \(Maxwell and Johnson 2000\).](#) Measurements were made on dark adapted (30 minutes) samples using pulse-amplitude modulated (PAM) fluorescence (PHYTO-PAM fitted with a PHYTO-ED emitter-detector unit (Heinz Walz GmbH)). The four channels of the PHYTO-PAM have different excitation wavelengths (470, 520, 645 and 665 nm) and the detector is protected by a long-pass filter, measuring emission at wavelengths > 710 nm. Data produced by excitation at 665 nm was used in this study. The instrument was set up with a measuring light pulse frequency of 2 (equivalent to photosynthetically active radiation (PAR) of $1 \mu\text{mol m}^{-2} \text{s}^{-1}$), a saturation pulse intensity setting of 10 and a pulse width of 200 ms. ϕ_{PSII} was calculated based on variable fluorescence (F_v), where F_o is the minimum fluorescence yield of the dark-adapted sample when all reaction centers II are open, and F_m is the maximum fluorescence yield when all reaction centers II are closed (Genty et al., 1989; Maxwell and Johnson, 2000):

$$\phi_{PSII} = F_v/F_m = (F_m - F_o)/F_m \quad (2)$$

Cell membrane permeability and therefore potential cell ‘leakiness’ was determined from counts of *Thalassiosira weissflogii* stained with SYTOX Green (Invitrogen, Thermo Fisher Scientific). SYTOX Green is a plasma membrane impermeable nucleic acid dye (Veldhuis et al., 2001; Franklin et al., 2012) that stains cells with a compromised plasma membrane, causing fluorescence with an emission peak of 523 nm when excited by a source at 450–490 nm. Samples (1 ml) were placed in sterile microcentrifuge tubes and incubated with 1.92 μM SYTOX Green stain for one hour in the dark at 20 °C. Stained sample (0.5 ml) was mixed with 1.5 ml of filter sterilized ASW and filtered through a 0.4 μm pore size polycarbonate filter (Nuclepore, Whatman) under low vacuum (< 150 mm Hg). Excess dye was removed by two rinses (1 ml) with filter sterilized ASW. The filters were mounted on glass slides using SlowFade Diamond Antifade Mountant (Invitrogen, Thermo Fisher Scientific). Cells (400 per sample) were counted using an epifluorescence microscope (Axioplan 2, Carl Zeiss MicroImaging) and categorized depending on how they stained (Thornton, 2014): intact plasma membrane (unstained), compromised plasma membrane (stained nucleus), and compromised plasma membrane with breakdown of internal membranes (stained throughout the cells).

2.3.3 Extracellular carbon pools

Two classes of exopolymer particles were measured in the MART. Transparent exopolymer particles (TEP) and Coomassie stainable particles (CSP) are ubiquitous in the ocean (Thornton, 2018) and are often associated with phytoplankton growth (Passow, 2002; Engel et al., 2015; Thornton, 2018) and death (Berman-Frank et al., 2007; Bouchard et al., 2011; Thornton and Chen, 2017). Exopolymer particles were collected by filtration (2 ml) under low vacuum onto 25 mm diameter 0.4 μm pore size polycarbonate filters (Nuclepore, Whatman, GE Healthcare Bio-Sciences) in triplicate. The filters were stained directly in the filter funnel with 1 ml of dye that was drawn through the filter slowly, followed by two rinses of 1 ml of UHP water to remove excess dye. TEP were stained with Alcian blue 8GX (Sigma-Aldrich), which is specific for acid polysaccharides. The Alcian blue solution was 0.02% (w/v) in 0.06% acetic acid (v/v) at pH 2.5 (Passow and Alldredge, 1995). CSP were stained on separate filters with Coomassie

245 Brilliant Blue G-250 (Sigma-Aldrich), which is specific for proteins (Long and Azam, 1996). Working Coomassie Brilliant Blue
G-250 solutions were prepared each day by making a 1/25 dilution of the stock solution (1 g Coomassie Brilliant Blue G-250 in
100 mL of UHP water) with 0.2 μm filtered ASW (Berges et al., 2001). This resulted in a 0.04% (w/v) working solution at pH of
7.4. Dry filters were mounted in a drop of immersion oil on a Cytoclear microscope slide (GE Water & Process Technologies;
Logan et al., 1994). A second drop of oil was placed on top of the filter, and it was covered with a glass coverslip. Prepared slides
250 were stored frozen (-20°C).

Exopolymer particles were quantified in terms of particle abundance (particles mL^{-1}) and concentration ($\text{mm}^2 \text{mL}^{-1}$) from
JPEG images of the filters, captured using an AxioCam ERc 5s color camera mounted on an Axioplan 2 microscope run from
Axiovision 4.8 software (Carl Zeiss MicroImaging). The image analysis method was based on Engel (2009), see Thornton and
Chen (2017) for details. Briefly, image analysis was conducted using ImageJ software (National Institutes of Health). Only the red
255 channel of the JPEG image was analysed, using the triangle method (Zack et al., 1977) to threshold the grayscale image into a
binary image where TEP or CSP were represented by black pixels.

Water samples were filtered through combusted glass fiber filters (GF/F, Whatman) and stored in combusted glass vials
at -20°C in the dark for analysis of fluorescent dissolved organic matter (FDOM). Samples were thawed slowly at room temperature
in the dark and placed in a 1 cm path quartz cuvette. Absorbance spectra and excitation-emission matrices (EEM) were measured
260 using a Horiba Aqualog fluorometer with excitation wavelengths from 240 to 500 nm in 2 nm increments. A high-purity water
sample (Starna Scientific Ltd.) was used to correct fluorescence and absorption spectra. Fluorescence spectra were corrected by
inner filter effect using the Aqualog software, and intensities were converted to Raman units (excitation at 350 nm, emission
intensity from 371 to 428 nm) (Kothawala et al., 2013; Lawaetz and Stedmon, 2009). FDOM was characterized using Coble's
peaks (Coble, 1996, 2007). These peaks in fluorescence spectra are associated with defined chemical properties of DOM, though
265 they cannot be assigned specific chemical formulae (Coble, 1996). We measured peak B (tyrosine-like, protein like; emission
intensity at 310 nm, with an excitation of 275 nm), peak T (tryptophan-like, protein-like; emission intensity at 340 nm, with an
excitation at 275), peak A (humic-like; peak emission intensity between at 380 to 460 nm, with an excitation at 260 nm); peak M
(humic-like; peak emission intensity between at 380 to 420 nm, with an excitation at 312 nm), peak C (humic-like; peak emission
intensity between at 420 to 480 nm; with an excitation at 350 nm).

270

2.3.4 Bacterial activity

Glucosidase activity was used as a proxy for bacterial metabolism during the MART experiment with *Thalassiosira weissflogii*.
Enzyme activity was measured using the fluorimetric method of Hoppe (1983) modified for a 96-well plate format (Marx et al.,
2001). The substrate analog for β -glucosidase activity was 4-methylumbelliferone- β -D-glucoside (Sigma-Aldrich), which is
275 cleaved by β -glucosidase to produce a fluorescent product, 4-methylumbelliferone (MUF). Each assay (200 μL) consisted of 180
 μL of water from the MART and 20 μL of substrate. The assay was intentionally not conducted in a biological buffer to maintain
the pH observed within the MART. Temperature within the plate reader varied < 1.5 °C between assays on different days. Substrate
concentration was 200 μM , which preliminary work showed to be saturating; therefore the enzyme velocities measured represent
 β -glucosidase activity. Fluorescence intensity was measured using a Spectramax Gemini EM spectrofluorometer run by SoftMax
280 Pro 4.8 software. The plate (opaque white 96 well (Grenier)) was mixed by vibration for 5 seconds prior to each measurement.
Fluorescence intensity was measured every 2.5 minutes for up to 12 hours, however only the first 120 minutes of data were used
to calculate rates as this was the period in which there was a linear increase in fluorescence, indicating that the substrate was not
limiting. The assay was calibrated using a dilution series of MUF standards, with calibrations measured several times over the

285 course of the MART experiment. Heat killed controls (MART water heat to 90-95 °C for 15 minutes) were run as well as ASW
and UHP water blanks. Samples from the MART were size fractionated by filtration to locate the enzyme activity; in addition to
whole water, the assay was run on water passed through GF/C filter (Whatman) with a nominal pore size of 1.2 µm, and double-
filtered through a GF/C filter followed by a 0.2 µm pore size syringe filter. Whole water samples contained phytoplankton, bacteria
and detritus. Passage through a GF/C filter was used to remove *T. weissflogii*, detritus and particle associated bacteria. Passage
through a 0.2 µm pore size filter was used to remove free-living bacteria in addition to *T. weissflogii* and detritus.

290

2.4 Artificial Seawater blank-blank MART

A MART experiment was conducted to determine the background conditions in a tank containing artificial seawater, but no
phytoplankton. This experiment represented a controlled blank, which was used to make comparisons to the experiments during
which phytoplankton were grown. The experiment was conducted over 5 days, using the same protocols as the experiments
inoculated with phytoplankton. The experiment was conducted in two phases; firstly, the tank was filled with ASW (no nutrient or
phytoplankton additions) and the aerosol were generated and sampled as described above on days 1 and 2. Secondly, L1 nutrients
were added on day 3 and the measurements were repeated on days 3 and 5. The blank measurements were broken down into these
two phases due to the compositional difference between ASW and ASW+L1, as L1 nutrients contain sources of macronutrients
(N, P and Si), trace metals, and also sources of organics (Guillard and Hargraves, 1993) in the form of Na₂EDTA (14.9 µM) and
vitamins (thiamine (0.3 µM), biotin (2.05 x 10⁻³ µM), and cyanocobalamin (3.69 x 10⁻⁴ µM). The concentrations of macronutrients
added were 60 µM NaNO₃, 20 µM NaH₂PO₄, and 60 µM Na₂SiO₃. Aerosol concentration and ice nucleation properties were
measured. Biological measurements were chlorophyll *a* concentration, bacteria counts, and TEP and CSP concentrations. Ice
nucleation measurements made on samples collected during this experiment were considered procedural blanks.

305 2.5 Field methods

Results from the MART experiments were compared with ice nucleation data collected *in situ* from the *R/V Atlantis* in the North
Atlantic during the second NASA North Atlantic Aerosols and Marine Ecosystems Study (NAAMES) research cruise in May 2016
(Wilbourn et al., 2020). The objective of NAAMES was to understand the ocean ecosystem-aerosol-cloud system of the western
subarctic Atlantic, including how plankton ecosystems influence the properties of remote marine aerosol and boundary layer clouds
(Behrenfeld et al., 2019, 2021). The data were collected at a single station (Station 4; 47° 39.360 N, 39° 11.398 W), after a storm
caused deep water entrainment with homogenous physical properties to well below 200 m on arrival at the station, followed by
shoaling resulting in a mixed layer < 25 m within 24 to 48 hours (Graff and Behrenfeld, 2018). Drifters and float data were used
to maintain the sampling station within the same water mass over the next 4 days (Graff and Behrenfeld, 2018). Della Penna and
Gaubert (2019) defined Station 4 as a subtropical (based on mean dynamic topography) anticyclonic eddy.


315 Detailed methods are provided in Wilbourn et al., (2020). Briefly, high performance liquid chromatography (HPLC) was
used to quantify phytoplankton pigments (Van Heukelem and Thomas, 2001) at the NASA Goddard Space Flight Center Ocean
Ecology Laboratory (Greenbelt, Maryland). Pigments were filtered onto combusted GF/F filters, stored at -80 °C, extracted in
methanol, and analyzed on a 4.6 x 150 mm HPLC Eclipse XDB column using an Agilent RR1200 HPLC system (Agilent
Technologies). In addition, small phytoplankton were enumerated and grouped using a cell sorting flow cytometer (BD Influx,
320 Becton Dickinson Biosciences) (Graff et al., 2012, 2015). The size ranges 0.5-1 µm, 1-3 µm, 3-50 µm corresponded to three groups
of microorganisms: *Synechococcus*, photosynthetic picoeukaryotes, and photosynthetic nanoeukaryotes. Cells were sorted based
on pigment fluorescence emissions at 692 nm and 530 nm, and forward and side scattering intensity.

Primary marine aerosol were generated from the surface of the ocean using the Sea Sweep aerosol generator (Bates et al., 2012, 2020). Aerosol sample collection using PIXE impactors, and ice nucleation measurements were conducted employing the technique described above for the mesocosm study. Air sample flow (1 Lpm) from the Sea Sweep was directed to the PIXE cascade impactor to collect 0.06-1 μm aerodynamic diameter on combusted aluminium foil. Aerosol samples were stored at $-80\text{ }^{\circ}\text{C}$ and transported to Texas A&M for ice nucleation measurements (Wilbourn et al., 2020) as described above.

3 Results

3.1 Phytoplankton culture experiments in a marine aerosol reference tank (MART)

Data collected during the 24 day MART experiment with *Synechococcus elongatus* are summarized in Fig. 2. Growth rates of *Synechococcus* per day and mean INP freezing temperatures are shown in Fig. 2a. In this experiment, growth rate was determined from changes in chlorophyll *a* concentration, a metric commonly used to track changes in phytoplankton biomass (Fig. 2b). Growth rate changed from days 2 to 4 was 1.34 day^{-1} , associated with a peak biomass of $34.1 \pm 0.2\ \mu\text{g chl. } a\ \text{L}^{-1}$ (mean \pm SD; $n = 3$). The chlorophyll *a* concentration declined between measurements made on days 4 and 8, resulting in negative growth rates during this period, indicative of a dying culture. There was a second period of growth during which the biomass of *Synechococcus* increased from mean values of 15.3 to $26.0\ \mu\text{g chl. } a\ \text{L}^{-1}$, but the growth rate was relatively low (0.21 day^{-1}) compared with the initial period of relatively fast growth (1.34 day^{-1}). Mean ice nucleation temperatures of aerosol collected from the MART were warmest during the initial period of high growth rates, with freezing temperatures of -23.9 ± 1.9 and $-18.0 \pm 2.3\text{ }^{\circ}\text{C}$ (mean \pm SD) on days 2 and 3, respectively. These temperatures were significantly warmer than homogeneous freezing of pure water droplets, indicating the presence of effective INPs in the aerosol. Mean ice nucleation temperatures for the remainder of the experiment were relatively cold, ranging from $-34.2 \pm 1.1\text{ }^{\circ}\text{C}$ (day 4) to $-30.7 \pm 1.1\text{ }^{\circ}\text{C}$ (day 8). Ice nucleation temperature measurements of aerosol collected in the blank MART experiment represent procedural blanks, as the samples were treated in the same way as samples from the MART experiments with phytoplankton. The mean ice nucleation temperature of aerosol collected from the headspace of a blank MART containing only ASW was $-34.9 \pm 0.9\text{ }^{\circ}\text{C}$, compared with $-31.3 \pm 0.9\text{ }^{\circ}\text{C}$ for ASW+L1 nutrients (Fig. 2a). As described in the methods, the addition of L1 nutrients included organics, mainly in the form of vitamins.



350 Figure 2. Growth of the cyanobacterium *Synechococcus elongatus* in a marine aerosol reference tank (MART) and associated ice nucleation. (a) Growth rates of *Synechococcus* (green bars) with time (indicated by the width of the green bars), plotted with freezing temperature of ice nucleating particles (INPs) (blue circles) over time (data points indicate mean \pm pooled SD). The freezing temperature of aerosol procedural blanks are shown for reference, where the black circle represents the freezing temperature of aerosol generated from artificial seawater medium (ASW) (mean \pm SD, $n = 7$) and the open circle represents the freezing temperature of aerosol generated from ASW + L1 nutrients (mean \pm SD, $n = 11$). Procedural blank measurements were made during a separate blank MART experiment and therefore their position on the x-axis does not indicate timing. (b). Chlorophyll *a* concentration was used as a measure of *Synechococcus* biomass and the quantum yield of photosystem II (ϕ_{PSII}) was used to indicate physiological status (mean \pm SD, $n = 3$). (c) The fraction of INPs frozen at different temperatures in aerosol collected from the MART. Each solid line represents data from a different day. Procedural blanks are shown as dotted lines for artificial seawater (ASW) and artificial sea water with added nutrients (ASW+L1).

360

Formatted: Space After: 10 pt, Line spacing: single


The quantum yield of photosystem II (ϕ_{PSII}) is a measure of how efficiently phytoplankton use light energy during photosynthetic photochemistry and it is frequently used as an indicator of general physiological status (Fig. 2b). The ϕ_{PSII} was relatively high during the first few days of growth in the MART, increasing from 0.41 ± 0.08 (day 2) to 0.48 ± 0.0 (day 5) (mean \pm SD; $n = 3$). The standard deviations of measurements of both ϕ_{PSII} and chlorophyll *a* were smaller than the data points in most measurements in Fig. 2b. Correlating with changes in chlorophyll concentrations, ϕ_{PSII} declined between days 6 and 8, and increased during the second period of growth from 0.17 ± 0.01 (day 8) to 0.45 ± 0.01 (day 11) (Fig. 2b). On days 21 and 22, ϕ_{PSII} was very low (< 0.1) indicating that the *Synechococcus* culture was dying, though chl. *a* concentration remained high (29.2 ± 0.1 $\mu\text{g chl. } a \text{ L}^{-1}$) (Fig. 2b). The decline in biomass and ϕ_{PSII} (days 5 to 8), followed by a second growth period and increasing ϕ_{PSII}

365

between days 8 and 11, was unexpected. There was no change in environmental conditions within the MART to explain the observed pattern.

The fraction of INPs frozen as a function of temperature is shown in Fig. 2c. The onset of freezing and the temperature of complete INP activation were both much warmer early in the experiment (days 2 and 3) than for aerosol collected on subsequent days. On day 2, INPs in the aerosol started to activate at -19.5°C and were completely activated at -26.5°C (Fig. 2c). Activation started at the warmer temperature of -14.0°C on day 3 and all INPs were activated at -22.0°C . In contrast, activation occurred at colder temperatures, between -26.6 and -33.5°C , for the later sampling days (Fig. 2c). There was no relationship between the biomass of *Synechococcus* in the tank and the concentration of aerosol in the overlying air (Fig. S1). Plots of fraction frozen against freezing temperature show that, except for warm freezing temperatures on days 2 and 3, the freezing temperatures of aerosol from the *Synechococcus* culture were between those of the procedural blanks; generally warmer than ASW on its own, but colder than ASW+L1. Therefore, except for days 2 and 3, there was no difference between the freezing temperature of SSA containing phytoplankton organic matter and the procedural blanks.

The diatom *Thalassiosira weissflogii* showed the same pattern as *Synechococcus elongatus*, with fast growth rates associated with ice nucleation at the warmest temperatures (Fig. 3a), with a rapid increase chlorophyll *a* concentration (38.7 to $188.9\ \mu\text{g L}^{-1}$ between days 1 and 3) and relatively high values of ϕ_{PSII} (> 0.5) (Fig. 3b). The standard deviations of measurements of both ϕ_{PSII} and chlorophyll *a* were smaller than the data points in most measurements in Fig. 3b. The onset of ice nucleation on days 2 and 3 were -19.1°C and -22.3°C , respectively (Fig. 3c). On these days, the fraction of INPs frozen reached 100% at -24°C (Fig. 3c). In contrast, the onset of nucleation was $< -27^{\circ}\text{C}$ for sampling days at slower growth rates later in the experiment (Fig. 3c). Fraction frozen curves showed that freezing occurred at temperatures significantly warmer than the procedural blanks at the beginning of the experiment (days 2 and 3) and at the end of the experiment (days 16 and 29). Several improvements to the experimental procedure were implemented for the *T. weissflogii* experiments. A higher temporal resolution of growth rate estimates was determined on a day-to-day basis using direct cell counts (Fig. 3d). The highest growth rate in the cultures was $3.0\ \text{day}^{-1}$ observed on day 1, but no samples for INP characterization were taken on that day. On days 2 and 3, growth rates were 1.0 and $0.5\ \text{day}^{-1}$, respectively, corresponding freezing temperatures of $-20.7 \pm 1.4^{\circ}\text{C}$ and $-23.0 \pm 0.7^{\circ}\text{C}$ (Fig. 3a). Total carbohydrate increased rapidly between days 2 ($13.9 \pm 1.1\ \mu\text{g ml}^{-1}$; mean \pm SD, $n = 3$) and 5 ($20.5 \pm 0.5\ \mu\text{g ml}^{-1}$), corroborating the pattern in biomass and growth rates observed from cell counts (Fig. 3d). Day 3 was the day on which the chl. *a* concentration was highest ($188.9 \pm 4.8\ \mu\text{g chl. } a\ \text{L}^{-1}$; mean \pm SD, $n = 3$) (Fig. 3b), with a cell count of $7.58 \times 10^4 \pm 4.39 \times 10^3\ \text{cells ml}^{-1}$ (mean \pm SD, $n = 3$) (Fig. 3d). Growth rates on days 7 and 9 were $0\ \text{day}^{-1}$, indicating stationary phase. Based on cell counts, there was a significant correlation between growth rate over the 24 hours before the sample was taken and mean nucleation temperature ($r = 0.93$, $p < 0.001$, $n = 20$). There was a correlation ($r = 0.917$, $p < 0.001$, $n = 13$) between ϕ_{PSII} and growth rate in *Thalassiosira weissflogii*, with high ϕ_{PSII} indicating photosynthetically efficient cells that were growing rapidly. High values of ϕ_{PSII} (> 0.44) were associated with ice nucleation at relatively warmer temperatures ($> -24^{\circ}\text{C}$) during the *T. weissflogii* experiment ~~both taxa~~. SYTOX Green staining was measured as a further indicator of the physiological status of the *Thalassiosira weissflogii* (Fig. 3e). It is a marker of cell permeability, in which the increase in staining indicates that cells are potentially leaking more dissolved organic matter (DOM) into the surrounding water. In the *T. weissflogii* experiment the proportion of cells that stained with the fluorescent probe increased rapidly as the concentration of diatoms decreased in the MART, from $< 5\%$ during the first seven days of culture, to $11 \pm 5\%$ (mean \pm SD) on day 9, and $92 \pm 2\%$ by day 13.



410 Figure 3. Growth of the diatom *Thalassiosira weissflogii* in a marine aerosol reference tank (MART) and associated ice nucleation. (a).
 Day-to-day growth rates of *T. weissflogii* determined based on changes in cell concentration (green bars) plotted with freezing
 temperature of ice nucleating particles (INPs) (blue circles) over time (data points indicate mean \pm pooled SD). Growth rates were
 measured for the first 13 days of the experiment. The freezing temperature of aerosol procedural blanks are shown for reference, where
 the black circle represents the freezing temperature of aerosol generated from artificial seawater medium (ASW) (mean \pm SD, n = 7)
 415 and the open circle represents the freezing temperature of aerosol generated from ASW + L1 nutrients (mean \pm SD, n = 11). Blank
 measurements were made during a separate blank MART experiment and therefore their position on the x-axis does not indicate timing.
 (b). Quantum yield of photosystem II (Φ_{PSII}) was used to indicate the physiological status (mean \pm SD, n = 3) of the diatom (open circles)
 over time, while changes in chlorophyll *a* show changes in biomass (green squares). (c). Fraction of INPs frozen and concentration of
 INPs in the atmosphere above the culture in the MART at different temperatures. The coloured **solid** lines represent different days.
 420 **Procedural blanks are shown as dashed lines for artificial seawater (ASW) and artificial sea water with added nutrients (ASW+L1).** (d).
 Changes in diatom biomass within the culture over time based on diatom cell abundance (mean \pm SD, n = 3) (blue circles) and total
 carbohydrate (open squares) (mean \pm SD, n = 3). (e) SYTOX Green staining as an indicator of cell leakiness and death over time. The

proportion of diatoms that were unstained and therefore had intact cells (purple triangles), partially stained with a stained nucleus (blue squares), or fully stained indicating dying or dead cells (red circles).

425

Concentrations of INPs in the headspace above both cultures are shown in Fig. 4. Concentration of INPs in both experiments were similar, despite cultures of *T. weissflogii* reaching a biomass (Fig. 3b) an order of magnitude greater than that of *S. elongatus* (Fig. 3a), as indicated by chlorophyll *a* concentration. The maximum concentration of INPs in both experiments was approximately 2×10^{-3} INP L⁻¹ air at -26 °C (*T. weissflogii*) and -29 °C (*S. elongatus*). INPs were observed at the warmest temperatures on day 3 in cultures of *S. elongatus*, with the onset of nucleation first detected (6×10^{-5} INPs L⁻¹) at -14 °C .

430

Information on additional variables which were assessed, but did not indicate a clear relationship to ice nucleation behavior is provided in the Supplement. Aerosol number concentrations in the MART were greater in the headspace over cultures (Fig. S1) compared with the headspace over blank seawater (Table S1), indicating that biomass and biological activity affected the production of aerosol. Aerosol number concentration was generally higher (mean = 2.21×10^6 L⁻¹) over the culture of *T. weissflogii* compared with *Synechococcus* (mean = 1.43×10^6 L⁻¹), reflecting the relatively higher biomass in the diatom culture, as indicated by chlorophyll *a* concentration in Figs. 2b and 3b above. Less aerosol were generated from a control MART tank containing only artificial seawater ($< 1 \times 10^6$ L⁻¹; Table S1). Aerosol number concentration increased in the control experiment after the addition of L1 nutrients, possibly to the addition of organic matter in the form of vitamins, and the growth of bacteria in the water (Table S1). In summary, there is no connection between aerosol concentration and ice nucleation activity indicating that the improvements in ice nucleating ability are not driven by the total aerosol available to catalyze freezing.

435

440

Results from the analysis of the extracellular material in the MART experiments, namely exopolymer particles (Transparent exopolymer particles (TEP) and Coomassie Staining Particles (CSP)) and fluorescent dissolved organic matter (FDOM) is provided in the supplementary material (Figures S2 and S3). DOM in the medium provides potential substrates for bacterial metabolism and growth. Measurements of β -glucosidase activity were used as a proxy for bacterial activity in the *T. weissflogii* experiment (Figure S4). β -glucosidase activity increased by two orders of magnitude over the course of the experiment, indicating that bacteria were playing a role in the transformation and remineralization of organic matter in the MART. Filtration removed glucosidase activity, indicating that the enzyme was associated with cells or aggregates of organic matter and there was no significant extracellular glucosidase in the culture medium. Diatom biomass (Fig. 3d) declined from day 8 onwards; the continued increase in glucosidase activity suggests that the enzyme activity was associated with heterotrophic bacteria and the remineralization of organic matter as the phytoplankton died.

445

450

Formatted: Font: Italic




Figure 4. Concentration of ice nucleating particles (INPs) in the headspace of a marine aerosol reference tank (MART) at different temperatures. Measurements were made over time on different days in batch cultures of (a) *Synechococcus elongatus* (cyanobacterium) and (b) *Thalassiosira weissflogii* (diatom).

455

3.2 Field observations from the North Atlantic

Field data from Wilbourn et al., (2020) were used to calculate phytoplankton growth rates to determine whether high growth rates were associated with ice nucleation at relatively warmer temperatures. Fig. 5a shows changes in phytoplankton biomass in the surface water of the North Atlantic Ocean, determined in this case, from pigments and flow cytometry. As Station 4 was a natural ecosystem there was grazing (Morison et al., 2019), which was absent during the MART experiments. Consequently, changes in biomass at Station 4 reflect the loss of biomass through grazing, as well as additions through phytoplankton growth. Net growth occurred at Station 4, with an increase in chlorophyll *a* concentration from 0.52 to 1.28 mg m⁻³ on days 1 and 4, respectively. Cell counts corroborated an accumulation in phytoplankton biomass, with concentrations of *Synechococcus* increasing from 2.34 x 10⁶ to 1.38 x 10⁷ cells L⁻¹. Changes in biomass between days were used to calculate net growth rates (Table 1) at Station 4. Total phytoplankton growth rate, based on change in chlorophyll *a*, was greatest between days 3 and 4, with a rate of 0.56 day⁻¹ (Fig. 5b). Growth rate of *Synechococcus* (Table 1) was greatest between days 3 and 4, though it is not known how much the rapid increase in *Synechococcus* contributed to the total increase in phytoplankton biomass. Measurements showed that mean ice nucleation temperatures decreased from day 1 to 3 on station 4 (Fig. 5b), and were warmest at -24.5 ± 0.9 °C (mean ± SD) on day 4, which was associated with the highest phytoplankton biomass (Fig. 5a) and highest net phytoplankton growth rate over the preceding 24 hours (Fig. 5b). The fraction of INPs as a function of temperature is shown in Fig. 5c. The onset ice nucleation temperature and the temperature of complete INPs activation was much warmer on day 4 (Fig. 5c). Taken together, our MART tank and NAAMES experiments shown that in two very different taxa grown in the laboratory, and in complex natural phytoplankton assemblages, the highest propensity to nucleate atmospheric ice crystals occurs during conditions of the fastest growth rates.

475




Figure 5. Phytoplankton growth and ice nucleation over 4 days at a Lagrangian station in the North Atlantic Ocean (24 to 27 May 2016). (a) Changes in phytoplankton biomass as indicated by pigment concentrations (colored stacked bars) and cell abundance (monochrome stacked bars). (b) Day-to-day growth rates determined from changes in chlorophyll *a* concentration and ice nucleation temperature of primary marine aerosol generated *in situ* from surface water (data points indicate mean \pm pooled SD) (c) Fraction of ice nucleating particles (INPs) frozen at different temperatures in primary marine aerosol generated *in situ*. Each coloured line represents aerosol collected on a different day. Field data replotted from Wilbourn et al. (2020), with the addition of growth rates.

480

485

Time period (days)	Growth rate (day ⁻¹)			
	chl. <i>a</i>	<i>Synechococcus</i>	picoeukaryotes	nanoeukaryotes
1-2	0.27	0.29	0.54	1.02
2-3	0.34	0.37	0.65	0.60
3-4	0.56	1.12	0.68	0.57

490 **Table 1. Net phytoplankton growth rates (day⁻¹) at a Lagrangian station tracked over 4 days (24 to 27 May 2016) in the North Atlantic Ocean. Growth rates calculated based on changes in biomass concentration over 24 hours. Chlorophyll *a* was used as a proxy for total phytoplankton biomass. Different components of the pico- and nanophytoplankton were counted using flow cytometry. Growth rates calculated using data from the North Atlantic Aerosol Marine Ecosystem Study (NAAMES) (Behrenfeld et al. 2019; Wilbourn et al. 2020).**

4 Discussion

500 Our results show that SSA generated from cultures of two contrasting taxa (*Thalassiosira weissflogii* (diatom) and *Synechococcus elongatus* (cyanobacterium)) promoted immersion mode ice nucleation at temperatures warmer than the homogeneous freezing of liquid water droplets. Several previous studies have shown that phytoplankton promote ice nucleation (Knopf et al., 2011; Alpert et al., 2011a, 2011b; Wilson et al., 2015; McCluskey et al., 2017, 2018; Wolf et al., 2019; Wilbourn et al., 2020). The unique finding of this study is that the physiological status of phytoplankton in the water affects the properties of SSA sufficiently to increase their ice nucleation temperature. Fast growing phytoplankton produced SSA containing INPs that catalysed ice nucleation at temperatures that were warmer than during other phases of growth, including stationary phase and death phase. This is a finding that was observed in two contrasting taxa and confirmed by a re-examination of field data from Wilbourn et al. (2020). In both the *Synechococcus* and *T. weissflogii* experiments, periods of elevated ice nucleation ability coincided with high Φ_{PSII} . Further work is needed to determine whether Φ_{PSII} could be developed as a reliable indicator of ice nucleation activity. The data presented here suggests that [the](#) physiology of fast-growing phytoplankton is associated with the production of a chemical signature (a specific compound, class of compounds, or even a change in the ratio of compounds) that affects the ice nucleation of aerosols at relatively warm temperatures.

It is well established that phytoplankton acclimate to different growth conditions (Geider et al., 1997, 1998; Moore et al., 2006), resulting in different chemical compositions (Geider and LaRoche, 2002; Dyhrman et al., 2012; Lin et al., 2016). While it is impossible to screen all the different biomolecules within a taxon for ice nucleation activity, observations of resource allocation and resource allocation models provide a conceptual framework to identify differences in the composition of phytoplankton growing under different conditions (Klausmeier et al., 2004; Liefer et al., 2019; Inomura et al., 2020). Klausmeier et al. (2004) explained the N:P stoichiometry of phytoplankton in terms of three strategies. ‘Survivalists’ are rich in resource acquisition machinery (pigments and proteins) and have a high N:P ratio; ‘generalists’ have a balance between resource acquisition and growth machinery and an N:P ratio at or near Redfield proportions; ‘bloomers’ are adapted for exponential growth and have a high proportion of growth machinery, resulting in a low N:P ratio (Arrigo, 2015; Klausmeier et al., 2004). Based on this model, rapidly growing phytoplankton associated with ice nucleation at relatively warm temperatures have the characteristics of ‘bloomers’. Rapidly growing phytoplankton have elevated cell quotas for P as cell processes required for rapid growth involve large quantities of nucleic acids. For example, growth requires P-rich ribosomes (composed of RNA and protein) for the synthesis of proteins. [This concept is also expressed by ecologists in terms of the growth rate hypothesis \(GRH\), where the C:P and N:P ratio of fast-growing organisms is low due to the high concentrations of ribosomal RNA required for protein synthesis \(Elser et al., 2000; Isanta-Navarro et al., 2022\), though the application of the GRH to all phytoplankton has been challenged \(Flynn et al., 2010\).](#)

530 Recent work with relatively pure samples of representative biomolecules from cells have shown significant differences in ice nucleation properties between compounds (Alsante et al., 2023; Steinke et al., 2020; Wolf et al., 2019). SSA collected from the MART tanks were a complex mixture of different biomolecules with different ice nucleation properties. These biomolecules were predominantly produced via photosynthesis and growth of the phytoplankton. Transformation of the phytoplankton organic matter by heterotrophic bacteria in the tank also played a role, as did background contamination of the ASW with organic matter. Determining how these components interact to affect immersion ice nucleation is a challenging and multifaceted problem. For example, concentration affects the immersion freezing temperature of the protein apoferritin, with higher concentrations resulting in freezing at warmer temperatures, due to protein aggregation (Cascajo-Casresana et al., 2020). The interactions of ice nucleating proteins with salts affect ice nucleation temperatures (Schwidetzky et al., 2021).

540 Our recent work shows that both DNA and RNA are moderately effective INPs, nucleating at a mean temperature of -20 ± 1 °C (Alsante et al., 2023). Given that DNA is reportedly enriched by a factor as high as 30,000 in artificially generated SSA using seawater from the North Atlantic Ocean (Rastelli et al., 2017), as well as direct metagenomic evidence of DNA in the atmosphere (Mayol et al., 2017; Lang-Yona et al., 2022), this represents a plausible marine INP. We did not measure the concentration of nucleic acids in the MART, but the GRH hypothesis and ‘bloomer’ strategy would suggest that nucleic acids were a relatively large proportion of the organic matter during the early stage of the cultures, when they were growing at the highest rates and affected ice nucleation at the warmest temperatures. However, nucleic acids would have accumulated in the cultures after the period of relatively warm ice nucleation as the total biomass increased, suggesting that ice nucleation at warm temperatures was not driven by the concentration of nucleic acids.

550 Another potential source of INP in fast growing phytoplankton are proteins associated with growth. The measured N:C ratio of phytoplankton increases with growth rate (Inomura et al., 2020), indicating that fast growing phytoplankton are synthesising more proteins and nucleic acids, storing more nitrogen, or both. Recent studies have shown that proteins can be effective INP (Cascajo-Casresana et al., 2020; Daily et al., 2022; Lukas et al., 2021; Roeters et al., 2021; Scwidetzky et al., 2021; Hartman et al., 2022). Proteins, like nucleic acids, would have accumulated in the cultures beyond the initial phase of fast growth rate as the phytoplankton biomass continued to increase. Therefore, the amount of protein in the water (and presumably the SSA generated from the water) was not a driver of the warm freezing temperatures observed during the early phase of growth. There are significant differences between proteins in their ability to act as INP (Cascajo-Casresana et al., 2020; Alsante et al., 2023). Some proteins are among the most efficient known INP. Examples of highly efficient protein INPs include InaZ from *Pseudomonas syringae*, which nucleates ice at temperatures as high as -2 °C (Maki et al., 1974; Roeters et al., 2021; Wex et al., 2015), and ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) (Alsante et al., 2023). RuBisCO initiates immersion mode ice nucleation at -6.8 °C, with complete freezing at -9 °C, and a mean freezing temperature of -7.9 ± 0.3 °C (Alsante et al., 2023). RuBisCO would have been abundant in the MART experiments as it is one of the most abundant proteins on Earth, which is ubiquitous in phytoplankton as it is essential to photosynthetic carbon fixation (Ellis et al., 1979; Raven 2013; Bar-On & Milo, 2019). In contrast, other proteins are significantly less efficient as INP; the mean immersion mode freezing temperature of alkaline phosphatase was -20.4 ± 1.3 °C in the same study (Alsante et al., 2023). These findings suggest that it is the presence of specific proteins, rather than bulk protein concentration, that determine the ice nucleation efficiency of organic matter.

565 A representative phytoplankton cell is composed of 25–50% protein, 5–50% polysaccharide, 5–20% lipids, 3–20% pigments and 20% nucleic acids (Emerson & Hedges, 2008). While we did not measure lipids, total carbohydrates were used as an indicator of biomass in the culture of *Thalassiosira weissflogii*. Carbohydrate concentration followed the same pattern as cell concentration, peaking after the measurement of most efficient ice nucleation. Carbohydrate breakdown became more significant as the culture aged, as indicated by measurements of glucosidase activity and the decrease in total carbohydrate concentration after

Formatted: Font: Italic

Formatted: Font: Italic

570 day 7. Wolf et al. showed that both lipids and carbohydrates from the cyanobacterium *Prochlorococcus* were efficient deposition
INP. Dreischmeier et al. (2017) showed that large polysaccharides derived from tree pollen acted as INP, whereas smaller
575 carbohydrates had anti-freeze properties. It was not possible to determine whether carbohydrates and lipids acted as a source of
INP in the MART tanks.

Differences in DOM concentration and composition could also play a role in explaining why ice nucleation temperature
changes with growth rate and physiological status. The proportion of fixed carbon released as DOM depends on both taxon and
physiological status (Thornton, 2014). Cell death also releases DOM into the surrounding water (Veldhuis et al., 2001; Thornton,
575 2014). The enrichment of SSA with organic matter has been related to stress and death in both field (O'Dowd et al., 2015) and
laboratory experiments (Wang et al., 2015). In addition to physiological stress, physical disruption of cells affect the release of
DOM and cell fragments from phytoplankton *in situ* through sloppy feeding by zooplankton (Møller et al., 2003; Møller, 2007)
and viral infection (Bratbak et al., 1993; Vardi et al., 2012; Kranzler et al., 2019). Wolf et al. (2019) measured the ice nucleation
biogenic material from *Prochlorococcus* cultures in the deposition mode, finding that small particles from lysed cultures were
580 more effective INPs than large particles. Morison et al. (2019) found significant rates of grazer mortality and low rates of viral
lysis at Station 4 in the Atlantic at the same time as our work, suggesting that these processes would have contributed cell fragments
and DOM to SSA sampled *in situ*. Grazing and viral lysis did not occur in the MART experiments. However, SYTOX Green
staining showed increased cell permeability during stationary and death phase of the *T. weissflogii* culture, indicating greater
potential to leak DOM into the surrounding medium. Evidence for DOM in both MART experiments came from measurements of
585 exopolymer particles and the presence of FDOM. FDOM in the MART showed that fluorescence peaks (peaks B and T; Coble,
1996) associated with proteins were high (Fig. S3b), both in terms of concentration and relative to other fluorescence peaks, during
the exponential growth of *Thalassiosira weissflogii* (Fig. 3~~2~~). However, Peaks B and T were not elevated during exponential
growth of *Synechococcus*. Based on our data, it was not possible to link ice nucleation activity with cell 'leakiness' and the
measured components of DOM.

590 After the initial phase of rapid growth during the *Synechococcus* MART experiment there was a decline in biomass, as
indicated by the halving of chl *a* concentration between days 4 and 8. This was followed by a second period of growth, which did
not correlate with relatively warm ice nucleation (Fig. 2). There are two possible explanations for this discrepancy. Firstly, the
second period of growth (0.21 day⁻¹) was not rapid compared with growth rates during the early phase of the culture (1.34 day⁻¹).
595 Secondly, the composition of the water had changed significantly since the initial phase of fast growth due to the death and
physiological stress of *Synechococcus* between days 4 and 8. In addition to the declines in chl *a* concentration and quantum yield
of photosystem II, there was also an increase in the amount of dissolved organic matter in the MART, as indicated by an increase
in FDOM concentration between measurements on days 1 and 6 (Fig. S3). While MART and mesocosm experiments often progress
with one distinct peak (such as in the experiment with *Thalassiosira weissflogii* (Fig. 3)) and compilation of growth curves in Lee
et al. (2015), there are examples where there is more than one peak in phytoplankton biomass over the course of an experiment
600 (Lee et al., 2015; McCluskey et al., 2017; Wang et al. 2015).

- A separate blank MART experiment was run to determine the effect of the combination of salts and potential background
contamination on the results. Ice nucleation measurements of these procedural blanks were higher than the homogeneous freezing
temperature (-38 °C; Kanji et al. 2017), but this was not surprising, and it is typical of procedural blank measurements (Alsante et
al., 2023; Irish et al., 2017; Wilbourn et al., 2020). Two opposing factors affected the freezing temperature of the procedural blanks:
605 salt depresses freezing temperature (Wilbourn et al. 2020; Perkins et al. 2020) and organic matter potentially act as an INP. The
data suggest that background organic matter was more significant than salt depression in determining the ice nucleation temperature
of phytoplankton-free medium. The procedural blanks suggest that the ASW contributed INP, both from the analytical grade salts

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Superscript

Formatted: Font: Italic

Formatted: Font: Italic

610 used to make the medium, and from the addition of nutrients (including vitamins and EDTA) to the growth medium. It is not possible to determine whether the INP activity of the medium was a direct effect of the organic matter background, or an indirect effect after the growth and transformation of organic matter by heterotrophic bacteria. These results show that ASW made with analytical grade salts offers no advantage over natural filtered seawater; both media will contain background organic matter contamination. The depression of freezing temperature by salts depends on the concentration and composition of salt, as well as the composition of potential INP interacting with the salt. Wilbourn et al. (2020) found that that decreasing the salinity of samples collected from the Atlantic Ocean from 30 g L⁻¹ NaCl to 3.75 g L⁻¹ increased heterogeneous ice nucleation freezing temperature by approximately 10 °C. However, interactions between salts and INP may be complex. Schwidetzky et al. (2021) found that some ions inhibited, while others facilitated, freezing in ice nucleating proteins from the bacterium *Pseudomonas syringae*. Further, the concentration of salts in solution at the time of freezing will not necessarily be the same as the original seawater generating the SSA. Wilbourn et al. (2020) proposed that SSA would activate as cloud condensation nuclei (CCN) before freezing, with the resulting water uptake diluting the salts in the sea spray by an estimated factor of approximately 27.

620 Number concentrations of INP were low (< 2 x 10⁻³ L⁻¹) during both MART experiments, even at temperature < -20 °C. Our concentrations were at the lower end of previous measurements from laboratory mesocosm and MART tanks (DeMott et al. 2016; McCluskey et al. 2017). Measurements of number concentrations of marine INP *in situ* are limited in spatial and temporal coverage, and vary by orders of magnitude. A summary of the results from several recent campaigns in the Southern Ocean and found the INP concentrations at -20 °C were 10⁻⁴ to 10⁻² per liter, with higher concentrations observed near the terrestrial influence of landmasses such as Australia (McFarquar et al., 2021). Welti et al. (2020) summarized published shipboard measurements of ice nucleation from the ocean, with a 10⁻⁴ to 10⁰ L⁻¹ range in INP concentration at -15 °C and a range in INP concentrations over 9 orders of magnitude (10⁻⁴ to 10⁵ L⁻¹) over the temperature range -40 to -5 °C. These studies included some sites close to land, where continental sources were likely the major source of INP.

630 Field measurements (Wilbourn et al., 2020) corroborated our laboratory results; the fastest phytoplankton growth was associated with the warmest ice nucleation temperatures. According to Della Penna and Gaube (2019), the water sampled on Days 1-4 was sampled from within the same anticyclonic eddy. Similarly, Graff and Behrenfeld (2018) assumed that the same water mass was sampled for all 4 days at Station 4 in their study of phytoplankton photoacclimation. Following those studies, our interpretation of the field data assumes that the samples used to calculate net growth were from the same water mass.

635 Specifically, our growth rates (Fig. 5b) represent growth over the 24 hours leading up to the aerosol sampling; the growth rate for day 4 (Fig. 5b) was based on the differences in morning chl. *a* measurements (sampled between 8:00 and 9:00 local time) taken on days 3 and 4. Morison et al. (2019) measured grazing for days 1-3 on Station 4; grazing rates were insignificant on day 1 followed by significant on days 2 and 3 (0.26 to 0.44 day⁻¹; based on chl. *a*). Their observations show that *in situ* growth rates were significantly higher than we estimated. Unfortunately, Morison does not provide grow rates for Day 4, out of concern that day 4 was potentially biologically different as the ship had drifted from the center to the periphery of the eddy.

640 Fast growing *Synechococcus* facilitated ice nucleation at relatively warm temperatures in the laboratory and *Synechococcus* was a significant component of the phytoplankton biomass at Station 4 in the Atlantic (Wilbourn et al., 2020). *Synechococcus* had the fastest net growth rate (1.12 day⁻¹) between days 3 and 4, leading up to the relatively warm ice nucleation temperatures observed on day 4 at Station 4. The proportion of *Synechococcus* cells was relatively low; it contributed 9.8 and 14.5 % of total cell numbers counted with flow cytometry on days 3 and 4, respectively (Fig. 3a). Consequently, it was not possible to determine the role of organic matter from *Synechococcus* in affecting ice nucleation temperature. Other components of the phytoplankton also grew relatively quickly between days 3 and 4, including picoeukaryotes (0.68 day⁻¹), which were the most abundant components of the small phytoplankton (between 79.7 and 83.4 % of cells counted using flow cytometry). Wilbourn et

Formatted: Font: (Default) +Headings (Times New Roman)

Formatted: Font: (Default) +Headings (Times New Roman)

Formatted: Font: (Default) +Headings (Times New Roman), 10 pt

Formatted: Font: (Default) +Headings (Times New Roman), 10 pt

Formatted: Font: (Default) +Headings (Times New Roman), 10 pt

Formatted: Font: Italic

Formatted: Font: (Default) +Headings (Times New Roman)

al., (2020) also measured the ice nucleation temperatures of *Synechococcus* isolated by flow cytometry on Day 4 at Station 4. There was no significant difference between the mean freezing temperature of *Synechococcus* and those of photosynthetic picoeukaryotes and photosynthetic nanoeukaryotes.

It should not be overlooked that heterotrophic microorganisms (bacteria and protists) were significant contributors to biomass at Station 4 (Bolaños et al., 2021). Bacteria play an important role in determining the fate of organic matter in the ocean (Kujawinski, 2011; Hasencz et al., 2020). Wang et al. (2015) proposed that the organic matter composition of SSA is affected not only by primary production, but also by the enzymatic degradation of organic matter by heterotrophic bacteria. Glucosidase activity was used as a proxy for bacterial activity in the MART culture of *T. weissflogii*; it increased by two orders of magnitude over the course of the experiment. This indicates that bacteria were playing a significant role in the remineralization and transformation of DOM in the MART tank, particularly as the culture aged during the death and stationary phases of culture. However, this had no observed influence on either the concentration of aerosols or their ice-nucleating ability. Hill et al. (2023) found a positive correlation between the concentration of heterotrophic nanoflagellates and ice nucleating entities in a mini-MART experiment using natural seawater. Heterotrophic nanoflagellates were not present in the simplified systems grown in the MART, therefore our experiment does not account for their effects. Heterotrophic nanoflagellates were a component of the grazing community on the North Atlantic, though they were not counted (Morison et al., 2020).

Organic matter in the surface ocean can be divided into pools based on age as it exists on a continuum from labile to ultra-recalcitrant (Hansell 2013). From the perspective of the generation of SSA, we can consider a constant background pool of recalcitrant organic matter with a lifetime longer than the mixing time of the ocean, and relatively labile pools of organic matter that turnover quickly and are affected by recent biological processes, such as phytoplankton growth, in the surface ocean (Brooks and Thornton, 2018). There is a growing paradigm that the production of organic-rich primary SSA is driven by relatively labile organic matter produced recent processes, such as phytoplankton blooms, in the surface ocean (O'Dowd et al. 2004, 2015; Rinaldi et al. 2013; Sciare et al. 2009). However, there is no consensus as some studies indicate that the organic composition of primary marine aerosol is relatively constant, indicating that it is determined by the relatively recalcitrant background pool of organic matter that is uncoupled from recent phytoplankton growth (Lewis et al. 2022; Quinn et al. 2014). INP represent a rare subset of atmospheric aerosol, with only 1 in 10^5 , or fewer, aerosol particles acting as INP (DeMott et al. 2010). This observation, coupled with the selective concentration of different pools of organic matter in the SML (Cunliffe et al., 2013; Thornton et al., 2016) and selectivity of the aerosolization process through bubble bursting (Aller et al., 2005, 2017; Rastelli et al. 2017), make it challenging to relate biological processes in the ocean with ice nucleation in the atmosphere.

Experiments with phytoplankton cultures can be controlled, and therefore are useful to understand the complex processes generating marine biogenic aerosol and INP (Hill et al., 2023; McCluskey et al., 2017, 2018; Prather et al. 2013; Wang et al. 2015). Recent work (Hill et al. 2023; McCluskey et al., 2017, 2018) suggest that the production of INP occurs during the collapse and decay of phytoplankton blooms during the later stages of a growth experiment. In contrast, our results suggest that it is during exponential growth in the early stages of a growth experiment, when the most effective ice nuclei are produced. These findings are not mutually exclusive, INP may be significant at more than one stage of a phytoplankton bloom, in terms of abundance, relatively warm freezing temperature, or both. It is a consensus that ice nucleation at temperatures significantly warmer than homogeneous freezing is driven by organic matter in SSA. Therefore, processes that affect the composition and abundance of organic matter will affect both the abundance and properties of INP. Different phytoplankton have different have significantly different traits in terms of cell size, growth rates, and chemical composition (Beardall et al., 2009; Edwards et al., 2012; Ho et al., 2003; Quigg et al., 2003) and therefore the choice of taxa may affect the results. To date, too few taxa of phytoplankton have been grown in these types of experiment to determine taxa-specific patterns in INP production with growth. Our experiments were different from most other

Formatted: Font: Not Bold

Formatted: Superscript

MART or large tank experiments as we used artificial seawater as the growth medium; most other experiments have used natural seawater from the Pacific Ocean (Hill et al., 2023; McCluskey et al., 2017, 2018; Prather et al. 2013; Wang et al. 2015). This water was filtered through a mesh screen to remove animals and the largest protists, but it still contains natural assemblages of heterotrophic bacteria, protists and viruses. It may be that processes such as viral lysis, grazing, and the processing of phytoplankton organic matter by heterotrophic bacteria are important in determining the properties of SSA and the INP that they contain. Some of these processes were absent from our MART experiments. Differences between experiments may also arise from the research objectives of the researchers; Hill et al. (2023) focussed their experiment on the ‘decay phase’ of the phytoplankton bloom and McCluskey et al. (2017) did not make ice nucleation measurements between days 2 and 8 of a MART experiment.

4 Conclusion

We initially anticipated that processes associated with the release of DOM and cell fragments into the water column associated with physiological stress, slow growth rates, and autocatalytic cell death would produce a larger number of more efficient INPs. Bloom collapse has been proposed as a major source of organic matter in SSA (O’Dowd et al., 2015). McCluskey et al., (2017) found an increase in the production of INPs, active between -25 and -15 °C, following the peak in chlorophyll concentration in a mesocosm experiment with natural seawater. They explained this consistent population of INPs as resulting from the collapse and decay of the phytoplankton blooms during the mesocosm experiments (McCluskey et al., 2017). Our data indicate the opposite; ice nucleation at relatively warm temperatures was associated with fast growth, healthy photosynthetic machinery (indicated by high values of Φ_{PSII}), and relatively few dying or leaking cells (indicated by SYTOX green staining).

Significantly, our results are the first to show that fast growing phytoplankton are a source of INPs that catalyse freezing at relatively warm temperatures. Insights into the relevant chemical properties of fast-growing cells will come from considering how phytoplankton allocate resources, with rapid growth requiring a relatively high cellular content of nucleic acids and specific proteins, compared with slow growth. Similar results were observed in two contrasting taxa of laboratory-grown phytoplankton and a taxonomically mixed population of phytoplankton in the ocean, indicating that the coupling between fast growth and efficient INPs is widespread phenomenon and not associated with any one taxonomic group. These results may have significant implications for the prediction of mixed phase cloud formation, life cycle, and precipitation on regional scales as rapidly growing phytoplankton form blooms over large areas of the ocean. Blooms such as the annual spring bloom in the North Atlantic Ocean may be a temporally and spatially predictable source of relatively active marine INPs in SSA.

715 Data availability

Data will become available in the Texas Data Repository (<https://dataverse.tdl.org/>) on publication.

Supplement

The supplement related to this article is available online.

Author contributions

SDB and DCOT designed the experiments with input from EKW, JM and ANA. EKW, JM, ANA, AW, and KM conducted the marine aerosol reference tank (MART) experiments, and analysed samples and data, in consultation with SDB and DCOT. GGB analysed samples and processed the data for the CDOM analysis. DCOT wrote the manuscript with assistance from SDB and ANA, and editorial input from EKW, JM, and GGB.

Competing interests

The authors declare no competing interests.

730 Acknowledgements

We thank Joseph Niehaus for his significant contribution to the construction and testing of the marine aerosol reference tank. We thank the Captain and crew of the *R. V. Atlantis*, and the NAAMES science team, for making the collection of field samples possible.

Financial Support

735 SDB and DCOT were supported by the National Science Foundation (United States) under Grant No. AGS-1539881. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. Funding for field sample collection was provided by NASA Earth Venture Suborbital-2 (EVS-2) Award #NNX15AE68G.

740 References

[Aller, J. Y., Kuznetsova, M. R., Jahns, C. J., and Kemp, P. F.: The sea surface microlayer as a source of viral and bacterial enrichment in marine aerosols, *Journal of Aerosol Science*, 36, 801-812, 10.1016/j.jaerosci.2004.10.012, 2005.](#)

745 [Aller, J. Y., Radway, J. C., Kilhau, W. P., Bothe, D. W., Wilson, T. W., Vaillancourt, R. D., Quinn, P. K., Coffman, D. J., Murray, B. J., and Knopf, D. A.: Size-resolved characterization of the polysaccharidic and proteinaceous components of sea spray aerosol, *Atmospheric Environment*, 154, 331-347, 10.1016/j.atmosenv.2017.01.053, 2017.](#)

[Alsante, A. N., Thornton, D. C. O., and Brooks, S. D.: Ice nucleation catalyzed by the photosynthesis enzyme RuBisCO and other abundant biomolecules, *Commun. Earth Environ.*, 4, 9, 10.1038/s43247-023-00707-7, 2023.](#)

750 [Alpert, P. A., Aller, J. Y., and Knopf, D. A.: Ice nucleation from aqueous NaCl droplets with and without marine diatoms, *Atmos. Chem. Phys.*, 11, 5539-5555, <https://doi.org/10.5194/acp-11-5539-2011>, 2011a.](#)

[Alpert, P. A., Aller, J. Y., and Knopf, D. A.: Initiation of the ice phase by marine biogenic surfaces in supersaturated gas and supercooled aqueous phases, *Phys. Chem. Chem. Phys.*, 13, 19882-19894, <https://doi.org/10.1039/c1cp21844a>, 2011b.](#)

[Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud-precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth Sci. Rev.*, 89, 13-41, <https://doi.org/10.1016/j.earscirev.2008.03.001>, 2008.](#)

760 [Arar, E. J. and Collins, G. B.: Method 445.0. *In vitro* determination of chlorophyll *a* and pheophytin *a* in marine and freshwater algae by fluorescence. U.S. Environmental Protection Agency, Cincinnati, Ohio, 1997.](#)

[Arrigo, K. R.: Marine microorganisms and global nutrient cycles, *Nature*, 437, 349-355, <https://doi.org/10.1038/nature04158>, 2005.](#)

Formatted: Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman

765

[Bar-On, Y. M. and Milo, R.: The global mass and average rate of rubisco, Proceedings of the National Academy of Sciences of the United States of America, 116, 4738-4743, 10.1073/pnas.1816654116, 2019.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

770

Bates, T. S., Quinn, P. K., Frossard, A. A., Russell, L. M., Hakala, J., Petaja, T., Kulmala, M., Covert, D. S., Cappa, C. D., Li, S. M., Hayden, K. L., Nuaaman, I., McLaren, R., Massoli, P., Canagaratna, M. R., Onasch, T. B., Sueper, D., Worsnop, D. R., and Keene, W. C.: Measurements of ocean derived aerosol off the coast of California, *J. Geophys. Res. D: Atmos.*, 117, 13, <https://doi.org/10.1029/2012jd017588>, 2012.

Formatted: Font: (Default) +Headings (Times New Roman)

775

Bates, T. S., Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L., Saliba, G., Lewis, S., Graff, J., Russell, L. M., and Behrenfeld, M. J.: Variability in marine plankton ecosystems are not observed in freshly emitted sea spray aerosol over the North Atlantic Ocean, *Geophysical Research Letters*, 47, 9, <https://doi.org/10.1029/2019gl085938>, 2020.

780

[Beall, C. M., Hill, T. C. J., DeMott, P. J., Koneman, T., Pikridas, M., Drewnick, F., Harder, H., Pohlker, C., Lelieveld, J., Weber, B., Iakovides, M., Prokes, R., Sciare, J., Andreae, M. O., Stokes, M. D., and Prather, K. A.: Ice-nucleating particles near two major dust source regions, *Atmos. Chem. Phys.*, 22, 12607-12627, 10.5194/acp-22-12607-2022, 2022.](#)

Formatted: Font: (Default) +Headings (Times New Roman), 10 pt

Formatted: Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt

785

[Beardall, J., Allen, D., Bragg, J., Finkel, Z. V., Flynn, K. J., Quigg, A., Rees, T. A. V., Richardson, A., and Raven, J. A.: Allometry and stoichiometry of unicellular, colonial and multicellular phytoplankton, *New Phytol.*, 181, 295-309, 10.1111/j.1469-8137.2008.02660.x, 2009.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

790

Berges, J. A., Franklin, D. J., and Harrison, P. J.: Evolution of an artificial seawater medium: Improvements in enriched seawater, artificial water over the last two decades, *J. Phycol.*, 37, 1138-1145, <https://doi.org/10.1046/j.1529-8817.2001.01052.x>, 2001.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) +Headings (Times New Roman)

795

Behrenfeld, M. J., Brooks, S. D., Gaube, P., and Mojica, K. D. A.: Editorial: Unraveling Mechanisms Underlying Annual Plankton Blooms in the North Atlantic and Their Implications for Biogenic Aerosol Properties and Cloud Formation, *Front. Mar. Sci.*, 8, 764035, <https://doi.org/10.3389/fmars.2021.764035>, 2021.

Berman-Frank, I., Rosenberg, G., Levitan, O., Haramaty, L., and Mari, X.: Coupling between autocatalytic cell death and transparent exopolymeric particle production in the marine cyanobacterium *Trichodesmium*, *Environ. Microbiol.*, 9, 1415-1422, <https://doi.org/10.1111/j.1462-2920.2007.01257.x>, 2007.

800

Behrenfeld, M. J., Moore, R. H., Hostetler, C. A., Graff, J., Gaube, P., Russell, L. M., Chen, G., Doney, S. C., Giovannoni, S., Liu, H. Y., Proctor, C., Bolalios, L. M., Baetge, N., Davie-Martin, C., Westberry, T. K., Bates, T. S., Bell, T. G., Bidle, K. D., Boss, E. S., Brooks, S. D., Cairns, B., Carlson, C., Halsey, K., Harvey, E. L., Hu, C. M., Karp-Boss, L., Kleb, M., Menden-Deuer, S., Morison, F., Quinn, P. K., Scarino, A. J., Anderson, B., Chowdhary, J., Crosbie, E., Ferrare, R., Haire, J. W., Hu, Y. X., Janz, S., Redemann, J., Saltzman, E., Shook, M., Siegel, D. A., Wisthaler, A., Martine, M. Y., and Ziemba, L.: The North Atlantic Aerosol and Marine Ecosystem Study (NAAMES): Science Motive and Mission Overview, *Front. Mar. Sci.*, 6, 122, <https://doi.org/10.3389/fmars.2019.00122>, 2019.

- 805 Bolaños, L. M., Choi, C. J., Worden, A. Z., Baetge, N., Carlson, C. A., and Giovannoni, S.: Seasonality of the Microbial Community Composition in the North Atlantic, *Front. Mar. Sci.*, **8**, 624164, <https://doi.org/10.3389/fmars.2021.624164>, 2021.
- Bouchard, J. N. and Purdie, D. A.: Effect of elevated temperature, darkness, and hydrogen peroxide treatment on oxidative stress and cell death in the bloom forming toxic cyanobacterium *Microcystis aeruginosa*, *J. Phycol.*, **47**, 1316-1325, <https://doi.org/10.1111/j.1529-8817.2011.01074.x>, 2011.
- 810
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., Zhang, X. Y.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 571 – 658, <https://doi.org/10.1017/CBO9781107415324.016>, 2014.
- 815
- Bratbak, G., Egge, J. K., and Heldal, M.: Viral mortality of the marine alga *Emiliania huxleyi* (Haptophyceae) and termination of algal blooms, *Mar. Ecol. Prog. Ser.*, **93**, 39-48, <https://doi.org/10.3354/meps093039>, 1993.
- 820
- Brooks, S. D. and Thornton, D. C. O.: Marine Aerosols and Clouds, *Annu. Rev. Mar. Sci.*, **10**, 289-313, <https://doi.org/10.1146/annurev-marine-121916-063148>, 2018.
- 825
- Coble, P. G.: Characterization of marine and terrestrial DOM in seawater using excitation emission matrix spectroscopy, *Mar. Chem.*, **51**, 325-346, [https://doi.org/10.1016/0304-4203\(95\)00062-3](https://doi.org/10.1016/0304-4203(95)00062-3), 1996.
- Coble, P. G.: Marine optical biogeochemistry: The chemistry of ocean color, *Chem. Rev.*, **107**, 402-418, <https://doi.org/10.1021/cr050350+>, 2007.
- 830
- Cascajo-Castresana, M., David, R. O., Iriarte-Alonso, M. A., Bittner, A. M., and Marcolli, C.: Protein aggregates nucleate ice: the example of apoferritin, *Atmos. Chem. Phys.*, **20**, 3291-3315, <https://doi.org/10.5194/acp-20-3291-2020>, 2020.
- Cochran, R. E., Ryder, O. S., Grassian, V. H., and Prather, K. A.: Sea Spray Aerosol: The Chemical Link between the Oceans, Atmosphere, and Climate, *Accounts Chem. Res.*, **50**, 599-604, <https://doi.org/10.1021/acs.accounts.6b00603>, 2017.
- 835
- Collier, K. N. and Brooks, S. D.: Role of organic hydrocarbons in atmospheric ice formation via contact freezing, *J. Phys. Chem. A*, **120**, 10169-10180, <https://doi.org/10.1021/acs.jpca.6b11890>, 2016.
- 840
- Croft, B., Martin, R. V., Moore, R. H., Ziemba, L. D., Crosbie, E. C., Liu, H. Y., Russell, L. M., Saliba, G., Wisthaler, A., Muller, M., Schiller, A., Gal, M., Chang, R. Y. W., McDuffie, E. E., Bilsback, K. R., and Pierce, J. R.: Factors controlling marine aerosol size distributions and their climate effects over the northwest Atlantic Ocean region, *Atmos. Chem. Phys.*, **21**, 1889-1916, <https://doi.org/10.5194/acp-21-1889-2021>, 2021.

845 [Cunliffe, M., Engel, A., Frka, S., Gasparovic, B., Guitart, C., Murrell, J. C., Salter, M., Stolle, C., Upstill-Goddard, R., and Wurl, P.: Sea surface microlayers: A unified physicochemical and biological perspective of the air-ocean interface, *Prog. Oceanogr.*, 109, 104-116, 10.1016/j.pocean.2012.08.004, 2013.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

Daily, M. I., Tarn, M. D., Whale, T. F., and Murray, B. J.: An evaluation of the heat test for the ice-nucleating ability of minerals and biological material, *Atmos. Meas. Tech.*, 15, 2635-2665, <https://doi.org/10.5194/amt-15-2635-2022>, 2022.

Della Penna, A. and Gaube, P.: Overview of (sub)mesoscale ocean dynamics for the NAAMES field program, *Front. Mar. Sci.*, 6, 384, <https://doi.org/10.3389/fmars.2019.00384>, 2019.

855 [DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 11217-11222, 10.1073/pnas.0910818107, 2010.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

DeMott, P. J., Hill, T. C. J., McCluskey, C. S., Prather, K. A., Collins, D. B., Sullivan, R. C., Ruppel, M. J., Mason, R. H., Irish, V. E., Lee, T., Hwang, C. Y., Rhee, T. S., Snider, J. R., McMeeking, G. R., Dhaniyala, S., Lewis, E. R., Wentzell, J. J. B., Abbatt, J., Lee, C., Sultana, C. M., Ault, A. P., Axson, J. L., Martinez, M. D., Venero, I., Santos-Figueroa, G., Stokes, M. D., Deane, G. B., Mayol-Bracero, O. L., Grassian, V. H., Bertram, T. H., Bertram, A. K., Moffett, B. F., and Franc, G. D.: Sea spray aerosol as a unique source of ice nucleating particles, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 5797-5803, <https://doi.org/10.1073/pnas.1514034112>, 2016.

865

Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A., and Smith, F.: Colorimetric method for determination of sugars and related substances, *Anal. Chem.*, 28, 350-356, 1956.

Dyrhman, S. T., Jenkins, B. D., Ryneerson, T. A., Saito, M. A., Mercier, M. L., Alexander, H., Whitney, L. P., Drzewianowski, A., Bulygin, V. V., Bertrand, E. M., Wu, Z. J., Benitez-Nelson, C., and Heithoff, A.: The transcriptome and proteome of the diatom *Thalassiosira pseudonana* reveal a diverse phosphorus stress response, *Plos One*, 7, <https://doi.org/10.1371/journal.pone.0033768>, 2012.

875 [Edwards, K. F., Thomas, M. K., Klausmeier, C. A., and Litchman, E.: Allometric scaling and taxonomic variation in nutrient utilization traits and maximum growth rate of phytoplankton, *Limnology and Oceanography*, 57, 554-566, 10.4319/lo.2012.57.2.0554, 2012.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

[Ellis, R. J.: Most abundant protein in the world, *Trends in Biochemical Sciences*, 4, 241-244, 10.1016/0968-0004\(79\)90212-3, 1979.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

880 [Elser, J. J., Sterner, R. W., Gorokhova, E., Fagan, W. F., Markow, T. A., Cotner, J. B., Harrison, J. F., Hobbie, S. E., Odell, G. M., and Weider, L. J.: Biological stoichiometry from genes to ecosystems, *Ecol. Lett.*, 3, 540-550, 10.1046/j.1461-0248.2000.00185.x, 2000.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

[Emerson, S.R., Hedges, J. I. *Chemical oceanography and the marine carbon cycle*. Cambridge University Press, New York, NY, 2008.](#)

Formatted: Line spacing: 1.5 lines

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: 10 pt

Engel, A.: 2009. Determination of marine gel particles, in: Practical Guidelines for the Analysis of Seawater, edited by Wurl, O.
885 CRC Press, Taylor & Francis Group, Boca Raton, Florida, USA, 125-142, <https://doi.org/10.1080/17451000903514220>, 2009.

Engel, A., Borchard, C., Loginova, A., Meyer, J., Hauss, H., and Kiko, R.: Effects of varied nitrate and phosphate supply on polysaccharidic and proteinaceous gel particle production during tropical phytoplankton bloom experiments, *Biogeosciences*, 12, 5647-5665, <https://doi.org/10.5194/bg-12-5647-2015>, 2015.

890

[Flynn, K. J., Raven, J. A., Rees, T. A. V., Finkel, Z., Quigg, A., and Beardall, J.: Is the growth rate hypothesis applicable to microalgae?, *Journal of Phycology*, 46, 1-12, 10.1111/j.1529-8817.2009.00756.x, 2010.](https://doi.org/10.1111/j.1529-8817.2009.00756.x)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt

Fornea, A. P., Brooks, S. D., Dooley, J. B., and Saha, A.: Heterogeneous freezing of ice on atmospheric aerosols containing ash,
895 soot, and soil, *J. Geophys. Res. D: Atmos.*, 114, <https://doi.org/10.1029/2009jd011958>, 2009.

Franklin, D. J., Airs, R. L., Fernandes, M., Bell, T. G., Bongaerts, R. J., Berges, J. A., and Malin, G.: Identification of senescence and death in *Emiliania huxleyi* and *Thalassiosira pseudonana*: Cell staining, chlorophyll alterations, and dimethylsulfoniopropionate (DMSP) metabolism, *Limnol. Oceanogr.*, 57, 305-317, <https://doi.org/10.4319/lo.2012.57.1.0305>,
900 2012.

Fuentes, E., Coe, H., Green, D., de Leeuw, G., and McFiggans, G.: Laboratory-generated primary marine aerosol via bubble-bursting and atomization, *Atmos. Meas. Tech.*, 3, 141-162, 10.5194/amt-3-141-2010, 2010a.

Fuentes, E., Coe, H., Green, D., de Leeuw, G., and McFiggans, G.: On the impacts of phytoplankton-derived organic matter on the
905 properties of the primary marine aerosol - Part 1: Source fluxes, *Atmos. Chem. Phys.*, 10, 9295-9317, <https://doi.org/10.5194/acp-10-9295-2010>, 2010b.

Geider, R. J., MacIntyre, H. L., and Kana, T. M.: Dynamic model of phytoplankton growth and acclimation: Responses of the balanced growth rate and the chlorophyll a:carbon ratio to light, nutrient-limitation and temperature, *Mar. Ecol. Prog. Ser.*, 148,
910 187-200, <https://doi.org/10.3354/meps148187>, 1997.

Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic regulatory model of phytoplanktonic acclimation to light, nutrients, and temperature, *Limnol. Oceanogr.*, 43, 679-694, <https://doi.org/10.4319/lo.1998.43.4.0679>, 1998.

915 Geider, R. J. and La Roche, J.: Redfield revisited: variability of C : N : P in marine microalgae and its biochemical basis, *Eur. J. Phycol.*, 37, 1-17, <https://doi.org/10.1017/s0967026201003456>, 2002.

Gantt, B. and Meskhidze, N.: The physical and chemical characteristics of marine primary organic aerosol: a review, *Atmos. Chem. Phys.*, 13, 3979-3996, <https://doi.org/10.5194/acp-13-3979-2013>, 2013.

920

Genty, B., Briantais, J. M., and Baker, N. R.: The relationship between the quantum yield of photosynthetic electron-transport and quenching of chlorophyll fluorescence, *Biochim. Biophys. Acta*, 990, 87-92, [https://doi.org/10.1016/s0304-4165\(89\)80016-9](https://doi.org/10.1016/s0304-4165(89)80016-9), 1989.

925 Graff, J. R. and Behrenfeld, M. J.: Photoacclimation responses in subarctic Atlantic phytoplankton following a natural mixing-restratification event, *Front. Mar. Sci.*, 5, 209, <https://doi.org/10.3389/fmars.2018.00209>, 2018.

Graff, J. R., Milligan, A. J., and Behrenfeld, M. J.: The measurement of phytoplankton biomass using flow-cytometric sorting and elemental analysis of carbon, *Limnol. Oceanogr. Methods*, 10, 910-920, <https://doi.org/10.4319/lom.2012.10.910>, 2012.

930

Graff, J. R., Westberry, T. K., Milligan, A. J., Brown, M. B., Dall'Olmo, G., van Dongen-Vogels, V., Reifel, K. M., and Behrenfeld, M. J.: Analytical phytoplankton carbon measurements spanning diverse ecosystems, *Deep Sea Res. Part I*, 102, 16-25, <https://doi.org/10.1016/j.dsr.2015.04.006>, 2015.

935 Guillard, R. R. L. and Hargraves, P. E.: *Stichochrysis immobilis* is a diatom, not a Chrysophyte, *Phycologia*, 32, 234-236, <https://doi.org/10.2216/i0031-8884-32-3-234.1>, 1993.

Guillard, R. R. L. and Sieracki, M. S.: Counting cells in cultures with the light microscope, in: *Algal Culturing Techniques*, edited by: Andersen R. A. Elsevier Academic Press, Burlington, MA, USA, 239-252, 2005.

940

[Hansell, D. A.: Recalcitrant Dissolved Organic Carbon Fractions, in: Annual Review of Marine Science, Vol 5, edited by: Carlson, C. A. and Giovannoni, S. J., Annual Review of Marine Science, Annual Reviews, Palo Alto, 421-445, 10.1146/annurev-marine-120710-100757, 2013.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

945 Hasencz, E. S., Jayarathne, T., Pendergraft, M. A., Santander, M. V., Mayer, K. J., Sauer, J., Lee, C., Gibson, W. S., Kruse, S. M., Malfatti, F., Prather, K. A., and Stone, E. A.: Marine bacteria affect saccharide enrichment in sea spray aerosol during a phytoplankton bloom, *ACS Earth Space Chem.*, 4, 1638-1649, <https://doi.org/10.1021/acsearthspacechem.0c00167>, 2020.

Hartmann, S., Ling, M., Dreyer, L. S. A., Zipori, A., Finster, K., Grawe, S., Jensen, L., Borck, S., Reicher, N., Drace, T., Niedermeier, D., Jones, N., Hoffmann, S. V., Wex, H., Rudich, Y., Boesen, T., and Santl-Temkiv, T.: Structure and protein-protein interactions of ice nucleation proteins drive their activity, *Front. Microbiol.*, 13, 872306, <https://doi.org/10.3389/fmicb.2022.872306>, 2022.

955 Harrison, P. J., Waters, R. E., and Taylor, F. J. R.: A broad-spectrum artificial seawater medium for coastal and open ocean phytoplankton, *J. Phycol.*, 16, 28-35, <https://doi.org/10.1111/j.1529-8817.1980.tb00724.x>, 1980.

[Hill, T. C. J., Malfatti, F., McCluskey, C. S., Schill, G. P., Santander, M. V., Moore, K. A., Rauker, A. M., Perkins, R. J., Celussi, M., Levin, E. J. T., Suski, K. J., Cornwell, G. C., Lee, C. S. P., Del Negro, P., Kreidenweis, S. M., Prather, K. A., and DeMott, P. J.: Resolving the controls over the production and emission of ice-nucleating particles in sea spray, *Environ. Sci. - Atmospheres*, 21, 10.1039/d2ea00154c, 2023.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

960 [Ho, T. Y., Quigg, A., Finkel, Z. V., Milligan, A. J., Wyman, K., Falkowski, P. G., and Morel, F. M. M.: The elemental composition of some marine phytoplankton, *Journal of Phycology*, 39, 1145-1159, 10.1111/j.0022-3646.2003.03-090.x, 2003.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

965 Hoose, C. and Mohler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817-9854, <https://doi.org/10.5194/acp-12-9817-2012>, 2012.

Hoppe, H. G.: Significance of exoenzymatic activities in the ecology of brackish water - measurements by means of methylumbelliferyl substrates, *Mar. Ecol. Prog. Ser.*, 11, 299-308, <https://doi.org/10.3354/meps011299>, 1983.

970

Hudson, J. G. and Noble, S.: Cumulus cloud and drizzle microphysics relationships with complete CCN spectra, *J. Geophys. Res. D: Atmos.*, 126, 20, <https://doi.org/10.1029/2021jd034966>, 2021.

975 Inomura, K., Omta, A. W., Talmy, D., Bragg, J., Deutsch, C., and Follows, M. J.: A mechanistic model of macromolecular allocation, elemental stoichiometry, and growth rate in phytoplankton, *Front. Microbiol.*, 11, 86, <https://doi.org/10.3389/fmicb.2020.00086>, 2020.

980 Irish, V. E., Elizondo, P., Chen, J., Chou, C., Charette, J., Lizotte, M., Ladino, L. A., Wilson, T. W., Gosselin, M., Murray, B. J., Polishchuk, E., Abbatt, J. P. D., Miller, L. A., and Bertram, A. K.: Ice-nucleating particles in Canadian Arctic sea-surface microlayer and bulk seawater, *Atmos. Chem. Phys.*, 17, 10583-10595, <https://doi.org/10.5194/acp-17-10583-2017>, 2017.

[Jsanta-Navarro, J., Prater, C., Peoples, L. M., Loladze, I., Phan, T., Jevasingh, P. D., Church, M. J., Kuang, Y., and Elser, J. J.: Revisiting the growth rate hypothesis: Towards a holistic stoichiometric understanding of growth. *Ecol. Lett.*, 25, 2324-2339, 10.1111/ele.14096, 2022.](#)

985

Jeffrey, S. W. and Humphrey, G. F.: New spectrophotometric equations for determining chlorophylls *a*, *b*, *c1* and *c2* in higher-plants, algae and natural phytoplankton, *Biochem. Physiol. Pflanzen*, 167, 191-194, [https://doi.org/10.1016/s0015-3796\(17\)30778-3](https://doi.org/10.1016/s0015-3796(17)30778-3), 1975.

990 Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Kramer, M.: Overview of Ice Nucleating Particles, in: *Ice Formation and Evolution in Clouds and Precipitation: Measurement and Modeling Challenges*, edited by: Baumgardner, D., McFarquhar, G. M., and Heymsfield, A. J., *Meteorological Monographs*, Amer Meteorological Society, Boston, <https://doi.org/10.1175/amsmonographs-d-16-0006.1>, 2017.

995 Klausmeier, C. A., Litchman, E., Daufresne, T., and Levin, S. A.: Optimal nitrogen-to-phosphorus stoichiometry of phytoplankton, *Nature*, 429, 171-174, <https://doi.org/10.1038/nature02454>, 2004.

Knopf, D. A., Alpert, P. A., Wang, B., and Aller, J. Y.: Stimulation of ice nucleation by marine diatoms, *Nat. Geosci.*, 4, 88-90, <https://doi.org/10.1038/ngeo1037>, 2011.

1000

Kothawala, D. N., Murphy, K. R., Stedmon, C. A., Weyhenmeyer, G. A., and Tranvik, L. J.: Inner filter correction of dissolved organic matter fluorescence, *Limnol. Oceanogr. Methods*, 11, 616-630, <https://doi.org/10.4319/lom.2013.11.616>, 2013.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

1005 Kranzler, C. F., Krause, J. W., Brzezinski, M. A., Edwards, B. R., Biggs, W. P., Maniscalco, M., McCrow, J. P., Van Mooy, B. A. S., Bidle, K. D., Allen, A. E., and Thamatrakoln, K.: Silicon limitation facilitates virus infection and mortality of marine diatoms, *Nat. Microbiol.*, 4, 1790-1797, <https://doi.org/10.1038/s41564-019-0502-x>, 2019.

1010 Kujawinski, E. B.: The impact of microbial metabolism on marine dissolved organic matter, *Annu. Rev. Mar. Sci.*, 3, 567-599, <https://doi.org/10.1146/annurev-marine-120308-081003>, 2011.

Lawaetz, A. J. and Stedmon, C. A.: Fluorescence intensity calibration using the Raman scatter peak of water, *Appl. Spectrosc.*, 63, 936-940, <https://doi.org/10.1366/000370209788964548>, 2009.

1015 [Lewis, S. L., Russell, L. M., Saliba, G., Quinn, P. K., Bates, T. S., Carlson, C. A., Baetge, N., Aluwihare, L. I., Boss, E., Frossard, A. A., Bell, T. G., and Behrenfeld, M. J.: Characterization of Sea Surface Microlayer and Marine Aerosol Organic Composition Using STXM-NEXAFS Microscopy and FTIR Spectroscopy. *ACS Earth and Space Chemistry*, 15, 10.1021/acsearthspacechem.2c00119, 2022.](#) **Formatted:** Font: (Default) Times New Roman, 10 pt
Formatted: Line spacing: 1.5 lines
Formatted: Font: (Default) Times New Roman, 10 pt

1020 Liefer, J. D., Garg, A., Fyfe, M. H., Irwin, A. J., Benner, I., Brown, C. M., Follows, M. J., Omta, A. W., and Finkel, Z. V.: The macromolecular basis of phytoplankton C:N:P under nitrogen starvation, *Front. Microbiol.*, 10, 763, <https://doi.org/10.3389/fmicb.2019.00763>, 2019.

Lin, S. J., Litaker, R. W., and Sunda, W. G.: Phosphorus physiological ecology and molecular mechanisms in marine phytoplankton, *J. Phycol.*, 52, 10-36, <https://doi.org/10.1111/jpy.12365>, 2016.

1025 Logan, B. E., Grossart, H. P., and Simon, M.: Direct observation of phytoplankton, TEP and aggregates on polycarbonate filters using brightfield microscopy, *J. Plankton Res.*, 16, 1811-1815, <https://doi.org/10.1093/plankt/16.12.1811>, 1994.

1030 Lohmann, U., Lüönd, F., and Mahrt, F.: *An Introduction to Clouds: From the Microscale to Climate*, Cambridge University Press, Cambridge, United Kingdom, 391 pp., 2016

Long, R. A. and Azam, F.: Abundant protein-containing particles in the sea, *Aquatic Microb. Ecol.*, 10, 213-221, <https://doi.org/10.3354/ame010213>, 1996.

1035 Lukas, M., Schwidetzky, R., Eufemio, R. J., Bonn, M., and Meister, K.: Toward understanding bacterial ice nucleation, *J. Phys. Chem. B*, 126, 1861-1867, <https://doi.org/10.1021/acs.jpcc.1c09342>, 2022.

1040 [Maki, L. R., Galyan, E. L., Changchi, M., and Caldwell, D. R.: Ice nucleation induced by *Pseudomonas syringae*, *Applied Microbiology* 28, 456-459. \[10.1128/aem.28.3.456-459.1974\]\(https://doi.org/10.1128/aem.28.3.456-459.1974\), 1974.](#) **Formatted:** Font: (Default) Times New Roman, 10 pt
Formatted: Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt, Italic
Formatted: Font: (Default) Times New Roman, 10 pt

Mansour, K., Decesari, S., Facchini, M. C., Belosi, F., Paglione, M., Sandrini, S., Bellacicco, M., Marullo, S., Santoleri, R., Ovadnevaite, J., Ceburnis, D., O'Dowd, C., Roberts, G., Sanchez, K., and Rinaldi, M.: Linking marine biological activity to aerosol chemical composition and cloud-relevant properties over the North Atlantic Ocean, *J. Geophys. Res. D: Atmos.*, 125, 19, <https://doi.org/10.1029/2019jd032246>, 2020.

- 1045 Marx, M. C., Wood, M., and Jarvis, S. C.: A microplate fluorimetric assay for the study of enzyme diversity in soils, *Soil Biol. Biochem.*, 33, 1633-1640, [https://doi.org/10.1016/s0038-0717\(01\)00079-7](https://doi.org/10.1016/s0038-0717(01)00079-7), 2001.
- Maxwell, K. and Johnson, G. N.: Chlorophyll fluorescence - a practical guide, *J. Exp. Bot.*, 51, 659-668, <https://doi.org/10.1093/jexbot/51.345.659>, 2000.
- 1050 Mayol, E., Arrieta, J. M., Jimenez, M. A., Martinez-Asensio, A., Garcias-Bonet, N., Dachs, J., Gonzalez-Gaya, B., Royer, S. J., Benitez-Barrios, V. M., Fraile-Nuez, E., and Duarte, C. M.: Long-range transport of airborne microbes over the global tropical and subtropical ocean, *Nat. Commun.*, 8, <https://doi.org/10.1038/s41467-017-00110-9>, 2017.
- 1055 McCluskey, C. S., Hill, T. C. J., Malfatti, F., Sultana, C. M., Lee, C., Santander, M. V., Beall, C. M., Moore, K. A., Cornwell, G. C., Collins, D. B., Prather, K. A., Jayarathne, T., Stone, E. A., Azam, F., Kreidenweis, S. M., and DeMott, P. J.: A dynamic link between ice nucleating particles released in nascent sea spray aerosol and oceanic biological activity during two mesocosm experiments, *J. Atmos. Sci.*, 74, 151-166, <https://doi.org/10.1175/jas-d-16-0087.1>, 2017.
- 1060 McCluskey, C. S., Hill, T. C. J., Sultana, C. M., Laskina, O., Trueblood, J., Santander, M. V., Beall, C. M., Michaud, J. M., Kreidenweis, S. M., Prather, K. A., Grassian, V., and DeMott, P. J.: A mesocosm double feature: insights into the chemical makeup of marine ice nucleating particles, *J. Atmos. Sci.*, 75, 2405-2423, <https://doi.org/10.1175/jas-d-17-0155.1>, 2018.
- Møller, E. F.: Production of dissolved organic carbon by sloppy feeding in the copepods *Acartia tonsa*, *Centropages typicus*, and
1065 *Temora longicornis*, *Limnol. Oceanogr.*, 52, 79-84, <https://doi.org/10.4319/lo.2007.52.1.0079>, 2007.
- Møller, E. F., Thor, P., and Nielsen, T. G.: Production of DOC by *Calanus finmarchicus*, *C. glacialis* and *C. hyperboreus* through sloppy feeding and leakage from fecal pellets, *Mar. Ecol. Prog. Ser.*, 262, 185-191, <https://doi.org/10.3354/meps262185>, 2003.
- 1070 Moore, C. M., Suggett, D. J., Hickman, A. E., Kim, Y. N., Tweddle, J. F., Sharples, J., Geider, R. J., and Holligan, P. M.: Phytoplankton photoacclimation and photoadaptation in response to environmental gradients in a shelf sea, *Limnol. Oceanogr.*, 51, 936-949, <https://doi.org/10.4319/lo.2006.51.2.0936>, 2006.
- Morison, F., Harvey, E., Franze, G., and Menden-Deuer, S.: Storm-induced predator-prey decoupling promotes springtime
1075 accumulation of North Atlantic phytoplankton, *Front. Mar. Sci.*, 6, 608, <https://doi.org/10.3389/fmars.2019.00608>, 2019.
- O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and Putaud, J. P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676-680, <https://doi.org/10.1038/nature02959>, 2004.
- 1080 O'Dowd, C., Ceburnis, D., Ovadnevaite, J., Bialek, J., Stengel, D. B., Zacharias, M., Nitschke, U., Connan, S., Rinaldi, M., Fuzzi, S., Decesari, S., Facchini, M. C., Marullo, S., Santoleri, R., Dell'Anno, A., Corinaldesi, C., Tangherlini, M., and Danovaro, R.: Connecting marine productivity to sea-spray via nanoscale biological processes: Phytoplankton dance or death disco?, *Sci. Rep.*, 5, 14883, <https://doi.org/10.1038/srep14883>, 2015.

1085 Palmer, T.: Short-term tests validate long-term estimates of climate change, *Nature*, 582, 185-186, <https://doi.org/10.1038/d41586-020-01484-5>, 2020.

Parsons, T. R., Maita, Y. and Lalli, C. M.: *A Manual of Chemical and Biological Methods for Seawater Analysis*. Pergamon Press, Oxford, United Kingdom, 172 pp., 1984.

1090

Passow, U. and Alldredge, A. L.: A dye-binding assay for the spectrophotometric measurement of transparent exopolymer particles (TEP), *Limnol. Oceanogr.*, 40, 1326-1335, <https://doi.org/10.4319/lo.1995.40.7.1326>, 1995.

1095 Passow, U.: Transparent exopolymer particles (TEP) in aquatic environments, *Prog. Oceanogr.*, 55, 287-333, [https://doi.org/10.1016/s0079-6611\(02\)00138-6](https://doi.org/10.1016/s0079-6611(02)00138-6), 2002.

Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M., Roberts, G. C., Russell, L. M., Ault, A. P., Baltrusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L., Hersey, S. P., Kim, M. J., Lambert, W. F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C., and Zhao, D. F.: Bringing the ocean into the laboratory to probe the chemical complexity of sea spray aerosol, *Proc. Natl. Acad. Sci. U.S.A.*, 110, 7550-7555, <https://doi.org/10.1073/pnas.1300262110>, 2013.

1105 [Perkins, R. J., de Vasquez, M. G. V., Beasley, E. E., Hill, T. C. J., Stone, E. A., Allen, H. C., and DeMott, P. J.: Relating Structure and Ice Nucleation of Mixed Surfactant Systems Relevant to Sea Spray Aerosol, *Journal of Physical Chemistry A*, 124, 8806-8821, \[10.1021/acs.jpca.0c05849\]\(https://doi.org/10.1021/acs.jpca.0c05849\), 2020.](#)

Formatted: Line spacing: 1.5 lines

1110 [Quigg, A., Finkel, Z. V., Irwin, A. J., Rosenthal, Y., Ho, T. Y., Reinfelder, J. R., Schofield, O., Morel, F. M. M., and Falkowski, P. G.: The evolutionary inheritance of elemental stoichiometry in marine phytoplankton, *Nature*, 425, 291-294, \[10.1038/nature01953\]\(https://doi.org/10.1038/nature01953\), 2003.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

Quinn, P. K., Bates, T. S., Schulz, K. S., Coffman, D. J., Frossard, A. A., Russell, L. M., Keene, W. C., and Kieber, D. J.: Contribution of sea surface carbon pool to organic matter enrichment in sea spray aerosol, *Nat. Geosci.*, 7, 228-232, <https://doi.org/10.1038/ngeo2092>, 2014.

1115 Quinn, P. K., Bates, T. S., Coffman, D. J., Upchurch, L., Johnson, J. E., Moore, R., Ziemba, L., Bell, T. G., Saltzman, E. S., Graff, J., and Behrenfeld, M. J.: Seasonal variations in western North Atlantic remote marine aerosol properties, *J. Geophys. Res. D: Atmos.*, 124, 14240-14261, <https://doi.org/10.1029/2019jd031740>, 2019.

1120 Rastelli, E., Corinaldesi, C., Dell'Anno, A., Lo Martire, M., Greco, S., Facchini, M. C., Rinaldi, M., O'Dowd, C., Ceburnis, D., and Danovaro, R.: Transfer of labile organic matter and microbes from the ocean surface to the marine aerosol: an experimental approach, *Sci Rep*, 7, 10, <https://doi.org/10.1038/s41598-017-10563-z>, 2017.

[Raven, J. A.: Rubisco: still the most abundant protein of Earth? *New Phytol.*, 198, 1-3, \[10.1111/nph.12197\]\(https://doi.org/10.1111/nph.12197\), 2013.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

1125 Redfield, A. C.: The biological control of chemical factors in the environment, *Am. Scientist*, 46, 205-221, 1958.

[Rinaldi, M., Fuzzi, S., Decesari, S., Marullo, S., Santoleri, R., Provenzale, A., von Hardenberg, J., Ceburnis, D., Vaishya, A., O'Dowd, C. D., and Facchini, M. C.: Is chlorophyll-a the best surrogate for organic matter enrichment in submicron primary marine aerosol?, *Journal of Geophysical Research-Atmospheres*, 118, 4964-4973, 10.1002/jgrd.50417, 2013.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

1130

Roeters, S. J., Golbek, T. W., Bregnhøj, M., Drace, T., Alamdari, S., Roseboom, W., Kramer, G., Santl-Temkiv, T., Finster, K., Pfaendtner, J., Woutersen, S., Boesen, T., and Weidner, T.: Ice-nucleating proteins are activated by low temperatures to control the structure of interfacial water, *Nat. Commun.*, 12, 1183, <https://doi.org/10.1038/s41467-021-21349-3>, 2021.

1135 Saliba, G., Chen, C. L., Lewis, S., Russell, L. M., Rivellini, L. H., Lee, A. K. Y., Quinn, P. K., Bates, T. S., Haentjens, N., Boss, E. S., Karp-Boss, L., Baetge, N., Carlson, C. A., and Behrenfeld, M. J.: Factors driving the seasonal and hourly variability of sea-spray aerosol number in the North Atlantic, *Proc. Natl. Acad. Sci. U.S.A.*, 116, 20309-20314, <https://doi.org/10.1073/pnas.1907574116>, 2019.

1140 Saliba, G., Chen, C. L., Lewis, S., Russell, L. M., Quinn, P. K., Bates, T. S., Bell, T. G., Lawler, M. J., Saltzman, E. S., Sanchez, K. J., Moore, R., Shook, M., Rivellini, L. H., Lee, A., Baetge, N., Carlson, C. A., and Behrenfeld, M. J.: Seasonal Differences and Variability of Concentrations, Chemical Composition, and Cloud Condensation Nuclei of Marine Aerosol Over the North Atlantic, *J. Geophys. Res. D: Atmos.*, 125, 24, <https://doi.org/10.1029/2020jd033145>, 2020.

1145 Sanchez, K. J., Roberts, G. C., Saliba, G., Russell, L. M., Twohy, C., Reeves, M. J., Humphries, R. S., Keywood, M. D., Ward, J. P., and McRobert, I. M.: Measurement report: Cloud processes and the transport of biological emissions affect Southern Ocean particle and cloud condensation nuclei concentrations, *Atmos. Chem. Phys.*, 21, 3427-3446, <https://doi.org/10.5194/acp-21-3427-2021>, 2021.

1150 Schneider, T., Kaul, C. M., and Pressel, K. G.: Possible climate transitions from breakup of stratocumulus decks under greenhouse warming, *Nat. Geosci.*, 12, 163-167, <https://doi.org/10.1038/s41561-019-0310-1>, 2019.

Schwidetzky, R., Lukas, M., YazdanYar, A., Kunert, A. T., Poschl, U., Domke, K. F., Frohlich-Nowoisky, J., Bonn, M., Koop, T., Nagata, Y., and Meister, K.: Specific ion-protein interactions influence bacterial ice nucleation, *Chem. Eur. J.*, 27, 7402-7407, <https://doi.org/10.1002/chem.202004630>, 2021.

[Sciare, J., Favez, O., Sarda-Estève, R., Oikonomou, K., Cachier, H., and Kazan, V.: Long-term observations of carbonaceous aerosols in the Austral Ocean atmosphere: Evidence of a biogenic marine organic source, *Journal of Geophysical Research-Atmospheres*, 114, 10, 10.1029/2009jd011998, 2009.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

1160

Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the

1165 role of aerosol-cloud interactions in the climate system, Proc. Natl. Acad. Sci. U.S.A., 113, 5781-5790,
https://doi.org/10.1073/pnas.1514043113, 2016.

Sellegrì, K., O'Dowd, C. D., Yoon, Y. J., Jennings, S. G., and de Leeuw, G.: Surfactants and submicron sea spray generation, J. Geophys. Res. D: Atmos., 111, 12, https://doi.org/10.1029/2005jd006658, 2006.

1170 [Steinke, I., Hiranuma, N., Funk, R., Hohler, K., Tullmann, N., Umo, N. S., Weidler, P. G., Mohler, O., and Leisner, T.: Complex plant-derived organic aerosol as ice-nucleating particles - more than the sums of their parts? Atmos. Chem. Phys., 20, 11387-11397, 10.5194/acp-20-11387-2020, 2020.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

1175 Stokes, M. D., Deane, G. B., Prather, K., Bertram, T. H., Ruppel, M. J., Ryder, O. S., Brady, J. M., and Zhao, D.: A Marine Aerosol Reference Tank system as a breaking wave analogue for the production of foam and sea-spray aerosols, Atmos. Meas. Tech., 6, 1085-1094, https://doi.org/10.5194/amt-6-1085-2013, 2013.

Thornton, D. C. O.: Diatom aggregation in the sea: mechanisms and ecological implications, Eur. J. Phycol., 37, 149-161, https://doi.org/10.1017/s0967026202003657, 2002.

1180

Thornton, D. C. O.: Dissolved organic matter (DOM) release by phytoplankton in the contemporary and future ocean, Eur. J. Phycol., 49, 20-46, https://doi.org/10.1080/09670262.2013.875596, 2014.

1185 Thornton, D. C. O.: Coomassie Stainable Particles (CSP): Protein Containing Exopolymer Particles in the Ocean, Front. Mar. Sci., 5, 206, https://doi.org/10.3389/fmars.2018.00206, 2018.

[Thornton, D. C. O., Brooks, S. D., and Chen, J.: Protein and Carbohydrate Exopolymer Particles in the Sea Surface Microlayer \(SML\). Frontiers in Marine Science, 3, 135, 10.3389/fmars.2016.00135, 2016.](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

1190 Thornton, D. C. O. and Chen, J.: Exopolymer production as a function of cell permeability and death in a diatom (*Thalassiosira weissflogii*) and a cyanobacterium (*Synechococcus elongatus*), J. Phycol., 53, 245-260, https://doi.org/10.1111/jpy.12470, 2017.

1195 Twohy, C. H., DeMott, P. J., Russell, L. M., Toohey, D. W., Rainwater, B., Geiss, R., Sanchez, K. J., Lewis, S., Roberts, G. C., Humphries, R. S., McCluskey, C. S., Moore, K. A., Selleck, P. W., Keywood, M. D., Ward, J. P., and McRobert, I. M.: Cloud-nucleating particles over the southern ocean in a changing climate, Earth's Future, 9, e2020EF001673, https://doi.org/10.1029/2020ef001673, 2021.

Vali, G.: Quantitative evaluation of experimental results on heterogeneous freezing nucleation of supercooled liquids, J. Atmos. Sci., 28, 402-409, https://doi.org/10.1175/1520-0469(1971)028<0402:Qeoera>2.0.Co;2, 1971.

1200

Vali, G.: Nucleation terminology, Bull. Am. Meteorol. Soc., 66, 1426-1427, 1985.

1205 Vali, G., DeMott, P. J., Mohler, O., and Whale, T. F.: Technical Note: A proposal for ice nucleation terminology, *Atmos. Chem. Phys.*, 15, 10263-10270, <https://doi.org/10.5194/acp-15-10263-2015>, 2015.

Van Heukelem, L. and Thomas, C. S.: Computer-assisted high-performance liquid chromatography method development with applications to the isolation and analysis of phytoplankton pigments, *J. Chromatogr. A*, 910, 31-49, [https://doi.org/10.1016/s0378-4347\(00\)00603-4](https://doi.org/10.1016/s0378-4347(00)00603-4), 2001.

1210 Van Mooy, B. A., Fredricks, H. F., Pedler, B. E., Dyhrman, S. T., Karl, D. M., Koblizek, M., Lomas, M. W., Mincer, T. J., Moore, L. R., Moutin, T., Rappe, M. S., and Webb, E. A.: Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity, *Nature*, 458, 69-72, <https://doi.org/10.1038/nature07659>, 2009.

1215 Vardi, A., Haramaty, L., Van Mooy, B. A. S., Fredricks, H. F., Kimmance, S. A., Larsen, A., and Bidle, K. D.: Host-virus dynamics and subcellular controls of cell fate in a natural coccolithophore population, *Proc. Natl. Acad. Sci. U.S.A.*, 109, 19327-19332, <https://doi.org/10.1073/pnas.1208895109>, 2012.

1220 Veldhuis, M. J. W., Kraay, G. W., and Timmermans, K. R.: Cell death in phytoplankton: correlation between changes in membrane permeability, photosynthetic activity, pigmentation and growth, *Eur. J. Phycol.*, 36, 167-177, <https://doi.org/10.1017/s0967026201003110>, 2001.

1225 Wang, X. F., Sultana, C. M., Trueblood, J., Hill, T. C. J., Malfatti, F., Lee, C., Laskina, O., Moore, K. A., Beall, C. M., McCluskey, C. S., Cornwell, G. C., Zhou, Y. Y., Cox, J. L., Pendergraft, M. A., Santander, M. V., Bertram, T. H., Cappa, C. D., Azam, F., DeMott, P. J., Grassian, V. H., and Prather, K. A.: Microbial control of sea spray aerosol composition: a tale of two blooms, *ACS Cent. Sci.*, 1, 124-131, <https://doi.org/10.1021/acscentsci.5b00148>, 2015.

1230 [Wex, H., Augustin-Bauditz, S., Boose, Y., Budke, C., Curtius, J., Diehl, K., Dreyer, A., Frank, F., Hartmann, S., Hiranuma, N., Jantsch, E., Kanji, Z. A., Kiselev, A., Koop, T., Mohler, O., Niedermeier, D., Nillius, B., Rosch, M., Rose, D., Schmidt, C., Steinke, I., and Stratmann, F.: Intercomparing different devices for the investigation of ice nucleating particles using Snomax \(R\) as test substance, *Atmos. Chem. Phys.*, 15, 1463-1485, \[10.5194/acp-15-1463-2015\]\(https://doi.org/10.5194/acp-15-1463-2015\), 2015.](#)

1235 Wilbourn, E. K., Thornton, D. C. O., Ott, C., Graff, J., Quinn, P. K., Bates, T. S., Betha, R., Russell, L. M., Behrenfeld, M. J., and Brooks, S. D.: Ice nucleation by marine aerosols over the North Atlantic Ocean in late spring, *J. Geophys. Res. D: Atmos.*, 125, e2019JD030913, <https://doi.org/10.1029/2019jd030913>, 2020.

Wilson, T. W., Ladino, L. A., Alpert, P. A., Breckels, M. N., Brooks, I. M., Browse, J., Burrows, S. M., Carslaw, K. S., Huffman, J. A., Judd, C., Kilhau, W. P., Mason, R. H., McFiggans, G., Miller, L. A., Najera, J. J., Polishchuk, E., Rae, S., Schiller, C. L., Si, M., Temprado, J. V., Whale, T. F., Wong, J. P. S., Wurl, O., Yakobi-Hancock, J. D., Abbatt, J. P. D., Aller, J. Y., Bertram, A. K., Knopf, D. A., and Murray, B. J.: A marine biogenic source of atmospheric ice-nucleating particles, *Nature*, 525, 234-238, <https://doi.org/10.1038/nature14986>, 2015.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines

Wolf, M. J., Coe, A., Dove, L. A., Zawadowicz, M. A., Dooley, K., Biller, S. J., Zhang, Y., Chisholm, S. W., and Cziczo, D. J.: Investigating the heterogeneous ice nucleation of sea spray aerosols using *Prochlorococcus* as a model source of marine organic matter, *Environ. Sci. Technol.*, 53, 1139-1149, <https://doi.org/10.1021/acs.est.8b05150>, 2019.

1245

Zack, G. W., Rogers, W. E., and Latt, S. A.: Automatic measurement of sister chromatid exchange frequency, *J. Histochem. Cytochem.*, 25, 741-753, <https://doi.org/10.1177/25.7.70454>, 1977.

Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of higher climate sensitivity in CMIP6 models, *Geophys. Res. Lett.*, 47, e2019GL085782, <https://doi.org/10.1029/2019gl085782>, 2020.

1250

[Zhao, X., Liu, X. H., Burrows, S. M., and Shi, Y.: Effects of marine organic aerosols as sources of immersion-mode ice-nucleating particles on high-latitude mixed-phase clouds, *Atmos. Chem. Phys.*, 21, 2305-2327. \[10.5194/acp-21-2305-2021\]\(https://doi.org/10.5194/acp-21-2305-2021\), 2021.](#)

1255

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), 10 pt

Formatted: Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines