

The role of temporal scales in extracting dominant meteorological drivers of major airborne pollutants

Miaoqing Xu¹, Jing Yang¹, Manchun Li², Xiao Chen¹, Qiancheng Lv¹, Qi Yao¹, Bingbo Gao³, Ziyue Chen^{1*}

5 ¹College of Global and Earth System Sciences, Beijing Normal University, 19 Xijiekou Street, Haidian, Beijing 100875, China

²School of Geography and Ocean Science, Nanjing University, Nanjing 210023, China

³College of Land Science and Technology, China Agricultural University, Beijing 100083, China

Correspondence to: Ziyue Chen (zychen@bnu.edu.cn)

10 **Abstract.** The influence of individual meteorological factors on different airborne pollutants has been massively conducted. However, few studies have considered the effect of temporal scales on the extracted pollutant-meteorology association. Based on Convergent Cross Mapping (CCM), we compared the influence of major meteorological factors on PM_{2.5}, PM₁₀ and O₃ concentrations in 2020 at the 3h and 24h scale respectively. In terms of the extracted dominant meteorological factor, the consistence between the analysis at 3h and 24h scale was relatively low, suggesting a large difference even from a qualitative
15 perspective. In terms of the mean ρ value, the effect of temporal scale on PM (PM_{2.5} and PM₁₀)-Meteorology association was consistent, yet largely different from the temporal-scale effect on O₃. Temperature was the most important meteorological factor for PM_{2.5}, PM₁₀ and O₃ across China at both 3h and 24 scale. For PM_{2.5} and PM₁₀, the extracted PM-temperature association at the 24h scale was stronger than that at the 3h scale. Meanwhile, for summer O₃, due to strong reactions between precursors, the extracted O₃-temperature association at the 3h scale was much stronger. Due to the discrete distribution, the
20 extracted association between all pollutants and precipitation was much weaker at the 3h scale. Similarly, the extracted PM-wind association was notably weaker at the 3h scale. Due to precursor transport, summertime O₃-wind association was stronger at the 3h scale. For atmospheric pressure, the pollutant-pressure association was weaker at the 3h scale except for summer, when interactions between atmospheric pressure and other meteorological factors were strong. From the spatial perspective, pollutant-meteorology association at 3h and 24h was more consistent in those heavily polluted regions. This research suggested
25 that temporal scales should be carefully considered when extracting natural and anthropogenic drivers for airborne pollution.

1 Introduction

Since 2013, PM_{2.5} induced haze events increased dramatically across China (Chen et al., 2020a; Wang et al., 2021a). To address this issue, a series of emission-cut policies were released and strictly implemented, leading to significantly reduced PM_{2.5} concentrations at the national scale (Wang et al., 2021b; Wang et al., 2022; Xiao et al., 2020). Conversely, with the
30 improvement of PM_{2.5} pollution, a soaring ground ozone level was observed since 2013, making composite airborne pollution

a rising challenge (Gong et al., 2017; Zheng et al., 2018). Against this background, a comprehensive understanding of their characteristics and driving factors is key for effectively predicting and managing composite airborne pollutants (Chen et al., 2018, 2019a, 2019c, 2020a).

35 The major influential factors for airborne pollutants are human factors, which closely relates to their compositions and formation (Cheng et al., 2017; Zhan et al., 2017), and meteorological factors, which closely relates to their dispersion (Chen et al., 2020; Guo et al., 2020; Zhang et al., 2020). Given the strong negative effects of airborne pollution on public health (Kelly et al., 2015; Gao et al., 2017; Yin et al., 2020) and crop yields (Zhou et al., 2018; Xu et al., 2021), massive studies have been conducted on the human and meteorological attribution of composite airborne pollution. For meteorological influencing factors, Yang et al. (2021) studied 284 major cities in China based on daily scales and found that $PM_{2.5}$ was mainly affected by wind, temperature and rainfall, while O_3 was mainly affected by temperature, relative humidity and sunshine duration. Wang et al. (2018) established 12 joint regression models and analyzed that the leading meteorological factors of $PM_{2.5}$ pollution in Zhejiang were temperature and wind speed based on the hour-scale data. For emission influencing factors, Wang et al. (2018) found that the emission influencing factor of $PM_{2.5}$ pollution in Zhejiang was NO_2 based on the analysis of hour-scale data. Zhai et al. (2019) estimated the correlation between $PM_{2.5}$ concentration and meteorological factors at the 10-day scale, found that the variation trend of $PM_{2.5}$ and SO_2 , NO_2 and CO was consistent, also SO_2 emission-control was the main driving factor for $PM_{2.5}$ variations. Despite massive studies conducted, notable inconsistency of dominant meteorological and anthropogenic drivers for airborne pollutants was observed between findings from previous studies. In addition to the variation of seasons and geographical locations, the temporal resolution of data sources can be another major reason for the distinct outputs. However, the influence of temporal scales in the attribution of airborne pollution has rarely been investigated.

In recent years, the research on pollutant-meteorology has been massively conducted since 2013 (Chen et al., 2020), yet some gaps remained. Due to the lack of high temporal-resolution data, previous studies were mainly conducted at the daily scale and many scholars may believe that the application of high-temporal-resolution data leads to a better extraction of pollutant-meteorology association. To fill this gap, we employed the data of major airborne pollutants, including $PM_{2.5}$, PM_{10} and O_3 , meteorological factors and some precursors across China with a temporal resolution of 3h and 24h respectively. By comparing the major drivers for airborne pollutants extracted using data sources with different temporal resolution, the role of temporal scales in the attribution of composite airborne pollution can be comprehensively understood. This research aims for an improved understanding of the mechanisms how different factors may affect airborne pollutants under various temporal scales and sheds useful light on a better management of composite airborne pollution through more effective emission-cut.

2 Methodology

2.1 Data sources

Per-3h meteorological data across China for January- December 2020 were obtained from the China Meteorological Administration. The meteorological variables used in this study included temperature, precipitation, wind direction, wind speed and atmospheric pressure, which were closely related to PM_{2.5}, PM₁₀ (Chen et al., 2020a) and O₃ concentrations (Chen et al., 2020b). For cities with more than one observation station, the average of records from multiple stations was employed. For a multi-scale comparison, the 24h meteorological data were produced by conducting average operation on the 3h meteorological data. The previous studies have proved that pollutant-meteorology association presented notable seasonal variations and thus if CCM has been conducted based a whole year, the p value was not significant at many cases and thus the comparison cannot be conducted. Therefore, in this research, we considered the experiments based on seasonal data respectively. For analyzing seasonal variations of pollutant-meteorology association, December, January, February were set as winter, March, April, May as Spring, June, July, August as Summer, September, October, November as Autumn.

Hourly concentration data of PM_{2.5}, PM₁₀ and O₃ during the same period were obtained from China National Environmental Monitoring Center, CNEMC. The meteorological data were matched according to cities and air pollutant stations, and the nearest station corresponding to each air pollution monitoring station was selected as its surrounding meteorological conditions. A total of 101 cities were successfully matched. For cities with more than one observation station, the average of records from multiple stations was employed. To match the temporal scale of meteorological data, the per-3h and per-24h pollutant data were produced by conducting average operation on the hourly concentration data.

2.2 Advanced Causation Model

Since 2013, when PM_{2.5} pollution was observed across China, research on airborne pollution has been massively conducted. Amongst a diversity of topics, research on the meteorological influences on major airborne pollutants (e.g., PM_{2.5} and O₃) has received growing emphasis. However, the major challenge for extracting and comparing the influence of individual meteorological factors lies in the complicated inner-interactions between multiple meteorological factors, which cause large uncertainties when applying traditional correlation analysis (Chen et al., 2020a). To address this issue, we employed an advanced causation model, Convergent Cross Mapping (CCM), to quantify the influence of each meteorological factor on PM_{2.5}, PM₁₀ and O₃. By removing the influence of disturbing factors, CCM (Sugihara et al., 2012) is capable of extracting reliable coupling between two variables in complicated ecosystems. CCM calculates the causal influence of the variable A on the target variable B as the ρ value, ranging from 0 to 1. Like the correlation coefficient, the ρ value can be used for comparing the influencing between multiple variables on the target variable.

Thanks to its advantage in effectively extracting the asymmetric, bidirectional association between two variables and identifying mirage correlation in complicated ecosystems with a diversity of variables, we have massively employed CCM to evaluate the influence of multiple meteorological factors on PM_{2.5} (Chen et al., 2017, 2018), O₃ (Cheng et al., 2019; Chen et al., 2020b) and NPP (Gao et al., 2022) and achieved reliable outputs. Based on a multi-model comparison experiment, our recent research (Chen et al., 2022) proved that CCM was the most suitable model for causation inference in complicated atmospheric environment. CCM is specifically designed to deal with the nonlinear relationship between two variables and is fully suitable for the nonlinear relationship between atmospheric factors. Compared with other mainstream statistical models, CCM was advantageous of identifying unique pollutant-meteorology association in local areas while maintaining general characteristics of pollutant-meteorology association across China. Furthermore, CCM generated meteorology-pollutant association were highly consistent with prior-knowledge. In this regard, for this research, we also employed CCM to quantify and compare the influence of temperature, precipitation, wind speed, wind direction and atmospheric pressure on PM_{2.5}, PM₁₀ and O₃ concentrations. CCM automatically considers all possible interaction forms and lag effects between the time series of two variables, which effectively reduces the influence of interference and avoids the influence of other factors. CCM is highly automatic to remove the uncertainty of manual setting and only the setting of several parameters is required: E (number of dimensions), τ (time lag) and b (number of nearest neighbors). For this research, τ , E, and b were set as 2 days, 3 and 4 according to previous studies (Chen et al., 2018; 2020b).

Based on the rarely employed 3h meteorological data sources, we compared the effects of temporal scales on the extracted pollutant-meteorology causation. Acknowledged, due to the data limitation at the 3h scale, which did not include humidity and sunshine duration, we could simply consider a limited number of meteorological factors (Temperature, Precipitation, Wind Speed, Wind Direction and Atmospheric Pressure), which was less than our previous studies based on meteorological data at the 24h scale, while some meteorological factors (e.g., humidity and sunshine duration) were missed in this research. However, since we compared the same set of these major meteorological factors at both 3h and 24h scale, the calculated consistence and difference could effectively reveal the potential effects of different temporal scales on the quantitative (the detailed ρ value) and qualitative (the dominant meteorological factor) findings of pollutant-meteorology association. Despite the limitation of number of meteorological factors, it caused limited influence on the temporal effects on pollutant-meteorology association. This is because CCM simply considers the causality between the target variable and one influencing variable, and removes the influence from other variables (Sugihara et al., 2012; Chen et al., 2020). Another limitation of this data was that this data set simply included one year's data and thus the inter-annual variation of temporal effects on pollutant-meteorology association could not be revealed. For this research, we majorly revealed the existence of strong temporal effects on pollutant-meteorology association, which can be fully supported by the one-year data with four seasons (four complete time series with more than 90 records (24h scale) and 720 records (3h scale) Meanwhile, the temporal variation of temporal effects on pollutant-meteorology association and its influencing factors should be further investigated in future studies, when the long time series data sets of 3h meteorological data become available.

3 Results

3.1 The comparison of dominant meteorological factors for PM_{2.5}, PM₁₀ and O₃ across China at 3h and 24h scale

Based on CCM, we calculated the dominant meteorological factors for seasonal O₃ (Table 1), PM_{2.5} (Table 2) and PM₁₀ (Table 3) concentrations at the 3h and 24h respectively. By comparing the extracted pollutant-meteorology association, we calculated the number of cities with the same meteorological factor at different temporal scales (Table 4). For all three airborne pollutants, the dominant meteorological factor at the 3h and 24h scale was the same in only around a third of cities, indicating the temporal scale played a large role in the analysis of pollutant-meteorology association. From the seasonal scale, the consistence between dominant meteorological factors extracted at 3h and 24h in autumn and winter was higher than that in spring and summer. This phenomenon indirectly suggested that meteorological influences on airborne pollutants were stronger in autumn and winter, and thus the role of dominant meteorological factor was highlighted.

Table 1 inserted here.

Table 2 inserted here.

Table 3 inserted here.

At the 3h scale, for O₃, the number of cities with precipitation as the dominant influencing factor was largest in winter, while the number of cities with temperature was largest in spring, summer and autumn; For PM_{2.5} and PM₁₀, the number of cities with temperature was largest in all seasons. As a comparison, at the 24h scale, for O₃, PM_{2.5} and PM₁₀, the number of cities with temperature as the dominant influencing factor was largest in spring, which was consistent with previous studies (Wang et al., 2018; Yang et al., 2021), summer and autumn while the number of cities with precipitation was largest in winter.

Table 4 inserted here.

For both the 3h and 24h scale, we could see temperature and precipitation exerted strong influences on O₃, PM_{2.5} and PM₁₀ in the majority of cities. However, the consistence of dominant factors between two temporal scales remained less than 50%. This may be attributed to the fact that the extraction of dominant meteorological factor amongst several factors was relatively qualitative and thus some subtle differences between different meteorological factors could not be revealed. Therefore, we

further presented the detailed comparison of the influence of individual meteorological factors on O₃, PM_{2.5} and PM₁₀ at 3h and 24h respectively.

3.2 The comparison of quantified influence of different meteorological factors on PM_{2.5}, PM₁₀ and O₃ across China at 3h and 24h scale

160 The detailed distribution of influence of individual meteorological factors on O₃, PM_{2.5} and PM₁₀ concentrations are presented
in Figure 1. Generally, meteorological influences on airborne pollutants presented a consistent trend between the 3h and 24h
scale, characterized with a generally similar violin shape. According to Figure 1, the violin shape and range of 3h pollutant-
meteorology was much sharper than the 24h pollutant-meteorology, indicating the 3h temporal scale was more sensitive to
165 meteorological factors, the mean of calculated ρ value across China also proved that temperature exerted a much stronger
influence on PM_{2.5}, PM₁₀ and O₃ than other factors. Furthermore, according to the violin shape of different pollutants, we found
that the pattern of PM_{2.5}-Meteorology and PM₁₀-Meteorology was generally consistent and largely different from the pattern
of O₃-Meteorology, indicating that meteorological influences on particulate matters and gaseous pollutants were different. The
major difference of pollutant-meteorology interactions at 3h and 24h was explained as follows:

170

Table 5 inserted here.

Figure 1 inserted here.

175 For all three airborne pollutants, temperature exerted a strongest influence across China in all seasons in terms of the largest
mean ρ . For PM_{2.5} and PM₁₀, the calculated influence of temperature at 24h scale was consistently larger than that at the 3h
scale. This may be attributed to the fact that the secondary reactions of the precursors of PM were relatively mild (Chen et al,
2016, 2020) and thus the temperature variation within 24h exerted a stronger influence than 3h temperature variation.
Meanwhile, the influence of temperature on O₃ presented a notable seasonal pattern. For the relatively cold season winter and
180 spring, when O₃ concentrations were relatively low, the influence of temperature at 24h scale was larger than that at the 3h
scale. For summer, when O₃ concentrations were the highest, the influence of temperature at 3h scale was much larger than
that at the 24h scale. This is mainly attributed to the fact that the high temperature in summer was the major trigger for quick
reactions between precursors and high O₃ concentrations. Therefore, short-term variations of temperature could strongly
influence O₃ concentrations in summer (Cheng et al., 2018, 2019).

185

For precipitation, since the distribution of precipitation in a day's time is not unified, and there may be no precipitation in
many 3h slots, the mean ρ of precipitation across China at the 3h scale was weaker than that at the 24h scale. As a comparison,
at the 24 scale, the occurrence of precipitation was significantly enhanced and thus the influence of precipitation on airborne

190 pollutants was much stronger. Across China, the precipitation intensity showed obvious seasonal variations, and most regions may have the maximum value in summer and minimum value in winter. Accordingly, the calculated ρ of precipitation on $PM_{2.5}$, PM_{10} and O_3 at 24h scale was remarkably larger than that at 3h scale.

195 Previous studies (Chen et al., 2017, 2018, 2020) proved that wind played a notable influence on PM. Similar to precipitation, the daily distribution of wind is not unified, and there may be calm wind conditions in many 3h slots. Therefore, the mean ρ of wind direction and wind speed on $PM_{2.5}$ and PM_{10} at 24h scale was notably larger than that at the 3h scale. Wind- O_3 interactions presented notable seasonal patterns. In the less-polluted Spring and Winter, the mean ρ of wind direction and wind speed at the 24h scale was larger than that at the 3h scale. In summer, when O_3 concentrations were relatively high, the mean ρ of wind direction and wind speed at the 3h scale was larger.

200 Atmospheric pressure mainly affects the transport and accumulation of pollutants by indirectly influencing other meteorological factors (e.g. wind and precipitation). Therefore, large uncertainties existed in the extracted pressure-pollutant causation. Generally, for $PM_{2.5}$, PM_{10} and O_3 , the mean ρ of atmospheric pressure across China at the 3h scale was weaker than that at the 24h scale, except for summer, when the interactions between atmospheric pressure and other meteorological factors were strong.

205 **3.3 The spatial patterns of dominant meteorological factors for $PM_{2.5}$, PM_{10} and O_3 across China at 3h and 24h scale**

As shown in Figure 2, 3, 4, the influence of meteorological factors on airborne pollutants has obvious seasonal variations and presented some regional similarity. For $PM_{2.5}$ (Figure 3) and PM_{10} (Figure 4), the dominant meteorological factor for Northern China was mainly wind, especially the heavily polluted winter, while the dominant meteorological factor for Yangtze River Delta and was mainly precipitation at both 3h and 24h scale. The dominant meteorological factor in Shandong Peninsula in spring and autumn, southern China in summer, northern and coastal areas in autumn, and northeast China in winter are also consistent at different temporal scales. For O_3 (Figure 2), especially the heavily polluted summer, temperature presented a prevailing role across the nation and was the dominant role for most cities. This output was consistent with our previous studies (Chen et al., 2018, 2019a), suggesting the general national trend of Pollutant-Meteorology association varied limitedly across temporal scales of research data, especially in those heavily polluted regions. Meanwhile, for those regions, where the airborne pollution was not severe and homogeneous, the temporal issues of meteorological influences on PM was notable and thus the dominant meteorological factor in these regions presented notable differences at 3h and 24h scale.

220 Generally, the distribution of the dominant elements at the 24h scale was more heterogeneous than that at the 3h scale. Regions with higher PM ($PM_{2.5}$ and PM_{10}) concentrations received more influences from temperature at the 3h scale than that at the 24h scale.

Figure 2 inserted here.

Figure 3 inserted here.

Figure 4 inserted here.

225

4 Discussion

Although previous studies (Tai et al., 2010; Hu et al., 2021b; Yousefian et al., 2021; Zhong et al., 2021) pointed out the difference of pollutant-meteorology association at different temporal scales, few studies have actually conducted the quantitative analysis due to the lack of data. This research suggested that the temporal effects on pollutant-meteorology association was significantly strong. While the obvious quantitative difference of the influence of individual factors on PM_{2.5}, PM₁₀ and O₃ (as shown in Figure 1), we found a very low consistence between extracted dominant meteorological factors (the consistence was less than 50% for all pollutants), indicating strong temporal effects even from a qualitative perspective. Based on the comparison of extracted pollutant-meteorology association at the 3h and 24h, there were no fixed spatiotemporal patterns of pollutant-meteorology association across temporal scales. However, we got several interesting and useful findings as follows. Firstly, we found the temporal effects of meteorological influences on different PM (e.g., PM_{2.5} and PM) were similar, yet notably different from that on gaseous pollutants (e.g., O₃). Secondly, there were notable difference of the temporal effects between different meteorological factors. The variation of pollutant-meteorology association for those factors with continuous observation record (e.g., Temperature) was notably different from those factors with discrete observation record (e.g. Precipitation) at 3h and 24h scale. Thirdly, the effects of temporal scales on pollutant-meteorology association varied significantly across seasons, characterized with notable difference between heavily polluted and less polluted seasons (e.g. The heavily polluted season for O₃ and PM was winter and summer respectively) . Despite a complicate pattern, we found that the heavier the pollution, the stronger pollutant-meteorology association was. Consequently, in the heavily polluted season, the short-term (e.g., 3h) variation of specific meteorological factors (e.g. Temperature) exerted a stronger influence on PM and O₃ than the daily variation. As a comparison, in the less polluted season, the daily accumulation of specific meteorological factors exerted a stronger influence on airborne pollutants than short-term (e.g., 3h) accumulation. While the general trend of pollutant-meteorology association was consistent with previous studies, the general ρ value was slightly smaller for this research. The underlying reason may be the reduced PM_{2.5} concentration in 2020 caused by the emission-cut during COVID-19. As explained in our previous studies (Chen et al., 2018), the higher PM_{2.5} concentration, the stronger meteorological influence on PM_{2.5} concentrations. Similar to our previous studies (Chen et al., 2017, 2018, 2022), we conducted the CCM analysis at the seasonal scale. This is because the large seasonal variation of pollutant-meteorology association may cause an insignificant output of CCM for an entire-year analysis, and cause large uncertainties.

235

240

245

250

This research suggested that the temporal scale played a complicated role and higher temporal-resolution did not guarantee a stronger pollutant-meteorology association. For instance, for hot seasons (e.g., summertime O₃), the reaction between O₃ precursors was strong and quick, and thus the 3h resolution could better feature the influence of temperature on O₃ concentrations. Meanwhile, the secondary reaction for PM_{2.5} was relatively slow (Chen et al. 2016), and the daily variation of temperature and PM_{2.5} concentrations presented a stronger association than the hourly variation of temperature and PM_{2.5} concentrations. Similarly, due to the discrete distribution, the daily influence of daily total precipitation on daily PM_{2.5} concentrations was also notably stronger than the influence of 3h precipitation on 3h PM_{2.5} concentrations. Furthermore, this type of uncertainty was not predicable across regions. Given the complicated effects of temporal scales on pollutant-meteorology association, scholars should properly choose the temporal-resolution of research data according to the aims, study sites, pollutant types and seasons. With the growing availability of long-term meteorological and pollutant data, multi-scale, instead of high-temporal-resolution, research is recommended to comprehensively understand the short- and long-term meteorological influences on different airborne pollutants.

For future research, the temporal effects of influence of meteorological factors (e.g., Humidity, Boundary layer height) on airborne pollutants should also be explored with the availability of new data sources. On the other, this research proved the important role of temporal scales in quantifying the influence of meteorological factors on airborne pollutants. Similarly, when inferring the association between precursors (NO₂, VOCs) and airborne pollutants, the temporal scales, which was rarely considered in previous studies, should also be comprehensively taken into account. The reaction rate between different precursors and the target pollutants in different regions and seasons could be better understood through multi-scale causation analysis. CCM is an ideal tool for quantifying the influence of individual meteorological factors on PM_{2.5} concentrations, as it can effectively remove the influence of other meteorological factors. Therefore, this research revealed a strong temporal effect on pollutant-meteorology association, from the perspective of the association of individual meteorological factors. However, admittedly, CCM is limited in establishing the overall effects of multiple meteorological factors on PM_{2.5} concentrations. Instead, other models, such as GAM (Generalized Additive Model), which work limited in extracting the association between PM_{2.5} and individual meteorological factors, are advantageous in extracting the overall influence of multiple meteorological factors on airborne pollutants (Gong et al., 2017; Zheng et al., 2018; Hu et al., 2021a). When such 3h meteorological data set become more easily available and includes a complete set of meteorological factors, we could also employ GAMs or CTMs to investigate the temporal effects on the combined effects of meteorological factors on airborne pollutants.

5 Conclusion

We employed CCM to compare the influence of major meteorological factors (Temperature, Precipitation, Wind Speed, Wind Direction and Atmospheric Pressure) on PM_{2.5}, PM₁₀ and O₃ concentrations in 101 cities across China at the 3h and 24h scale.

285 The result revealed a strong effect of temporal scale on the pollutant-meteorology association from different perspective. In terms of the extracted dominant meteorological factor, the consistence between the analysis at 3h and 24h scale was relatively low (the consistence for all pollutants was less than 50%), suggesting a large difference even from a qualitative perspective. In terms of the mean ρ value, the effect of temporal scale on the influence of individual meteorological factors on Particulate Matter (PM_{2.5} and PM₁₀) was consistent, which was largely different from the temporal-scale effect on gaseous pollutants.

290 Temperature was the most important meteorological factor for PM_{2.5}, PM₁₀ and O₃ across China at both 3h and 24 scale. For PM_{2.5} and PM₁₀, the secondary reaction of which was relatively slow, the extracted PM-temperature association at the 24h scale was stronger than that at the 3h scale. Meanwhile, for summer O₃, due to the quick and strong reactions between precursors, the extracted O₃-temperature association at the 3h scale was much stronger than that at the 24h scale. Due to the discrete distribution, the extracted association between all pollutants and precipitation was much weaker at the 3h scale.

295 Similarly, the extracted PM-wind association was notably weaker at the 3h scale. Due to the transport of precursors, summertime O₃-wind association was stronger at the 3h scale. For atmospheric pressure, the pollutant-pressure association was weaker at the 3h scale except for summer, when the interactions between atmospheric pressure and other meteorological factors were strong. From the spatial perspective, the influence of meteorological factors on airborne pollutants has obvious seasonal variations and presented some regional similarity, pollutant-meteorology association at 3h and 24h was more


300 consistent at those heavily polluted regions. This research provides a comprehensive understanding for the effect of temporal scale on pollutant-meteorology association and sheds useful light on better extracting the natural and anthropogenic drivers for airborne pollution.

References

- Chen, Z., et al. 2022. Causation inference in complicated atmospheric environment. *Environmental Pollution*, 303, 119057.
- 305 Chen, Z., et al. 2020a. Influence of meteorological conditions on PM_{2.5} concentrations across China: A review of methodology and mechanism. *Environment international*, 139, 105558.
- Chen, Z., et al. 2020b. Understanding the causal influence of major meteorological factors on ground ozone concentrations across china. *Journal of Cleaner Production*, 242, 118498.
- Chen, Z., et al. 2019a. Spatial self-aggregation effects and national division of city-level pm_{2.5} concentrations in china based
- 310 on spatio-temporal clustering. *Journal of Cleaner Production*, 207(10), 875-881.
- Chen, Z., et al. 2019b. Understanding long-term variations of meteorological influences on ground ozone concentrations in beijing during 2006–2016. *Environmental Pollution*, 245(FEB.), 29-37.
- Chen, Z., et al. 2019c. The control of anthropogenic emissions contributed to 80% of the decrease in pm 2.5 concentrations in beijing from 2013 to 2017. *Atmospheric Chemistry and Physics*, 19(21), 13519-13533.
- 315 Chen, Z., et al. 2018. Understanding meteorological influences on pm_{2.5} concentrations across china: a temporal and spatial perspective. *Atmospheric Chemistry & Physics Discussions*, 1-30.

- Chen, Z., et al. 2017. Detecting the causality influence of individual meteorological factors on local pm2.5 concentration in the jing-jin-ji region. *Rep*, 7, 40735.
- Chen, Z., et al. 2016. Understanding temporal patterns and characteristics of air quality in Beijing: A local and regional perspective. *Atmospheric environment*, 127, 303-315.
- 320 Cheng, N., et al. 2017. Spatio-temporal variations of pm2.5 concentrations and the evaluation of emission reduction measures during two red air pollution alerts in beijing. *Scientific Reports*, 7(1), 8220.
- Cheng, N., et al. 2018. Ground ozone concentrations over Beijing from 2004 to 2015: Variation patterns, indicative precursors and effects of emission-reduction. *Environmental Pollution*, 237, 262-274.
- 325 Cheng, N., et al. 2019. Ground ozone variations at an urban and a rural station in Beijing from 2006 to 2017: Trend, meteorological influences and formation regimes. *Journal of Cleaner Production*, 235, 11-20.
- Gao, B., et al. 2022. Temporally or spatially? Causation inference in Earth System Sciences. *Science Bulletin*, 67(3), 232-235.
- Gao, J., et al. 2017. Haze, public health and mitigation measures in China: A review of the current evidence for further policy response. *Science of the Total Environment*, 578, 148-157.
- 330 Guo, J., et al, 2020. The climatology of lower tropospheric temperature inversions in China from radiosonde measurements: roles of black carbon, local meteorology, and large-scale subsidence. *Journal of Climate*, 33 (21): 9327–9350.
- Gong, X., et al. 2017. Quantifying o3 impacts in urban areas due to wildfires using a generalized additive model. *Environmental Science and Technology*, 51(22), 13216.
- Hu, C., et al. 2021a. Understanding the impact of meteorology on ozone in 334 cities of China. *Atmospheric Environment*, 335 248, 118221.
- Hu, M., et al. 2021b. Spatial-temporal heterogeneity of air pollution and its relationship with meteorological factors in the Pearl River Delta, China. *Atmospheric Environment*, 254, 118415.
- Kelly, F. J., Fussell, J. C. 2015. Air pollution and public health: emerging hazards and improved understanding of risk. *Environmental geochemistry and health*, 37(4), 631-649.
- 340 Sugihara, G., et al. 2012. Detecting causality in complex ecosystems. *Science*, 338(6106), 496-500.
- Tai, A. P., Mickley, L. J., Jacob, D. J. 2010. Correlations between fine particulate matter (PM2. 5) and meteorological variables in the United States: Implications for the sensitivity of PM2. 5 to climate change. *Atmospheric environment*, 44(32), 3976-3984.
- Wang, B., et al. 2018. Long term causality analyses of industrial pollutants and meteorological factors on PM2. 5 concentrations in Zhejiang Province. In 2018 5th International Conference on Information Science and Control Engineering (ICISCE) ,301-305. IEEE.
- 345 Wang, X., et al. 2021a. Dominant synoptic patterns associated with the decay process of PM 2.5 pollution episodes around Beijing. *Atmospheric Chemistry and Physics*, 21(4), 2491-2508.
- Wang, N., et al. 2021b. Air quality during COVID-19 lockdown in the Yangtze River Delta and the Pearl River Delta: Two different responsive mechanisms to emission reductions in China. *Environmental science & technology*, 55(9), 5721-5730.
- 350

- Wang, Z., et al. 2021. The estimation of hourly PM_{2.5} concentrations across China based on a Spatial and Temporal Weighted Continuous Deep Neural Network (STWC-DNN). *ISPRS Journal of Photogrammetry and Remote Sensing*, 190, 38-55.
- Xiao, Q., et al. 2020. Changes in spatial patterns of PM_{2.5} pollution in China 2000–2018: Impact of clean air policies. *Environment international*, 141, 105776.
- 355 Xu, M., et al. 2021. Estimating the impact of ground ozone concentrations on crop yields across China from 2014 to 2018: A multi-model comparison. *Environmental Pollution*, 283, 117099.
- Yang, Z., et al. 2017. Nonlinear and lagged meteorological effects on daily levels of ambient PM_{2.5} and O₃: Evidence from 284 Chinese cities. *Journal of Cleaner Production*, 278, 123931.
- Yin, P., J. et al. 2020. Higher risk of cardiovascular disease associated with smaller size-fractioned particulate matter, *Environmental Science & Technology Letters*, 7(2), 95-101.
- 360 Yousefian, F., et al. 2021. Temporal variations of ambient air pollutants and meteorological influences on their concentrations in tehran during 2012–2017. *ISEE Conference Abstracts*, 2021(1).
- Zhai, S., et al. 2019. Fine particulate matter (PM_{2.5}) trends in China, 2013–2018: Separating contributions from emission influencing emissions and meteorology. *Atmospheric Chemistry and Physics*, 19(16), 11031-11041.
- 365 Zhan, D., et al. 2017. Spatiotemporal variations and driving factors of air pollution in China. *International Journal of Environmental Research and Public Health*, 14(12), 1538.
- Zhang, Y. et al., 2020. Vertical wind shear modulates particulate matter pollutions: A perspective from Radar wind profiler observations in Beijing, China. *Remote Sensing*, 12(3), 546.
- Zheng, B., et al. 2018. Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions. *Atmospheric Chemistry and Physics*, 18(19), 14095-14111.
- 370 Zhong, Q., et al. 2021. PM_{2.5} reductions in Chinese cities from 2013 to 2019 remain significant despite the inflating effects of meteorological conditions. *One Earth*, 4(3), 448-458.
- Zhou, L., Chen, X., Tian, X. 2018. The impact of fine particulate matter (PM_{2.5}) on China's agricultural production from 2001 to 2010. *Journal of Cleaner Production*, 178, 133-141.



375

Figure 1: The violin chart of the ρ value of individual meteorological factors on PM_{2.5}, PM₁₀ and O₃ across China at 3h and 24h scale.




Figure 2: The dominant meteorological factor for O₃ concentrations across China at 3h and 24h scale.





Figure 3: The dominant meteorological factor for PM_{2.5} concentrations across China at 3h and 24h scale.



385 **Figure 4: The dominant meteorological factor for PM₁₀ concentrations across China at 3h and 24h scale.**

386 **Table 1: The number of cities with this meteorological factor as the dominant meteorological factor for O₃.**

O ₃	3h				24h			
	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter
Temperature	64	78	75	42	59	38	58	33
Precipitation	15	9	8	43	21	18	15	47
Atmospheric pressure	7	5	4	3	8	8	4	3
Wind Direction	6	4	5	1	8	23	14	7
Wind Speed	9	5	9	12	5	14	10	11

387

Table 2: The number of cities with this meteorological factor as the dominant meteorological factor for PM_{2.5}.

PM _{2.5}	3h				24h			
	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter
Temperature	62	60	79	59	44	43	61	30
Precipitation	7	9	8	19	22	19	14	35
Atmospheric pressure	12	8	4	2	3	8	6	5
Wind Direction	12	12	8	6	22	16	13	13
Wind Speed	8	12	2	15	10	15	7	18

390 **Table 3: The number of cities with this meteorological factor as the dominant meteorological factor for PM₁₀.**

PM ₁₀	3h				24h			
	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter
Temperature	65	53	73	56	45	34	55	31
Precipitation	19	11	13	24	20	34	20	36
Atmospheric pressure	9	10	4	2	10	8	7	4
Wind Direction	4	13	7	4	13	14	10	14
Wind Speed	4	14	4	15	13	11	9	16

Table 4: The number of cities with the same dominant factor at both 3h and 24h scale.

	spring	summer	autumn	winter
O ₃ -meteorological elements	32	42	58	53
PM _{2.5} -meteorological elements	36	42	62	42
PM ₁₀ -meteorological elements	42	29	56	43

Table 5: The mean ρ of individual meteorological factors on PM_{2.5}, PM₁₀ and O₃ across China.

		Temperature		Precipitation		Atmospheric pressure		Wind Direction		Wind Speed	
		3h	24h	3h	24h	3h	24h	3h	24h	3h	24h
O ₃	Spring	0.213	0.283	0.050	0.140	0.058	0.070	0.028	0.048	0.030	0.042
	Summer	0.238	0.114	0.013	0.042	0.055	0.017	0.049	0.049	0.065	0.037
	Autumn	0.218	0.210	0.013	0.032	0.032	0.032	0.038	0.034	0.039	0.034
	Winter	0.133	0.198	0.100	0.191	0.058	0.035	0.045	0.058	0.052	0.062
PM _{2.5}	Spring	0.095	0.128	0.027	0.059	0.030	0.034	0.018	0.048	0.015	0.030
	Summer	0.079	0.108	0.012	0.040	0.016	0.013	0.018	0.036	0.018	0.032
	Autumn	0.143	0.182	0.016	0.029	0.018	0.025	0.019	0.045	0.017	0.028
	Winter	0.120	0.140	0.045	0.090	0.020	0.035	0.027	0.044	0.045	0.064
PM ₁₀	Spring	0.106	0.129	0.031	0.068	0.030	0.034	0.014	0.039	0.015	0.036
	Summer	0.081	0.125	0.012	0.049	0.022	0.016	0.019	0.030	0.016	0.028
	Autumn	0.158	0.220	0.016	0.041	0.022	0.045	0.021	0.040	0.021	0.041
	Winter	0.109	0.127	0.046	0.082	0.020	0.029	0.028	0.050	0.043	0.063