

The Impact of Aerosols on the Stratiform Clouds over Southern West Africa: A Large-Eddy Simulation Study

Lambert Delbeke¹, Chien Wang¹, Pierre Tulet¹, Cyrielle Denjean², Maurin Zouzoua³, Nicolas Maury⁴
and Adrien Deroubaix⁵

¹Laboratoire d'Aérodynamique, [Université de Toulouse, Université Paul Sabatier](#), CNRS, [UT3, IRD](#), Toulouse, France

²CNRM, [Université de Toulouse, Météo-France](#), CNRS, Toulouse, France

³Laboratoire Atmosphères, Milieux, Observations Spatiales, IPSL, CNRS, Guyancourt, France

⁴CNRM, Université de Toulouse, [Météo-France](#), CNRS, Toulouse; now at LMD/IPSL, Paris, France

⁵[IUP, Institute of Environmental Physics, University of Bremen, Bremen](#), and Max Plank Institute [for Meteorology, Hamburg](#), Germany

Correspondence to: Chien Wang (chien.wang@aero.obs-mip.fr), Cyrielle Denjean (cyrielle.denjean@meteo.fr)

Abstract. Low level stratiform clouds (LLSCs), covering a large area, appear frequently during the wet monsoon season in southern West Africa. This region is also the place where different types of aerosols coexist, including biomass burning aerosols coming from Central and South Africa and anthropogenic aerosols emitted by local activities. We investigate the semi-direct and indirect effects of these aerosols on the diurnal cycle of LLSCs by constructing a case study based on ~~aircraft-based~~ [airborne](#) and ground-based observations from the Dynamic-Aerosol-Chemistry-Cloud-Interaction in West Africa (DACCIWA) field campaign. This case is modelled using a Large Eddy Simulation (LES) model with fine scale resolution and in-situ aerosol measurements including size distribution and chemical composition. The model has successfully reproduced the observed life cycle of the LLSC, from stratus formation to stabilization during the night, to upward development after sunrise until break-up of cloud deck in afternoon. Various sensitivity simulations using different measured aerosol profiles ~~based on measurements~~ also suggest that aerosols can affect the cloud life cycle through both the indirect and semi-direct effect. Despite precipitation produced by the modeled cloud is nearly negligible, clouds lifetime is still sensitive to the aerosol concentration. As expected, modeled cloud microphysical features including cloud droplet number concentration, mean radius, and thus cloud reflectivity are all controlled by aerosol concentration. However, it is found that the difference in cloud reflectivity is not always the only factor in determining the variation of the incoming solar radiation at ground and cloud life cycle specifically beyond sunrise. Instead, the difference in cloud-void space brought by dry air entrainment from above and thus the speed of consequent evaporation - also influenced by aerosol concentration, is ~~an another~~ [another](#) important factor to consider. Results have shown that clouds in the case with lower aerosol concentration and larger droplet size appear to be less affected by entrainment and convection. In addition, we found that an excessive atmospheric heating up to 12 K day^{-1} produced by absorbing black carbon aerosols (BC) in our modeled cases can also affect the life cycle of modeled clouds. Such a heating is found to lower the height of cloud top ~~and stabilize the cloud layer~~, resulting a less extent in vertical development and accelerating clouds break-up. The semi-direct effect impacts on indirect effect by reducing cloud reflectivity particularly in case of polluted environment. Finally, semi-direct effect is

found to contribute positively to the indirect radiative forcing due to a decreased cloud-void space, and negatively by causing thinner clouds that would break-up faster in late afternoon, all depending on the phase in stratiform cloud diurnal cycle.

1. Introduction

Low-level stratiform clouds (LLSCs) are important to Earth's radiative budget through the reflection of solar radiation due to their high albedo (Hartmann et al., 1992; Chen et al., 2000) and large cloud deck covering Earth's surface more than any other cloud type (Eastman and Warren, 2014). LLSCs often occupy the upper few hundred meters in the planetary boundary layer (PBL), and their persistent appearance relies on a stable PBL that is normally associated with a large-scale subsidence above PBL under a ~~high-pressure~~high-pressure system. LLSCs are often formed over cooler subtropical and mid-latitude oceans, constantly covering more than 50% of these areas (Wood, 2012). During the West African monsoon season, LLSCs frequently form over continental southern West Africa (SWA) in the night and would likely break up in the early afternoon of the following day there (Schrage and Fink, 2012; Schuster et al., 2013). LLSCs are, under polluted conditions characterized by numerous and small cloud droplets, increasing the cloud albedo, suppressing drizzle, and extending lifetime (Twomey, 1957; Haywood and Boucher, 2000; Liu et al., 2014; Carslaw et al., 2017). The presence of LLSCs impacts the radiative budget of atmospheric boundary layer and surface and also affect the diurnal cycle of the convective boundary layer and thus the regional climate (Knippertz et al., 2011; Hannak et al., 2017). However, the diurnal cycle of LLSCs is still poorly represented in weather and climate models, especially over SWA, because the processes behind the variability of LLSCs cover remain elusive (Knippertz et al., 2011; Hannak et al., 2017; Hill et al., 2018).

Stratiform clouds are sensitive to aerosol properties (concentration, chemistry) and vertical distribution. This is because that aerosol can directly scatter or absorb solar radiation (direct effect or aerosol-radiation effect), or by serving as cloud nuclei, influence cloud microphysical structure and thus reflectance or lifetime (indirect aerosol effects or radiative effect of aerosol-cloud interaction plus cloud adjustment) (Boucher et al., 2013). The heating associated with aerosol absorption would be able to perturb atmospheric thermodynamic stability and thus dynamical processes as well (semi-direct effect) (Hansen et al., 1998). All these effects can modify the energy budget and thus the status of the planetary boundary layer where the stratiform clouds form. Aerosols inside stratiform clouds are also modified by physico-chemical processes which can influence the aerosol concentration (Wood, 2012). Interactions between aerosols and clouds, and their effects on radiation, precipitation, and regional circulations, remain one of the largest uncertainties in understanding and projecting climate change. Indeed, the indirect effect of aerosol forcing is still difficult to estimate (IPCC 2021) and climate models struggle to minimize such uncertainties (Li et al., 2022). Some aerosol ~~particules~~constituents such as black carbon absorb a substantial amount of shortwave radiation, which results in rapid atmospheric thermodynamic adjustments. This semi-direct aerosol radiative effect can be positive or negative depending on the relative distribution of the aerosol with respect to clouds. Several previous studies were conducted to investigate aerosol-clouds interactions of LLSCs using high-resolution LES models but mainly over ocean (Ackerman et al., 2004; Sandu et al., 2008; Twohy et al., 2013; Flossmann and Wobrock, 2019) where surface fluxes are having little diurnal variation, moisture is mainly provided by evaporation from

80 sea surface to maintain the stratiform cloud layer. By contrast over land, moisture supply is dependent on the characteristics of the surface (Wood, 2012).

During the West Africa Monsoon (WAM), aerosols can come from both local and remote sources to SWA. Large amount of Biomass Burning Aerosols (BBA) are transported from southern and Central African towards SWA during the summer monsoon (Haslett *et al.*, 2019). These air masses are further
85 loaded with additional aerosols from anthropogenic emissions upon reaching the highly urbanized regions near the coast. (Chatfield *et al.*, 1998; Sauvage *et al.*, 2005; Mari *et al.*, 2008; Murphy *et al.*, 2010; Reeves *et al.*, 2010; Menut *et al.*, 2018; Haslett *et al.*, 2019). A significant quantity of wind-blown mineral dust aerosols emitted from the Sahara and Sahel throughout the year with a peak in springtime (Marticorena and Bergametti, 1996) can also reach SWA far south often in June (Knippertz *et al.*, 2017).
90 Local sources of aerosols in SWA are related to anthropogenic activities near the coast from where polluted plumes transport inland (Deroubaix *et al.*, 2019). These emissions are supposed to increase with the expected growth~~ing~~ of the population (Lioussse *et al.*, 2014). These different sources of aerosols give a complex mix of species with a high loading, having a serious impact on human health (Bauer *et al.*, 2019) and also on the diurnal cycle of ~~winds,~~ LLSCs as well as precipitation over SWA (Taylor *et al.*,
95 2019).

The DACCIWA project was designed to better characterize cloud-aerosol-precipitation interactions in SWA (Knippertz *et al.*, 2015). The measurement campaign provides a comprehensive set of ground-based and airborne~~-based~~ measurements of clouds and aerosols in June-July 2016. (Knippertz *et al.*, 2017; Kalthoff *et al.*, 2018; Flamant *et al.*, 2018) and model analyses. Measurements were conducted at three
100 supersites, Savè (Benin), Kumasi (Ghana) and Ile-Ife (Nigeria) (Fig.1),~~and~~ and coordinated with three research aircrafts: the French ATR-42 operated by SAFIRE (Service des Avions Français Instrumentés pour la Recherche en Environnement), the British Twin Otter operated by British Antarctic Survey, and the German Falcon aircraft operated by DLR (Deutsches Zentrum für Luft und Raumfahrt). Additional radiosoundings were launched from Savè with high temporal frequency, which specifically benefits the
105 monitoring of the LLSCs diurnal evolution.




Figure 1. Map of southern West Africa with Savè, Kumasi and Ile-Ife locations and the flight track (blue line) of the ATR-42 the 3 July 2016 with NASA Suomi NPP/VIIRS true color corrected reflectance (<https://worldview.earthdata.nasa.gov/>).

110 Based on observations and parcel modeling, Taylor *et al.* (2019) and Denjean *et al.* (2020a) showed
 that the majority of cloud condensation nuclei and absorbing aerosols were from ubiquitous long-range
 transported BBA, causing a polluted background which limits the effect of local pollution on cloud
 properties and aerosol radiative effects. Modeling studies have suggested that light-absorbing aerosols
 from combustion sources like BBA and anthropogenic sources can impact on the formation, evolution,
 115 and precipitation of LLSCs, especially over south-eastern Atlantic. Using COSMO-ART model in a
 simulation for 2-3 July 2016 case, Deetz *et al.* (2018) found that under the influence of the Maritime
 Inflow (MI, cold air) from Guinean Gulf, stratus-stratocumulus transition areis susceptible to the aerosol
 direct effect, resulting in a spatial shift in the MI front and a temporal shift of the cloud transition. Over
 SWA and influenced by anthropogenic emission sources, the break-up time of LLSCs can be delayed by
 120 one hour and daily precipitation rate can decrease by 7.5% according to Deroubaix *et al.* (2022). Moreover,
 semi-direct and indirect effects were studied together by varying respective magnitude or foreing
 emissions of anthropogenic aerosols, but they were not examined separately. Haslett *et al.* (2019) denote
 that cloud droplets number concentration increases up to 27 % due to transported BBA using COSMO-
 ART, and-making cloud and rain less sensitive to future increase in anthropogenic emissions on regional

125 scale. The impact of sedimentation on LLSCs has been studied by Dearden *et al.* (2018) using the Met
Office NERC Cloud model (MONC) who highlight that sedimentation of cloud droplets, determined by
droplet size, could affect liquid water path by removing droplets from the entrainment zone, or by
lowering the cloud base and creating more heterogeneous cloud structure~~cloud droplets size has an effect~~
~~on liquid water path~~. Menut *et al.* (2019) showed with WFR-CHIMERE that a decrease of anthropogenic
130 emissions along the SWA coast leadled to a northward shift of the monsoonal precipitation and the
increase of surface wind speed over arid region in the Sahel, resulting in aan increase of mineral dust
emission. These previous modeling studies all highlighted in a regional scale and take care in majority
of only certain aerosol chemical ~~composition~~compositions, but however, they do did not take into account
all aerosol species detected during the field campaign especially black carbon (BC). Pedruzo-Bagazgoitia
135 *et al.* (2020) analyzed the stratocumulus-cumulus transition at fine scale (a dozen of kilometer sidelong)
using a LES at high resolution (50x50 m²) but without considering aerosols effects.

The aim of this study is to understand the relative impacts of local and transported aerosols on the
life cycle of LLSCs during the monsoon period over SWA by using observational data obtained from the
well-documented DACCIWA field campaign alongside a high-resolution LES model including
140 interactive aerosol module that is able to represent the complex aerosol compositions. This modelling
case is ~~focused on local scale and fine resolution over SWA which made it also~~ one of ~~the a~~ few studies
that model and ~~analyse~~analyzed stratiform cloud diurnal cycle over land ~~when a majority take place~~
~~over rather than~~ ocean. For this purpose, using observational data we firstly identified a reference case for
modeling, that is a LLSCs case observed on July 3, ~~2016~~2016, at Savè site. The short description of
145 observations, data, and the model as well as configurations of different simulations will be presented in
the Method section after the Introduction. Then an analysis will be driven to understand and validate the
reference case compared to measurements. Then, to assess the aerosol effects on stratiform clouds and
~~based on using~~ observed aerosol data, we have constructed several different aerosol profiles which differ
in term of aerosol size distribution and chemistry for sensitivity modeling studies. A first sensitivity
150 analysis will be driven to ~~asses~~assess the impact of aerosol concentration and consequently the indirect
effect on LLSC diurnal cycle. Another analysis will be focused on the impact of aerosol optical properties
by switching off aerosol (semi) direct contributions to the radiative budget to exhibit the relative changes
imposed by direct and semi-direct effects. In the DACCIWA framework, such analysis is a first and
differs from other modelling studies by performing this set of scenarios and configurations in order to
155 better investigate on indirect and semi-direct effects of aerosols from biomass burning and local
anthropogenic sources. Finally, this study will conclude by a summary of findings.

2. Methods

2.1 Observations

160 The relevant measurements of the DACCIWA field campaign used to select our LLSC case and to
configure the model simulations are described as follows:

i) Radiosondes were launched with the MODEM system every 1 to 1.5 hour between 17:00 and
11:00 UTC at the supersite of Savè in Benin (local time of Benin is UTC+1). This site~~It~~ is located at
185 km from the coast and 166 m above sea level (~~a.m.s.l~~) where the area is rather flat, and the
165 vegetation is mainly composed of small trees and shrubs. Temperature, pressure, relative humidity, and

wind vertical profiles in the lower atmosphere ~~reaching a maximum height of around~~(up to 1500 m ~~a.g.l~~ (above ground level) were measured with a 1 s temporal resolution (4-5 m of vertical resolution) (Derrien et al., 2016). These sondes use two ~~ballons~~balloons of different volumes to reach a preset time of ascent and after the cutting of the larger balloon, the second allows to retrieve the sonde for another use (Legain *et al.*, 2013).

ii) At the supersite of Savè, meteorological parameters were measured using different instruments. A CHM15k Ceilometer was deployed by the Karlsruher Institut für Technologie (KIT) to measure the cloud base height continuously with a 1 min resolution and a 15 m vertical resolution. Three cloud base heights are recorded from the backscatter profiles produced by the lidar with a wavelength of 1064 nm and a 5-7 kHz rate (Handwerker *et al.*, 2016). The cloud cover was monitored every day by using a MOBOTIX S15 cloud camera, installed by ~~UPS~~(Université Paul Sabatier (~~UPS~~) team, to obtain pictures in visible and IR every 2 min. The aperture angles for the IR channel corresponds to a 158 m x 114 m area at a height of 200 m. ~~P and pictures are coded in RGB components over 256 colors. A pixel is colored depending on the emissivity of the sky and so its brightness temperature. So a low cloud base is seen as red and a clear sky is seen as blue.~~ A microwave radiometer (humidity and temperature profiler HATPRO-G4 from Radiometer Physics GmbH) was installed by KIT to measure brightness temperature to retrieve absolute humidity, liquid water path, and air temperature. The surface ~~heat and radiation~~ fluxes ~~and net radiations~~ were measured with an energy balance station deployed over grass and bushes. Additional measurements were also soil heat flux, air density and turbulence parameters as well as sensible and latent heat flux.

iii) The aircraft campaign took place from 29 June to 16 July 2016. It was a collaborative work between three research aircraft but in this study only the ATR-42 is selected as it flew around Savè between 10:00 and 11:00 UTC and probed the cloud layer. The cloud droplet size distribution was measured with a cloud droplet probe (CDP) (Taylor *et al.*, 2019). The chemical composition for non-refractive compounds was measured with an Aerodyne compact Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS) ~~analyzed~~(Brito et al., 2018). The black carbon (BC) mass concentration was measured with a single particle soot photometer (SP2) (Denjean *et al.*, 2020b). The aerosol number size distribution was measured with a custom-built scanning mobility sizer spectrometer (SMPS, 20–485 nm), an ultra-high sensitivity aerosol spectrometer (UHSAS, 0.04–1 μm), and an optical particle counter (OPC GRIMM model 1.109, 0.3–32 μm) corrected for the complex refractive index provided in Denjean et al. (2020a). The total concentration number of particles larger than 10 nm was measured by condensation particle counter (CPC, model MARIE). ~~A suite of instruments measured also meteorological~~ Meteorological variables such ~~145~~ as temperature, humidity, pressure, and winds ~~speed and direction were also measured by a suite of airborne instruments.~~ Gas ~~A gas~~ concentration analyzer ~~was used to measure~~ ~~some certain chemical gases including like~~ CO_2 , CH_4 , and CO .

2.2 Description of the studied case

Our study analyzes the diurnal cycle of LLSCs based on the case study of 3 July 2016 at the Savè supersite (Fig. 2). The cloud deck formed during the night, at around 02:00 UTC, close to ~~the appearance of the core of the Nocturnal-Low-Level Jet (NLLJ), which could have a maximum speed around 6 m s^{-1} (Kalthoff *et al.*, 2018) where associated with a maximum~~ the cooling is maximum (Lohou *et al.*, 2020).

and could reach a maximum speed of around 6 m s^{-1} (Kalthoff et al., 2018). At formation, the cloud base and top heights ~~are~~ were located around $310 \pm 30 \text{ m}$ and $640 \pm 100 \text{ m}$, respectively, and ~~were~~ maintained due to the cloud top radiative cooling and cold advection (Dione et al., 2019).

210 The diurnal cycle of LLSCs over SWA typically involves four phases: stable phase, jet phase, stratus phase, and convective phase (Dione et al., 2019; Lohou et al., 2020). The stable phase begins just after sunset and is characterized by a weak monsoon flow and the cessation of buoyancy-driven turbulence (generated by surface heating) within the atmospheric boundary layer (ABL)-PBL (Zouzoua et al., 2021). The jet phase corresponds to the settlement of key drivers of cooler air advection. Maritime inflow (MI),
215 cold and slightly humid air from the Guinean coast, reaches Savè at the end of the afternoon (between 16:00 UTC and 20:00 UTC) then the NLLJ formation occurs (Adler et al., 2019). The stratus phase begins with LLSC formation when the advective cooling continuously increases the relative humidity (RH) until saturation is reached between 22:00 and 06:00 UTC. Turbulent mixing ~~maintains cooling below the cloud base~~ beneath the NLLJ and alongside strong radiative cooling at the cloud top leads to the persistence of
220 a thick stratus layer (Schuster et al., 2013; Babić et al., 2019). The LLSCs life cycle ends during the final convective phase which begins when the ABL-PBL develops vertically due to solar heating at the surface alongside a weak radiative cooling at cloud top (e.g., Ghonima et al., 2016). By using dataset from Savè supersite Zouzoua et al. (2021) identified three scenarios of evolution depending on the LLSCs coupling to the surface at sunrise. The coupling was assessed by the departure between the Cloud Base Height
225 (CBH) and the Lifting Condensation Level (LCL).




Figure 2. 3 July cloud evolution with the representation of the Cloud Base Height (CBH), the Cloud Top Height (CTH), LCL and ATR-42 flight track near Savè.

230 The LLSCs observed on 3 July 2016 follow the four aforementioned phases and evolves by scenario C described by Zouzoua et al. (2021) as seen in Figure 2; the cloud is coupled to the surface at sunrise (06:30 UTC) and its base rises with growing ABL-PBL until its break-up in the late afternoon (around
235 16:00 UTC). This cloud deck stands longer (2-3 hours more) compared to other LLSCs observed during

the campaign. ~~Co~~The co-located radar at Savè supersite allows the detection of light precipitation from higher clouds during the first hours of the convective phase while no precipitation was detected by the surface rain gauge. Thus, this late LLSC break-up could be explained by a significant increase in its liquid water content (LWC) caused by evaporation of this light precipitation (Zouzoua *et al.*, 2021).
240 Nevertheless, our focus of this study is on the diurnal cycle of LLSC as influenced by aerosols alongside planetary boundary layer dynamics rather than examining the above hypothesis appeared to be related to a process beyond the local scale. Therefore, our model setting is made to specifically eliminate the influence of mid-cloud layer for the purpose.

On 3 July 2016, the ATR-42 flew around Savè supersite and probed the boundary layer around 10:00
245 UTC. The measurements confirmed an important aerosol number concentration with a maximum of around 3500 cm⁻³ mainly located in the Aikten mode. The ATR-42 detected an export of pollution from Lomé (a coast city) which could explain the higher loading of aerosol in the Aikten mode (Denjean *et al.*, 2020a). The aerosol chemical composition was mainly dominated by organics (55.3%), followed by sulfates (24.5%), ammoniac (11.2%) and nitrates (6.2%) while little BC mass was detected around Savè
250 (2.8%). However, these data are directly extracted from DACCIWA ~~database~~database, and the size distribution has to be corrected from the aerosol refractive index to avoid bias. For this purpose, Denjean *et al.* (2020a) provided corrected profiles for different typical aerosol population encountered during the DACCIWA campaign.

255 2.3 Meso-NH Model

In this study, we have simulated the observed case using the French model Meso-NH (Lac *et al.*, 2018). Meso-NH is a non-hydrostatic atmospheric research model that has been applied to studies in scales, ranging from synoptic to turbulent. ~~Employed-Deployed~~ in a limited area, the model uses advanced numerical techniques like monotonic advection schemes for scalar transport and ~~fourth-order~~fourth order
260 WENO advection scheme for momentum (Jiang and Shu, 1996). Sub-grid turbulence is parametrized using turbulence kinetic energy (TKE) based on Deardorff turbulent mixing length. ~~The advection scheme used for horizontal and vertical velocities is CEN4TH, a A 4th-fourth order advection scheme CEN4TH, eentredcentered~~ on space and time, is applied with a Runge-Kutta ~~eentredcentered~~ 4th order temporal scheme for momentum advection. Aerosol and chemistry are also well represented. Here, Meso-NH
265 version 5.4.2 (~~www.mesonh.aero.obs-mip.fr~~) is used and the relevant component modules and parametrizations for this study are described as follows:

LIMA (Liquid Ice Multiple Aerosol) is a complete two-moment scheme (Vié *et al.*, 2016) predicting both the mass mixing ratio and the number concentration of aerosol species. Based on ICE3-ICE4 schemes (Caniaux *et al.*, 1994; Pinty and Jabouille, 1998; Lascaux *et al.*, 2006) and the two-moment
270 warm microphysical scheme C2R2 from (Cohard and Pinty, 2000), LIMA predicts the number concentration of cloud droplets, raindrops and pristine ice crystals. It includes a prognostic representation of aerosol population using a superimposition of several aerosol modes with each mode defined by its chemical composition, size distribution and aerosols can act as a Cloud Condensation Nuclei (CCN) or an Ice Freezing Nuclei (IFN). For boundary layer cloud LESSs, a pseudo-prognostic approach for
275 supersaturation was developed (Thouaron *et al.*, 2012) to limit the droplet concentration production and so it would represent cloud-top supersaturation better. A variant to C2R2, called KHKO, was developed

by Geoffroy *et al.* (2008) for clouds producing drizzle following Khairoutdinov and Kogan (2000) parametrization. These clouds are low precipitating warm clouds, and not sufficiently thick to produce heavy rain. The precipitating hydrometeors are drizzle only and their diameter are of the order of several dozens of micrometers. These modifications for KHKO were brought inside LIMA warm phase in order to better represent drizzle ~~completing LIMA.~~

ECMWF radiation module, originated from ECMWF and based on two-stream methods, calculates the atmospheric heating rate and the net surface radiative forcing. Longwave radiation scheme used is Rapid Radiation Transfer Model (RRTM; Mlawer *et al.*, (1997)), based on the correlated k-distribution method. It integrates 16 bands and 140 g points (Morcrette, 2002). The shortwave scheme uses the photon path distribution method (Fouquart and Bonnel, 1980) in six spectral bands. Fluxes are calculated independently in clear and cloudy portion before being aggregated. The liquid cloud effective radius is computed from the liquid water content with the Martin *et al.* (1994) parametrization.

ORILAM (Organic Inorganic Lognormal Aerosols Model) is an aerosol module coupled to Meso-NH and connected to LIMA (Tulet *et al.*, 2005). It describes the size distribution and the chemical composition of aerosols using two lognormal functions for the Aitken and accumulations modes. These modes are internally mixed and for each of them the model computes the evolution of the primary species (black carbon and primary organic carbon), three inorganic ions (NO_3^- , NH_4^+ , SO_4^{2-}) and condensed water. ORILAM includes Second Organic Aerosols (SOA) (Tulet *et al.*, 2006) but are not taken into account in this study. Three moments (zeroth, third and sixth) are considered for each mode to compute the evolution of total number, median ~~diameter~~diameter, and geometric standard deviation. The size distribution can evolve through a particle coagulation process with both intramodal and intermodal calculations. It can also evolve through condensation and merging between modes. ORILAM includes the CCN activation scheme of Abdul-Razzak and Ghan (2004) in order to replace the one of LIMA to calculate the number of activated CCN. The others LIMA parametrizations in warm phase like the calculation of drizzle remain active. The use of ORILAM needs to activate the gas phase chemistry scheme of Meso-NH (Tulet *et al.*, 2003; Mari *et al.*, 2004) using the EXQSSA solver. Inorganic chemistry system (EQSAM, Metzger *et al.* (2002)) solves the chemical composition of sulfate nitrate-water-ammonium aerosols based on thermodynamics equilibrium. For secondary organic aerosols, the thermodynamic equilibrium uses the MPMPO scheme from Griffin *et al.* (2003). ORILAM ~~computes directly~~directly computes the evolution of aerosol extinction, SSA and asymmetry factors that are coupled online with the radiation scheme of Meso-NH for the 6 short wavelengths from the aerosol chemical composition and size parameters (Aouizerats *et al.*, 2010).

SURFEX is a standardized surface module containing surface schemes externalized of Meso-NH (Masson *et al.*, 2013). Each grid point can be split into four tiles: land, town, sea and inland water (lake, rivers). In case of a shrubs typical surface, the interactions between soil, biological and atmosphere are calculated by ISBA parametrization (Noilhan and Planton, 1989). It represents the effect of vegetation and bare soil. Several evapotranspiration formulations are available for plants and for simulating the CO_2 fluxes. Soil is represented as a bucket of two or three layer. The land tile can be separated in up to 19 subtiles following the type of vegetation.

2.4 Model settings

Based on observations and the capability of the model, a reference case (REF) was first designed to simulate through LES. The reference case serves as a base to reproduce the major features of the observed LLSCs diurnal cycle particularly under an observed aerosol profile. It also serves as a reference for further sensitivity simulations with different aerosol configurations to study the impacts of aerosol composition alongside abundance on LLSCs.

The domain is a 3D box of 9.6 km x 9.6 km x 2 km in size with a horizontal resolution of 40 m x 40 m. The vertical resolution is 10 m between 0 m and 1200 m then 40 m above until 2 km of altitude. Such high resolution is able to resolve explicitly the biggest turbulent eddies. A periodic boundary condition on the horizontal directions is applied and an absorbing layer is set at 1.8 km. A thermodynamic perturbation is deployed to activate turbulence at the beginning of the simulation at 23:00 UTC of 2 July and the spin-up is 1h. A subsidence profile is applied following Bellon and Stevens (2013) scheme $w_{subs}(z) = -w_0(1 - e^{-\frac{z}{z_w}})$, $w_{subs}(z) = -w_0[1 - \exp(-z/z_w)]$, with $w_0 = 15 \text{ mm s}^{-1}$ and $z_w = 250 \text{ m}$. This subsidence profile is applied during the entire simulation to keep a nearly constant cloud top height during the stratus phase and to better control the convective phase. The surface energy and water fluxes are simulated by SURFEX ISBA scheme parametrized by data from Savè supersite measurements with the typical vegetation around of Savè which consists of shrubs, crops or taller trees assuming a flat surface corresponding to the area around Savè. A time-step of 2s is used which appears to be adequate to study accurately the LLSCs diurnal variations particularly involving aerosol and cloud microphysics. Note that the radiation scheme is called every 10 minutes. Note that previous studies regarding nocturnal stratus-stratocumulus suggested that a vertical resolution as fine as 5 meters near the cloud top would be necessary for reproducing the cloud top entrainment and thus cloud macrophysical structures (Stevens et al., 2005). Since the nocturnal-diurnal life cycle in our case involves a dynamically evolving cloud top (particularly in the daytime), it makes it difficult to prescribe a highlight zone for finer resolution. Our fast-testing results did not suggest an alarming difference between the run with 10 m and 5 m vertical resolution (not shown). Therefore, the current vertical resolution and the time step are selected to well cover all possible cloud tops during the simulation time and to provide the best economic computational performance for aerosol-cloud interaction with a fully coupled chemistry model.




Figure 3. REF aerosol chemical mass compositions (a) and size distribution fitted into 2 modes described in Table 1 (b) used in REF.

| | $N_a (cm^{-3})$ | σ | D (nm) |
|--------|-----------------|----------|--------|
| Mode 1 | 654 | 1.49 | 63.98 |
| Mode 2 | 1530 | 1.53 | 190.97 |

Table 1. REF aerosol size distribution described by two modes configured by three parameters (number concentration N_a , standard deviation σ and diameter D).


REF case is configured using the ~~radiosounding profiler~~ radiosondes of 2 July at 23:00 UTC for temperature, ~~humidity~~ humidity, and horizontal wind components (U, V). The simulation is then controlled by tendency profiles of temperature, ~~and~~ humidity, and horizontal wind applied homogeneously on the domain each hour. These tendency profiles are based on the ~~hourly radiosondes~~ ~~radiosoundings profiles~~ launched on 3 July between 00:00 and 11:00 UTC. After 11:00 UTC, the next tendency profiles were designed based on the measurements of the microwave radiometer, the analysis of ~~resulted~~ surface incoming solar radiative flux, cloud thickness and cover. Note that, despite these best possible efforts in configuring a set of observation-constrained tendency profiles to reproduce observed cloud field, it is difficult to eliminate the possibility that such profiles could reflect certain local thermodynamic effects however small they are. In practice, our principal is to make the profiles to be able to force the modeled clouds reproduce observed quantities of major features such as cloud top, base, LWP, surface incoming solar radiation, among others, in the REF case. This would serve the best purpose for us to address the major issue of this study, i.e., the role of different aerosol profiles in the diurnal cycle of modeled LLSCs.

We decided to use a "background" distribution as the aerosol profile for REF simulation. This profile, described in Denjean *et al.* (2020a), actually reflects the influence of aged BBA on clouds with minor

influence of local anthropogenic sources. The aerosol number size distribution is dominated by a particle accumulation mode centered at 190 nm and a smaller Aiken mode centered at 64 nm as seen in Figure 3b. This profile exhibits a high loading of aerosols with a maximum of 1400 cm⁻³ detected in the accumulation mode. The aerosol chemical composition was dominated by organics (52.2%) followed by sulfates (25.3%), ammonium (8.9%), BC (10.7%) and nitrates (2.8%). The configuration of ORILAM ~~have~~has been initialized using the REF aerosol chemical composition and number size distribution given in Table 1 and Figure 3b by fitting the SMPS profiles into two lognormal modes using the "py-smps" package (Hagan *et al.*, 2022) with each mode having the same chemical composition.

385 **3. Analysis of REF Results**

3.1 Simulated LLSCs evolution



390 **Figure 4.** Simulated LLSCs deck evolution compared to Savè ceilometer and radar measurements, vertical colored bars attribute at each altitude level a modeled cloud presence probability. Here mean CBH and CTH represents cloud base and cloud top height, respectively.

395 The simulation of the REF scenario reproduces the formation of the observed LLSCs deck on 3 July 2016 as shown in Figure 4. The formation of clouds leads, as described in section 2.2, to the end of the

jet phase. The domain mean CBH estimated from the mixing ratio of cloud droplets follows the ceilometer's measurements during the stratus phase between 2:00 and 10:00 UTC, varying between 100 and 200 m of altitude. The simulated mean Cloud Top Height (CTH) evolves from 400 to 550 m of altitude, in range of the values from 500 to 580 m detected by the radar. During the convective phase, the model results differ slightly from the observations. Note that ~~To~~ to analyze the cloud cover profile over the domain, the Cloud Presence Probability (CPP) at each altitude level model layer, differing from cloud fraction that is often defined as a column metrics, -is calculated as a percentage of by taking all cloud pixels which with exceed 0.05 g kg⁻¹ in a total condensed water mixing ratio exceeding 0.05 g kg⁻¹ as a cloud pixel at each at the given model layer altitude level (Fig. 4).

Nevertheless, the mean simulated CBH and CTH are overestimated compared to ceilometer and radar values in some period particularly late morning and afternoon. The simulated CBH can differ from ceilometer one by 150 m of altitude at 11:00 UTC. The CTH is often overestimated by 100 m. Between 15:00 and 16:00 UTC, the simulated mean CBH follows approaches again the detected values ceilometer readings (600 to 950 m) (no radar values are available to validate the simulated CTH). As mentioned in section 2.1, the ceilometer is a lidar while the radar values are derived from reflectivity vertical profiles which have a 30 m of resolution. The differences between the model and the observation between 13:00 and 16:00 UTC are likely due to the domain averaging of the simulated values, the tendency profiles established from corrected radiosonde values, the ceilometer values limited to only one vertical direction, the vertical resolution of radar profiles, or the limitation of radar in detecting hydrometeors.




Figure 5. Comparison between modeled liquid water path (LWP, $g\ kg^{-1}\ m$) and the images from Savè cloud camera at 06:00 (top), 12:00 (middle) and 16:00 UTC (bottom).

420


~~To analyze the cloud cover over the domain, Cloud Presence Probability (CPP) at each altitude level is calculated by taking all pixels which exceed $0.05\ g\ kg^{-1}$ in total condensed water mixing ratio as a cloud pixel at each altitude level. Liquid water path (LWP) at each column (Fig. 5), calculated based on cloud pixels, brings a view on the horizontal organization and homogeneity of the cloud deck. During the stratus phase, the CPP is nearly equal to 100% between CBH and CTH giving a homogeneous cloud deck.~~

425 The top panel of Figure 5 gives a comparison of the cloud organization in the model and from cloud observations with visible-sky camera (visible range).

At 06:00 UTC, ~~it can be seen that cloud~~ deck covers the entire domain as seen in both modeled result and in observations (note the clear-distinct cloud rolls in model results). Between 10:00 and 13:00 UTC, the CPP ~~at the mid distance from~~ layers between mean CBH and CTH decreases from near 100% to 90%. Near the two averaged values, CPP decreases more to reach near 60% and 80% at CBH and CTH, respectively. This leads to a more-less inhomogeneous cloud deck confirmed by the LWP map and the observation of the cloud-sky camera at 12:00 UTC shown in the middle row of Figure 5. Indeed, more ~~no-cloud~~ pixels begin to appear between clouds and ~~the~~ sunlight is seen through the cloud deck by the camera. Finally, the CPP continues to decrease until the end of the convection phase ~~by~~ reaching with a maximum ~~of near~~ barely reaching 80%, and a value around mean CBH and CTH ~~it reaches as low as~~ 20% and 40%, respectively. This demonstrates the break-up of the cloud deck during convection and the ~~decreasing of~~ cloud thickness thinning. The bottom panels of Figure 5 show clearly the dissipation of a large number of clouds alongside substantially thinning of the others ~~break-up of clouds~~ at 16:00 UTC PM. The LWP map (Fig. 5b) shows numerous thin clouds corresponding to those seen by the camera of Savè.

Figure 6a shows the comparison between the SW radiation flux at the surface (SWRADSURF), averaged over the modeled domain and measurements performed by the energy balance station. Observed values are fitted following the locally Weighted Scatterplot Smoothing (LOWESS) method (Cleveland, 1979), which is a non-parametric regression method performing weighted local linear fits. The temporal evolution of the modeled SWRADSURF follows ~~well~~ the observations well although some biases can be observed. After 06:00 UTC the solar radiation reaches the ground and as the cloud deck thickness and covering show little variations, the radiative flux increases gradually by reaching near 200 W m^{-2} at the end of the stratus phase (10:00 UTC). As ~~the~~ clouds deck becomes inhomogeneous during the convective phase (10:00 to 16:00 UTC), the solar flux reaches a maximum of 300 W m^{-2} , which is a bit less than the fitted 350 W m^{-2} value. Finally, when the clouds break-up, more solar radiation can reach the surface. After this period during which model and observations agree well, from 15:00 UTC the mean curve decreases to 200 W m^{-2} while the fit curve is near 320 W m^{-2} due to an overestimation of the cloud thickness by the model. At 16:00 UTC, both modeled and measurement values are very close at around 280 W m^{-2} . ~~But, as the LLSCs breakup later than common clouds during DACCIWA campaign (more than 2-3 hours), it implies that modeled SWRADSURF is low.~~ Generally, the modeled maximum values are higher than the ones detected by the Savè instrument. For example, at 10:00 UTC, the balance detected a peak of 300 W m^{-2} while the model value reached near 400 W m^{-2} . ~~Such difference is reduced during the convective phase.~~

Figure 6b and 6c shows that the evolution of the modeled latent and sensible heat fluxes ~~well~~ reproduced those measured by the instrument well as shown in the middle and bottom row respectively of Figure 6. During the night, the sensible heat flux was negative then increased to 0 W m^{-2} close to the sunrise time (6:00 UTC), indicating a reduction of the cooling close to the ground (Dione *et al.*, 2019). Between 9:00 and 14:00 UTC, the modeled two heat fluxes followed the measured trends though overestimated by almost 70 and 18 W m^{-2} . Then the mean modeled curves go below the fitted observed curves at 15:00 UTC and finally decrease to almost 0 W m^{-2} after 18:00 UTC. The difference between modeled and observed latent and sensible heat fluxes may be due to the different area covered by the measurements and the model and the prescribed subgrid-scale distributions of cloud droplets.



470 **Figure 6.** Comparison between Savè surface observation and REF simulation for SW radiation flux at surface
 (SWRADSURF,a), sensible heat flux (H ,b) and latent heat flux (LE ,c) all expressed in $W m^{-2}$ at the surface. The variation of
 REF for each parameter indicates the range of possible values these parameters can take.

In Summary, the REF simulation has successfully reproduced the major observations obtained by the
 475 instruments at Savè on 3 July 2016. The modeled cloud thickness and coverage represent well the cloud
 situation with some inaccuracies due to the lack of data and insufficient correction of the tendency profiles
 applied all along the simulation to control temperature and humidity every hour. The modeled heating of
 the ground by solar radiation also follows the measurements of the energy balance of Savè and maximum

variation on the domain are a bit overestimated. The sensible and latent heat fluxes detected at Savè have also been well captured by the model.

3.2 Thermodynamic, dynamical, cloud microphysical, and radiative analyses


Thermodynamic, dynamical, and radiative processes and their interaction with cloud microphysics are among the key factors in determining the life cycle of LLSCs. Here we discuss the evolution of these processes simulated by the model in the REF case in order to better understand the reasons behind model-observation consistency or discrepancy. The discussion will be emphasized in three periods. The first period is the transition between jet and stratus phase (between 00:00 and 04:00 UTC) to observe how clouds are formed. The second period is the stratus phase between 06:00 and 10:00 UTC because of the stability of the cloud layer observed by instruments of Savè. The third period is the convective phase between 12:00 and 17:00 UTC to study how the properties of LLSCs evolve during the break-up stage.

3.2.1 Transition jet-status phase

The formation of clouds is controlled by the temperature and humidity tendency profiles established from the radiosonde measurements made on the 3rd of July 2016. Figure 7 gives domain-averaged profiles of the simulated temperature (T) and relative humidity. As explained in section 2.2, maritime inflow already reached the site, increasing that increased humidity and created creating NLLJ. Temperature decreases from 24°C to 23°C at ground from 00:00 to 04:00 UTC and from 24°C to 21°C near 400 m of altitude. The advection of cold and slightly humid air leads to the increase of RH as expected reaching 100% at 02:00 UTC at 100 m. After this time, RH exceeds saturation between 100 and 500 m of altitude. The inversion occurs around 325 m and 500 m respectively at 02:00 UTC and at 04:00 UTC. The NLLJ is well represented as the mean wind speed (w_s) before cloud formation is more greater than 7 m s^{-1} . After cloud formation of cloud, the NLLJ core corresponds near to the mean cloud base height (Adler et al., 2019; Babic et al., 2019; Lohou et al., 2020). The turbulence during this period is shear-driven due to this NLLJ which yields a well-mixed the sub-cloud layer. The TKE is high above ground (0.5 to $0.2 \text{ m}^2 \text{ s}^{-2}$), then decreases to near zero along altitude above rough 200 meters at 00:00 UTC. After 02:00 UTC, TKE increases at the level of the CTH (350 and 500 m) and decreases at the center of clouds ($0.04 \text{ m}^2 \text{ s}^{-2}$), indicating this area is less turbulent than the extremities of the cloud layer.

Cloud droplet number concentration or CDNC (N_c) is determined by the supersaturation in an updraft and the concentration of aerosols that activate at their supersaturation. In Figure 7e, simulated aerosol concentrations were the highest close to the ground and decreased with altitude up to around 2 km. This simulated aerosol profile is similar to those observed by airborne measurements during DACCIWA (Taylor et al., 2019; Denjean et al., 2020a) with locally emitted aerosols transported above the boundary layer due to a combination of land-sea surface temperature gradients, orography-forces circulation and the diurnal cycle of the wind along the coastline (Deroubaix et al., 2019; Flamant et al., 2018). The simulated clouds microphysics reflectreflects polluted conditions with N_c reaching 1750 droplets droplet cm^{-3} and r_c around $5 \mu\text{m}$ which is not enough to form even drizzle (size between 0.2 mm and 0.5 mm, (Pruppacher et al., 1998; Sandu et al., 2008)). These values are in the range of those measured inland at the same altitude by Taylor et al. (2019) during DACCIWA. Median of simulated CDNC was 500

520 *droplets cm⁻³* at the beginning of cloud formation and reached *1750 droplets cm⁻³* latter, most likely due to the continuous activation of aerosol into cloud droplets.



525 **Figure 7.** Profiles from left to right of temperature (T, a), relative humidity (RH, b), equivalent potential temperature (θ_e , c), horizontal wind speed (w_{sws} , d), aerosol number concentration (N_a , dashed curve, e), cloud droplets number concentration (N_c , plain curve, e), cloud droplet radius (r_c , f), turbulent kinetic energy (TKE, g), longwave heating rate (LWHR, dashed curve, h) and shortwave heating rate (SWHR, plain curve, h) at 00:00, 02:00 and 04:00 UTC. Dashdot horizontal lines represent mean cloud base height (CBH) and dotted horizontal lines the mean cloud top height (CTH).

530 The emission of thermal radiation by the clouds during the stratus phase creates a cooling at the cloud top as demonstrated by the profiles evolution of the Long-Wave Heating Rate (LWHR) at Figure 7h. The more numerous the cloud droplets are ~~the cloud droplets~~ the stronger the cooling is, as shown in ~~the figure~~ Fig. 7h that Short-Wave Heating Rate (SWHR) can reach $-50 K day^{-1}$. This strong longwave emission is able to reduce the thermal production of turbulence above at the cloud top, ~~and contribute to the stability of air masses at the cloud top,~~ deepening the temperature inversion ~~above cloud top~~. A stabilized cloud top layer by radiative cooling and a NLLJ core contributing to the shear-driven turbulence below the cloud base allows the well mixing of the cloud layer, making the LCL to correspond to the LLCSS base as seen at Figure 2 (Adler *et al.*, 2019; Lohou *et al.*, 2020). ~~In fact, modeled cloud layers are very stable before the sunrise as observed at Savè during the DACCIWA campaign.~~

540


3.2.2 Stratus phase

The stratus phase starts just after the sunrise. To maintain stratus in almost the same state as in the previous period needs certain proper temperature and humidity conditions as shows ~~at-in~~ Figure 8. Indeed, the ground temperature ~~at the ground~~ is still at 23°C at 06:00 and 08:00 UTC, and 20°C at the mean CTH

545 (500 and 550 m respectively). RH profiles indicate that supersaturation still exists between the range of CBH and CTH, allowing droplets condensation to continue. Air masses are quite stable-well-mixed between ground and CTH within PBL during this time as θ_e is near equal to 347 K and the inversion layer is settled where θ_e is reaching 350–351 K. The horizontal wind speed between the ground and the cloud base decreases, which indicates the reducing of the NLLJ core (nearly 2 m s^{-1}) and its rises in altitude due to the turbulent mixing induced by the LW cooling at the cloud top during the night. The turbulence between ground and cloud center decreases to $0.03 \text{ m}^2 \text{ s}^{-2}$ then finally increases slightly to $0.04 \text{ m}^2 \text{ s}^{-2}$ at the mean CTH. The TKE is a bit stronger at 08:00 UTC, reaching $0.05 \text{ m}^2 \text{ s}^{-2}$ in the cloud layer which is explained by an increase of the vertical wind speed.

555 The aerosol concentration at 06:00 and 08:00 UTC is around 2000 cm^{-3} up to 500 m then it decreases along altitude. This concentration is still high to allow the formation of 1100–1200 *droplets* cm^{-3} between CBH and CTH. The concentration of cloud droplets leads to a maximum droplet radius of $6 \mu\text{m}$, which is still not enough to form drizzle. The cloud layer has an albedo close to 1 due to the high droplet concentration. The presence of light absorbing aerosol causes the shortwave that causes the Short-Wave Heating Rate (SWHR) amplified at the cloud top by semi-direct effect. At 08:00 UTC, the SWHR and LWHR are equal to 27 K day^{-1} and -70 K day^{-1} , respectively.

560




565 **Figure 8.** Profiles from left to right of temperature (T, a), relative humidity (RH, b), equivalent potential temperature (θ_e , c), horizontal wind speed (w_s , d), aerosol number concentration (N_a , dashed curve, e), cloud droplets number concentration (N_c , plain curve, e), cloud droplet radius (r_c , f), turbulent kinetic energy (TKE, g), longwave heating rate (LWHR, dashed curve, h) and shortwave heating rate (SWHR, plain curve, h) at 06:00, 08:00 and 10:00 UTC. Dashdot horizontal lines represent mean cloud base height (CBH) and dotted horizontal lines the mean cloud top height (CTH).

570 At 10:00 UTC, the cloud layer starts to rise significantly, with CBH and CTH reaching 340 and 660 m, respectively. Moreover, more-stronger solar radiations-irradiance can reach the ground (220 W m^2), leading to the heating of the surface and the increasing of the sensible and latent heat fluxes as seen in Figure 6. It also increases the temperature near ground to $24 \text{ }^\circ\text{C}$ and at the cloud top the temperature is

20 °C. Supersaturation occurs obviously between cloud base and cloud top and the inversion layer is observed above the cloud top between 660 and 750 m. The NLLJ core is no longer present as the horizontal wind speed decreased to 0.5–0.75 $m s^{-1}$. However, the turbulent kinetic energy increases to 0.1 $m^2 s^{-2}$ ~~almost to the~~ throughout the vertical layer from 50 meter above the ground to a level just below the cloud top. This enhancement of turbulence is expected to increase entrainment ~~from above~~ entering the cloud from above as well. Clouds are having a maximum of 1000 cm^{-3} and a maximum droplet radius reaching near 6 μm . Given these values, the cloud albedo is near equal to 1 and the SWHR increases to 45 $K day^{-1}$. It almost compensates the LWHR value of 65 $K day^{-1}$. During the 3rd of July, the radar detected light precipitation from higher clouds but this simulation doesn't model this type of clouds. So, the cloud deck stands longer due to the set-up of tendency profiles especially the one for humidity which keeps supersaturation at stratus layer until 10:00 UTC.

585 3.2.3 Convective phase

This phase is extending from 12:00 to 17:00 UTC in 3 July 2016 case when the SW radiation flux at surface is maximized at 300 $W m^{-2}$ (Figure 6), leading to a more intense heating from the surface. During this period, the temperature at ground evolves from 25 to 27 °C as seen at Figure 9. The temperature near the cloud top is 20°C at 12:00 and 14:00 UTC. At the break-up time (16:00 UTC), the temperature is lower than 18°C at the CTH. The formation of clouds is possible due to RH exceeding 100% in upper altitude, as the convection of humid air masses causes the CBH and CTH to rise from 450 to 925 m and from 760 to 1100 m, respectively. Moreover, at the break-up (16:00 UTC), the equivalent potential temperature decreases above 450 m of altitude, indicating air masses become more unstable with altitude. The horizontal wind speed is weak at the beginning of the phase with 0.5 $m s^{-1}$ at ground level but increases along time to reach 1 $m s^{-1}$ at ground and 3 $m s^{-1}$ around 700m. This increase coincides the dissipation of the LLSCs and indicates ~~indicates the end of clouds break-up and~~ the arrival of the marine inflow.



600 **Figure 9.** Profiles from left to right of temperature (T, a), relative humidity (RH, b), equivalent potential temperature (θ_e , c),
 | horizontal wind speed (w_s , d), aerosol number concentration (N_a , dashed curve, e), cloud droplets number concentration
 | (N_c , plain curve, e), cloud droplet radius (r_c , f), turbulent kinetic energy (TKE, g), longwave heating rate (LWHR, dashed curve,
 | h) and shortwave heating rate (SWHR, plain curve, h) at 12:00, 14:00, 16:00 and 17:00 UTC. Dashed-dot horizontal
 | lines represent mean cloud base height (CBH) and dotted horizontal lines the mean cloud top height (CTH).

605

The turbulence profiles evolve along altitude during the convection phase, reaching $0.075 \text{ m}^2 \text{ s}^{-2}$ in
 center of clouds and almost zero near 850 m at 12:00 UTC. This profile evolves at 14:00 UTC to reach
 | $0.25 \text{ m}^2 \text{ s}^{-2}$ and zero at cloud center and 1000 m, indicating a reinforcement of turbulence due to an
 | elevation of dynamical production via the vertical wind speed increase. Finally at break-up time,
 610 turbulence near the ground decrease but at cloud level the TKE has a value of $1.5 \text{ m}^2 \text{ s}^{-2}$ showing a strong
 turbulence layer. This turbulent layer further moves to a upper altitude after 17:00 UTC.

The aerosol distribution varies along with the dynamical situation. The maximum aerosols
 concentration reaches 1800 cm^{-3} below 800 m and 1700 cm^{-3} below 1000 m at 12:00 and 17:00 UTC
 615 respectively. Again, with so numerous aerosols, the cloud droplets concentration as a domain mean has a
 maximum value of $900 \text{ droplets cm}^{-3}$ at 12:00 UTC. This value decreases along time as more clouds
 | dissipated. At the break-up of clouds, reduced cloud coverage allows more solar radiation to reach the
 ground. The maximum value of SWHR drastically changes from 45 K day^{-1} at 12:00 UTC (almost
 compensating cloud top cooling) to about 10 K day^{-1} at 16:00 UTC. The cloud top cooling is near constant
 at the end of convection phase with -45 K day^{-1} .

620

4. Sensitivity study to examine the influence of different aerosol profiles on LLSC diurnal cycle

625 Previous studies have indicated that the life cycle of stratus or stratocumulus within planetary
boundary layer depends on the subtle balance among several critical while interconnected forcings
including surface heat fluxes, cloud top and base radiative profiles, and thus turbulent mixing (e.g.,

Stevens *et al.*, 2005; Dussen *et al.*, 2014, Ghonima *et al.*, 2016). Apparently, our simulation results of the REF case support previous findings particularly for cases over land with surface sensible heat playing a significant role. Nevertheless, the role of aerosols in such a life cycle have rarely explored in-depth. Given the critical role of aerosols in determining cloud macro- and microphysical features and thus radiation, this is a must-addressed issue to advance our understanding of the LLSC life cycle. A unique component of our study is the deployment of an interactive aerosol and atmospheric chemistry module in this observation-constrained modeling effort. In the following section we will discuss roles of aerosol variations in both number concentration and chemical composition in influencing the diurnal cycle of observed LLSCs.

4.1 Aerosol profiles used in sensitivity simulations

The result of REF simulation has demonstrated that the Meso-NH model is able to reproduce many observed features of the July 3 LLSC case ~~even though Meso-NH model has some~~ despite certain biases. Moreover, the dynamical, thermodynamic and aerosol parameters are reasonably well simulated by the model. It is well acknowledged that aerosol, from both anthropogenic activities and biomass burning emissions, may influence cloud formation directly through absorbing solar radiation and indirectly by serving as CCN. To identify the aerosol sources and key processes of aerosol in such cloud enhancement, we tested the respective influence of anthropogenic sources and of aerosol semi-direct effects. For this purpose, we have configured two different aerosol scenarios based on observations during the field campaign (Figure A1 and Table 2), then applied them in a set of sensitivity simulations that would be otherwise the same as the configuration of REF simulation.

| Case | | $N_a (cm^{-3})$ | σ | D (nm) |
|-------|--------|-----------------|---------------------|--------|
| POL | Mode 1 | 17100 | 1.54 | 55.19 |
| | Mode 2 | 2650 | 20142.14 | 101.83 |
| CLEAN | Mode 1 | 65 | 1.49 | 63.98 |
| | Mode 2 | 153 | 1.53 | 190.97 |

Table 2. POL and CLEAN aerosols size distribution described by two modes configured following three parameters (number concentration, standard deviation and diameter).

To investigate the impacts of anthropogenic and biomass burning sources on clouds, three additional numerical experiments were performed in addition to REF (Table 2): (1) an experiment with strong anthropogenic pollution influence based on the aerosol chemical composition observed by Brito *et al.* (2018) and Denjean *et al.* (2020a) in urban plumes originating from the polluted cities of Lomé, Accra and Abidjan, which is called POL; (2) an experiment designed to underestimate aerosol emission by ~~deriving~~ dividing REF aerosol concentration by 10, called CLEAN; and (3) an experiment without (with) aerosol semi-direct effect, called ADEOFF (ADEON).

4.2 Impact of aerosol loads on micro- and macrophysical properties of low-level clouds

660 Figure 10 compares three experiments conducted with enhanced anthropogenic emissions (POL), underestimated aerosol emissions (CLEAN) and background conditions (REF) to provide a quantitative estimation of aerosol loads on radiation and LLCs. In these simulations, both the semi-direct and the indirect effects are taken into account, which act simultaneously on cloud formation and evaporation. The POL case is mostly similar to REF. Before the sunrise, the mean CBH and CTH and cloud presence probability of both cases are almost the same until 08:00 UTC (see Figure 4 and Figure A2a and A2b). After this time, the POL mean CBH is 10 m inferior to the reference reaching 340 m at 10:00 UTC while the mean CTH is mostly 10 superior even reaching 940 m instead of 920 m at 15:00 UTC. The cloud presence probabilities of both cases are also largely the same until the break-up stage. For example, the cloud extends to 670 m thick (from 630 to 1300 m above the ground) at 16 UTC instead of 540 m as in REF. On the other hand, POL and REF have produced clearly different cloud droplet number concentrations alongside mean radius throughout the ~~life-time~~lifetime of modeled clouds (Figure 10a and 10b). At the cloud formation (02:00 UTC), despite having similar liquid water content (LWC) around 0.35 g m^{-3} at 250 m in both cases, N_c^{POL} reaches 333 droplets cm^{-3} and r_c^{POL} 6.45 μm instead of 653 droplets cm^{-3} and 5.1 μm for REF case. This is explained by the differences in the characterization of the aerosols between the two scenarios and the vertical wind speed as Abdul-Razzak and Ghan (2000) include vertical wind speed in their activation scheme. At 02:00 UTC this parameter is less than 0.30 m s^{-1} and allow POL and REF clouds droplets concentration and radius to evolve ~~this way~~under this condition. This trend is reversed ~~the droplets number concentration and radius~~ at 06:00 UTC ~~these numbers~~when the droplets number concentration and radius are equal to 1208 droplets cm^{-3} and 6.43 μm for POL, and 1305 droplets cm^{-3} ~~with a radius of~~and 6.12 μm for REF, respectively. After 08 UTC and until the clouds break-up, N_c^{POL} is superior to N_c^{REF} by reaching a maximum difference of 1425 droplets cm^{-3} at 14:00 UTC. Their respective radius radii are 4.42 μm and 5.18 μm while the liquid water content profiles are quite the same as near 0.47 g m^{-3} at 750 m. These results are in good agreement with the ACPIM parcel model simulation done by Taylor et al. (2019) where CDNC varies in a range of 500–1400 droplets cm^{-3} depending on the inland or offshore (offshore + local emissions) aerosols origin.

685 The difference in cloud macrophysical features such as CBH and CTH between CLEAN and REF ~~are~~is visible though largely limited to a few tens of meters. However, their differences in other macrophysical features including cloud coverage and microphysical features are rather significant. Indeed, from formation to break-up of stratiform clouds, N_c^{CLEAN} is inferior to N_c^{REF} and r_c^{CLEAN} is superior to r_c^{REF} . At 02:00 UTC, N_c^{CLEAN} has a maximum value of 181 droplets cm^{-3} for a radius of 7.58 μm instead of 653 droplets cm^{-3} and 5.1 μm for N_c^{REF} and r_c^{REF} respectively with the same liquid water content value (0.35 g m^{-3}). Between 02:00 UTC and 08:00 UTC, r_c^{CLEAN} increases to reach at the latter time 12.55 μm . After 08:00 UTC, r_c^{CLEAN} decreases slowly showing a maximum value of 10.97 μm at 14:00 UTC. It has to be notified for this time that LWC^{CLEAN} reaches near 0.45 g m^{-3} instead of 0.49 g m^{-3} for LWC^{REF} . This change in the available volume of liquid water with such droplet size explains the decreasing of N_c^{CLEAN} while the aerosol number concentration remains stable around 200–250 cm^{-3} .





Figure 10. Evolution of cloud droplets concentration N_c (top) and cloud droplets radius r_c (middle) with the scenarios given and designated by letter a (REF), b (POL) and c (CLEAN). Bottom panel gives the evolution of mean domain SWRADSURF differences between POL / CLEAN and REF.

As demonstrated from above discussions that modeled cloud microphysical features respond to the variation of aerosol number concentration as expected, i.e., higher aerosol concentration leads to higher cloud droplet number concentration (POL > REF > CLEAN) while smaller mean droplet radius (POL < REF < CLEAN) and hence a higher cloud reflectivity (POL > REF > CLEAN). However, interestingly, as shown in the bottom-panel of Fig. 10c, the response of the incoming solar radiation (SWRADSURF) at ground does not follow always such an expectation. In fact, SWRADSURF appears to be higher in POL than REF from sunrise to 13:00 UTC, and the values in both runs are also clearly higher than that in

710 CLEAN. This tendency is only reversed after 13:00 UTC when solar flux reaches its peak until the break-up stage.



715 **Figure 11.** Liquid water path (LWP, $g\ kg^{-1}\ m$) in POL (left column), REF (mid-column), and CLEAN (right column) **ADEON** runs at 10:00 UTC (top row), 14:00 UTC (middle row), and 16:00 UTC (bottom row).

720 Figure 11 shows that the major reason behind the above-described trend of SWRADSURF is the difference in cloud coverage in competing with the effect brought by different cloud reflectivity of various runs, especially before the noon time when zenith angle is still low high. After sunrise, the cloud top starts to rise and cloud layer becomes thicker. In the meantime, this upward development brings a downward entrainment of dry air from the temperature inversion layer zone above the cloud top and causes evaporation in the cloud. For a cloud with a large quantity of very small droplets as in POL and REF, the evaporation rate of droplets would exceed more rapidly than that in CLEAN case, thus cloud-void or thin cloud layer would form much easier than in the latter case. As shown in Fig. 11 and Table 3, cloud layer
725 in CLEAN is slightly denser than those in POL and REF while cloud-void or thin cloud pixels account a

substantially lower ratio within the domain. ~~When direct solar flux is relatively low~~ Thus, before noontime, cloud reflectivity seems ~~to becoming-become~~ the secondary factor comparing to the cloud-void space in determining the value of SWRADSURF. As a result, SWRADSURF in CLEAN is significantly lower than REF then POL until zenith angle becomes ~~higher-lower~~ closer to noontime. ~~This-The lower SWRADSURF in CLEAN~~ would also have reduced the turbulent mixing as well as delayed the convection. At 14:00 UTC, difference in cloud thickness and cloud-void space still exists while becomes relatively smaller among the three different runs (Fig. 11 and Table 3), cloud reflectivity now becomes the primary reason to cause a different SWRADSURF as shown in Fig. 10 (bottom panel). Interestingly, modeled clouds in POL and REF appear to dissipate earlier and much faster than in CLEAN in the break-up stage (Fig. 11, bottom panel), ~~the very low number of cloud pixels with different thickness in the domain seems having brought some variations in SWRADSURF.~~

| | LWP 10 UTC | PCP 10 UTC | LWP 14 UTC | PCP 14 UTC | LWP 16 UTC | PRP 16 UTC |
|-------|------------|------------|------------|------------|------------|------------|
| POL | 14.87 | 12.79 | 10.98 | 42.17 | 1.96 | 99.66 |
| REF | 15.67 | 10.11 | 11.34 | 42.69 | 2.74 | 99.67 |
| CLEAN | 16.98 | 6.95 | 11.79 | 44.93 | 4.12 | 94.47 |

Table 3. Domain averaged liquid water path (LWP; $g\ kg^{-1}\ m$) and ~~(b)~~ poor-cloud pixel percentage (PCP, defined by the percentage of pixels where LWP < $10\ g\ kg^{-1}\ m$; percentage) in three different runs.

To summarize, as expected, aerosol concentration is a major factor in controlling the cloud microphysical features by changing the simulated droplet number concentration and radius for clouds with similar liquid water content. However, despite this well-known Twomey effect the incoming solar radiation at ground did not decrease due to additional CDNC. Instead, cloud macrophysical features and in particular cloud-void space caused by dry entrainment from inverse layer above the cloud is the dominant factor impacting the incoming solar radiation at ground. Cloud macrophysical properties determine the break-up speed of modeled clouds and therefore, the life cycle of the modeled LLSCs. It is worth indicating though, another factor that might contribute to the cloud life cycle, i.e., the atmospheric heating caused by the semi-direct optical effect of absorbing aerosol component such as black carbon has not been analyzed up to this moment and will be discussed in the following section.

4.3 Impact of aerosol semi-direct effect on low-level clouds

The semi-direct effect of aerosols on LLSCs that represents the modifications of the LLSCs properties and atmospheric dynamics due to absorption of SW radiation by absorbing aerosol, ~~This effect~~ has been estimated ~~here~~ by conducting three additional experiments constructed accordingly ~~as-in the~~ same ~~way~~ as the original experiments (hereafter ADEON including REF, POL, and CLEAN) but excluding aerosol direct effects (named ADEOFF) and then comparing the results between each of the paired runs. ~~Indeed~~ ~~Apparently~~, BC is the major species behind the semi-direct effect in our case study, ~~as proved by scenario REF_NOBC as seen in Figure A2c. This scenarios retakes the characteristics of REF in term of size distribution and total mass but replace BC mass part by a equivalent mix of sulfate,~~

ammonium, nitrate and organic carbon. By running ADEON and ADEOFF configurations for REF_NOBC and comparing cloud evolution and cloud presence probability, it can be seen that there is few minor differences between these two configurations.





Figure 12. Evolution of the difference of the mean CBH and CTH (a), SWHR (b) and TKE (c) between the simulation runs with and without aerosol direct effect (ADEON-ADEOFF) for REF, POL and CLEAN.



770 **Figure 13.** Mean difference surface SW radiative flux (SWRADSURF) between Clear-Sky (CS) and cloudy scenarios giving
 775 the flux dissipated by clouds in ADEON (a) and ADEOFF (b) configurations. SWRADSURF difference between ADEON
 and ADEOFF configuration for the three scenarios (c).

The changes in ~~cloud-vertical-dispersion~~ cloud top and base, SWHR and TKE due to aerosol
 775 absorption and potential feedbacks are shown in Figure 12. The results demonstrates that light-absorbing
 BC aerosols can cause a substantial atmospheric heating accompanied by a substantial warming tendency

near LLSCs-top (Figure 12b). The domain averaged heating due to BC aerosols (SWHR difference) in CLEAN case is rather insignificant in comparison with the two other cases as 1.30 K day^{-1} (and a cooling -2.25 K day^{-1} above due to cloud top change) at 14:00 UTC, whereas it reaches 12.16 K day^{-1} ($-13.14 \text{ K day}^{-1}$) for POL, 7.71 K day^{-1} (-9.24 K day^{-1}) in REF, respectively. Accordingly, in ADEON runs, more water vapor tends to condense onto cloud droplets under the higher relative humidity in the lower PBL and decreasing turbulent mixing (Figure 12c, with a maximum decreasing of $-0.18 \text{ m}^2 \text{ s}^{-2}$ for POL) lead to a decrease of the cloud top height, limiting entrainment, and also reduced SW reflection due to BC in-cloud absorption. The cloud top height reduction in two polluted cases POL and REF is quite significant as shown in Figure 12a, where CTH in POL and REF has decreased by up to 100 and 70 meters due to the presence of BC, respectively. On the other hand, CBH is also increased about 20 meters in both cases before break-up. In comparison, CTH and CBH appear to be less affected in CLEAN run due to its low BC content. Before break-up, in-cloud TKE below the heating layer has been reduced in some extent (Fig.12c). On the other hand, due to a lower cloud top in polluted cases, boundary layer top with active turbulent exchange would also be lowered. The effect of BC absorption in lowering modeled clouds in POL and REF (implying a reduced upward development) is likely another factor to slow down their break-up as discussed before.

We find that the semi-direct effect can manifest both an enhancing and a weakening contribution to the (negative) indirect radiative forcing (Lohmann and Feichter, 2001; Koch and Del Genio, 2010a; Huang *et al.*, 2014; Yamaguchi *et al.*, 2015; Stjern *et al.*, 2017; Kreidenweis *et al.*, 2019). At 14:00 UTC, the flux difference between ADEON and ADEOFF at ground reaches -33 W m^{-2} and -75 W m^{-2} for REF and POL, respectively (Fig. 13c). This is explained by a decreased void space in ADEON runs that allows less solar radiations-irradiance to attain the surface despite the cloud layer being thinner (Fig. 42e-13c and A3), hence the semi-direct effect contributes positively to the enhancement of (negative) indirect radiative forcing in this case. But at 16:00 UTC the flux difference between ADEON and ADEOFF becomes positive with values for REF and POL as 32 W m^{-2} and 66 W m^{-2} , respectively. As the clouds break-up slower in ADEOFF during this stage due to being thicker-clouds (Fig. A3), more clouds inside the domain with increased thickness causes less-weaker SW radiations-irradiance reaching ground. In this case, the semi-direct effect weakens the semidirect radiative forcing.

The above results have demonstrated the important role of solar absorption by aerosols in determining the life cycle of LLSCs. The atmospheric heating by light absorbing BC would limit the elevation of cloud layertop, especially during the break-up stage (Koch and Del Genio, 2010b; Zhang and Zuidema, 2019). Such heating can also decrease cloud-void space then delay break-up until late afternoon, especially for clouds with higher cloud droplet number concentration in polluted environment such as in POL and REF runs. This study case also exhibits either a positive (e.g., decreasing cloud-void space) or a negative (e.g., accelerating break-up in late afternoon due to a thinner cloud) contribution of the semi-direct effect to the indirect radiative forcing.

5. Conclusions

A characteristic case of LLSCs over SWA have-has been simulated with Meso-NH model in high-resolution Large-Eddy Sim-ulationSimulation configuration constrained by DACCIWA measurements. The model has successfully reproduced observed life cycle alongside key macro and microphysical

features as well as surface radiative and heat fluxes. To determine the impact of aerosols on the LLSCs diurnal cycle, sensitivity simulations using several different aerosol profiles have also been conducted. 820 These aerosol profiles contain different number concentrations and chemical compositions in order to reflect the situations associated with various aerosol populations encountered during the campaign.

The results from various sensitivity simulations suggest that both aerosol concentration and chemical composition can effectively influence the LLSCs life cycle. The impact of the aerosol concentration, as reflected from a comparison among simulations using aerosol profiles with different number 825 concentrations, is initiated from resultant cloud microphysical features in particular the cloud droplet number concentration and mean droplet size. Such a difference created by different aerosol number concentration also affect cloud reflectivity as expected. Interestingly, we have found that the difference in cloud reflectivity caused by different aerosol concentration does not always dominate the surface incoming solar radiation and thus cloud development after sunrise due to another competing factor: the 830 cloud-void space caused by the air ~~entrainment-entrained~~ from the inversion layer above cloud top dominates the variation of surface incoming solar radiation before noontime. Clouds influenced by higher aerosol concentrations and thus higher cloud droplet number concentration ~~while-and~~ smaller droplet sizes are found to evaporate more easily and thus impose more cloud-void spaces. For the same reason, clouds with higher droplet concentration are likely to break-up earlier.

In addition, our sensitivity runs including versus excluding aerosol direct radiative effects have also 835 demonstrated the impact specifically of solar absorption by black carbon on the cloud life cycle. The excessive atmospheric heating ~~by black carbon up-reaches-reaching~~ 12 K day^{-1} in our modeled cases ~~is found to by black carbon can~~ lower the cloud top height and reduce dry entrainment. Working with above-indicated aerosol concentration effect this ~~heating and its consequences~~ might delay break-up until late 840 afternoon, while beyond that modeled clouds in polluted cases with higher aerosol concentrations and BC included would break-up faster due to thinner cloud layer. Therefore, semi-direct effect can contribute positively to the indirect radiative forcing (negative in quantity) due to decreased cloud-void space, or negatively by causing thinner cloud layer and thus a faster cloud break-up in late afternoon, all depending on the phase in stratiform cloud diurnal cycle.

Our study has demonstrated that the life cycle and thus the radiative forcing of LLSCs over land 845 area of SWA can be substantially influenced by aerosols from both long-range transported biomass burning plumes and from local urban emissions. In fact, more aerosol profiles had been collected during the DACCIWA campaign besides the ones used in this study. Future research works could reveal the aerosol impact under an even broader range of aerosol properties and to examine the 850 temporal variations of LLSCs radiative effects evolved with different large-scale meteorological conditions with different associated air mass. More analysis on different cloud cases in SWA would also be able to assess or refute current results on semi-direct effect.

855
Code and data availability. The data obtained during the DACCIWA campaign at the Savè supersite ~~alongside all other data used in this study~~ are publicly available on the SEDOO database

(~~<http://baobab.sedoo.fr/DACCIWA/>~~)(~~<http://baobab.sedoo.fr/DACCIWA/>~~). The Meso-NH code is maintained and updated by LAERO and CNRM, it is freely available for download at <http://mesonh.aero.obs-mip.fr/mesonh52/>. ~~All data used in this study are publicly available on the AERIS Data and Service Center, which can be found at <http://baobab.sedoo.fr/DACCIWA/>.~~

Author contributions. LD and CW designed the simulations and LD conducted model simulations and data analyses. LD and CW wrote this paper with contribution from all co-authors. CW advised and helped to better understand the different aspects of this research work. PT advised and trained LD for Meso-NH and ORILAM module use. CD processed and provided the aerosol profiles used in previous simulations and NM was part of this work. MZ helped to select the study case and advised during the study case construction and analysis. AD brought a critical eye to this work.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study is supported by L'Agence National de la Recherche (ANR) of France under "Programme d'Investissements d'Avenir" (ANR-18-MPGA-003 EUROACE) and co-funded by University Toulouse III Paul Sabatier. The computation of this work was performed using HPC resources of French GENCI-IDRIS (Grant A0110110967 and A0090110967) and French Regional Computations center CALMIP. ~~The first author LD thanks the Laboratoire d'Aérodynamique, Université de Toulouse, France, for funding and hosting the his Ph.D. research activities. A special thank to all people whose work was involved in the measurement and processing of DACCIWA campaign data especially over the Savè supersite. First Author LD also thanks the MesoNH team, especially Quentin Rodier-, Juan Escobar, and Philippe Wautelet from CNRM, France, for its-their advises on using Meso-NH use-, Benoit Vié and Marie Mazoyer (CNRM) are thanked for their help to handle and modify microphysical scheme LIMA-, A thank to Quentin Libois (CNRM) for bringing its understanding of explaining the details of Meso-NH's radiative schemes. Many thanks for all co-authors for their help and time given to this article-, and a special thanks specifically to Fabienne Lohou (LAERO) for her introduction of DACCIWA campaign alongside her guidance in using relevant data products. A special thanks the authors to all people whose work was involved in the measurement and processing of DACCIWA campaign data especially over the Savè supersite. Many constructive comments and suggestions from Dr. Mónica Zamora Zapata and an anonymous reviewer have made a substantial impact on our effort to improve the quality of the manuscript.~~

Appendix A








Figure A1. Mass composition (a,c) and size distribution provided by (Denjean et al., 2020a) and fitted into 2 modes described in Table 2 (b,d) for scenarios POL (top), CLEAN (bottom).

900





905 **Figure A2.** Mean LLSCs deck evolution of POL (a) and CLEAN (b) cases with the representation of REF's one to make comparison and REF_NOBC_ADEON and ADEOFF runs (c), vertical colored bars for POL/CLEAN (left) and REF (right) attribute at each altitude level a cloud presence density for both cases at each hour.



915 **Figure A3.** Liquid water path (LWP, $g\ kg^{-1}\ m$) in POL (left column), REF (mid-column), and CLEAN (right column) ADEOFF runs at 10:00 UTC (top row), 14:00 UTC (middle row), and 16:00 UTC (bottom row).

References

- Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types, *Journal of Geophysical Research: J. Geophys. Res.-Atmospheres*, 105, 6837–6844, <https://doi.org/https://doi.org/10.1029/1999JD901161>, 2000.
- 920 Abdul-Razzak, H. and Ghan, S. J.: Parameterization of the influence of organic surfactants on aerosol activation, *J. Geophys. Res.-AtmosJournal of Geophysical Research: Atmospheres*, 109, <https://doi.org/https://doi.org/10.1029/2003JD004043>, 2004.
- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03174>, 2004.
- 925 Adler, B., Babic, K., Kalthoff, N., Lohou, F., Lothon, M., Dione, C., Pedruzo-Bagazgoitia, X., and Andersen, H.: Nocturnal low-level clouds in the atmospheric boundary layer over southern West Africa: an observation-based analysis of conditions and processes, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 19, 663–681, <https://doi.org/10.5194/acp-19-663-2019>, 2019.
- 930 Aouizerats, B., Thouron, O., Tulet, P., Mallet, M., Gomes, L., and Henzing, J. S.: Development of an online radiative module for the computation of aerosol optical properties in 3-D atmospheric models: validation during the EUCAARI campaign, *Geoscientific Model Development*, 3, 553–564, <https://doi.org/10.5194/gmd-3-553-2010>, 2010.
- Babic, K., Adler, B., Kalthoff, N., Andersen, H., Dione, C., Lohou, F., Lothon, M., and Pedruzo-Bagazgoitia, X.: The observed diurnal cycle of low-level stratus clouds over southern West Africa: a case study, *Atmos. Chem. Phys Atmospheric Chemistry and Physics*, 19, 1281–1299, <https://doi.org/10.5194/acp-19-1281-2019>, 2019.
- 935 Bauer, S. E., Im, U., Mezuman, K., and Gao, C. Y.: Desert Dust, Industrialization, and Agricultural Fires: Health Impacts of Outdoor Air Pollution in Africa, *J. Geophys Res.-AtmosJournal of Geophysical Research: Atmospheres*, 124, 4104–4120, <https://doi.org/https://doi.org/10.1029/2018JD029336>, 2019.
- Bellon, G. and Stevens, B.: Time Scales of the Trade Wind Boundary Layer Adjustment, *Journal of the Atmospheric Sci. enees*, 70, 1071 – 1083, <https://doi.org/10.1175/JAS-D-12-0219.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2013.
- [Boucher, O., D. Randall, P. Artaxo, C. Bretherton, G. Feingold, P. Forster, V.-M. Kerminen, Y. Kondo, H. Liao, U. Lohmann, P. Rasch, S.K. Satheesh, S. Sherwood, B. Stevens and X.Y. Zhang: Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change \[Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley \(eds.\)\]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.](#)
- 945 Brito, J., Freney, E., Dominutti, P., Borbon, A., Haslett, S. L., Batenburg, A. M., Colomb, A., Dupuy, R., Denjean, C., Burnet, F., Bourriane, T., Deroubaix, A., Sellegri, K., Borrmann, S., Coe, H., Flamant, C., Knippertz, P., and Schwarzenboeck, A.: Assessing the role of anthropogenic and biogenic sources on PM₁ over southern West Africa using aircraft measurements, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 18, 757–772, <https://doi.org/10.5194/acp-18-757-2018>, 2018.
- [Brooks, B., Catoire, V., Knippertz, P., and Coe, H.: Aerosol influences on low level clouds in the West African monsoon, Atmospheric Chemistry and Physics](#), 19, 8503–8522, <https://doi.org/10.5194/acp-19-8503-2019>, 2019.
- 955 Caniaux, G., Redelsperger, J.-L., and Lafore, J.-P.: A Numerical Study of the Stratiform Region of a Fast-Moving Squall Line. Part I: General Description and Water and Heat Budgets, *Journal of Atmospheric Sci. enees*, 51, 2046 – 2074, [https://doi.org/10.1175/15200469\(1994\)051<2046:ANSOTS>2.0.CO;2](https://doi.org/10.1175/15200469(1994)051<2046:ANSOTS>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 1994.
- 960 Carslaw, K. S., Gordon, H., Hamilton, D. S., Johnson, J. S., Regayre, L. A., Yoshioka, M., and Pringle, K. J.: Aerosols in the Pre-industrial Atmosphere, *Current Climate Change Reports*, 3, 1–15, <https://doi.org/10.1007/s40641-017-0061-2>, 2017.
- Chatfield, R. B., Vastano, J. A., Li, L., Sachse, G. W., and Connors, V. S.: The Great African Plume from biomass burning: Generalizations from a three-dimensional study of TRACE A carbon monoxide, *J. Geophys Res.-AtmosJournal of Geophysical Research: Atmospheres*, 103, 28059–28077, <https://doi.org/https://doi.org/10.1029/97JD03363>, 1998.

- 965 Chen, T., Rossow, W. B., and Zhang, Y.: Radiative Effects of Cloud-Type Variations, *Journal of Climate*, 13, 264 – 286, [https://doi.org/10.1175/1520-0442\(2000\)013<0264:REOCTV>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0264:REOCTV>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2000.
- Cleveland, W. S.: Robust Locally Weighted Regression and Smoothing Scatterplots, *Journal of the American Statistical Association*, 74, 829–836, <https://doi.org/10.1080/01621459.1979.10481038>, 1979.
- 970 Cohard, J.-M. and Pinty, J.-P.: A comprehensive two-moment warm microphysical bulk scheme. I: Description and tests, *Quarterly Journal of the Royal Meteorological Society*, 126, 1815–1842, <https://doi.org/https://doi.org/10.1002/qj.49712656613>, 2000.
- Dearden, C., Hill, A., Coe, H., and Choularton, T.: The role of droplet sedimentation in the evolution of low-level clouds over southern West Africa, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 18, 14253–14269, <https://doi.org/10.5194/acp-18-14253-2018>, 2018.
- 975 Deetz, K., Vogel, H., Knippertz, P., Adler, B., Taylor, J., Coe, H., Bower, K., Haslett, S., Flynn, M., Dorsey, J., Crawford, I., Kottmeier, C., and Vogel, B.: Numerical simulations of aerosol radiative effects and their impact on clouds and atmospheric dynamics over southern West Africa, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 18, 9767–9788, <https://doi.org/10.5194/acp-18-9767-2018>, 2018.
- 980 Denjean, C., Bourriane, T., Burnet, F., Mallet, M., Maury, N., Colomb, A., Dominutti, P., Brito, J., Dupuy, R., Sellegri, K., Schwarzenboeck, A., Flamant, C., and Knippertz, P.: Overview of aerosol optical properties over southern West Africa from DACCWA aircraft measurements, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 20, 4735–4756, <https://doi.org/10.5194/acp-20-4735-2020>, 2020a.
- Denjean, C., Brito, J., Libois, Q., Mallet, M., Bourriane, T., Burnet, F., Dupuy, R., Flamant, C., and Knippertz, P.: Unexpected Biomass Burning Aerosol Absorption Enhancement Explained by Black Carbon Mixing State, *Geophysical Research Letters*, 47, e2020GL089055, <https://doi.org/https://doi.org/10.1029/2020GL089055>, e2020GL089055 2020GL089055, 2020b.
- Deroubaix, A., Menut, L., Flamant, C., Brito, J., Denjean, C., Dreiling, V., Fink, A., Jambert, C., Kalthoff, N., Knippertz, P., Ladkin, R., Mailler, S., Maranan, M., Pacifico, F., Piguet, B., Siour, G., and Turquety, S.: Diurnal cycle of coastal anthropogenic pollutant transport over southern West Africa during the DACCWA campaign, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 19, 473–497, <https://doi.org/10.5194/acp-19-473-2019>, 2019.
- 990 Deroubaix, A., Menut, L., Flamant, C., Knippertz, P., Fink, A. H., Batenburg, A., Brito, J., Denjean, C., Dione, C., Dupuy, R., Hahn, V., Kalthoff, N., Lohou, F., Schwarzenboeck, A., Siour, G., Tuccella, P., and Voigt, C.: Sensitivity of low-level clouds and precipitation to anthropogenic aerosol emission in southern West Africa: a DACCWA case study, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 22, 3251–3273, <https://doi.org/10.5194/acp-22-3251-2022>, 2022.
- 995 Derrien, S., Bezombes, Y., Bret, B., Gabella, O., Jarnot, C., Medina, P., Piques, E., Delon, C., Dione, C., Campistron, B., Durand, P., Jambert, C., Lohou, F., Lothon, M., Pacifico, F., and Meyerfeld, Y.: DACCWA field campaign, Savè super-site, UPS instrumentation, 10.6096/DACCWA.1618, 2016.
- 1000 Dione, C., Lohou, F., Lothon, M., Adler, B., Babic, K., Kalthoff, N., Pedruzo-Bagazgoitia, X., Bezombes, Y., and Gabella, O.: Low-level stratiform clouds and dynamical features observed within the southern West African monsoon, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 19, 8979–8997, <https://doi.org/10.5194/acp-19-8979-2019>, 2019.
- Eastman, R. and Warren, S. G.: Diurnal Cycles of Cumulus, Cumulonimbus, Stratus, Stratocumulus, and Fog from Surface Observations over Land and Ocean, *Journal of Climate*, 27, 2386 – 2404, <https://doi.org/10.1175/JCLI-D-13-00352.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2014.
- 1005 Flamant, C., Deroubaix, A., Chazette, P., Brito, J., Gaetani, M., Knippertz, P., Fink, A. H., de Coetlogon, G., Menut, L., Colomb, A., Denjean, C., Meynadier, R., Rosenberg, P., Dupuy, R., Dominutti, P., Duplissy, J., Bourriane, T., Schwarzenboeck, A., Ramonet, M., and Totems, J.: Aerosol distribution in the northern Gulf of Guinea: local anthropogenic sources, long-range transport, and the role of coastal shallow circulations, *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 18, 12363–12389, <https://doi.org/10.5194/acp-18-12363-2018>, 2018.
- 1010 Flossmann, A. I. and Wobrock, W.: Cloud Processing of Aerosol Particles in Marine Stratocumulus Clouds, *Atmosphere*, 10, <https://doi.org/10.3390/atmos10090520>, 2019.

- Fouquart, Y. and Bonnel, B.: Computations of solar heating of the earth's atmosphere – A new parameterization, *Beitrag zur Physik der Atmosphäre*, 53, 35-62, pp-35-62, 1980.
- 1015 Geoffroy, O., Brenguier, J.-L., and Sandu, I.: Relationship between drizzle rate, liquid water path and droplet concentration at the scale of a stratocumulus cloud system, *Atmospheric Chemistry and Physics*, 8, 4641–4654, <https://doi.org/10.5194/acp-8-4641-2008>, 2008.
- Ghonima, M., T. Heus, J. R. Norris, and J. Kleissl, 2016: Factors controlling stratocumulus cloud lifetime over coastal land, *J. Atmos. Sci.*, 73, 2961-2983.
- 1020 Griffin, R. J., Nguyen, K., Dabdub, D., and Seinfeld, J. H.: A Coupled Hydrophobic-Hydrophilic Model for Predicting Secondary Organic Aerosol Formation, *Journal of Atmospheric Chemistry*, 44, 171–190, <https://doi.org/10.1023/A:1022436813699>, 2003.
- Hagan, H. D., Thompson, B., Palmo, J., and Ruskowski, A.: py-smps, <https://github.com/quant-aq/py-smps>, 2022.
- 1025 Handwerker, J., Scheer, S., and Gamer, T.: DACCIWA field campaign, Savè super-site, Cloud and precipitation, <https://doi.org/10.6096/dacciwa.1686>, 2016.
- Hannak, L., Knippertz, P., Fink, A. H., Kniffka, A., and Pante, G.: Why Do Global Climate Models Struggle to Represent Low-Level Clouds in the West African Summer Monsoon?, *Journal of Climate*, 30, 1665 – 1687, <https://doi.org/10.1175/JCLI-D-16-0451.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2017.
- 1030 Hansen, J., M. Sato, R. Ruedy, A. Lacis, and V. Oinas, Global warming in the twenty-first century: An alternative scenario. *PNAS*, 97, 9875-9880, 1998.
- Hartmann, D. L., Ockert-Bell, M. E., and Michelsen, M. L.: The Effect of Cloud Type on Earth's Energy Balance: Global Analysis, *Journal of Climate*, 5, 1281 – 1304, [https://doi.org/10.1175/1520-0442\(1992\)005<1281:TEOCTO>2.0.CO;2](https://doi.org/10.1175/1520-0442(1992)005<1281:TEOCTO>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 1992.
- 1035 Haslett, S. L., Taylor, J. W., Evans, M., Morris, E., Vogel, B., Dajuma, A., Brito, J., Batenburg, A. M., Borrmann, S., Schneider, J., Schulz, C., Denjean, C., Bourriane, T., Knippertz, P., Dupuy, R., Schwarzenböck, A., Sauer, D., Flamant, C., Dorsey, J., Crawford, I., and Coe, H.: Remote biomass burning dominates southern West African air pollution during the monsoon, *Atmospheric Chemistry and Physics*, 19, 15217–15234, <https://doi.org/10.5194/acp-19-15217-2019>, 2019.
- 1040 Haywood, J. and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review, *Reviews of Geophysics*, 38, 513–543, <https://doi.org/10.1029/1999RG000078>, 2000.
- Hill, P. G., Allan, R. P., Chiu, J. C., Bodas-Salcedo, A., and Knippertz, P.: Quantifying the Contribution of Different Cloud Types to the Radiation Budget in Southern West Africa, *Journal of Climate*, 31, 5273 – 5291, <https://doi.org/10.1175/JCLI-D-17-0586.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2018.
- 1045 Huang, J., Wang, T., Wang, W., Li, Z., and Yan, H.: Climate effects of dust aerosols over East Asian arid and semiarid regions, *J. Geophys. Res.-Atmospheric Research: Atmospheres*, 119, 11,398–11,416, <https://doi.org/10.1002/2014JD021796>, 2014.
- 1050 Jiang, G.-S. and Shu, C.-W.: Efficient Implementation of Weighted ENO Schemes, *Journal of Computational Physics*, 126, 202–228, <https://doi.org/10.1006/jcph.1996.0130>, 1996.
- Kalthoff, N., Lohou, F., Brooks, B., Jegede, G., Adler, B., Babic, K., Dione, C., Ajao, A., Amekudzi, L. K., Aryee, J. N. A., Ay-’oola, M., Bessardon, G., Danuor, S. K., Handwerker, J., Kohler, M., Lathon, M., Pedruzo-Bagazgoitia, X., Smith, V., Sunmonu, L., Wieser, A., Fink, A. H., and Knippertz, P.: An overview of the diurnal cycle of the atmospheric boundary layer during the West African monsoon season: results from the 2016 observational campaign, *Atmospheric Chemistry and Physics*, 18, 2913–2928, <https://doi.org/10.5194/acp-18-2913-2018>, 2018.
- 1055 Khairoutdinov, M. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Monthly Weather Review*, 128, 229 – 243, [https://doi.org/10.1175/1520-0493\(2000\)128<0229:ANCPPI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0229:ANCPPI>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2000.

- Knippertz, P., Fink, A. H., Schuster, R., Trentmann, J., Schrage, J. M., and Yorke, C.: Ultra-low clouds over the southern West African monsoon region, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL049278>, 2011.
- Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Liousse, C., Mari, C., Allan, R. P., Brooks, B., Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCIWA Project: Dynamics–Aerosol–Chemistry–Cloud Interactions in West Africa, *Bull. Am. Meteor. Soc.*, 96, 1451–1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2015.
- Knippertz, P., Fink, A. H., Deroubaix, A., Morris, E., Tocquer, F., Evans, M. J., Flamant, C., Gaetani, M., Lavaysse, C., Mari, C., Marsham, J. H., Meynadier, R., Affo-Dogo, A., Bahaga, T., Brosse, F., Deetz, K., Guebsi, R., Latifou, I., Maranan, M., Rosenberg, P. D., and Schlueter, A.: A meteorological and chemical overview of the DACCIWA field campaign in West Africa in June–July 2016, *Atmos. Chem. Phys.*, 17, 10893–10918, <https://doi.org/10.5194/acp-17-10893-2017>, 2017.
- Koch, D. and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and synthesis, *Atmos. Chem. Phys.*, 10, 7685–7696, <https://doi.org/10.5194/acp-10-7685-2010>, 2010a.
- Koch, D. and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and synthesis, *Atmos. Chem. Phys.*, 10, 7685–7696, <https://doi.org/10.5194/acp-10-7685-2010>, 2010b.
- Kreidenweis, S. M., Petters, M., and Lohmann, U.: 100 Years of Progress in Cloud Physics, Aerosols, and Aerosol Chemistry Research, *Meteorological Monographs*, 59, 11.1–11.72, <https://doi.org/10.1175/AMSMONOGRAPHSD-18-0024.1>, 2019.
- Lac, C., Chaboureau, J.-P., Masson, V., Pinty, J.-P., Tulet, P., Escobar, J., Leriche, M., Barthe, C., Aouizerats, B., Augros, C., Aumond, P., Auguste, F., Bechtold, P., Berthet, S., Bielli, S., Bosseur, F., Caumont, O., Cohard, J.-M., Colin, J., Couvreur, F., Cuxart, J., Delautier, G., Dauhut, T., Ducrocq, V., Filippi, J.-B., Gazen, D., Geoffroy, O., Gheusi, F., Honnert, R., Lafore, J.-P., Lebeau-pin Brossier, C., Libois, Q., Lunet, T., Mari, C., Maric, T., Mascart, P., Mogé, M., Molinié, G., Nuissier, O., Pantillon, F., Peyrillé, P., Pergaud, J., Perraud, E., Pianezze, J., Redelsperger, J.-L., Ricard, D., Richard, E., Riette, S., Rodier, Q., Schoetter, R., Seyfried, L., Stein, J., Suhre, K., Taufour, M., Thouron, O., Turner, S., Verrelle, A., Vié, B., Visentin, F., Vionnet, V., and Wautelet, P.: Overview of the Meso-NH model version 5.4 and its applications, *Geoscientific Model Development*, 11, 1929–1969, <https://doi.org/10.5194/gmd-11-1929-2018>, 2018.
- Lascaux, F., Richard, E., and Pinty, J.-P.: Numerical simulations of three different MAP IOPs and the associated microphysical processes, *Quart. J. Roy. Meteor. Soc.*, 132, 1907–1926, <https://doi.org/10.1256/qj.05.197>, 2006.
- Legain, D., Bousquet, O., Douffet, T., Tzanos, D., Moulin, E., Barrie, J., and Renard, J.-B.: High-frequency boundary layer profiling with reusable radiosondes, *Atmospheric Measurement Techniques*, 6, 2195–2205, <https://doi.org/10.5194/amt-6-2195-2013>, 2013.
- Li, J., Carlson, B. E., Yung, Y. L., Lv, D., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, V., Kahn, R. A., Zhang, P., Dubovik, O., Ding, A., Laci, A. A., Zhang, L., and Dong, Y.: Scattering and absorbing aerosols in the climate system, *Nature Reviews Earth & Environment*, 3, 363–379, <https://doi.org/10.1038/s43017-022-00296-7>, 2022.
- Liousse, C., Assamoi, E., Criqui, P., Granier, C., and Rosset, R.: Explosive growth in African combustion emissions from 2005 to 2030, *Environ. Res. Lett.*, 9, 035003, <https://doi.org/10.1088/1748-9326/9/3/035003>, ~~publisher: IOP Publishing~~, 2014.
- Liu, Y., Jia, R., Dai, T., Xie, Y., and Shi, G.: A review of aerosol optical properties and radiative effects, *Journal of Meteorological Research*, 28, 1003–1028, <https://doi.org/10.1007/s13351-014-4045-z>, 2014.
- Lohmann, U. and Feichter, J.: Can the direct and semi-direct aerosol effect compete with the indirect effect on a global scale?, *Geophysical Research Letters*, 28, 159–161, <https://doi.org/10.1029/2000GL012051>, 2001.
- Lohou, F., Kalthoff, N., Adler, B., Babic, K., Dione, C., Lohon, M., Pedruzo-Bagazgoitia, X., and Zouzoua, M.: Conceptual model of diurnal cycle of low-level stratiform clouds over southern West Africa, *Atmos. Chem. Phys.*, 20, 2263–2275, <https://doi.org/10.5194/acp-20-2263-2020>, 2020.

- 110 Mari, C., Evans, M. J., Palmer, P. I., Jacob, D. J., and Sachse, G. W.: Export of Asian pollution during two cold front episodes of the TRACE-P experiment, *J. Geophys. Res.-Atmos.*~~Journal of Geophysical Research: Atmospheres~~, 109, <https://doi.org/https://doi.org/10.1029/2003JD004307>, 2004.
- Mari, C. H., Cailley, G., Corre, L., Saunois, M., Attié, J. L., Thouret, V., and Stohl, A.: Tracing biomass burning plumes from the Southern Hemisphere during the AMMA 2006 wet season experiment, *Atmos. Chem. Phys.*~~Atmospheric Chemistry and Physics~~, 8, 3951–3961, <https://doi.org/10.5194/acp-8-3951-2008>, 2008.
- 115 Marticorena, B. and Bergametti, G.: Two-year simulations of seasonal and interannual changes of the Saharan dust emissions, *Geophys. Res. Lett.*~~Geophysical Research Letters~~, 23, 1921–1924, <https://doi.org/https://doi.org/10.1029/96GL01432>, 1996.
- Martin, G. M., Johnson, D. W., and Spice, A.: The Measurement and Parameterization of Effective Radius of Droplets in Warm Stratocumulus Clouds, *Journal of Atmospheric Sciences*, 51, 1823 – 1842, [https://doi.org/10.1175/15200469\(1994\)051<1823:TMAPOE>2.0.CO;2](https://doi.org/10.1175/15200469(1994)051<1823:TMAPOE>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 1994.
- 120 Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essauouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Scity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geoscientific Model Development*, 6, 929–960, <https://doi.org/10.5194/gmd-6-929-2013>, 2013.
- Menut, L., Flamant, C., Turquety, S., Deroubaix, A., Chazette, P., and Meynadier, R.: Impact of biomass burning on pollutant surface concentrations in megacities of the Gulf of Guinea, *Atmos. Chem. Phys.*~~Atmospheric Chemistry and Physics~~, 18, 2687–2707, <https://doi.org/10.5194/acp18-2687-2018>, 2018.
- 130 Menut, L., Tuccella, P., Flamant, C., Deroubaix, A., and Gaetani, M.: The role of aerosol–radiation–cloud interactions in linking anthropogenic pollution over southern west Africa and dust emission over the Sahara, *Atmos. Chem. Phys.*~~Atmospheric Chemistry and Physics~~, 19, 14657–14676, <https://doi.org/10.5194/acp-19-14657-2019>, 2019.
- Metzger, S., Dentener, F., Pandis, S., and Lelieveld, J.: Gas/aerosol partitioning: 1. A computationally efficient model, *J. Geophys. Res.-Atmos.*~~Journal of Geophysical Research: Atmospheres~~, 107, ACH 16–1–ACH 16–24, <https://doi.org/https://doi.org/10.1029/2001JD001102>, 2002.
- 135 Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.-Atmos.*~~Journal of Geophysical Research: Atmospheres~~, 102, 16663–16682, <https://doi.org/https://doi.org/10.1029/97JD00237>, 1997.
- 140 Morcrette, J.-J.: The Surface Downward Longwave Radiation in the ECMWF Forecast System, *Journal of Climate*, 15, 1875 – 1892, [https://doi.org/10.1175/1520-0442\(2002\)015<1875:TSDLRI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1875:TSDLRI>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2002.
- Murphy, J. G., Oram, D. E., and Reeves, C. E.: Measurements of volatile organic compounds over West Africa, *Atmos. Chem. Phys.*~~Atmospheric Chemistry and Physics~~, 10, 5281–5294, <https://doi.org/10.5194/acp-10-5281-2010>, 2010.
- 145 Noilhan, J. and Planton, S.: A Simple Parameterization of Land Surface Processes for Meteorological Models, *Monthly Weather Review*~~Rev.~~, 117, 536 – 549, [https://doi.org/10.1175/1520-0493\(1989\)117<0536:ASPOLS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<0536:ASPOLS>2.0.CO;2), ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 1989.
- Pedruzo-Bagazgoitia, X., de Roode, S. R., Adler, B., Babic, K., Dione, C., Kalthoff, N., Lohou, F., Lothon, M., and Vilà-Guerau de Arellano, J.: The diurnal stratocumulus-to-cumulus transition over land in southern West Africa, *Atmos. Chem. Phys.*~~Atmospheric Chemistry and Physics~~, 20, 2735–2754, <https://doi.org/10.5194/acp-20-2735-2020>, 2020.
- 150 Pinty, J.-P. and Jabouille, P.: A mixed-phased cloud parameterization for use in a mesoscale non-hydrostatic model: simulations of a squall line and of orographic precipitation, 1998.
- Pruppacher, H. R., Klett, J. D., and Wang, P. K.: Microphysics of Clouds and Precipitation, *Aerosol Science and Technology*, 28, 381–382, <https://doi.org/10.1080/02786829808965531>, ~~publisher: Taylor & Francis, eprint: https://doi.org/10.1080/02786829808965531~~, 1998.
- 155 Reeves, C. E., Formenti, P., Afif, C., Ancellet, G., Attié, J.-L., Bechara, J., Borbon, A., Cairo, F., Coe, H., Crumeyrolle, S., Fierli, F., Flamant, C., Gomes, L., Hamburger, T., Jambert, C., Law, K. S., Mari, C., Jones, R. L., Matsuki, A., Mead,

- 160 M. I., Methven, J., Mills, G. P., Minikin, A., Murphy, J. G., Nielsen, J. K., Oram, D. E., Parker, D. J., Richter, A., Schlager, H., Schwarzenboeck, A., and Thouret, V.: Chemical and aerosol characterisation of the troposphere over West Africa during the monsoon period as part of AMMA, *Atmos. Chem. Phys.* ~~Atmospheric Chemistry and Physics~~, 10, 7575–7601, <https://doi.org/10.5194/acp-10-7575-2010>, 2010.
- Sandu, I., Brenguier, J.-L., Geoffroy, O., Thouron, O., and Masson, V.: Aerosol Impacts on the Diurnal Cycle of Marine Stratocumulus, *Journal of the J. Atmos. pherie Sci_enees*, 65, 2705 – 2718, <https://doi.org/10.1175/2008JAS2451.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2008.
- 165 Sauvage, B., Thouret, V., Cammas, J.-P., Gheusi, F., Athier, G., and Nédélec, P.: Tropospheric ozone over Equatorial Africa: regional aspects from the MOZAIC data, *Atmos. Chem. Phys.* ~~Atmospheric Chemistry and Physics~~, 5, 311–335, <https://doi.org/10.5194/acp-5-311-2005>, 2005.
- Schrage, J. M. and Fink, A. H.: Nocturnal Continental Low-Level Stratus over Tropical West Africa: Observations and Possible Mechanisms Controlling Its Onset, *Mon. thly Weather Rev_iew*, 140, 1794 – 1809, <https://doi.org/10.1175/MWR-D-11-00172.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2012.
- 170 Schuster, R., Fink, A. H., and Knippertz, P.: Formation and Maintenance of Nocturnal Low-Level Stratus over the Southern West African Monsoon Region during AMMA 2006, *Journal_ J_of the Atmos. pherie Sci_enees*, 70, 2337 – 2355, <https://doi.org/10.1175/JAS-D-120241.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society~~, 2013.
- 175 ~~Stevens, B., Moeng, C.-H., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J.-C., Jiang, H., Khairoutdinov, M., Kirkpatrick, M. P., Lewellen, D. C., Lock, A., Müller, F., Stevens, D. E., Whelan, E., and Zhu, P.: Evaluation of Large-Eddy Simulations via Observations of Nocturnal Marine Stratocumulus, *Mon. Weather Rev_.* 133, 1443–1462, <https://doi.org/10.1175/MWR2930.1>, 2005.~~
- 180 Stjern, C. W., Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, , Andrews, T., Boucher, O., Faluvegi, G., Iversen, T., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivíe, D., Richardson, T., Shawki, D., Shindell, D., Smith, C. J., Takemura, T., and Voulgarakis, A.: Rapid Adjustments Cause Weak Surface Temperature Response to Increased Black Carbon Concentrations, *J. Geophys Res.-Atmos* ~~Journal of Geophysical Research: Atmospheres~~, 122, 11,462–11,481, <https://doi.org/https://doi.org/10.1002/2017JD027326>, 2017.
- 185 Taylor, J. W., Haslett, S. L., Bower, K., Flynn, M., Crawford, I., Dorsey, J., Choularton, T., Connolly, P. J., Hahn, V., Voigt, C., Sauer, D., Dupuy, R., Brito, J., Schwarzenboeck, A., Bourriane, T., Denjean, C., Rosenberg, P., Flamant, C., Lee, J. D., Vaughan, A. R., Hill, P. G., ~~Brooks, B., Catoire, V., Knippertz, P., and Coe, H.: Aerosol influences on low-level clouds in the West African monsoon, *Atmos. Chem. Phys.* 19, 8503–8522, <https://doi.org/10.5194/acp-19-8503-2019>, 2019.~~
- 190 ~~Brooks, B., Catoire, V., Knippertz, P., and Coe, H.: Aerosol influences on low-level clouds in the West African monsoon, *Atmospheric Chemistry and Physics*, 19, 8503–8522, <https://doi.org/10.5194/acp-19-8503-2019>, 2019.~~
- Thouron, O., Brenguier, J.-L., and Burnet, F.: Supersaturation calculation in large eddy simulation models for prediction of the droplet number concentration, *Geoscientific Model Development*, 5, 761–772, <https://doi.org/10.5194/gmd-5-761-2012>, 2012.
- 195 Tulet, P., Crassier, V., Solmon, F., Guedalia, D., and Rosset, R.: Description of the Mesoscale Nonhydrostatic Chemistry model and application to a transboundary pollution episode between northern France and southern England, *J. Geophys Res.-Atmos* ~~Journal of Geophysical Research: Atmospheres~~, 108, ACH 5–1–ACH 5–11, <https://doi.org/https://doi.org/10.1029/2000JD000301>, 2003.
- Tulet, P., Crassier, V., Cousin, F., Suhre, K., and Rosset, R.: ORILAM, a three-moment lognormal aerosol scheme for mesoscale atmospheric model: Online coupling into the Meso-NH-C model and validation on the Escompte campaign, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2004JD005716>, ~~eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2004JD005716>~~, 2005.
- 200 Tulet, P., Grini, A., Griffin, R. J., and Petitcol, S.: ORILAM-SOA: A computationally efficient model for predicting secondary organic aerosols in three-dimensional atmospheric models, *Journal of Geophysical Research:-J. Geophys Res.-Atmos* ~~Atmospheres~~, 111, <https://doi.org/https://doi.org/10.1029/2006JD007152>, 2006.
- 205 Twohy, C. H., Anderson, J. R., Toohey, D. W., Andrejczuk, M., Adams, A., Lytle, M., George, R. C., Wood, R., Saide, P., Spak, S., Zuidema, P., and Leon, D.: Impacts of aerosol particles on the microphysical and radiative properties of

- stratocumulus clouds over the southeast Pacific Ocean, *Atmos. Chem. Phys.* ~~Atmospheric Chemistry and Physics~~, 13, 2541–2562, <https://doi.org/10.5194/acp-13-2541-2013>, 2013.
- Twomey, S.: PRECIPITATION BY DIRECT INTERCEPTION OF CLOUD-WATER, *Weather*, 12, 120–122, <https://doi.org/https://doi.org/10.1002/j.1477-8696.1957.tb00453.x>, 1957.
- Vié, B., Pinty, J.-P., Berthet, S., and Leriche, M.: LIMA (v1.0): A quasi two-moment microphysical scheme driven by a multimodal population of cloud condensation and ice freezing nuclei, *Geoscientific Model Development*, 9, 567–586, <https://doi.org/10.5194/gmd-9-567-2016>, 2016.
- Wood, R.: Stratocumulus Clouds, *Monthly Weather Review*, 140, 2373 – 2423, <https://doi.org/10.1175/MWR-D-11-00121.1>, ~~place: Boston MA, USA Publisher: American Meteorological Society,~~ 2012.
- Yamaguchi, T., Feingold, G., Kazil, J., and McComiskey, A.: Stratocumulus to cumulus transition in the presence of elevated smoke layers, *Geophysical Research Letters*, 42, 10,478–10,485, <https://doi.org/https://doi.org/10.1002/2015GL066544>, 2015.
- Zhang, J. and Zuidema, P.: The diurnal cycle of the smoky marine boundary layer observed during August in the remote southeast Atlantic, *Atmos. Chem. Phys.* ~~Atmospheric Chemistry and Physics~~, 19, 14493–14516, <https://doi.org/10.5194/acp-19-14493-2019>, 2019.
- Zouzoua, M., Lohou, F., Assamoi, P., Lothon, M., Yoboue, V., Dione, C., Kalthoff, N., Adler, B., Babic, K., Pedruzo-Bagazgoitia, X., and Derrien, S.: Breakup of nocturnal low-level stratiform clouds during the southern West African monsoon season, *Atmos. Chem. Phys.* ~~Atmospheric Chemistry and Physics~~, 21, 2027–2051, <https://doi.org/10.5194/acp-21-2027-2021>, 2021.