

The Impact of Aerosols on the Stratiform Clouds over Southern West Africa: A Large-Eddy Simulation Study

Lambert Delbeke¹, Chien Wang¹, Pierre Tulet¹, Cyrielle Denjean², Maurin Zouzoua³, Nicolas Maury⁴
5 and Adrien Deroubaix⁵

¹Laboratoire d'Aérodynamique, Université de Toulouse, CNRS, UT3, IRD, Toulouse, France

²CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

³Laboratoire Atmosphères, Milieux, Observations Spatiales, IPSL, CNRS, Guyancourt, France

⁴CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse; now at LMD/IPSL, Paris, France

10 ⁵IUP, Institute of Environmental Physics, University of Bremen, Bremen, and Max Plank Institute for Meteorology, Hamburg, Germany

Correspondence to: Chien Wang (chien.wang@aero.obs-mip.fr), Cyrielle Denjean (cyrielle.denjean@meteo.fr)

Abstract. The low level stratiform clouds (LLSCs) covering a large area appear frequently during the wet monsoon season in southern West Africa. This region is also a place where different types of aerosols
15 coexist, including the biomass burning aerosols coming from Central and South Africa and the aerosols emitted by local anthropogenic activities. We investigate the indirect and semi-direct effects of these aerosols on the life cycle of LLSCs by conducting a case study based on the airborne and ground-based observations from the field campaign of Dynamic-Aerosol-Chemistry-Cloud-Interaction in West Africa (DACCIIWA). This case is modeled using a Large Eddy Simulation (LES) model with fine resolution and
20 in-situ aerosol measurements including size distribution and chemical composition. The model has successfully reproduced the observed life cycle of the LLSC, from stratus formation to stabilization during the night, and to the upward development after sunrise until break-up of cloud deck in the late afternoon. Additional sensitivity simulations using different measured aerosol profiles also suggest that aerosols can affect the cloud life cycle through both the indirect and semi-direct effect. As expected,
25 modeled cloud microphysical features including cloud droplet number concentration, mean radius, and thus cloud reflectivity are all controlled by aerosol concentration. However, it is found that the variation in cloud reflectivity induced by different aerosol profiles is not always the only factor in determining the incoming solar radiation at ground and thus cloud life cycle after sunrise. Instead, the difference in cloud fraction brought by dry air entrainment from above and thus the speed of consequent evaporation - also
30 influenced by aerosol concentration, is another important factor to consider. Clouds influenced by higher aerosol concentrations and thus having higher number concentration and smaller size of cloud droplets are found to evaporate more easily and thus impose a lower cloud fraction. In addition, by comparing the simulations including versus excluding we find that an excessive atmospheric heating up to $12 K day^{-1}$ produced by absorbing black carbon aerosols (~~BC~~), we find that the semi-direct effect resulted from this heating is to lower the cloud top as well as liquid water path while increase cloud fraction, thus to delay the intense cloud break-up until the later afternoon when convection is further strengthened~~in our modeled cases lowers the height of cloud top and liquid water path, resulting a weaker extent in vertical development while a higher cloud fraction and delaying intense cloud break-up before later afternoon.~~

~~While the thinner clouds resulted from such a heating, on the other hand, would break up faster in late afternoon when convection is further strengthened.~~


1. Introduction

Low-level stratiform clouds (LLSCs) have a higher albedo and a larger cloud cover than many other types of clouds (Hartmann *et al.*, 1992; Chen *et al.*, 2000; Eastman and Warren, 2014). Their reflection of solar radiation is ~~thus~~ important to Earth's radiative budget. LLSCs often occupy the upper few hundred meters in the planetary boundary layer (PBL). Their appearance can be persistent when ~~they are~~ associated with a high-pressure system with a large-scale subsidence that ~~normally brings a stable~~ stabilizes the PBL ~~with a large-scale subsidence above. These clouds~~ LLSCs are often formed over cooler subtropical and mid-latitude oceans, constantly covering more than 50% of these areas (Wood, 2012). During the West African monsoon season, LLSCs can also form frequently over continental southern West Africa (SWA) in the night, then usually break up in the early afternoon of the following day (Schrage and Fink, 2012; Schuster *et al.*, 2013). Under a polluted condition, LLSCs are characterized by numerous and small cloud droplets, increasing the cloud albedo, suppressing drizzle, and extending the cloud lifetime (Twomey, 1957; Haywood and Boucher, 2000; Liu *et al.*, 2014; Carslaw *et al.*, 2017). The presence of LLSCs impacts on the radiative budget of the atmosphere, surface fluxes, the diurnal cycle of the convective boundary layer, and thus the regional climate (Knippertz *et al.*, 2011; Hannak *et al.*, 2017). However, the processes behind the life cycle of LLSCs particularly over SWA remain elusive, hence the representation of these clouds in weather and climate models is still poor (Knippertz *et al.*, 2011; Hannak *et al.*, 2017; Hill *et al.*, 2018).

Stratiform clouds are sensitive to aerosol concentration, chemical composition, and vertical distribution. This is because that aerosol can directly scatter or absorb solar radiation (the direct effect or aerosol-radiation effect), or by serving as cloud nuclei, influence cloud microphysical structure and thus reflectance or lifetime (the indirect aerosol effects or radiative effect of aerosol-cloud interaction plus cloud adjustment) (Boucher *et al.*, 2013). Specifically, the heating associated with aerosol absorption of solar radiation would be able to perturb the thermodynamic stability and thus dynamical processes in the atmosphere as well (the semi-direct effect) (Hansen *et al.*, 1998), and serve as a positive or negative addition to the indirect effect depending on the relative distribution of the aerosol with respect to clouds (*e.g.*, Johnson *et al.*, 2004; Feingold *et al.*, 2005). All these effects can modify the energy budget and thus the status of the planetary boundary layer where the stratiform clouds form and evolve. Aerosols inside stratiform clouds can also be modified by aqueous physio-chemical processes, further altering the forcing strength of aerosol population, whether remaining inside droplets or being released through evaporation, due to their modified morphology and chemical composition (Wood, 2012). Interactions between aerosols and clouds, and their effects on radiation, precipitation, and regional circulations, remain one of the largest uncertainties in understanding and projecting climate change. Indeed, it is still difficult to estimate the indirect effect of aerosols and thus to minimize the uncertainty associated with this effect in the climate models ~~the indirect effect of aerosol is still difficult to estimate~~ (Boucher *et al.*, 2013; Li *et al.*, 2022), ~~making any effort to minimize the associated uncertain in the climate models demanding~~ (Li *et al.*, 2022). ~~Several previous studies had were conducted to investigate~~ aerosol-cloud interactions in LLSCs using high-resolution Large-Eddy Simulation (LES) models, Many of these studies were though

80 ~~mainly on the~~ cases over ocean (e.g., Ackerman *et al.*, 2004; Sandu *et al.*, 2008; Twohy *et al.*, 2013; Flossmann and Wobrock, 2019), where latent heat flux at the surface plays a more important role in the life cycle of LLSCs than sensible heat, while the latter dominates surface fluxes often have insignificant diurnal variation and latent heat alongside moisture from evaporation at sea surface dominate, differing from in the cases over land (Wood, 2012; Ghonima *et al.*, 2014).

85 During the West Africa Monsoon (WAM), aerosols can come from both local and remote sources to SWA. A large amount of Biomass Burning Aerosols (BBA) can be transported from southern and central African towards SWA during the summer monsoon (Haslett *et al.*, 2019). The air masses transporting BBA are also loaded with additional aerosols from anthropogenic emissions upon reaching the highly urbanized regions near the coast (Chatfield *et al.*, 1998; Sauvage *et al.*, 2005; Mari *et al.*, 2008; Murphy
90 *et al.*, 2010; Reeves *et al.*, 2010; Menut *et al.*, 2018; Haslett *et al.*, 2019). A significant quantity of mineral dust aerosols emitted from the Sahara and Sahel throughout the year with a peak in springtime (Marticorena and Bergametti, 1996) can also reach SWA, often in June (Knippertz *et al.*, 2017). Local aerosol sources in SWA are related to anthropogenic activities near the coast (projected to increase with growing population, Lioussé *et al.*, 2014), from where polluted plumes would transport inland
95 (Deroubaix *et al.*, 2019). These different emission sources lead to a complex mix of aerosol constituents in the aera, having a serious impact on human health (Bauer *et al.*, 2019), and possibly complicating the aerosol impacts on the life cycle of LLSCs as well as precipitation over SWA (Taylor *et al.*, 2019).



100 **Figure 1.** Map of southern West Africa with Savè, Kumasi and Ile-Ife locations and the flight track (blue line) of the ATR-42 the 3 July 2016 with NASA Suomi NPP/VIIRS true color corrected reflectance (<https://worldview.earthdata.nasa.gov/>).

The DACCIWA project was designed to better characterize cloud-aerosol-precipitation interactions in SWA (Knippertz *et al.*, 2015). The measurement campaign conducted in June-July 2016 has provided a comprehensive set of ground-based and airborne measurements of clouds and aerosols (Knippertz *et al.*, 2017; Kalthoff *et al.*, 2018; Flamant *et al.*, 2018). The measurements were conducted at three supersites, Savè (Benin), Kumasi (Ghana) and Ile-Ife (Nigeria) (Fig. 1) and coordinated with three research aircrafts: the French ATR-42 operated by SAFIRE (Service des Avions Français Instrumentés pour la Recherche en Environnement), the British Twin Otter operated by British Antarctic Survey, and the German Falcon aircraft operated by DLR (Deutsches Zentrum für Luft und Raumfahrt). Additional radiosoundings were launched from Savè with high temporal frequency, which specifically benefits the monitoring of the LLSCs evolution.

115 DACCIWA campaign has also inspired many modeling studies. Based on the observations from DACCIWA and a parcel model, Taylor *et al.* (2019) and Denjean *et al.* (2020a) showed that most cloud condensation nuclei and absorbing aerosols observed during DACCIWA campaign were from ubiquitous long-range transported BBA, causing a polluted background which limits the effect of local pollution on cloud properties and aerosol radiative effects. Using COSMO-ART model in a simulation of 2-3 July

2016 case, Deetz *et al.* (2018) found that under the influence of the cold air brought by the Maritime Inflow (MI) from Guinean Gulf, stratus-stratocumulus transition is susceptible to the aerosol direct effect, resulting in a spatial shift in the MI front and a temporal shift of the cloud transition. Influenced by anthropogenic emission sources, the break-up time of LLSCs over SWA can be delayed by one hour and daily precipitation rate can decrease by 7.5% according to Deroubaix *et al.* (2022). Moreover, the joint rather than separate impact of the semi-direct and indirect effects of aerosol were also studied with varying magnitude of anthropogenic aerosol emissions by Haslett *et al.* (2019) using COSMO-ART model. The study indicated that cloud droplets number concentration could increase up to 27 % due to transported BBA, making cloud and rain less sensitive to further increase in regional anthropogenic emissions. The impact of sedimentation on LLSCs was indicated by previous studies (*e.g.*, Bretherton *et al.*, 2007). This issue has also been addressed in a modeling of DACCIWA case by Dearden *et al.* (2018) using the Met Office NERC Cloud model (MONC), who demonstrated that sedimentation of cloud droplets, determined by droplet size, could affect liquid water path by removing droplets from the entrainment zone, or by lowering the cloud base and creating more heterogeneous cloud structure. Menut *et al.* (2019) showed in a WRF-CHIMERE simulation that a decrease of anthropogenic emissions along the SWA coast could lead to a northward shift of the monsoonal precipitation and an increase of surface wind speed over arid region in the Sahel, resulting in an increase of mineral dust emission. Pedruzo-Bagazgoitia *et al.* (2020) analyzed the stratocumulus-cumulus transition at a fine scale (a dozen of kilometer sidelong) using a LES at high resolution (50x50 m²), though aerosol effects were not being taken into consideration. These previous modeling studies all highlighted in a regional scale. Majority of them, however, only addressed the response of LLSCs to the aerosol abundance rather than aerosol effects associated with different chemical compositions (such as the semi-direct effect of black carbon) by taking the advantage of measurement data did not well use the enrich information of aerosol chemical compositions (*e.g.*, black carbon, or BC) obtained during the field campaign.

The aim of this study is to improve our understanding of the impacts associated with both local and transported aerosols on the life cycle of LLSCs during the monsoon period over SWA. In doing so, observational data obtained from the well-documented DACCIWA field campaign have been used to constrain a high-resolution LES model incorporated with an interactive aerosol module that is able to represent the complex aerosol compositions besides size distributions. This modeling effort is also among a few studies that model and analyze stratiform cloud nocturnal-diurnal life cycle over land rather than ocean. A description of observations, data, and the model as well as configurations of different simulations is presented in the Method section after the Introduction. Then the results of an analysis aiming to understand and evaluate the modeled reference case against measurements are discussed. Thereafter, the results from several sensitivity simulations are presented. These sensitivity simulations use various observed aerosol profiles with different size distributions and chemical compositions and are designed to examine the indirect and semi-direct effects of aerosols on the life cycle of modeled LLSCs, making this study the first such modeling attempt within the framework of DACCIWA campaign. The last section of the paper summarizes major research findings of this study.

2. Methods

2.1 Observational data

We have used certain measurements of DACCIWA field campaign to configure the model simulations our LLSC case. These data are described as follows.


160 i) Radiosonde data: During DACCIWA campaign, radiosondes were launched with the MODEM system every 1 to 1.5 hour between 17:00 and 11:00 UTC (the local time of Benin is UTC+1) at the supersite of Savè in Benin. This site is located 185 km from the coast and 166 m above sea level. The area is rather flat, and the vegetation is mainly composed of small trees and shrubs. Vertical profiles from ground to 1500 m altitude of temperature, pressure, relative humidity, and wind were measured
165 with a 1 s temporal interval (4 - 5 m of vertical resolution) (Derrien *et al.*, 2016). These sondes were obtained using two balloons of different volumes to reach a preset time of ascent, and after the cutting of the larger balloon, the second one would be used to retrieve the sonde for another use (Legain *et al.*, 2013).

ii) Ground-based measurements: At the supersite of Savè, a CHM15k Ceilometer was deployed by
170 the Karlsruher Institut für Technologie (KIT) to measure the cloud base height continuously with a 1 min interval and a 15 m vertical resolution, based on the backscatter profiles produced by the lidar with a wavelength of 1064 nm and a 5-7 kHz rate (Handwerker *et al.*, 2016). The cloud cover was monitored every day by using a MOBOTIX S15 cloud camera, installed by Université Paul Sabatier (UPS) team, to obtain pictures in visible and IR every 2 min. The aperture angles for the IR channel corresponds to a
175 158 m x 114 m area at a height of 200 m and the pictures are coded in RGB components. A microwave radiometer (the humidity and temperature profiler HATPRO-G4 from Radiometer Physics GmbH) was installed by KIT to measure the brightness temperature to retrieve absolute humidity, liquid water path, and air temperature. The surface heat and radiation fluxes were measured with an energy balance station deployed over grass and bushes. Additional measurements include soil heat flux, air density, and
180 turbulence parameters.

iii) Airborne measurements: The aircraft campaign took place from 29 June to 16 July 2016, conducted collaboratively by three research aircrafts (see Introduction). In this study, only data from the ATR-42 were used as it flew around Savè between 10:00 and 11:00 UTC and probed the cloud layer. The cloud droplet size distribution was measured with a cloud droplet probe (CDP) (Taylor *et al.*,
185 2019). The chemical composition for non-refractive compounds was measured with the Aerodyne compact Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS) (Brito *et al.*, 2018). The black carbon (BC) mass concentration was measured with a single particle soot photometer (SP2) (Denjean *et al.*, 2020b). The aerosol size distribution was measured with a custom-built scanning mobility sizer spectrometer (SMPS, 20–485 nm), an ultra-high sensitivity aerosol spectrometer (UHSAS, 0.04–1 μm),
190 and an optical particle counter (OPC GRIMM model 1.109, 0.3–32 μm) corrected for the complex refractive index provided in Denjean *et al.* (2020a). The total number concentration of particles larger than 10 nm was measured by a condensation particle counter (CPC, model MARIE). Meteorological variables such as temperature, humidity, pressure, and wind speed and direction were also measured by a suite of airborne instruments. A gas concentration analyzer was used to measure certain chemical
195 gases including CO₂, CH₄, and CO.

2.2 Description of the studied case

Our study analyzes the life cycle of LLSCs based on the observed case of 3 July 2016 at the Savè supersite (Fig. 2). The cloud deck formed during the night, at around 02:00 UTC, close to the appearance of the core Nocturnal-Low-Level Jet (NLLJ), which could have a maximum speed around 6 m s^{-1} (Kalthoff *et al.*, 2018), associated with a maximum cooling (Lohou *et al.*, 2020). At formation, the cloud had its base and top located around $310 \pm 30 \text{ m}$ and $640 \pm 100 \text{ m}$, respectively, and was maintained by the cloud top radiative cooling and cold advection (Dione *et al.*, 2019).



205 **Figure 2.** 3 July cloud evolution with the representation of the Cloud Base Height (CBH), the Cloud Top Height (CTH), LCL and ATR-42 flight track near Savè.

The life cycle of LLSCs over SWA typically involves four phases: the stable phase, jet phase, stratus phase, and convective phase (Dione *et al.*, 2019; Lohou *et al.*, 2020). The stable phase begins just after sunset and is characterized by a weak monsoon flow and the cessation of buoyancy-driven turbulence within the PBL generated by surface heating (Zouzoua *et al.*, 2021). The jet phase corresponds to the settlement of key drivers of cooler air advection. Maritime Inflow (MI), a cold and slightly humid air mass from the Guinean coast, often reaches Savè at the end of the afternoon (between 16:00 UTC and 20:00 UTC), then is followed by the NLLJ formation (Adler *et al.*, 2019). The stratus phase begins with LLSC formation when advective cooling continuously increases the relative humidity (RH) until air reaches saturation between 22:00 and 06:00 UTC. The turbulent mixing beneath the NLLJ alongside a strong radiative cooling at the cloud top leads to the persistence of a thick stratus layer (Schuster *et al.*, 2013; Babic *et al.*, 2019). The LLSCs life cycle ends during the final convective phase, which begins when the PBL develops vertically due to solar heating at the surface alongside a weak radiative cooling at cloud top (*e.g.*, Ghonima *et al.*, 2016). By using dataset from Savè supersite, Zouzoua *et al.* (2021) identified three scenarios of evolution depending on the LLSCs coupling to the surface at sunrise. The

coupling was assessed by the departure between the Cloud Base Height (CBH) and the Lifting Condensation Level (LCL).

225 The LLSCs observed on 3 July 2016 follow the four aforementioned phases and evolve by scenario C described by Zouzoua *et al.* (2021) as seen in Figure 2. The cloud is coupled to the surface at sunrise (06:30 UTC), its base rises with growing PBL until break-up occurs in the late afternoon around 16:00 UTC. The cloud deck of July 3 case stands longer (2-3 hours more) comparing to other LLSCs observed during the campaign. The co-located Ka band mobile, dual-polarization Doppler radar (8.5 mm, 35.5 MHz) at Savè supersite detected light drizzle precipitation from higher clouds in a rather short period
230 during the first hours of the convective phase, while no precipitation was detected by the surface rain gauge. Thus, the late LLSC break-up could be explained hypothetically by the cooling alongside moistening brought by the evaporation of this light precipitation, which could enhance the liquid water path of the beneath LLSC (Zouzoua *et al.*, 2021). Nevertheless, our focus of this study is on the life cycle of LLSC as influenced by aerosols alongside planetary boundary layer dynamics rather than examining
235 the above hypothesis, which is likely related to a process beyond the local scale. Therefore, our model setting is made to specifically eliminate the influence of mid-cloud layer for this purpose.

On 3 July 2016, the ATR-42 flew around Savè supersite and probed the boundary layer around 10:00 UTC. The airborne instruments detected aerosol size distributions with a maximum number concentration around 3500 cm^{-3} mainly in the Aikten mode. The ATR-42 also detected an export of polluted air mass
240 from Lomé (a coast city), which could explain the measured high aerosol concentration in the Aikten mode (Denjean *et al.*, 2020a). The measured aerosol chemical composition was mainly dominated by organics (55.3%), followed by sulfates (24.5%), ammoniac (11.2%), and nitrates (6.2%), while only a small amount of BC mass was detected around Savè (2.8%). However, the measured aerosol size distributions were found to need a correction based on aerosol refractive index to avoid bias. For this
245 purpose, Denjean *et al.* (2020a) provided corrected profiles for various types of aerosol populations measured during the DACCIWA campaign. Our modeling has thus used corrected rather than “raw” measurements.

2.3 Meso-NH Model

250 In this study, we have simulated the observed case using the Meso-NH model (Lac *et al.*, 2018). Meso-NH is a non-hydrostatic atmospheric research model that has been applied to studies in different scales ranging from synoptic to turbulent. Deployed in a limited area, the model uses advanced numerical techniques like monotonic advection schemes for scalar transport, and fourth order advection scheme for momentum (Jiang and Shu, 1996). Sub-grid turbulence is parametrized using turbulence kinetic energy
255 (TKE) based on Deardorff turbulent mixing length. In this study, a fourth order advection scheme CEN4TH, centered on space and time, is applied with a Runge-Kutta centered 4th order temporal scheme for momentum advection (Lunet *et al.*, 2017). Aerosol and chemistry are also well represented. Here, Meso-NH version 5.4.2 is used and the relevant component modules and parametrizations for this study are described as follows.

260 The aerosol-cloud framework of Meso-NH version used in this study is LIMA (Liquid Ice Multiple Aerosol). LIMA includes a complete two-moment scheme predicting both the mass mixing ratio and the number concentration of aerosol species (Vié *et al.*, 2016), using a superimposition of several aerosol

265 modes, with each mode defined by its chemical composition and size distribution. Aerosols can act as
either Cloud Condensation Nuclei (CCN) or Ice Forming Nuclei (IFN). Based on the ICE3-ICE4 ice
microphysics schemes (Caniaux *et al.*, 1994; Pinty and Jabouille, 1998; Lascaux *et al.*, 2006) and the two-
moment warm microphysical scheme C2R2 (Cohard and Pinty, 2000), LIMA also predicts the mass as
well as number concentration of cloud droplets, raindrops, and pristine ice crystals. For modeling
boundary layer cloud in LES mode, a pseudo-prognostic approach for correcting the diagnostically
derived supersaturation was developed (Thouren *et al.*, 2012) to limit the droplet concentration
270 production and to better represent cloud-top supersaturation. A variant to C2R2, called KHKO, was
developed by Geoffroy *et al.* (2008) for clouds producing drizzle (differentiated from cloud droplet with
a radius larger than 25 μm) following Khairoutdinov and Kogan (2000) parametrization. The-Only the
sedimentation of drizzle is calculated in the scheme. The KHKO alongside necessary modifications has
been brought in the LIMA warm phase framework in order to better represent drizzle in thin and low
275 precipitating warm clouds. Therefore, we have adopted this version of LIMA in our modeling.

To better represent aerosols, we have used the aerosol module ORILAM (Organic Inorganic
Lognormal Aerosols Model) in this study and coupled it with Meso-NH to interconnect the cloud
microphysics module with LIMA (Tulet *et al.*, 2005). ORILAM describes the size distribution and
chemical composition of aerosols using two lognormal functions respectively for the Aitken and
280 accumulations mode. These modes are internally mixed. For each of them, the model computes the
evolution of the primary species (black carbon and primary organic carbon), three inorganic ions (NO_3^- ,
 NH_4^+ , SO_4^{2-}), and condensed water. ORILAM includes a Secondary Organic Aerosols (SOA) module
(Tulet *et al.*, 2006) that is, however, not included in this study. Three moments (the zeroth, third, and
sixth) are considered for each mode to compute the evolution of total number, median diameter, and
285 geometric standard deviation. Note that the choice of the 6th moment is numerical since it allows one to
calculate the coagulation coefficients explicitly and to facilitate the integration of the aerosol solver. The
size distribution can evolve through both intra- and intermodal particle coagulation. It can also evolve
through condensation and merging between modes. ORILAM includes the CCN activation scheme of
Abdul-Razzak and Ghan (2004) (though without the correction of Thouren *et al.*, 2012) in order to replace
290 the one of LIMA to calculate the number of activated CCN. The use of ORILAM needs to activate the
gas phase chemistry scheme of Meso-NH (Tulet *et al.*, 2003; Mari *et al.*, 2004) using the EXQSSA solver.
ORILAM has a module for gas-particle thermodynamic equilibrium (EQSAM for inorganics and
MPMPO for organics) that allows the model to calculate the contents of inorganic and organic
compositions including water within aerosols (*e.g.*, Metzger *et al.*, 2002; Griffin *et al.*, 2003). The solver
295 combines moment 0 (integrated number) and 3 (integrated new volume resulted from the hygroscopic
growth) to calculate the new dimensional distribution (Tulet *et al.*, 2005, 2006). ORILAM directly
computes the evolution of aerosol extinction, single scattering albedo (SSA), and asymmetry factor that
are coupled online with the radiation scheme of Meso-NH for the 6 short wavelengths from the aerosol
chemical composition and size parameters (Aouizerats *et al.*, 2010).

300 ECMWF radiation module is adopted in this study. Based on the two-stream method, this module
calculates the atmospheric heating rate and then net surface radiative forcing. Longwave radiation scheme
used is the Rapid Radiation Transfer Model (RRTM; Mlawer *et al.*, 1997), based on the correlated k-
distribution method. It integrates 16 bands and 140 g points (Morcrette, 2002). The shortwave scheme

305 uses the photon path distribution method (Fouquart and Bonnel, 1980) in six spectral bands. Fluxes are calculated independently in clear and cloudy portions before being aggregated. The liquid cloud effective radius is computed from the liquid water content and droplet number concentration with the Martin *et al.* (1994) parametrization [for continental condition](#).

310 The surface model used in our modeling is the SURFEX, which is a standardized surface module containing surface schemes externalized of Meso-NH (Masson *et al.*, 2013). With SURFEX, each grid point can be split into four tiles: land, town, sea, and inland water (lake, rivers). In case of a shrubs surface, the interactions between soil, biological processes, and the atmosphere are calculated by ISBA parametrization (Noilhan and Planton, 1989). Several evapotranspiration formulations are available for simulating plants and CO₂ fluxes. Soil is represented as a bucket of two or three layers. The land tile can be separated into as many as 19 subtiles following the type of vegetation.

315 **2.4 Model settings**

320 Based on the observations and the capability of the model, a reference case (REF) was first designed to simulate using the LES. The reference case serves as a base for the model to reproduce the major features of the observed LLSCs life cycle particularly using an observed aerosol profile. It also serves as a comparison base for further sensitivity simulations with different aerosol configurations to study the impacts of aerosol composition ~~alongside-and~~ abundance on LLSCs.

325 The model domain is a 3D box of 9.6 km x 9.6 km x 2 km in size, with a horizontal resolution of 40 m x 40 m. Note that the radiation module still proceeds its calculations above 2 km using prescribed profiles. The vertical resolution is 10 m between 0 m and 1200 m then 40 m above to explicitly resolve the important turbulent eddies. A periodic boundary condition on the horizontal directions is applied and a “sponge layer” is set between 1.8 and 2 km height to absorb wave reflection. A thermodynamic perturbation is deployed to activate turbulence at the beginning of the simulation at 23:00 UTC of 2 July and the spin-up is 1h (though observed clouds formed around 02:00 UTC). A subsidence profile is applied following Bellon and Stevens (2013) scheme $w_{subs}(z) = -w_0[1 - \exp(-z/z_w)]$, with $w_0 = 15 \text{ mm s}^{-1}$ and $z_w =$
330 250 m. This subsidence profile is applied during the entire simulation to keep a nearly constant cloud top height during the stratus phase and to better control the convective phase. The surface energy and water fluxes are simulated by SURFEX ISBA scheme, parametrized using data from Savè supersite, with the typical vegetation consisting of shrubs, crops, or taller trees, assuming a flat surface in the area around Savè. A time-step of 2s is used, which appears to be adequate based on testing runs to study the LLSCs
335 nocturnal-diurnal variations particularly involving aerosol and cloud microphysics. The radiation scheme is called every 10 minutes. Note that previous studies regarding nocturnal stratus-stratocumulus suggested that a vertical resolution as fine as 5 meters near the cloud top would be necessary for reproducing the cloud top entrainment and thus cloud macrophysical structures (Stevens *et al.*, 2005). However, the nocturnal-diurnal life cycle in our case involves a dynamically evolving cloud top from 400 to 1200 m,
340 particularly in the daytime, making it a difficult task to prescribe a highlighted zone for finer resolution. Our fast-testing results, on the other hand, did not suggest any significant difference between the run with 10 m and 5 m vertical resolution (not shown). Therefore, the current vertical resolution and the time step are selected to well cover all possible cloud tops during the simulation and to provide a best economic computational performance for modeling aerosol-cloud interaction with a fully coupled chemistry model.




Figure 3. Aerosol chemical mass compositions (a) and size distribution fitted into 2 modes described in Table 1 (b) used in REF.

350

	$N_a (cm^{-3})$	σ	D (nm)
Mode 1	654	1.49	63.98
Mode 2	1530	1.53	190.97

355

Table 1. REF aerosol size distribution described by two modes configured by three parameters (number concentration N_a , standard deviation σ and diameter D).

360

REF case is configured using the radiosondes of 2 July at 23:00 UTC for temperature, humidity, and horizontal wind components (U, V). The simulation is then forced by tendency profiles of temperature, humidity, and horizontal wind applied homogeneously on the domain each hour. These tendency profiles are based on the hourly radiosondes launched on 3 July between 00:00 and 11:00 UTC. After 11:00 UTC, the next tendency profiles were derived based on the measurements of the microwave radiometer, the analysis of surface incoming solar radiative flux, and the cloud thickness and cover. Note that, despite these best possible efforts in configuring a set of observation-constrained tendency profiles to reproduce the observed cloud field, it is difficult to eliminate the possibility that such profiles could reflect certain local thermodynamic effects however small they are. In practice, our main goal is to make the profiles to be able to force the model to reproduce observed quantities of major features such as cloud top, base, liquid water path (LWP), surface incoming solar radiation, among others, in the REF case. This would serve the best purpose for us to address the major issue of this study, *i.e.*, the role of different aerosol profiles in the life cycle of modeled LLSCs.


370

We use a "background" distribution as the aerosol profile for REF simulation. This profile, derived from the corrected original measurements as described in Denjean *et al.* (2020a), reflects the influence of aged BBA on clouds with a minor influence from local anthropogenic sources. The aerosol size distribution is dominated by a particle accumulation mode centered at 190 nm and a smaller Aiken mode centered at 64 nm (Figure 3b). This profile exhibits a high loading of aerosols with a maximum of 1400 cm⁻³ in the accumulation mode. The aerosol chemical composition is dominated by organics (52.2%), followed by sulfates (25.3%), ammonium (8.9%), BC (10.7%), and nitrates (2.8%). The configuration of ORILAM has been initialized using the REF aerosol chemical composition and size distribution given in Table 1 and Figure 3b by fitting the SMPS profiles into two lognormal modes using the "py-smmps" package (Hagan *et al.*, 2022), with each mode having the same chemical composition.

3. Analysis of REF Results

3.1 Simulated the life cycle of LLSCs


The simulation of the REF scenario reproduces the formation of the observed LLSCs on 3 July 2016 as shown in Figure 4. The formation of clouds leads to, as described in section 2.2, the end of the jet phase. The domain mean CBH, derived from the modeled mixing ratio of cloud droplets, follows the ceilometer's measurements during the stratus phase between 02:00 and 10:00 UTC, varying between 100 and 300 m of altitude. The simulated domain mean CTH evolves from 400 to 650 m of altitude during the same period, well within the range from 500 to 580 m detected by the radar. The modeled domain mean CBH and CTH, however, overestimate the measurements of ceilometer and radar, respectively, during some periods in late morning and afternoon. The difference between the simulated and ceilometer detected CBH can be as large as 150 m, *e.g.*, at 11:00 UTC. While modeled CTH is often higher than radar measurements by 100 m. Between 15:00 and 16:00 UTC, the simulated domain mean CBH approaches again the ceilometer readings from 600 to 950 m (no radar values are available to evaluate the simulated CTH). As mentioned in section 2.1, the ceilometer is a vertically pointing lidar, its detected values come from the vertical profiles of reflectivity with a 30 m of resolution. The differences between the model and the observation could come from the different representation between the simulated result (a domain average) and ceilometer detection (limited to only vertical direction), in addition to the vertical resolution of observed profiles. The same could also apply to the difference between modeled and radar detected CTH, in addition to the limitation of radar in detecting hydrometeors. Nevertheless, certain model weaknesses likely associated with a lack of hourly radiosondes during the afternoon period as an observational constrain would contribute to these discrepancies as well.



410 **Figure 4.** Simulated LLSCs deck evolution compared to Savè ceilometer and radar measurements, vertical color bars attribute cloud fraction in percentage at each altitude level. Here simulated CBH and CTH represent domain-averaged cloud base and cloud top height, respectively. Different phases might have overlaps; therefore, their marks only serve a reference purpose here.


To analyze the cloud cover profile over the domain, the cloud fraction (CF) at each model layer is calculated as the occupation percentage of the cloud pixels with a total condensed water mixing ratio exceeding 0.05 g kg^{-1} at that given layer (Fig. 4). Note that this cloud fraction differs from the cloud fraction defined as a column metric. In addition, Liquid Water Path (LWP) at each column (Fig. 5), calculated based on column integrated cloud water mixing ratio, brings a view on the horizontal organization and homogeneity of the cloud deck. During the stratus phase, the CF is nearly equal to 100% between CBH and CTH (Fig. 4), suggesting a more homogeneous cloud deck consistent with cloud observations with sky camera (visible range) (Fig. 5, the top panels). Notably, peak LWP values between 06:00 and 12:00 UTC are quite close while domain-mean values differ (Fig. 5). In comparison, both peak and domain mean LWP are sharply lower at 16:00 UTC due to cloud break-up and dissipation. Between 10:00 and 13:00 UTC, CF of the layers between domain mean CBH and CTH starts to decrease from near 100% to 90%, while CF at CBH and CTH decreases more substantially to reach near 60% and 80%, respectively. This leads to a less homogeneous cloud deck confirmed by the LWP map and the observation of the sky camera at 12:00 UTC (the middle row of Figure 5). Indeed, more cloud-free pixels begin to appear, hence sunlight is seen through the cloud deck by the cloud camera. Finally, CF continues to

430 decrease until the end of the convection phase with a maximum barely reaching 80%, and the values around domain mean CBH and CTH level are as low as 20% and 40%, respectively (Fig. 4). This demonstrates the break-up of the cloud deck during convection and the cloud thinning. The bottom panels of Figure 5 clearly show the dissipation of many cloud blocks alongside substantially thinning of the remaining ones at 16:00 UTC.



435

Figure 5. Comparison between modeled liquid water path (LWP, $g\ kg^{-1}\ m$) and the images from Savè cloud camera at 06:00 (top), 12:00 (middle) and 16:00 UTC (bottom).



440 **Figure 6.** Comparison between Savè surface observation and REF simulation for SW radiation flux at surface (SWRADSURF, a), sensible heat flux (H, b) and latent heat flux (LE, c) all expressed in $W m^{-2}$ at the surface. The variation of REF for each parameter indicates the range of possible values these parameters can take.

Figure 6a shows the comparison between the modeled domain-average shortwave (SW) radiation flux at the surface (SWRADSURF) and the corresponding measurements performed by the energy balance station. Observed values are fitted following the LOcally Weighted Scatterplot Smoothing (LOWESS) method (Cleveland, 1979). The temporal evolution of the modeled SWRADSURF follows the observations rather well despite some biases. The solar radiation reaches the ground around 06:00 UTC and increases gradually thereafter by reaching near 200 W m^{-2} at the end of the stratus phase (10:00 UTC). As cloud deck becomes inhomogeneous during the convective phase (10:00 to 16:00 UTC), the modeled surface solar flux reaches a maximum of 300 W m^{-2} , which is a bit lower than the fitted 350 W m^{-2} value from measurements. When the clouds break up further, more solar radiation can reach the surface, and model and observation agree well thereafter with an exception at 15:00 UTC, where the mean modeled curve decreases to 200 W m^{-2} while the fitted observation curve is near 320 W m^{-2} due to an overestimation of the cloud thickness by the model. At 16:00 UTC, both modeled and measurement values are very close around 280 W m^{-2} . Generally, the modeled maximum values are higher than the ones detected by the Savè ground instrument.

Figure 6b and 6c show that the evolutions of the modeled domain-mean latent and sensible heat flux reproduce those measured by the instrument rather well. During the night, the sensible heat flux is negative then increases to 0 W m^{-2} close to the sunrise time (06:00 UTC), indicating a reduction of the cooling close to the ground (Dione *et al.*, 2019). Between 09:00 and 14:00 UTC, the modeled sensible and latent heat flux follow the measured trends though with a clear temporal offset, leading to an overestimate of almost 70 and 18 W m^{-2} , respectively. Then the modeled curves go below the fitted observations at 15:00 UTC until after 18:00 UTC. The difference between modeled and observed heat fluxes may be again due to the different representations, as modeled quantities are domain-mean values while measurements were made at a single point.


In Summary, the REF simulation has ~~successfully~~ reproduced all the major observations at Savè on 3 July 2016 reasonably well. For example, the modeled cloud thickness and coverage reflect the measured cloud macrophysical status despite some discrepancies, likely due to a lack of hourly radiosonde data to constrain the tendency profiles particularly in the afternoon hours. The modeled solar radiation at ground also follows the measurements very well except for certain overestimates. In addition, the sensible and latent heat fluxes measured at Savè have also been well captured by the model despite certain temporal offsets.

3.2 Thermodynamic, dynamical, cloud microphysical, and radiative analyses

Thermodynamic, dynamical, and radiative processes and their interaction with cloud microphysics are among the key factors in determining the life cycle of LLSCs. Here we discuss the evolutions of these processes simulated by the model in the REF case to better understand the reasons behind model-observation consistency or discrepancy. The discussion will be emphasized on three periods: the transition between jet and stratus phase when cloud forms (between 00:00 and 04:00 UTC), the stratus phase between 06:00 and 10:00 UTC, and the convective phase between 12:00 and 17:00 UTC corresponding to the break-up stage of LLSCs.

3.2.1 Transition jet-status phase

485 Figure 7 displays the modeled domain-average profiles of selected macro- and microphysical features for the transition of jet to status phase, when maritime inflow already reached the site. As expected, the advection of cold and slightly humid air leads to an increase of relative humidity (RH) to reach 100% at 02:00 UTC at 100 m above ground. After this time, RH exceeds saturation between 100 and 500 m of altitude. The inversion occurs around 325 m and 500 m respectively at 02:00 UTC and at 04:00 UTC. 490 The NLLJ is well represented in modeled results as the mean wind speed (w_s) before cloud formation is greater than 7 m s^{-1} . After cloud formation, the NLLJ core nearly corresponds to the mean cloud base height (Adler *et al.*, 2019; Babic *et al.*, 2019; Lohou *et al.*, 2020). The turbulence during this period is shear-driven due to this NLLJ, yielding a well-mixed sub-cloud layer. The turbulent kinetic energy (TKE) is high above ground (0.05 to $0.1 \text{ m}^2 \text{ s}^{-2}$), then decreases to near zero above rough 200 meters at 00:00 UTC. At 02:00 and 04:00 UTC, TKE increases at the level of CTH (350 and 500 m, respectively) and decreases at the center of clouds (near zero and $0.04 \text{ m}^2 \text{ s}^{-2}$), indicating this area is less turbulent than the extremities of the cloud layer. 495



500 **Figure 7.** Profiles from left to right of temperature (T, a), relative humidity (RH, b), equivalent potential temperature (θ_e , c), horizontal wind speed (w_s , d), aerosol number concentration (N_a , dashed curve, e), cloud droplets number concentration (N_c , plain curve, e), mean cloud droplet radius (r_c , f), turbulent kinetic energy (TKE, g), longwave heating rate (LWHR, dashed curve, h) and shortwave heating rate (SWHR, plain curve, h) at 00:00, 02:00 and 04:00 UTC. The horizontal dashed dot lines represent mean cloud base height (CBH) and dotted horizontal lines the mean cloud top height (CTH).

505 Cloud droplet number concentration or CDNC (N_c) is determined by the supersaturation in an updraft and the number of aerosols that can activate at this supersaturation. In Figure 7e, simulated aerosol concentration is the highest close to the ground then decreases with altitude up to around 2 km, similar to the airborne measurements during DACCWA (Taylor *et al.*, 2019; Denjean *et al.*, 2020a; Deroubaix *et al.*, 2019; Flamant *et al.*, 2018). The simulated cloud microphysical features reflect a polluted condition

510 as N_c reaches above $1200 \text{ droplet cm}^{-3}$ and mean cloud droplet radius r_c around $5 \mu\text{m}$ that is not enough to form drizzle (larger than $25 \mu\text{m}$ as defined in the model; typical size reaching the ground can be between 0.2 mm and 0.5 mm , ref. Pruppacher *et al.*, 1998; Sandu *et al.*, 2008). These modeled values are in the range of corresponding measurements at the same altitude by Taylor *et al.* (2019).


The emission of thermal radiation by the clouds during the stratus phase creates a cooling at the cloud
515 top as demonstrated by the evolution of modeled Long-Wave Heating Rate (LWHR) profiles at Figure 7h. For LLSCs at this stage with many low LWP ~~bloekscolumns~~ (e.g., with $\text{LWP} < 25 \text{ g/m}^2$; Petters *et al.*, 2012), the more numerous the cloud droplets are the stronger the cooling is (e.g., Petters *et al.*, 2012), as shown in Fig. 7h that LWHR can reach -50 K day^{-1} . This strong longwave emission can reduce the thermal production of turbulence above the cloud top, deepening the temperature inversion. A stabilized
520 cloud top layer by radiative cooling and a NLLJ core contributing to the shear-driven turbulence below the cloud base leads to a well-mixed cloud layer, making the LCL to correspond to the LLCs base as seen in Fig. 2 (Adler *et al.*, 2019; Lohou *et al.*, 2020).

3.2.2 Stratus phase

525 The stratus phase starts just after the sunrise. Maintaining stratus in almost the same state throughout this phase needs a stable ground temperature and moisture supply. As shows in (Figure 8), between 06:00 and 08:00 UTC the ground temperature varies little around 23°C , supersaturation still exists between CBH and CTH, and air masses are quite well-mixed within the boundary layer as θ_e is near constantly at 347 K (Fig. 8c). The horizontal wind speed between the ground and the cloud base decreases from the
530 magnitude in the previous transition phase (Fig. 8d), indicating a weakening NLLJ core (nearly 2 m s^{-1}). TKE value between ground and cloud center decreases from its previous magnitude to $0.03 \text{ m}^2 \text{ s}^{-2}$, while increases slightly to $0.04 \text{ m}^2 \text{ s}^{-2}$ at the mean CTH. At 08:00 UTC, TKE reaches $0.05 \text{ m}^2 \text{ s}^{-2}$ in the cloud layer, owing to an increase of surface solar heating (Fig. 8g).

The aerosol concentration from 06:00 and 08:00 UTC is around 2000 cm^{-3} below 500 m altitude,
535 then decreases along altitude, which is high enough to sustain a CDNC of $1100\text{--}1200 \text{ droplets cm}^{-3}$ between CBH and CTH as shown in Fig. 8e. The maximum layer-mean droplet radius is about $6 \mu\text{m}$, still not enough to form a significant drizzle. The cloud layer has an albedo close to 1 due to the high CDNC. The presence of light absorbing aerosol amplifies the Short-Wave Heating Rate (SWHR) at the cloud top. At 08:00 UTC, the maximum SWHR and LWHR are about 25 K day^{-1} and -60 K day^{-1} , respectively (Fig.
540 8h).

At 10:00 UTC, the cloud layer starts to rise significantly, with CBH and CTH reaching 340 and 660 m , respectively (Fig. 8). Moreover, stronger solar irradiance reaches the ground (220 W m^{-2}), leading to the heating of the surface and the increasing of the sensible and latent heat fluxes as seen in Figure 6. It also increases the surface temperature to 24°C and at the cloud top to 20°C (Fig. 8a). The NLLJ core is
545 no longer present at this moment. TKE increases to $0.1 \text{ m}^2 \text{ s}^{-2}$ throughout the vertical layer from 50 meter above the ground to a level just below the cloud top (Fig. 8g). This enhancement of turbulence is expected to increase entrainment entering the cloud from above as well. The SWHR increases to 45 K day^{-1} , almost compensates the LWHR cooling of 62 K day^{-1} .



550 **Figure 8.** Profiles from left to right of temperature (T , a), relative humidity (RH, b), equivalent potential temperature (θ_e , c), horizontal wind speed (w_s , d), aerosol number concentration (N_a , dashed curve, e), cloud droplets number concentration (N_c , plain curve, e), cloud droplet radius (r_c , f), turbulent kinetic energy (TKE, g), longwave heating rate (LWHR, dashed curve, h) and shortwave heating rate (SWHR, plain curve, h) at 06:00, 08:00 and 10:00 UTC. Dashdot horizontal lines represent mean cloud base height (CBH) and dotted horizontal lines the mean cloud top height (CTH).

555

3.2.3 Convective phase

This phase extends from 12:00 to 17:00 UTC on 3 July 2016. During this period, surface SW radiation flux is maximized at 300 W m^{-2} (Figure 6), leading to the highest surface heating of the day and an increase of the ground temperature from 25 to 27 °C (Fig. 9a). Convection of humid air masses causes the CBH and CTH to rise from 450 to 925 m and from 760 to 1100 m, respectively. Moreover, at 16:00 UTC, the equivalent potential temperature decreases above 450 m of altitude, indicating an unstable air mass there. The horizontal wind speed is weak at the beginning of the phase with 0.5 m s^{-1} at ground level but increases along time to reach 1 m s^{-1} and from 1 to 3 m s^{-1} around 700m altitude. This increase coincides the dissipation of the LLSCs and indicates the arrival of the marine inflow.

565 TKE value below the cloud base is higher or similar to that inside the cloud from 12:00 to 14:00 UTC, showing a well-mixed PBL (Fig. 9g). From 16:00 UTC, TKE decreases near the ground but increases at cloud level to a value of $0.15 \text{ m}^2 \text{ s}^{-2}$, showing a strong turbulence layer within the vertically lifted while thinner cloud due to enhanced convection.

570 The aerosol distribution varies along with the dynamical situation, with a maximum concentration reaching 1700 cm^{-3} within the PBL. The domain mean CDNC has a maximum value of $900 \text{ droplets cm}^{-3}$ at 12:00 UTC. This value then decreases along time as more clouds dissipate (Fig. 9e). After clouds become thinner and start to break, reduced CF allows more solar radiation to reach the ground. The maximum value of SWHR changes from 45 K day^{-1} at 12:00 UTC (almost compensating the longwave

cooling at cloud top) to about $10 K day^{-1}$ at 16:00 UTC. The cloud top LW cooling is near constant at the
 575 end of convection phase with $-45 K day^{-1}$ (Fig. 9h)




Figure 9. Profiles from left to right of temperature (T, a), relative humidity (RH, b), equivalent potential temperature (θ_e , c),
 580 horizontal wind speed (w_s , d), aerosol number concentration (N_a , dashed curve, e), cloud droplets number concentration (N_c , plain curve, e), cloud droplet radius (r_c , f), turbulent kinetic energy (TKE, g), longwave heating rate (LWHR, dashed curve, h) and shortwave heating rate (SWHR, plain curve, h) at 12:00, 14:00, 16:00 and 17:00 UTC. Dash-dot horizontal lines represent mean cloud base height (CBH) and dotted horizontal lines the mean cloud top height (CTH).

4. Sensitivity Study to Examine the Influence of Different Aerosol Profiles on LLSC Life Cycle

585 Previous studies have indicated that the life cycle of stratus or stratocumulus within the planetary boundary layer depends on a subtle balance among several critical while interconnected forcings, including surface heat fluxes, cloud top and base radiative profiles, and thus turbulent mixing (*e.g.*, Stevens *et al.*, 2005; Dussen *et al.*, 2014, Ghonima *et al.*, 2016). Our simulation results of the REF case support previous findings particularly for the cases over land, where the surface sensible heat plays a significant role. Nevertheless, the role of aerosols in such a life cycle have rarely been examined in depth.
 590 Given the critical role of aerosols in determining cloud macro- and microphysical features and thus radiation, this is an important issue to address to advance our understanding of the LLSC life cycle. A unique component of our study is the deployment of an interactive aerosol and atmospheric chemistry module in the observation-constrained LES modeling framework. The REF simulation has demonstrated
 595 that this model is capable to reproduce many observed dynamical, thermodynamic, and aerosol features of the July 3 LLSC case despite certain biases. Thus, we have designed additional sensitivity simulations, and using the results of REF run as a base to further isolate the aerosol impacts on LLSC life cycle through: (1) the difference in cloud droplet number concentrations resulted from aerosol profiles that

600 differ in both number concentration and chemical composition; and (2) the semi-direct effects from absorption of black carbon aerosols. In the following sections, we discuss the modeling configurations alongside outcomes of these two sets of simulations.

4.1 Impact of different aerosol profiles on micro- and macrophysical properties of LLSCs

605 We have firstly configured two sensitivity simulations with observation-based aerosol profiles differing from the one used in REF run (Figure A1 and Table 2). The first simulation uses an aerosol profile that reflects an influence of heavy anthropogenic pollution, obtained based on the aerosol chemical composition and size distribution observed by Brito *et al.* (2018) and Denjean *et al.* (2020a) within urban plumes originated from cities of Lomé, Accra and Abidjan, hereafter referred as POL. The second is a simulation that uses a clean aerosol profile derived by dividing REF aerosol concentration by 10, called
 610 CLEAN. These two sensitivity simulations are otherwise configured the same as the REF simulation. Comparing to REF case, aerosol profile of POL has a slightly higher peak number concentration but in a different mode. In addition, sulfate mass ratio in POL aerosol profile is much higher than that in REF profile, while organic carbon mass ratios are quite close in both profiles. REF, POL, and CLEAN runs simulate the July 3 case with different aerosol number concentrations and chemical compositions as
 615 reflected in their size distributions. Therefore, these simulations are expected to produce different CDNCs alongside dynamical consequences. Comparison between their results could provide us with information about the aerosol impacts on LLSC life cycle through abundance.

Case		$N_a (cm^{-3})$	σ	D (nm)
POL	Mode 1	17100	1.54	55.19
	Mode 2	2650	2.14	101.83
CLEAN	Mode 1	65	1.49	63.98
	Mode 2	153	1.53	190.97

620 **Table 2.** Aerosol size distribution parameters for POL and CLEAN runs including number concentration, standard deviation, and diameter for two aerosol modes.

Indeed, POL and REF have produced clearly different cloud microphysical features including droplet number concentrations alongside mean radius throughout the lifetime of modeled clouds (Fig. 10a and 10b). At the time of cloud formation (02:00 UTC), despite having a similar liquid water content (LWC) around $0.35 g m^{-3}$ at 250 m in both cases, N_c^{POL} reaches 333 *droplets cm⁻³* and r_c^{POL} 6.45 μm instead of
 625 653 *droplets cm⁻³* and 5.1 μm for REF case, indicating a result of differences mainly in Mode 2 aerosol numbers between the two scenarios (at 02:00 UTC the updraft near cloud base is rather weak at less than $0.30 m s^{-1}$ in both cases). This trend is about to reverse at 06:00 UTC when the CDNC and radius are equal to 1208 *droplets cm⁻³* and 6.43 μm in POL, and 1305 *droplets cm⁻³* and 6.12 μm in REF, respectively.
 630 After 08 UTC and until the cloud break up, N_c^{POL} is much higher than N_c^{REF} with a maximum difference of 1425 *droplets cm⁻³* at 14:00 UTC. Their respective radii are 4.42 μm and 5.18 μm while the liquid


635 water content profiles are quite the same as near 0.47 g m^{-3} at 750 m. The difference between POL and REF in CDNC after sunrise suggests that the activation favors the POL profile with higher sulfate content when updraft is strengthened. The above results of CDNC are in good agreement with the ACPIM parcel model simulation done by Taylor *et al.* (2019) where CDNC varies in a range of 500–1400 *droplets cm*⁻³ depending on the inland or offshore (offshore + local emissions) aerosols origin.

640 The difference between CLEAN and REF in cloud microphysical features are also significant. As expected, from formation to break-up of the clouds, N_c^{CLEAN} is lower than N_c^{REF} and r_c^{CLEAN} is larger than r_c^{REF} . At 02:00 UTC, N_c^{CLEAN} has a maximum value of 181 *droplets cm*⁻³ and r_c^{CLEAN} of 7.58 μm , in comparison to 653 *droplets cm*⁻³ and 5.1 μm for N_c^{REF} and r_c^{REF} respectively with the same liquid water content value (0.35 g m^{-3}). r_c^{CLEAN} further increases to 12.55 μm at 08:00 UTC, then decreases slowly to a maximum value of 10.97 μm at 14:00 UTC with LWC^{CLEAN} reaches near 0.45 g m^{-3} instead of 0.49 g m^{-3} for LWC^{REF} , likely due to an increased activation ratio of aerosols after sunrise. Despite a relatively larger droplet size in CLEAN than POL and REF case, there is no clear sign of significant drizzles even during the convection stage (Fig. 10). ~~Nevertheless, Note that sedimentation thus evaporation of larger droplets (smaller than drizzles though) from entrainment zone and cloud base could likely create a thermodynamic perturbation as well (e.g., Stevens *et al.*, 1998; Jiang *et al.*, 2002). Consistent with certain previous findings (e.g., Bretherton *et al.*, 2007), in a LES simulations using passive aerosol profile for July 4-5 DACCIWA case, Dearden *et al.* (2018) found that the sedimentation would remove larger cloud droplets from the entrainment zone thus, through a feedback, lead to a cloud deck with higher LWP while smaller CF than the case where such a sedimentation is completely excluded. Since the sedimentation of non-drizzle droplets is not included in our model, our results cannot be used to directly address this issue. This could imply a similar contrast between CLEAN and the two polluted cases in our simulations, by simply assuming the total drizzle sedimentation amount is proportional to the droplet size (i.e., inversely to the CDNC), though the quantity of such a perturbation seems rather small here, not to mention the more sophisticated feedback involved in our case introduced by the dynamic aerosol-cloud interaction in our model.~~

645
650
655

660 As demonstrated from above discussions that modeled cloud microphysical features generally respond to the variation of aerosol number concentration as expected, *i.e.*, higher aerosol concentration leads to higher cloud droplet number concentration (POL > REF > CLEAN) while smaller mean droplet radius (POL < REF < CLEAN) and hence a higher cloud reflectivity (POL > REF > CLEAN). Though exception does exist. For example, differences in the aerosol size distribution and chemical composition between REF and POL could lead to an outcome opposite to the general expectation particularly under a weak updraft. As shown in Fig. 10c, the response of the incoming solar radiation at ground (SWRADSURF) does not always follow such an expectation in cloud microphysics and thus reflectivity in responding to aerosol variation. In fact, SWRADSURF appears to be higher in POL than REF from sunrise to 13:00 UTC, and the values in both runs are also clearly higher than that in CLEAN. This tendency is only reversed after 13:00 UTC when solar flux reaches its peak until the break-up stage.

665



670

Figure 10. Evolution of cloud droplets concentration N_c (top) and cloud droplets radius r_c (middle) with the scenarios given and designated by letter a (REF), b (POL) and c (CLEAN). Bottom panel gives the evolution of mean domain SWRADSURF differences between POL or CLEAN and REF.

675




Figure 11. Liquid water path (LWP, $g\ kg^{-1}\ m$) in POL (left column), REF (mid-column), and CLEAN (right column) runs at 10:00 UTC (top row), 14:00 UTC (middle row), and 16:00 UTC (bottom row).

680 Figure 11 shows that the major reason behind the above-described trend of SWRADSURF is the
 difference in cloud fraction in competing with the effect brought by different cloud reflectivity in various
 runs, especially before noon when zenith angle is still high. After sunrise, the cloud top starts to rise and
 cloud layer becomes thicker. In the meantime, this upward development brings a downward entrainment
 of dry air from the temperature inversion zone above the cloud top and causes evaporation in the cloud.
 685 For a cloud with a large quantity of very small droplets as in POL and REF, the evaporation rate of
 droplets would exceed that in CLEAN case, thus more cloud-void spaces or a thinner cloud layer would
 form ~~much~~ easier than in the latter case. Note that a similar macrophysical response to aerosol

concentration variation (in a simple high versus low setting) was also suggested in a marine cloud case though with a coarse vertical resolution of 50 m (Wang *et al.*, 2003). As shown in Fig. 11 and Table 3, cloud layer in CLEAN is slightly denser than those in POL and REF while cloud-void or thin cloud pixels account for a substantially lower ratio within the domain. Thus, before noontime, cloud reflectivity seems to become the secondary factor comparing to cloud fraction in determining the value of SWRADSURF. As a result, SWRADSURF in CLEAN is significantly lower than REF and POL until zenith angle becomes lower closer to noontime. The lower SWRADSURF in CLEAN would also have reduced the turbulent mixing as well as delayed the convection that would cause extensive cloud break-up. At 14:00 UTC, differences in cloud thickness and cloud-void space still exist but become relatively smaller among the three different runs (Fig. 11 and Table 3), cloud reflectivity now becomes the primary reason to cause a different SWRADSURF as shown in Fig. 10 (bottom panel). ~~Interestingly, modeled clouds in POL and REF appear to dissipate earlier and much faster than in CLEAN in the break-up stage late afternoon, largely due to their smaller droplet sizes~~ (Fig. 11, bottom panel).

	LWP 10 UTC	PCP 10 UTC	LWP 14 UTC	PCP 14 UTC	LWP 16 UTC	PCP 16 UTC
POL	14.87	12.79	10.98	42.17	1.96	99.66
REF	15.67	10.11	11.34	42.69	2.74	99.67
CLEAN	16.98	6.95	11.79	44.93	4.12	94.47

Table 3. Domain averaged liquid water path (LWP; $g\ kg^{-1}\ m$) and poor-cloud pixel percentage (PCP, defined by the percentage of pixels where LWP < 10 $g\ kg^{-1}\ m$; percentage) in three different runs.

Looking into various temporally varying metrics of LWP in different model runs, we find that in general, LWP is inversely proportional to CDNC, as LWP in POL < LWP in REF < LWP in CLEAN, and this is applied to different metrics of LWP (Fig. 12, ref. ADEON curves; Table 3) as well. However, in comparison, the peak LWP varies less significantly in CLEAN case, while peak LWPs in two other runs decrease with domain averaged quantities in convection stage. ~~There were different opinions regarding the mechanisms behind such an inverse relation between LWP and CDNC (e.g., Ackerman *et al.*, 2004; Bretherton *et al.*, 2007), not to mention that most such hypotheses were proposed based on the cases of marine low clouds that might not be directly applied to the cases over land.~~ In our analysis, the difference in turbulent mixing driven by the surface radiative heating, as influenced by different microphysical features in various cases, seems having played a critical role in resulting the inverse relation between LWP and CDNC. The situation of cloud fraction (CF) is somewhat more complicated. As shown in Table 3 and Fig. A3, CF relation with CDNC varies in different stages. An inverse relation between CF and CDNC generally stands in the earlier and later period of the convection stage. This is primarily due to the faster evaporation of clouds with higher CDNC driven by entrainment in the former period (note the controlling role of CF in determining the surface incoming solar radiation and thus turbulence in this stage), or by strong convection in the latter. In the middle of the convection stage (13:00-15:00 UTC), the above relation, however, would reverse or become insignificant, owing to a weaker turbulent mixing in polluted cases since the cloud reflectivity becomes the dominant factor in controlling the surface

incoming solar radiation as discussed previously. Therefore, an analysis throughout the entire LLSC life cycle is very important to understand the response of CF alongside LWP to aerosol variation. Note that the atmospheric heating caused by absorbing black carbon aerosol is already included in this series of sensitivity simulations, though its impacts on the above result will be discussed later based on another set of sensitivity runs.




Figure 12. Domain averaged LWP (LWP mean), maximum LWP (LWP max), and domain averaged LWP over pixels where $LWP > 10 \text{ g kg}^{-1} \text{ m}$ in ADEON and ADEOFF runs in REF (upper panel), POL (middle panel), and CLEAN (lower panel) cases, derived using hourly model outputs.

To summarize, as expected, aerosol concentration is a major factor in controlling the cloud microphysical features by determining the simulated droplet number concentration and radius with similar liquid water content. However, our results suggest that cloud reflectivity as a function of CDNC is not necessarily a dominant factor to solely determine the surface incoming solar radiation. Instead, the response of cloud macrophysical features such as cloud fraction as well as LWP to the variation caused by dry entrainment from inversion layer above the cloud is also a competing factor in determining the incoming solar radiation at ground. Our sensitivity simulations utilize different aerosol profiles that reflect the variations in both aerosol concentration and chemical composition based on observations, the results

indicate a critical role of cloud microphysical response to aerosol in deciding the LWP and CF response. The overall negative response of LWP to aerosol concentration derived here agrees with several previous studies (e.g., [Aekerman-Wang et al., 2003](#); Jiang and Feingold, 2006). While the case for CF response is more complicated, varying in different stages in cloud life cycle. It is worth indicating though, another factor that might contribute to the cloud life cycle, *i.e.*, the atmospheric heating caused by the semi-direct effect of absorbing aerosol component such as black carbon has not been analyzed up to this moment and will be discussed in the following section.




4.2 Impact of aerosol semi-direct effect on LLSCs


The semi-direct effect of aerosols, resulted from SW radiation absorption by absorbing aerosol, could affect atmospheric dynamics surrounding LLSCs and thus their life cycle. To examine this effect, we have designed three additional sensitivity simulations, configured accordingly in the same way as their original experiments POL, REF, and CLEAN (hereafter ADEON of REF, POL, and CLEAN, respectively) but excluding aerosol direct effects (named ADEOFF). Therefore, comparison between the ADEOFF runs and their paired original ADEON runs provides information regarding the isolated impacts of the semi-direct effect on the LLSC life cycle for cases with different aerosol profiles. Apparently, BC is the major species behind the semi-direct effect in our case. The changes in cloud top and base, SWHR, and TKE due to aerosol absorption and associated feedbacks are shown in Figure 13. The results demonstrate that light-absorbing BC aerosols can cause a substantial atmospheric heating accompanied by a warming tendency near the top of LLSCs (Fig. 13b). At 14:00 UTC, the domain averaged heating due to BC aerosols (difference in SWHR between ADEON and ADEOFF) and a consequent cooling just above the cloud due mostly to the cloud top change are 12.16 K day^{-1} and $-13.14 \text{ K day}^{-1}$ in POL, and 7.71 K day^{-1} and -9.24 K day^{-1} in REF, respectively. In comparison, the atmospheric heating and associated cooling of 1.30 K day^{-1} and -2.25 K day^{-1} in CLEAN case are clearly insignificant. Accordingly, in ADEON runs, ~~more water vapor tends to condense onto cloud droplets under the higher relative humidity in the lower PBL and decreasing~~ turbulent mixing in PBL has been reduced (Fig. 13c, with a maximum decrease of $-0.18 \text{ m}^2 \text{ s}^{-2}$ for POL), leading to a decrease of the cloud top height, limiting entrainment and reducing incoming solar radiation at surface due to BC in-cloud absorption. The cloud top height reduction due to the semi-direct effect in two polluted cases POL and REF is quite substantial as shown in Figure 13a, where CTH in POL and REF has decreased by up to 100 and 70 meters due to the presence of BC, respectively. On the other hand, CBH has also increased about 20 meters in both cases before break-up, suggesting a thinner cloud layer owing to the semi-direct effect. In comparison, CTH, CBH, and thus cloud vertical extent ~~appear to be~~ less affected in CLEAN run due to its low BC content. Before break-up, in-cloud TKE just below the top of the heating layer has been reduced in some extent (Fig. 13c). On the other hand, due to a lower cloud top in the polluted cases, planetary boundary layer height would also be lowered. The effect of BC absorption in lowering modeled cloud top and thinning cloud layer in POL and REF (implying a reduced upward development) is likely another factor to slow down their break-up as discussed before.

The impact of the semi-direct effect on other critical macrophysical features such as cloud fraction and LWP can be also seen from the model results. For instance, LWP particularly the maximum LWP is clearly lower in the ADEON runs of the two polluted cases (REF and POL) (Fig. 12). In addition, an




increase of cloud fraction due to the semi-direct effect can be seen throughout the convection stage until 15:00 UTC when massive-intense cloud break-up occurs (Fig. A3). All these imply a critical role of the semi-direct effect on cloud radiation.


785





790 **Figure 13.** Evolution of the differences of the mean CBH and CTH (a), SWHR (b), and TKE (c) between the simulation runs with and without aerosol direct effect (ADEON-ADEOFF) for REF, POL and CLEAN.





795

Figure 14. The domain-mean differences of Mean difference surface SW radiative flux (SWRADSURF) between Clear Sky (CS) and cloudy scenarios giving the flux dissipated by clouds in ADEON (a) and ADEOFF (b) configurations. SWRADSURF difference between ADEON and ADEOFF configuration for the three scenarios (c).

800

We find that the semi-direct effect can both enhance and weaken the (negative) indirect radiative forcing as also indicated by some previous works (Lohmann and Feichter, 2001; Koch and Del Genio, 2010a; Huang *et al.*, 2014; Yamaguchi *et al.*, 2015; Stjern *et al.*, 2017; Kreidenweis *et al.*, 2019; Zhang and Zuidema, 2019). In the convection stage before 15:00 UTC, the difference in SWRADSURF between ADEON and ADEOFF is negative, reaching $-33 W m^{-2}$ and $-75 W m^{-2}$ for REF and POL at 14:00 UTC, respectively (Fig. 14e). This can be explained by an increase in cloud fraction in ADEON runs (Fig. A3, Table 3) that allows less solar irradiance to attain the surface despite the cloud layer being thinner, not to mention that solar irradiance itself has already been reduced due to BC absorption (Fig. 12, 14e and A2). Note that the different chemical compositions between POL and REF also lead to a quantitatively different effect. Hence, the semi-direct effect contributes positively to the enhancement of (negative) indirect radiative forcing in this case. On the other hand, at 16:00 UTC, the flux difference between ADEON and ADEOFF becomes positive with values for REF and POL as $32 W m^{-2}$ and $66 W m^{-2}$, respectively. As the clouds break up more slowly in ADEOFF during this stage due to thicker cloud layers (Fig. A2 and A3), more clouds inside the domain with increased thickness causes weaker SW irradiance reaching the ground. In other words, the semi-direct effect makes the cloud dissipate faster in the later convective stage. In this case, the semi-direct effect weakens the indirect radiative forcing.

805

810

815

The above results have demonstrated the important role of solar absorption by aerosols in determining the life cycle of LLSCs. Note that our modeling configurations are based on the aerosol profiles that are relatively well-mixed throughout the PBL then with concentration gradually decreasing

820 along altitude above PBL. Certain previous sensitivity experiments suggested that the location of BC
layer within or above PBL could have different impacts on the development of convection, entrainment,
and thus life cycle of the low clouds within PBL. For instance, Johnson *et al.* (2004) suggested that
without considering the indirect effect of aerosols, BC existing within boundary layer would lower LWP
by nearly 20% in a marine low stratocumulus case, where the cloud response is less sensitive to the
825 change in surface shortwave heating comparing to the situation in our case. Feingold *et al.* (2005) found
that smoke plumes containing BC near the surface would reduce the cloudiness through both the
atmospheric heating and the weakening effect on surface heat fluxes by BC. These results though
obtained with somewhat different model configurations than ours (*e.g.*, coarser vertical resolution,
different surface, etc.) are in a qualitative agreement with our findings. Nevertheless, the unique
830 configuration of our model allows us to quantitatively examine the semi-direct effect with varying aerosol
chemical compositions and thus extent of aerosol absorption. This has led us to reveal further insights of
the complicated interplays among various aerosol effects besides their individual impacts on the life cycle
of LLSCs.

5. Conclusions

835 An observed LLSC case over southern West Africa has been simulated with Meso-NH model in a
Large-Eddy Simulation configuration constrained by the measurements from DACCWA field campaign.
The model has successfully reproduced the observed nocturnal-to-diurnal life cycle alongside key macro-
and microphysical features as well as surface radiative and heat fluxes. To determine the impact of
aerosols on the modeled life cycle of LLSCs, sensitivity simulations using several different aerosol
840 profiles as well as the ones adopting these profiles but excluding the aerosol direct radiation effect
have also been conducted. These aerosol profiles contain different size distributions and chemical
compositions, reflecting the situations associated with various aerosol populations encountered during
the field campaign.

The results from sensitivity simulations suggest that both aerosol size distribution and chemical
845 composition can effectively influence the LLSCs life cycle. The impact of the aerosol size distribution,
as reflected from a comparison among simulations using aerosol profiles with different number
concentrations and modal distributions, is initiated from resultant cloud microphysical features in
particular the cloud droplet number concentration and mean droplet size. Such a difference created by
different aerosol size distributions also affect cloud reflectivity as expected. ~~Interestingly, w~~We have
850 found that the difference in cloud reflectivity caused by different aerosol concentration does not always
dominate the surface incoming solar radiation and thus cloud development after sunrise. This is due to a
competing factor: the difference in cloud fraction resulted from different evaporation speed of cloud
droplets (a function of CDNC) due to the dry air entrained from the inversion layer above cloud top,
which specifically dominates the variation of surface incoming solar radiation before noontime. Clouds
855 influenced by higher aerosol concentrations and thus having higher number concentration and smaller
sizes of cloud droplets ~~are found to~~ evaporate more easily and ~~thus impose~~this can lead to a lower cloud
fraction. For the same reason, clouds with higher droplet concentration are likely to break up earlier.

In addition, our sensitivity runs including versus excluding aerosol direct radiative effects have also
demonstrated the impact specifically of solar absorption by black carbon on the cloud life cycle. The

860 excessive atmospheric heating reaching 12 K day^{-1} introduced by black carbon in our modeled cases is found to be able to lower the cloud top height as well as liquid water path, reduce dry entrainment, and increase cloud fraction. Working with the cloud fraction response to aerosol size distribution, this heating and its consequences might delay break-up of the LLSCs until late afternoon. All these would enhance the aerosol indirect effect. On the other hand, the modeled clouds in polluted cases with higher aerosol
865 concentrations and BC content would break up faster in late afternoon due to their thinner cloud layers. In this case the semi-direct effect would weaken the indirect effect.

Our study has demonstrated that the life cycle and thus the radiative forcing of LLSCs over land area of SWA can be substantially influenced by aerosols from both long-range transported biomass burning plumes and from local anthropogenic emissions. In fact, more aerosol profiles had been
870 collected during DACCIWA campaign besides the ones used in this study. Future research works could reveal the aerosol impact under an even broader range of aerosol properties and to examine the temporal variations of LLSCs radiative effects evolving with different large-scale meteorological conditions with different associated air mass. More analysis on different cloud cases in SWA would also be able to assess or refute current results on semi-direct effect.

875

Code and data availability. The data obtained during DACCIWA campaign at the Savè supersite alongside all other data used in this study are publicly available on the SEDOO database
880 (<http://baobab.sedoo.fr/DACCIWA/>). The Meso-NH code is maintained and updated by LAERO and CNRM, it is freely available for download at <http://mesonh.aero.obs-mip.fr/mesonh52/>.

Author contributions. LD and CW designed the simulations and LD conducted model simulations and data analyses. LD and CW wrote this paper with contribution from all other co-authors. CW advised and helped LD to better understand the different aspects of this research work. PT advised and trained LD to
885 use Meso-NH and ORILAM module. CD processed and provided the aerosol profiles used in previous simulations and NM was part of this work. MZ helped to select the study case and advised during the study case construction and analysis. AD brought a critical eye to this work.

890 **Competing interests.** The authors declare that they have no conflict of interest.

Acknowledgements. This study is supported by L'Agence National de la Recherche (ANR) of France under "Programme d'Investissements d'Avenir" (ANR-18-MPGA-003 EUROACE) and co-funded by the University of Toulouse III Paul Sabatier. The computation of this work was performed using HPC
895 resources of French GENCI-IDRIS (Grant A0110110967 and A0090110967) and French Regional Computations center CALMIP. LD thanks the Laboratoire d'Aérodologie, Université de Toulouse, France, for funding and hosting his Ph.D. research activities. LD also thanks the Meso-NH team, especially Quentin Rodier, Juan Escobar, and Philippe Wautelet, for their advises on using Meso-NH, Benoit Vié and Marie Mazoyer for their help to handle and modify microphysical scheme LIMA, Quentin Libois for
900 explaining the details of Meso-NH's radiative schemes, and specifically Fabienne Lohou (LAERO) for

her introduction of DACCIWA campaign alongside her guidance in using relevant data products. A special thanks the authors to all people whose work was involved in the measurement and processing of DACCIWA campaign data especially over the Savè supersite. Many constructive comments and suggestions from Dr. Mónica Zamora Zapata and an anonymous reviewer as well as the handling editor, 905 Dr. Graham Feingold have made a substantial impact on our effort to improve the quality of the manuscript.




Figure A1. Mass composition (a,c) and size distribution provided by (Denjean et al., 2020a) and fitted into 2 modes described in Table 2 (b,d) for scenarios POL (top), CLEAN (bottom).




Figure A2. Liquid water path (LWP, $g\ kg^{-1}\ m$) in POL (left column), REF (mid-column), and CLEAN (right column) ADEOFF runs at 10:00 UTC (top row), 14:00 UTC (middle row), and 16:00 UTC (bottom row).




Figure A3. Domain averaged cloud fraction for AODON (solid lines) and AODOFF (dotted lines) of REF (red), POL (green), and CLEAN (blue) cases, derived using hourly model outputs. The cloud fraction here is a column quantity, defined as pixels where $LWP > 5 \text{ g kg}^{-1} \text{ m}$.

930

References

- Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types, *J. Geophys. Res.-Atmos.*, 105, 6837–6844, <https://doi.org/https://doi.org/10.1029/1999JD901161>, 2000.
- 935 Abdul-Razzak, H. and Ghan, S. J.: Parameterization of the influence of organic surfactants on aerosol activation, *J. Geophys. Res.-Atmos.*, 109, <https://doi.org/https://doi.org/10.1029/2003JD004043>, 2004.
- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03174>, 2004.
- 940 Adler, B., Babic, K., Kalthoff, N., Lohou, F., Lothon, M., Dione, C., Pedruzo-Bagazgoitia, X., and Andersen, H.: Nocturnal low-level' clouds in the atmospheric boundary layer over southern West Africa: an observation-based analysis of conditions and processes, *Atmos. Chem. Phys.*, 19, 663–681, <https://doi.org/10.5194/acp-19-663-2019>, 2019.
- Aouizerats, B., Thouron, O., Tulet, P., Mallet, M., Gomes, L., and Henzing, J. S.: Development of an online radiative module for the computation of aerosol optical properties in 3-D atmospheric models: validation during the EUCAARI campaign, *Geoscientific Model Development*, 3, 553–564, <https://doi.org/10.5194/gmd-3-553-2010>, 2010.
- 945 Babic, K., Adler, B., Kalthoff, N., Andersen, H., Dione, C., Lohou, F., Lothon, M., and Pedruzo-Bagazgoitia, X.: The observed diurnal cycle of low-level stratus clouds over southern West Africa: a case study, *Atmos. Chem. Phys.*, 19, 1281–1299, <https://doi.org/10.5194/acp-19-1281-2019>, 2019.
- Bauer, S. E., Im, U., Mezuman, K., and Gao, C. Y.: Desert Dust, Industrialization, and Agricultural Fires: Health Impacts of Outdoor Air Pollution in Africa, *J. Geophys. Res.-Atmos.*, 124, 4104–4120, <https://doi.org/https://doi.org/10.1029/2018JD029336>, 2019.
- 950

- Bellon, G. and Stevens, B.: Time Scales of the Trade Wind Boundary Layer Adjustment, *J. Atmos. Sci.*, 70, 1071 – 1083, <https://doi.org/10.1175/JAS-D-12-0219.1>, 2013.
- 955 Boucher, O., D. Randall, P. Artaxo, C. Bretherton, G. Feingold, P. Forster, V.-M. Kerminen, Y. Kondo, H. Liao, U. Lohmann, P. Rasch, S.K. Satheesh, S. Sherwood, B. Stevens, and X.Y. Zhang: Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 960 Bretherton, C. S., Blossy, F. N., and Uchida, J. : Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo, *Geophys. Res. Lett.*, 34, L03813, doi:10.1029/2006GL027648, 2007.
- Brito, J., Freney, E., Dominutti, P., Borbon, A., Haslett, S. L., Batenburg, A. M., Colomb, A., Dupuy, R., Denjean, C., Burnet, F., Bourriane, T., Deroubaix, A., Sellegri, K., Borrmann, S., Coe, H., Flamant, C., Knippertz, P., and Schwarzenboeck, A.: Assessing the role of anthropogenic and biogenic sources on PM₁ over southern West Africa using aircraft measurements, *Atmos. Chem. Phys.*, 18, 757–772, <https://doi.org/10.5194/acp-18-757-2018>, 2018.
- 965 Caniaux, G., Redelsperger, J.-L., and Lafore, J.-P.: A Numerical Study of the Stratiform Region of a Fast-Moving Squall Line. Part I: General Description and Water and Heat Budgets, *J. Atmos. Sci.*, 51, 2046 – 2074, [https://doi.org/10.1175/15200469\(1994\)051<2046:ANSOTS>2.0.CO;2](https://doi.org/10.1175/15200469(1994)051<2046:ANSOTS>2.0.CO;2), 1994.
- 970 Carslaw, K. S., Gordon, H., Hamilton, D. S., Johnson, J. S., Regayre, L. A., Yoshioka, M., and Pringle, K. J.: Aerosols in the Pre-industrial Atmosphere, *Current Climate Change Reports*, 3, 1–15, <https://doi.org/10.1007/s40641-017-0061-2>, 2017.
- Chatfield, R. B., Vastano, J. A., Li, L., Sachse, G. W., and Connors, V. S.: The Great African Plume from biomass burning: Generalizations from a three-dimensional study of TRACE A carbon monoxide, *J. Geophys. Res.-Atmos.*, 103, 28059–28077, <https://doi.org/https://doi.org/10.1029/97JD03363>, 1998.
- 975 Chen, T., Rossow, W. B., and Zhang, Y.: Radiative Effects of Cloud-Type Variations, *J. Clim.*, 13, 264 – 286, [https://doi.org/10.1175/1520-0442\(2000\)013<0264:REOCTV>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0264:REOCTV>2.0.CO;2), 2000.
- Cleveland, W. S.: Robust Locally Weighted Regression and Smoothing Scatterplots, *Journal of the American Statistical Association*, 74, 829–836, <https://doi.org/10.1080/01621459.1979.10481038>, 1979.
- Cohard, J.-M. and Pinty, J.-P.: A comprehensive two-moment warm microphysical bulk scheme. I: Description and tests, *Quar. J. Roy. Meteorol. Soc.*, 126, 1815–1842, <https://doi.org/https://doi.org/10.1002/qj.49712656613>, 2000.
- 980 Dearden, C., Hill, A., Coe, H., and Choularton, T.: The role of droplet sedimentation in the evolution of low-level clouds over southern West Africa, *Atmos. Chem. Phys.*, 18, 14253–14269, <https://doi.org/10.5194/acp-18-14253-2018>, 2018.
- 985 Deetz, K., Vogel, H., Knippertz, P., Adler, B., Taylor, J., Coe, H., Bower, K., Haslett, S., Flynn, M., Dorsey, J., Crawford, I., Kottmeier, C., and Vogel, B.: Numerical simulations of aerosol radiative effects and their impact on clouds and atmospheric dynamics over southern West Africa, *Atmos. Chem. Phys.*, 18, 9767–9788, <https://doi.org/10.5194/acp-18-9767-2018>, 2018.
- Denjean, C., Bourriane, T., Burnet, F., Mallet, M., Maury, N., Colomb, A., Dominutti, P., Brito, J., Dupuy, R., Sellegri, K., Schwarzenboeck, A., Flamant, C., and Knippertz, P.: Overview of aerosol optical properties over southern West Africa from DACCIIWA aircraft measurements, *Atmos. Chem. Phys.*, 20, 4735–4756, <https://doi.org/10.5194/acp-20-4735-2020>, 2020a.
- 990 Denjean, C., Brito, J., Libois, Q., Mallet, M., Bourriane, T., Burnet, F., Dupuy, R., Flamant, C., and Knippertz, P.: Unexpected Biomass Burning Aerosol Absorption Enhancement Explained by Black Carbon Mixing State, *Geophys. Res. Lett.*, 47, e2020GL089055, <https://doi.org/https://doi.org/10.1029/2020GL089055>, e2020GL089055, 2020b.
- 995 Deroubaix, A., Menut, L., Flamant, C., Brito, J., Denjean, C., Dreiling, V., Fink, A., Jambert, C., Kalthoff, N., Knippertz, P., Ladkin, R., Mailler, S., Maranan, M., Pacifico, F., Pigué, B., Siour, G., and Turquety, S.: Diurnal cycle of coastal anthropogenic pollutant transport over southern West Africa during the DACCIIWA campaign, *Atmos. Chem. Phys.*, 19, 473–497, <https://doi.org/10.5194/acp-19-473-2019>, 2019.
- Deroubaix, A., Menut, L., Flamant, C., Knippertz, P., Fink, A. H., Batenburg, A., Brito, J., Denjean, C., Dione, C., Dupuy, R., Hahn, V., Kalthoff, N., Lohou, F., Schwarzenboeck, A., Siour, G., Tuccella, P., and Voigt, C.: Sensitivity of low-

- level clouds and precipitation to anthropogenic aerosol emission in southern West Africa: a DACCIWA case study, *Atmos. Chem. Phys.*, 22, 3251–3273, <https://doi.org/10.5194/acp-22-3251-2022>, 2022.
- Derrien, S., Bezombes, Y., Bret, B., Gabella, O., Jarnot, C., Medina, P., Piques, E., Delon, C., Dione, C., Campistron, B., Durand, P., Jambert, C., Lohou, F., Lothon, M., Pacifico, F., and Meyerfeld, Y.: DACCIWA field campaign, Savè super-site, UPS instrumentation, 10.6096/DACCIWA.1618, 2016.
- 1005 Dione, C., Lohou, F., Lothon, M., Adler, B., Babic, K., Kalthoff, N., Pedruzo-Bagazgoitia, X., Bezombes, Y., and Gabella, O.: Low-level stratiform clouds and dynamical features observed within the southern West African monsoon, *Atmos. Chem. Phys.*, 19, 8979–8997, <https://doi.org/10.5194/acp-19-8979-2019>, 2019.
- Eastman, R. and Warren, S. G.: Diurnal Cycles of Cumulus, Cumulonimbus, Stratus, Stratocumulus, and Fog from Surface Observations over Land and Ocean, *J. Clim.*, 27, 2386 – 2404, <https://doi.org/10.1175/JCLI-D-13-00352.1>, 2014.
- 1010 Feingold, G., Jiang, H. L., and Harrington, J. Y.: On smoke suppression of clouds in Amazonia, *Geophys. Res. Lett.*, 32, 804, doi: 10.1029/2004GL021369, 2005.
- Flamant, C., Deroubaix, A., Chazette, P., Brito, J., Gaetani, M., Knippertz, P., Fink, A. H., de Coetlogon, G., Menut, L., Colomb, A., Denjean, C., Meynadier, R., Rosenberg, P., Dupuy, R., Dominutti, P., Duplissy, J., Bourriane, T., Schwarzenboeck, A., Ramonet, M., and Totems, J.: Aerosol distribution in the northern Gulf of Guinea: local anthropogenic sources, long-range transport, and the role of coastal shallow circulations, *Atmos. Chem. Phys.*, 18, 12363–12389, <https://doi.org/10.5194/acp-18-12363-2018>, 2018.
- 1015 Flossmann, A. I. and Wobrock, W.: Cloud Processing of Aerosol Particles in Marine Stratocumulus Clouds, *Atmosphere*, 10, <https://doi.org/10.3390/atmos10090520>, 2019.
- Fouquart, Y. and Bonnel, B.: Computations of solar heating of the earth’s atmosphere – A new parameterization, *Beitrag zur Physik der Atmosphäre*, 53, 35-62, 1980.
- 1020 Geoffroy, O., Brenguier, J.-L., and Sandu, I.: Relationship between drizzle rate, liquid water path and droplet concentration at the scale of a stratocumulus cloud system, *Atmos. Chem. Phys.*, 8, 4641–4654, <https://doi.org/10.5194/acp-8-4641-2008>, 2008.
- Ghonima, M., Heus, T., Norris, J. R., and J. Kleissl, J.: Factors controlling stratocumulus cloud lifetime over coastal land, *J. Atmos. Sci.*, 73, 2961-2983, 2016.
- 1025 Griffin, R. J., Nguyen, K., Dabdub, D., and Seinfeld, J. H.: A Coupled Hydrophobic-Hydrophilic Model for Predicting Secondary Organic Aerosol Formation, *J. Atmos. Chem.*, 44, 171–190, <https://doi.org/10.1023/A:1022436813699>, 2003.
- Hagan, H. D., Thompson, B., Palmo, J., and Ruskowski, A.: py-smps, <https://github.com/quant-aa/py-smps>, 2022.
- 1030 Handwerker, J., Scheer, S., and Gamer, T.: DACCIWA field campaign, Savè super-site, Cloud and precipitation, <https://doi.org/10.6096/daccciwa.1686>, 2016.
- Hannak, L., Knippertz, P., Fink, A. H., Kniffka, A., and Pante, G.: Why Do Global Climate Models Struggle to Represent Low-Level Clouds in the West African Summer Monsoon?, *J. Clim.*, 30, 1665 – 1687, <https://doi.org/10.1175/JCLI-D-16-0451.1>, 2017.
- 1035 Hansen, J., M. Sato, R. Ruedy, A. Lacis, and V. Oinas, Global warming in the twenty-first century: An alternative scenario. *PNAS*, 97, 9875-9880, 1998.
- Hartmann, D. L., Ockert-Bell, M. E., and Michelsen, M. L.: The Effect of Cloud Type on Earth’s Energy Balance: Global Analysis, *J. Clim.*, 5, 1281 – 1304, [https://doi.org/10.1175/1520-0442\(1992\)005<1281:TEOCTO>2.0.CO;2](https://doi.org/10.1175/1520-0442(1992)005<1281:TEOCTO>2.0.CO;2), 1992.
- 1040 Haslett, S. L., Taylor, J. W., Evans, M., Morris, E., Vogel, B., Dajuma, A., Brito, J., Batenburg, A. M., Borrmann, S., Schneider, J., Schulz, C., Denjean, C., Bourriane, T., Knippertz, P., Dupuy, R., Schwarzenböck, A., Sauer, D., Flamant, C., Dorsey, J., Crawford, I., and Coe, H.: Remote biomass burning dominates southern West African air pollution during the monsoon, *Atmos. Chem. Phys.*, 19, 15217–15234, <https://doi.org/10.5194/acp-19-15217-2019>, 2019.
- Haywood, J. and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review, *Rev. Geophys.*, 38, 513–543, <https://doi.org/https://doi.org/10.1029/1999RG000078>, 2000.
- 1045 Hill, P. G., Allan, R. P., Chiu, J. C., Bodas-Salcedo, A., and Knippertz, P.: Quantifying the Contribution of Different Cloud Types to the Radiation Budget in Southern West Africa, *J. Clim.*, 31, 5273 – 5291, <https://doi.org/10.1175/JCLI-D-17-0586.1>, 2018.

- 1050 Huang, J., Wang, T., Wang, W., Li, Z., and Yan, H.: Climate effects of dust aerosols over East Asian arid and semiarid regions, *J. Geophys. Res.-Atmos.*, 119, 11,398–11,416, <https://doi.org/https://doi.org/10.1002/2014JD021796>, 2014.
- Jiang, G.-S. and Shu, C.-W.: Efficient Implementation of Weighted ENO Schemes, *J. Comp. Phys.*, 126, 202–228, <https://doi.org/https://doi.org/10.1006/jcph.1996.0130>, 1996.
- Jiang, H., Cotton, W. R., and Feingold, G.: Simulations of aerosol-cloud-dynamical feedbacks resulting from entrainment of aerosol into the marine boundary layer during the Atlantic Stratocumulus Transition Experiment, *J. Geophys. Res.*, 107, 4813, doi:10.1029/2001JD001502, 2002.
- 1055 Jiang, H., and Feingold, G.: Effect of aerosol on warm convective clouds: Aerosol-cloud-surface flux feedbacks in a new coupled large eddy model, *J. Geophys. Res.*, 111, doi:10.1029/2005JD006138, 2006.
- Johnson, B. T., Shine, K. P., and Forster, P. M.: The semi-direct aerosol effect: Impact of absorbing aerosols on marine stratocumulus, *Q. J. R. Meteorol. Soc.*, 130, 1407-1422, 2004.
- 1060 Kalthoff, N., Lohou, F., Brooks, B., Jegede, G., Adler, B., Babic, K., Dione, C., Ajao, A., Amekudzi, L. K., Aryee, J. N. A., Ay'oola, M., Bessardon, G., Danuor, S. K., Handwerker, J., Kohler, M., Lathon, M., Pedruzo-Bagazgoitia, X., Smith, V., Sunmonu, L., Wieser, A., Fink, A. H., and Knippertz, P.: An overview of the diurnal cycle of the atmospheric boundary layer during the West African monsoon season: results from the 2016 observational campaign, *Atmos. Chem. Phys.*, 18, 2913–2928, <https://doi.org/10.5194/acp-18-2913-2018>, 2018.
- 1065 Khairoutdinov, M. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Mon. Weather Rev.*, 128, 229 – 243, [https://doi.org/10.1175/1520-0493\(2000\)128<0229:ANCPPI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0229:ANCPPI>2.0.CO;2), 2000.
- Knippertz, P., Fink, A. H., Schuster, R., Trentmann, J., Schrage, J. M., and Yorke, C.: Ultra-low clouds over the southern West African monsoon region, *Geophys. Res. Lett.*, 38, <https://doi.org/https://doi.org/10.1029/2011GL049278>, 2011.
- 1070 Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Liousse, C., Mari, C., Allan, R. P., Brooks, B., Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCWA Project: Dynamics–Aerosol–Chemistry–Cloud Interactions in West Africa, *Bull. Amer. Meteor. Soc.*, 96, 1451 – 1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, 2015.
- 1075 Knippertz, P., Fink, A. H., Deroubaix, A., Morris, E., Tocquer, F., Evans, M. J., Flamant, C., Gaetani, M., Lavaysse, C., Mari, C., Marsham, J. H., Meynadier, R., Affo-Dogo, A., Bahaga, T., Brosse, F., Deetz, K., Guebsi, R., Latifou, I., Maranan, M., Rosenberg, P. D., and Schlueter, A.: A meteorological and chemical overview of the DACCWA field campaign in West Africa in June–July 2016, *Atmos. Chem. Phys.*, 17, 10893–10918, <https://doi.org/10.5194/acp-17-10893-2017>, 2017.
- Koch, D. and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and synthesis, *Atmos. Chem. Phys.*, 10, 7685–7696, <https://doi.org/10.5194/acp-10-7685-2010>, 2010a.
- 1080 Koch, D. and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and synthesis, *Atmos. Chem. Phys.*, 10, 7685–7696, <https://doi.org/10.5194/acp-10-7685-2010>, 2010b.
- Kreidenweis, S. M., Petters, M., and Lohmann, U.: 100 Years of Progress in Cloud Physics, Aerosols, and Aerosol Chemistry Research, *Meteorological Monographs*, 59, 11.1 – 11.72, <https://doi.org/10.1175/AMSMONOGRAPHS-D-18-0024.1>, 2019.
- 1085 Lac, C., Chaboureau, J.-P., Masson, V., Pinty, J.-P., Tulet, P., Escobar, J., Leriche, M., Barthe, C., Aouizerats, B., Augros, C., Aumond, P., Auguste, F., Bechtold, P., Berthet, S., Bielli, S., Bosseur, F., Caumont, O., Cohard, J.-M., Colin, J., Couvreux, F., Cuxart, J., Delautier, G., Dauhut, T., Ducrocq, V., Filippi, J.-B., Gazen, D., Geoffroy, O., Gheusi, F., Honnert, R., Lafore, J.-P., Lebeau-pin Brossier, C., Libois, Q., Lunet, T., Mari, C., Maric, T., Mascart, P., Mogé, M., Molinié, G., Nuissier, O., Pantillon, F., Peyrillé, P., Pergaud, J., Perraud, E., Pianezze, J., Redelsperger, J.-L., Ricard, D., Richard, E., Riette, S., Rodier, Q., Schoetter, R., Seyfried, L., Stein, J., Suhre, K., Taufour, M., Thouron, O., Turner, S., Verrelle, A., Vié, B., Visentin, F., Vionnet, V., and Wautelet, P.: Overview of the Meso-NH model version 5.4 and its applications, *Geoscientific Model Development*, 11, 1929–1969, <https://doi.org/10.5194/gmd-11-1929-2018>, 2018.
- 1095 Lascaux, F., Richard, E., and Pinty, J.-P.: Numerical simulations of three different MAP IOPs and the associated microphysical processes, *Quart. J. Roy. Meteorol. Soc.*, 132, 1907–1926, <https://doi.org/https://doi.org/10.1256/qj.05.197>, 2006.

- 1100 Legain, D., Bousquet, O., Douffet, T., Tzanos, D., Moulin, E., Barrie, J., and Renard, J.-B.: High-frequency boundary layer profiling with reusable radiosondes, *Atmospheric Measurement Techniques*, 6, 2195–2205, <https://doi.org/10.5194/amt-6-2195-2013>, 2013.
- Li, J., Carlson, B. E., Yung, Y. L., Lv, D., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, V., Kahn, R. A., Zhang, P., Dubovik, O., Ding, A., Lacis, A. A., Zhang, L., and Dong, Y.: Scattering and absorbing aerosols in the climate system, *Nature Reviews Earth & Environment*, 3, 363–379, <https://doi.org/10.1038/s43017-022-00296-7>, 2022.
- 1105 Liousse, C., Assamoi, E., Criqui, P., Granier, C., and Rosset, R.: Explosive growth in African combustion emissions from 2005 to 2030, *Environ. Res. Lett.*, 9, 035003, <https://doi.org/10.1088/1748-9326/9/3/035003>, 2014.
- Liu, Y., Jia, R., Dai, T., Xie, Y., and Shi, G.: A review of aerosol optical properties and radiative effects, *Journal of Meteorological Research*, 28, 1003–1028, <https://doi.org/10.1007/s13351-014-4045-z>, 2014.
- Lohmann, U. and Feichter, J.: Can the direct and semi-direct aerosol effect compete with the indirect effect on a global scale?, *Geophys. Res. Lett.*, 28, 159–161, <https://doi.org/https://doi.org/10.1029/2000GL012051>, 2001.
- 1110 Lohou, F., Kalthoff, N., Adler, B., Babic, K., Dione, C., Lathon, M., Pedruzo-Bagazgoitia, X., and Zouzoua, M.: Conceptual model of diurnal cycle of low-level stratiform clouds over southern West Africa, *Atmos. Chem. Phys.*, 20, 2263–2275, <https://doi.org/10.5194/acp-20-2263-2020>, 2020.
- Lunet, T., Lac, C., Auguste, F., Visentin, F., Masson, V., and Escobar, J.: Combination of WENO and Explicit Runge- Kutta methods for wind transport in Meso-NH model, *Mon. Weather Rev.*, 145, 3817–3838, <https://doi.org/10.1175/MWR-D616-0343.12017>, 2017.
- 1115 Mari, C., Evans, M. J., Palmer, P. I., Jacob, D. J., and Sachse, G. W.: Export of Asian pollution during two cold front episodes of the TRACE-P experiment, *J. Geophys. Res.-Atmos.*, 109, <https://doi.org/https://doi.org/10.1029/2003JD004307>, 2004.
- Mari, C. H., Cailley, G., Corre, L., Saunois, M., Attié, J. L., Thouret, V., and Stohl, A.: Tracing biomass burning plumes from the Southern Hemisphere during the AMMA 2006 wet season experiment, *Atmos. Chem. Phys.*, 8, 3951–3961, <https://doi.org/10.5194/acp-8-3951-2008>, 2008.
- Marticorena, B. and Bergametti, G.: Two-year simulations of seasonal and interannual changes of the Saharan dust emissions, *Geophys. Res. Lett.*, 23, 1921–1924, <https://doi.org/https://doi.org/10.1029/96GL01432>, 1996.
- 1120 Martin, G. M., Johnson, D. W., and Spice, A.: The Measurement and Parameterization of Effective Radius of Droplets in Warm Stratocumulus Clouds, *J. Atmos. Sci.*, 51, 1823 – 1842, [https://doi.org/10.1175/1520-0469\(1994\)051<1823:TMAPOE>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<1823:TMAPOE>2.0.CO;2), 1994.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geoscientific Model Development*, 6, 929–960, <https://doi.org/10.5194/gmd-6-929-2013>, 2013.
- 1130 Menut, L., Flamant, C., Turquety, S., Deroubaix, A., Chazette, P., and Meynadier, R.: Impact of biomass burning on pollutant surface concentrations in megacities of the Gulf of Guinea, *Atmos. Chem. Phys.*, 18, 2687–2707, <https://doi.org/10.5194/acp18-2687-2018>, 2018.
- Menut, L., Tuccella, P., Flamant, C., Deroubaix, A., and Gaetani, M.: The role of aerosol–radiation–cloud interactions in linking anthropogenic pollution over southern west Africa and dust emission over the Sahara, *Atmos. Chem. Phys.*, 19, 14657–14676, <https://doi.org/10.5194/acp-19-14657-2019>, 2019.
- 1140 Metzger, S., Dentener, F., Pandis, S., and Lelieveld, J.: Gas/aerosol partitioning: 1. A computationally efficient model, *J. Geophys. Res.-Atmos.*, 107, ACH 16–1–ACH 16–24, <https://doi.org/https://doi.org/10.1029/2001JD001102>, 2002.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.-Atmos.*, 102, 16663–16682, <https://doi.org/https://doi.org/10.1029/97JD00237>, 1997.
- 1145 Morcrette, J.-J.: The Surface Downward Longwave Radiation in the ECMWF Forecast System, *J. Clim.*, 15, 1875 – 1892, [https://doi.org/10.1175/1520-0442\(2002\)015<1875:TSDLRI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1875:TSDLRI>2.0.CO;2), 2002.

- Murphy, J. G., Oram, D. E., and Reeves, C. E.: Measurements of volatile organic compounds over West Africa, *Atmos. Chem. Phys.*, 10, 5281–5294, <https://doi.org/10.5194/acp-10-5281-2010>, 2010.
- Noilhan, J. and Planton, S.: A Simple Parameterization of Land Surface Processes for Meteorological Models, *Mon. Weather Rev.*, 117, 536 – 549, [https://doi.org/10.1175/1520-0493\(1989\)117<0536:ASPOLS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<0536:ASPOLS>2.0.CO;2), 1989.
- 1150 Pedruzo-Bagazgoitia, X., de Roode, S. R., Adler, B., Babic, K., Dione, C., Kalthoff, N., Lohou, F., Lothon, M., and Vilà-Guerau de Arellano, J.: The diurnal stratocumulus-to-cumulus transition over land in southern West Africa, *Atmos. Chem. Phys.*, 20, 2735–2754, <https://doi.org/10.5194/acp-20-2735-2020>, 2020.
- Petters, J. L., Harrington, J. Y., and Clothiaux, E. E.: Radiative–dynamical feedbacks in low liquid water path stratiform clouds, *J. Atmos. Sci.*, 69, 1498–1512, <https://doi.org/10.1175/JAS-D-11-0169.1>, 2012
- 1155 Pinty, J.-P. and Jabouille, P.: A mixed-phased cloud parameterization for use in a mesoscale non-hydrostatic model: simulations of a squall line and of orographic precipitation, *Proc. Conf. on Cloud Physics*, 217–220. 1998.
- Pruppacher, H. R., Klett, J. D., and Wang, P. K.: Microphysics of Clouds and Precipitation, *Aerosol Science and Technology*, 28, 381–382, <https://doi.org/10.1080/02786829808965531>, 1998.
- 1160 Reeves, C. E., Formenti, P., Afif, C., Ancellet, G., Attié, J.-L., Bechara, J., Borbon, A., Cairo, F., Coe, H., Crumeyrolle, S., Fierli, F., Flamant, C., Gomes, L., Hamburger, T., Lambert, C., Law, K. S., Mari, C., Jones, R. L., Matsuki, A., Mead, M. I., Methven, J., Mills, G. P., Minikin, A., Murphy, J. G., Nielsen, J. K., Oram, D. E., Parker, D. J., Richter, A., Schlager, H., Schwarzenboeck, A., and Thouret, V.: Chemical and aerosol characterisation of the troposphere over West Africa during the monsoon period as part of AMMA, *Atmos. Chem. Phys.*, 10, 7575–7601, <https://doi.org/10.5194/acp-10-7575-2010>, 2010.
- 1165 Sandu, I., Brenguier, J.-L., Geoffroy, O., Thouron, O., and Masson, V.: Aerosol Impacts on the Diurnal Cycle of Marine Stratocumulus, *J. Atmos. Sci.*, 65, 2705 – 2718, <https://doi.org/10.1175/2008JAS2451.1>, 2008.
- Sauvage, B., Thouret, V., Cammas, J.-P., Gheusi, F., Athier, G., and Nédélec, P.: Tropospheric ozone over Equatorial Africa: regional aspects from the MOZAIC data, *Atmos. Chem. Phys.*, 5, 311–335, <https://doi.org/10.5194/acp-5-311-2005>, 2005.
- 1170 Schrage, J. M. and Fink, A. H.: Nocturnal Continental Low-Level Stratus over Tropical West Africa: Observations and Possible Mechanisms Controlling Its Onset, *Mon. Weather Rev.*, 140, 1794 – 1809, <https://doi.org/10.1175/MWR-D-11-00172.1>, 2012.
- Schuster, R., Fink, A. H., and Knippertz, P.: Formation and maintenance of nocturnal low-level stratus over the Southern West African monsoon region during AMMA 2006, *J. Atmos. Sci.*, 70, 2337 – 2355, <https://doi.org/10.1175/JAS-D-120241.1>, 2013.
- 1175 Stevens, B., Cotton, W. R., Feingold, G., and Moeng, C.-H.: Large-eddy simulations of strongly precipitating, shallow, stratocumulus-topped boundary layers, *J. Atmos. Sci.*, 55, 3616–3638, 1998.
- 1180 Stevens, B., Moeng, C.-H., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J.-C., Jiang, H., Khairoutdinov, M., Kirkpatrick, M. P., Lewellen, D. C., Lock, A., Müller, F., Stevens, D. E., Whelan, E., and Zhu, P.: Evaluation of large-eddy simulations via observations of nocturnal marine stratocumulus, *Mon. Weather Rev.*, 133, 1443–1462, <https://doi.org/10.1175/MWR2930.1>, 2005.
- 1185 Stjern, C. W., Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, , Andrews, T., Boucher, O., Faluvegi, G., Iversen, T., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivić, D., Richardson, T., Shawki, D., Shindell, D., Smith, C. J., Takemura, T., and Voulgarakis, A.: Rapid adjustments cause weak surface temperature response to increased black carbon concentrations, *J. Geophys. Res.-Atmos.*, 122, 11,462–11,481, <https://doi.org/https://doi.org/10.1002/2017JD027326>, 2017.
- Taylor, J. W., Haslett, S. L., Bower, K., Flynn, M., Crawford, I., Dorsey, J., Choularton, T., Connolly, P. J., Hahn, V., Voigt, C., Sauer, D., Dupuy, R., Brito, J., Schwarzenboeck, A., Bourriane, T., Denjean, C., Rosenberg, P., Flamant, C., Lee, J. D., Vaughan, A. R., Hill, P. G., Brooks, B., Catoire, V., Knippertz, P., and Coe, H.: Aerosol influences on low-level clouds in the West African monsoon, *Atmos. Chem. Phys.*, 19, 8503–8522, <https://doi.org/10.5194/acp-19-8503-2019>, 2019.
- 1190 Thouron, O., Brenguier, J.-L., and Burnet, F.: Supersaturation calculation in large eddy simulation models for prediction of the droplet number concentration, *Geoscientific Model Development*, 5, 761–772, <https://doi.org/10.5194/gmd-5-761-2012>, 2012.
- 1195

- Tulet, P., Crassier, V., Solmon, F., Guedalia, D., and Rosset, R.: Description of the mesoscale nonhydrostatic chemistry model and application to a transboundary pollution episode between northern France and southern England, *J. Geophys. Res.-Atmos.*, 108, ACH 5–1–ACH 5–11, <https://doi.org/https://doi.org/10.1029/2000JD000301>, 2003.
- 1200 Tulet, P., Crassier, V., Cousin, F., Suhre, K., and Rosset, R.: ORILAM, a three-moment lognormal aerosol scheme for mesoscale atmospheric model: Online coupling into the Meso-NH-C model and validation on the Escompte campaign, *J. Geophys. Res.-Atmos.*, 110, <https://doi.org/10.1029/2004JD005716>, 2005.
- Tulet, P., Grini, A., Griffin, R. J., and Petitcol, S.: ORILAM-SOA: A computationally efficient model for predicting secondary organic aerosols in three-dimensional atmospheric models, *J. Geophys. Res.-Atmos.*, 111, <https://doi.org/10.1029/2006JD007152>, 2006.
- 1205 Twohy, C. H., Anderson, J. R., Toohey, D. W., Andrejczuk, M., Adams, A., Lytle, M., George, R. C., Wood, R., Saide, P., Spak, S., Zuidema, P., and Leon, D.: Impacts of aerosol particles on the microphysical and radiative properties of stratocumulus clouds over the southeast Pacific Ocean, *Atmos. Chem. Phys.*, 13, 2541–2562, <https://doi.org/10.5194/acp-13-2541-2013>, 2013.
- Twomey, S.: PRECIPITATION BY DIRECT INTERCEPTION OF CLOUD-WATER, *Weather*, 12, 120–122, <https://doi.org/https://doi.org/10.1002/j.1477-8696.1957.tb00453.x>, 1957.
- 1210 Van der Dussen, J. J., de Roode, S. R., and Siebesma, A. P.: Factors controlling rapid stratocumulus cloud thinning, *J. Atmos. Sci.*, 71, 655–664, <https://doi.org/10.1175/JAS-D-13-0114.1>, 2014.
- Vié, B., Pinty, J.-P., Berthet, S., and Leriche, M.: LIMA (v1.0): A quasi two-moment microphysical scheme driven by a multimodal population of cloud condensation and ice freezing nuclei, *Geoscientific Model Development*, 9, 567–586, <https://doi.org/10.5194/gmd-9-567-2016>, 2016.
- 1215 Wang, S., Wang, Q., and Feingold, G.: Turbulence, Condensation, and Liquid Water Transport in Numerically Simulated Nonprecipitating Stratocumulus Clouds, *J. Atmos. Sci.*, 60, 262–278, [https://doi.org/10.1175/1520-0469\(2003\)060<0262:TCALWT>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)060<0262:TCALWT>2.0.CO;2), 2003.
- Wood, R.: Stratocumulus Clouds, *Mon. Weather Rev.*, 140, 2373 – 2423, <https://doi.org/10.1175/MWR-D-11-00121.1>, 2012.
- 1220 Yamaguchi, T., Feingold, G., Kazil, J., and McComiskey, A.: Stratocumulus to cumulus transition in the presence of elevated smoke layers, *Geophys. Res. Lett.*, 42, 10,478–10,485, <https://doi.org/https://doi.org/10.1002/2015GL066544>, 2015.
- Zhang, J. and Zuidema, P.: The diurnal cycle of the smoky marine boundary layer observed during August in the remote southeast Atlantic, *Atmos. Chem. Phys.*, 19, 14493–14516, <https://doi.org/10.5194/acp-19-14493-2019>, 2019.
- 1225 Zouzoua, M., Lohou, F., Assamoi, P., Lothon, M., Yoboue, V., Dione, C., Kalthoff, N., Adler, B., Babic, K., Pedruzo-Bagazgoitia, X., and Derrien, S.: Breakup of nocturnal low-level stratiform clouds during the southern West African monsoon season, *Atmos. Chem. Phys.*, 21, 2027–2051, <https://doi.org/10.5194/acp-21-2027-2021>, 2021.