

Measurement report: Size-resolved mass concentration of equivalent black carbon-containing particle larger than 700 nm and its role in radiation

Weilun Zhao¹, Ying Li^{2,3}, Gang Zhao⁴, Song Guo⁴, Nan Ma⁵, Shuya Hu⁴, Chunsheng Zhao¹

¹Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing 100871, China

²Department of Ocean Science and Engineering, Southern University of Science and Technology, Shenzhen 518055, China

³Southern Marine Science and Engineering Guangdong Laboratory, Guangzhou 511458, China

⁴State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

⁵Institute for Environmental and Climate Research, Jinan University, Guangzhou 511443, China

Correspondence to: Chunsheng Zhao (zcs@pku.edu.cn)

Abstract. Black carbon (BC) mass size distribution (BCMSD) is crucial in both environment and climate system due to BC's intense size-dependent absorption of solar radiation. BC-containing particles of size larger than 700 nm ($BC_{>700}$) could contribute to larger than half of bulk BC mass concentration. Unfortunately, previous methods concentrated on BC-containing particles less than 700 nm because of technical limitation. The contribution of BC to absorption and radiative effect could be underestimated without consideration of $BC_{>700}$. In this study, equivalent BCMSD (eBCMSD) from 150 nm up to 1.5 μm was measured at high time resolution of 1 h for the first time by an aerodynamic aerosol classifier in tandem with an aethalometer in two field campaigns over eastern China, namely Changzhou located in the Yangtze River Delta and Beijing located in the North China Plain. The results revealed that the pattern of eBCMSD in Changzhou (Beijing) was mostly bimodal (unimodal) peaking at 240 and 1249 nm (427 nm). The peak diameter of eBCMSD in Changzhou did not shift significantly with increasing pollution (240 to 289 nm). In contrast, the peak diameter of eBCMSD in Beijing shifted towards larger size from 347 to 527 nm with increasing pollution, indicating the aging process in urban site was different from that in regional background site. eBCMSD in both Changzhou and Beijing had significant diurnal cycle with smaller (bigger) value of eBCMSD during daytime (nighttime). Equivalent $BC_{>700}$ ($eBC_{>700}$) was ubiquitous and varied significantly with different locations and pollution levels. The campaign-averaged contribution of $eBC_{>700}$ to bulk eBC mass concentration ($m_{eBC,bulk}$), bulk absorption coefficient ($\sigma_{ab,bulk}$) as well as estimated direct radiative forcing of eBC (DRF_{eBC}) in Changzhou and Beijing were 27.8 (20.9 ~ 36.5) % and 24.1 (17.5 ~ 34.2) %, 19.6 (15.8 ~ 24.6) % and 25.9 (19.6 ~ 33.7) %, as well as 20.5 (18.4 ~ 22.2) % and 21.0 (16.3 ~ 26.1) %, respectively. $m_{eBC,bulk}$, $\sigma_{ab,bulk}$ as well as DRF_{eBC} of $eBC_{>700}$ in Changzhou (Beijing) varied by 3.6 (5.1) times from 0.11 (0.07) to 0.40 (0.36) $\mu\text{g m}^{-3}$, 3.2 (5.5) times from 0.54 (0.63) to 1.75 (3.45) Mm^{-1} as well as 2.4 (4.7) times from 0.1 (0.1) to 0.24 (0.47) W m^{-2} , respectively, with the aggravation of pollution. The contribution of $eBC_{>700}$ to $m_{eBC,bulk}$ and $\sigma_{ab,bulk}$ had significant diurnal cycle with higher (lower) fraction during daytime (nighttime) in both Changzhou and Beijing.

32 A case study indicated that the contribution of $e_{BC>700}$ to $m_{eBC,bulk}$, $\sigma_{ab,bulk}$ and DRF_{eBC} could reach up to 50 %, 50 % and
33 40 %, respectively. It was highly recommended to consider whole size range of BC-containing particles in the model
34 estimation of BC radiative effect.

35 **1 Introduction**

36 Black carbon (BC) is the strong light-absorbing carbonaceous particle (Bond and Bergstrom, 2006) from incomplete
37 combustion of fossil fuel or biomass (Bond et al., 2004). Absorption of BC reduces atmospheric visibility (Moosmuller et al.,
38 2009) and has warming effect on the climate system (Bond, 2001). BC radiative effect had considerable uncertainties.
39 Estimated BC radiative effects from different models did not even converge to same order of magnitude (Bond et al.,
40 2013;Szopa et al., 2021).

41 Previous estimation of BC radiative effect was based on bulk BC mass concentration ($m_{BC,bulk}$) from emission inventory
42 and prescribed mass absorption cross section (MAC) (Bond et al., 2013). Both $m_{BC,bulk}$ and MAC was influenced by BC mass
43 size distribution (BCMSD), which was one of the BC microphysical properties. BC radiative effect was highly sensitive to
44 BCMSD (Matsui et al., 2018), and BCMSD could result in obvious variation in aerosol radiative forcing (Zhao et al., 2019).
45 BCMSD depended on the emission source essentially. For example, the peak diameter of freshly emitted BCMSD from fossil
46 fuel was generally smaller than that from biomass burning (Berner et al., 1984;Artaxo et al., 1998;Schwarz et al., 2008). After
47 BC was emitted to the ambient environment, BCMSD was influenced by aging process, during which BC optical properties
48 underwent remarkable changes (Zhang et al., 2008). For instance, BC could be coated by other non-BC materials during
49 atmospheric transport. The existence of non-BC coating enhanced BC absorption and this phenomenon was termed as “lensing
50 effect” (Fuller et al., 1999). The accurate quantification of lensing effect was a critical challenge in estimating BC radiative
51 effect (Liu et al., 2017). and the information of BCMSD was required to resolve the influence of “lensing effect” on BC
52 radiative forcing.

53 Guo (2016) reported that elemental carbon (EC, Petzold et al. (2013)) containing particle larger than 2.1 μm accounted for
54 27.6 ~ 35.2 % of bulk EC mass concentration ($m_{EC,bulk}$). Wang et al. (2017) reported that EC-containing particle larger than
55 1.1 μm accounted for 40.6 ~ 65.5 % of $m_{EC,bulk}$. Wang et al. (2022) indicated EC-containing particle larger than 1 μm
56 contributed to 50 ~ 54 % of $m_{EC,bulk}$. Therefore, BC-containing particle larger than 1 μm contributed to significant part of total
57 BC mass. Wang et al. (2022) found that these large carbon-containing particles were super-aggregated BC particles with fractal
58 structure or BC-containing particles with massive coating from secondary processes. Chakrabarty et al. (2014) found that the
59 optical property of these super BC agglomerates could be significant. It should be noted that current characterization of BC-
60 containing particle larger than 1 μm could be only achieved through offline microscopy analysis (Chakrabarty et al., 2014) or
61 EC mass size distribution (ECMSD) measurement by off-line thermo/optical organic carbon/elemental carbon analysis of
62 size-segregated filter-based samples (Chow et al., 2001). The resulting time-resolution of ECMSD was 24 ~ 48 h. Considering
63 that the typical time scale of BC aging was 4 ~ 18 h (Peng et al., 2016), current measured ECMSD could not resolve

64 atmospheric aging of BC-containing particles larger than 1 μm . Actually, current method capable of measuring BC-containing
65 particle on time scale of BC aging was limited to size less than 700 nm, namely laser-induced incandescence technique
66 (Schwarz et al., 2006). The characterization of BC-containing particles larger than 700 nm ($\text{BC}_{>700}$) during atmospheric aging
67 was still unclear. The contribution of $\text{BC}_{>700}$ to absorption and BC radiative forcing was lack of study.

68 In this study, equivalent BC (eBC, Petzold et al. (2013)) mass size distribution (eBCMSD) up to 1.5 μm was measured with
69 a time resolution of 1 h to study the evolution of equivalent $\text{BC}_{>700}$ (e $\text{BC}_{>700}$) as well as the contribution of e $\text{BC}_{>700}$ to bulk
70 eBC mass concentration ($m_{\text{eBC,bulk}}$), bulk absorption coefficient ($\sigma_{\text{ab,bulk}}$) and eBC direct radiative forcing. The size referred to
71 the particle size, not the BC core size, in this study. eBCMSD was determined by an aerosol aerodynamic classifier (AAC,
72 Cambustion, UK, Tavakoli and Olfert (2013)) in tandem with an aethalometer (model AE33, Magee, USA, Drinovec et al.
73 (2015), AAC – AE33) based on the method proposed by Zhao et al. (2022). eBCMSD was measured in two different locations
74 of eastern China to study the spatial difference of e $\text{BC}_{>700}$. Direct radiative forcing of eBC (DRF_{eBC}) was estimated by the
75 Santa Barbara DISORT (discrete ordinates radiative transfer) Atmospheric Radiative Transfer (SBDART) model (Ricchiazzi
76 et al., 1998).

77 The structure of this study was organized as follows. Section 2 introduced the field measurement, instrumental setup, and
78 details about estimation of DRF_{eBC} . Section 3 discussed the evolution as well as mass, absorption and radiation contribution
79 of e $\text{BC}_{>700}$ based on the field measurement. Section 4 came to the conclusions.

80 **2 Methods**

81 **2.1 Field measurement**

82 The AAC-AE33 system was first applied to a field measurement in Changzhou, Jiangsu Province, China (119°36'E, 31°43'
83 N), situated at the Yangtze River Delta, from May 17th to June 3rd in 2021 (Summer). Changzhou was between two meta cities,
84 namely Nanjing (82 km to the northwest) and Shanghai (187 km to the southeast). There were no emission sources around
85 the measurement site in Changzhou. Thus, pollution around the site was dominated by regional transportation and the
86 measurement site in Changzhou was a typical regional background site (Zhao et al., 2022).

87 Then, the AAC-AE33 was deployed in Beijing, China (116°18'E, 39°59'N), located in the North China Plain, from October
88 29th 2021 to January 25th 2022 (Winter). The measurement site was near 2 busy streets, namely Zhongguancun Street to the
89 west and Chengfu Road to the south. Therefore, the measurement site in Beijing was representative of urban environment
90 Zhao et al. (2019).

91 **2.2 Instrumental setup**

92 The instrumental setup for eBCMSD measurement was illustrated in detail by Zhao et al. (2022) and introduced here briefly.
93 As shown in Fig. 1, a PM_{10} inlet (16.67 L min^{-1}) was used to sample ambient aerosol particles. Then particles passed through
94 a silica gel diffusion drier, where relative humidity (RH) was decreased to less than 30 %, before sampled by the AAC-AE33.
95 AAC-AE33 measured size-resolved absorption coefficient ($\sigma_{\text{ab,size-resolved}}$) at a flow rate of 3 L min^{-1} in Changzhou and 2 L

min⁻¹ in Beijing, respectively. AAC was set to scan 12 logarithmically equally distributed aerodynamic sizes ranging from 200 nm to 1.5 μm in Changzhou and 150 nm to 1.5 μm in Beijing, respectively. It should be pointed out that particle diameter (D_p) was aerodynamic size in this study. Particles of each scanned size were sampled for 5 min, so the time resolution of $\sigma_{ab, \text{size-resolved}}$ came to 1 h. The measured $\sigma_{ab, \text{size-resolved}}$ at wavelength of 880 nm by AE33 was used to derive eBCMSD because BC was the major contributor of aerosol absorption at 880 nm (Ramachandran and Rajesh, 2007).

The principle of retrieving σ_{ab} from AE33 was proposed by Hansen et al. (1984) and described here briefly. Aerosol-laden flow with flow rate of F flew into AE33, where the aerosol particles were collected on a region with area of S of a filter. The filter was illuminated by light sources at specific wavelengths. Part of the light transmitted through the particle-laden (particle-free) area of the filter and the transmitted light intensity was denoted as $I(I_0)$. The light attenuation ATN was defined as

$$\text{ATN} = -100 \cdot \ln\left(\frac{I}{I_0}\right). \quad (1)$$

Assume ATN changed by ΔATN in time interval of Δt , the attenuation coefficient σ_{ATN} was defined as

$$\sigma_{\text{ATN}} = \frac{S}{100 \cdot F} \cdot \frac{\Delta\text{ATN}}{\Delta t}. \quad (2)$$

The light attenuation was actually caused by both absorption of particles and scattering of particles as well as the filter, which was called multi-scattering effect. The multi-scattering effect was corrected based on the study of Zhao et al. (2020), where a parameter $C_f = 2.9$ was introduced to derive σ_{ab} :

$$\sigma_{ab} = \frac{\sigma_{\text{ATN}}}{C_f}. \quad (3)$$

Besides multi-scattering effect, loading effect was also required to be corrected, namely, the change in ATN was not linearly dependent on aerosol loading. Drinovec et al. (2015) developed “dual-spot” technique to correct loading effect, which was adopted by this study.

MAC was required to convert absorption coefficient to eBC mass concentration. The size-dependent MAC was modeled based on the scheme proposed by Zhao et al. (2021) at wavelength of 880 nm, which required size-resolved particle number concentration ($N_{\text{size-resolved}}$). Concisely, took 700nm of D_p as an example, the number fraction of BC-containing particle (f_{BC}) was assumed a fixed parameter (0.35), and the number concentration of BC-containing particle (N_{BC}) at D_p of 700 nm could be derived by

$$N_{\text{BC}} = f_{\text{BC}} \cdot N_{\text{size-resolved}} \cdot \Delta\log D_p, \quad (4)$$

where $\Delta\log D_p$ was the logarithmic width of the D_p size bin. The fixed- f_{BC} assumption led to ~ 3 % uncertainty in derived m_{eBC} . We assumed that all BC-containing particles at D_p of 700 nm had the same core size D_{BC} . An optimal D_{BC} was found so that calculated absorption ($\sigma_{ab, \text{calc}}$) matched measured absorption ($\sigma_{ab, \text{meas}}$), namely

$$\sigma_{ab, \text{calc}} = \rho_{\text{BC}} \frac{\pi}{6} D_{\text{BC}}^3 \cdot \text{MAC}_{\text{Mie}} \cdot N_{\text{BC}} = \sigma_{\text{size-resolved}} \cdot \Delta\log D_p = \sigma_{ab, \text{meas}}, \quad (5)$$

where ρ_{BC} was density of BC and assumed to be a fixed value (1.8 g cm⁻³), MAC_{Mie} was Mie-calculated MAC at D_p of 700

126 nm and the optimal D_{BC} . eBCMSD at D_p of 700 nm could be calculated by

$$127 \quad \text{eBCMSD}_{D_p=700\text{nm}} = \frac{\sigma_{\text{ab,size-resolved}}}{\text{MAC}_{\text{Mie}}}. \quad (6)$$

128 The assumption on the MAC led to $\sim 24\%$ uncertainty on derived m_{eBC} . It should be noted that dust was not considered in
 129 this study. $N_{\text{size-resolved}}$ was measured by a scanning mobility particle sizer (SMPS, TSI, USA) at 0.3 L min^{-1} as well as an
 130 aerodynamic particle sizer (APS, TSI, USA) at 5 L min^{-1} in Changzhou and an AAC in tandem with condensation particle
 131 counter (CPC, TSI, USA, AAC – CPC, Johnson et al. (2018)) at 1 L min^{-1} in Beijing, respectively. AAC-AE33 measured
 132 $\sigma_{\text{ab,size-resolved}}$ and determined eBCMSD synchronously. Therefore, the contribution of $\text{eBC}_{>700}$ to both bulk absorption and
 133 $m_{\text{eBC,bulk}}$ could be quantified simultaneously.

134 In this study, the bulk mass concentration of eBC-containing particle ($m_{\text{eBC,bulk}}$) was defined as

$$135 \quad m_{\text{eBC,bulk}} = \int_{200 \text{ nm}}^{1500 \text{ nm}} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p, \quad (7)$$

136 where $\frac{dm_{\text{eBC}}}{d\log D_p}$ was eBCMSD, and the lower limit of integral was 200 nm in both Changzhou and Beijing for the convenience

137 of comparison. The bulk mass concentration of $\text{eBC}_{>700}$ ($m_{\text{eBC,bulk},>700}$) was defined as

$$138 \quad m_{\text{eBC,bulk},>700} = \int_{700 \text{ nm}}^{1500 \text{ nm}} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p. \quad (8)$$

139 The contribution of $\text{eBC}_{>700}$ to $m_{\text{eBC,bulk}}$ ($f_{m,>700}$) was defined as

$$140 \quad f_{m,>700} = \frac{m_{\text{eBC,bulk},>700}}{m_{\text{eBC,bulk}}} \times 100 \%. \quad (9)$$

141 It should be noted that BC-containing particle of D_p lower than 200 nm and greater than 1500 nm was not considered in this
 142 study, which leads to discrepancy between true $f_{m,>700}$ ($\hat{f}_{m,>700}$) and estimated $f_{m,>700}$. By simple mathematical analysis, it
 143 could be proofed that $\hat{f}_{m,>700}$ in the range of

$$144 \quad \frac{1}{2} \left(f_{m,>700} + \frac{\int_{1500 \text{ nm}}^{+\infty} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p}{\int_{200 \text{ nm}}^{1500 \text{ nm}} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p} \right) < \hat{f}_{m,>700} < f_{m,>700} + \frac{\int_{1500 \text{ nm}}^{+\infty} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p}{\int_{200 \text{ nm}}^{1500 \text{ nm}} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p}, \quad (10)$$

145 where $\int_{200 \text{ nm}}^{1500 \text{ nm}} \frac{dm_{\text{eBC}}}{d\log D_p} d\log D_p$ was actually $m_{\text{eBC,bulk}}$ in this study.

146 Similarly, the bulk absorption coefficient ($\sigma_{\text{ab,bulk}}$) was defined as

$$147 \quad \sigma_{\text{ab,bulk}} = \int_{200 \text{ nm}}^{1500 \text{ nm}} \frac{d\sigma_{\text{ab}}}{d\log D_p} d\log D_p, \quad (11)$$

148 where $\frac{d\sigma_{\text{ab}}}{d\log D_p}$ was $\sigma_{\text{ab,size-resolved}}$. The bulk absorption coefficient of $\text{eBC}_{>700}$ ($\sigma_{\text{ab,bulk},>700}$) was defined as

$$149 \quad \sigma_{\text{ab,bulk},>700} = \int_{700 \text{ nm}}^{1500 \text{ nm}} \frac{d\sigma_{\text{ab}}}{d\log D_p} d\log D_p. \quad (12)$$

150 The contribution of $\text{eBC}_{>700}$ to $\sigma_{\text{ab,bulk}}$ ($f_{\text{ab},>700}$) was defined as

$$151 \quad f_{\text{ab},>700} = \frac{\sigma_{\text{ab,bulk},>700}}{\sigma_{\text{ab,bulk}}} \times 100 \%. \quad (13)$$

152 **2.3 Estimation of direct radiative forcing of equivalent black carbon**

The direct radiative effect was one of the BC characteristics that arouse extensive concerns. The SBDART model was employed to study the characteristics of DRF_{eBC} . Specifically, the instantaneous DRF_{eBC} was estimated at the top of atmosphere (TOA) under the cloud-free condition. Wavelengths from 250 nm to 4 μm were simulated in this study. Direct radiative forcing of aerosol ($DRF_{aerosol}$) was defined as (Zhao et al., 2018):

$$DRF_{aerosol} = (F_{aerosol,\downarrow} - F_{aerosol,\uparrow}) - (F_{clearsky,\downarrow} - F_{clearsky,\uparrow}), \quad (14)$$

where $F_{aerosol,\downarrow}$ ($F_{aerosol,\uparrow}$) was downward (upward) radiative irradiance flux at TOA with aerosol, and $F_{clearsky,\downarrow}$ ($F_{clearsky,\uparrow}$) was downward (upward) radiative irradiance flux at TOA without aerosol. Direct radiative forcing of aerosol without eBC ($DRF_{aerosol,noneBC}$) was defined as:

$$DRF_{aerosol,noneBC} = (F_{aerosol,noneBC,\downarrow} - F_{aerosol,noneBC,\uparrow}) - (F_{clearsky,\downarrow} - F_{clearsky,\uparrow}), \quad (15)$$

where $F_{aerosol,noneBC,\downarrow}$ ($F_{aerosol,noneBC,\uparrow}$) was downward (upward) radiative irradiance flux at TOA with aerosol except eBC.

The DRF_{eBC} was defined as the difference between $DRF_{aerosol}$ and $DRF_{aerosol,noneBC}$:

$$DRF_{eBC} = (F_{aerosol,\downarrow} - F_{aerosol,\uparrow}) - (F_{aerosol,noneBC,\downarrow} - F_{aerosol,noneBC,\uparrow}). \quad (16)$$

Similarly, the direct radiative forcing of $eBC_{>700}$ ($DRF_{eBC,>700}$) was defined as:

$$DRF_{eBC,>700} = (F_{aerosol,\downarrow} - F_{aerosol,\uparrow}) - (F_{aerosol,noneBC,>700,\downarrow} - F_{aerosol,noneBC,>700,\uparrow}), \quad (17)$$

where $F_{aerosol,noneBC,>700,\downarrow}$ ($F_{aerosol,noneBC,>700,\uparrow}$) was downward (upward) radiative irradiance flux at TOA with aerosol except $eBC_{>700}$. The contribution of $eBC_{>700}$ to DRF_{eBC} ($f_{DRF,>700}$) was defined as

$$f_{DRF,>700} = \frac{DRF_{eBC,>700}}{DRF_{eBC}} \times 100 \%. \quad (18)$$

SBDART simulation required information of surface albedo, vertical profiles of meteorological parameters and aerosol optical parameters. Surface albedo was acquired from Moderate Resolution Imaging Spectroradiometer (MODIS)/Terra surface reflectance data with temporal and spatial resolution of 1 d and 0.05° (MOD09CMG). The gridded data around the measurement site was averaged to represent surface albedo of the measurement site.

The vertical profile of meteorological parameters included vertical profile of pressure, temperature, water vapor and ozone, which were obtained from the fifth generation ECMWF (European Center for Medium Range Weather Forecasts) reanalysis data for global climate and weather (ERA5). The ERA5 data had temporal and spatial resolution of 1 h and 0.25° with 38 vertical layers from surface to about 50 km above surface. At each layer, the gridded data around the measurement site was also averaged to represent meteorological parameters of the measurement site. The time resolution of meteorological parameters was averaged to daily to match that of surface albedo.

The vertical profile of aerosol optical parameters included the vertical profile of bulk aerosol extinction coefficient ($\sigma_{ext,bulk}$), single scattering albedo (SSA) and asymmetry factor (g) at different wavelengths, which were parameterized based on the study of Zhao et al. (2019) and described here briefly. The bulk aerosol particle number concentration (N_{bulk}) was parameterized according to aircraft study by Liu et al. (2009). Dry $N_{size-resolved}$ at different heights had the same shape after

184 normalized by corresponding N_{bulk} . The parameterization of $m_{\text{eBC,bulk}}$ and eBCMSD was the same as N_{bulk} and dry $N_{\text{size-resolved}}$.
 185 As for mixing state, 51% of eBC mass was assumed externally mixed and the rest of eBC mass was assumed internally mixed
 186 with core-shell geometry (Ma et al., 2012) in each size bin. For the case of aerosol without eBC-containing particle (larger
 187 than 700 nm), eBCMSD (larger than 700 nm) was set to 0. The aerosol optical parameters varying with height-dependent RH
 188 were calculated by Mie scattering theory and κ -Kohler theory (Petters and Kreidenweis, 2007) assuming hygroscopic growth
 189 parameter of 0.22 (Tan et al., 2019). The refractive indices of eBC, water and non-eBC material without water were assumed
 190 $1.8 + 0.54i$ (average among the wavelengths of 450, 550 and 700 nm) (Kuang et al., 2015), $1.33 + 10^{-7}i$ (wavelength of 550
 191 nm) and $1.53 + 10^{-7}i$ (wavelength of 550 nm) (Wex et al., 2002), respectively. The refractive index of non-eBC material mixed
 192 with water after hygroscopic growth was derived by volume-weighted rule (Wex et al., 2002). In short, real-time measured
 193 eBCMSD and $N_{\text{size-resolved}}$ were used as boundary condition at ground level to construct parameterized vertical aerosol profile.
 194 When calculating aerosol optical parameters at each altitude, mixing state of BC-containing particle was assumed to be the
 195 same at each altitude, each time and each D_p . With the above information, the vertical profiles of $\sigma_{\text{ext,bulk}}$, SSA and g could be
 196 calculated based on Mie theory. The time resolution of aerosol optical parameters was averaged to daily to match that of
 197 surface albedo.

198 **3 Results and discussion**

199 **3.1 Case study**

200 A pollution episode took place from October 31st, 2021 to November 6, 2021 in Beijing, which was used for case study to
 201 illustrate the large variability of eBC_{>700}. The geometric mean diameter (\bar{D}_p) of eBCMSD was defined as

$$202 \log \bar{D}_p = \frac{\int \log D_p \frac{dm_{\text{eBC}}}{d \log D_p} d \log D_p}{\int \frac{dm_{\text{eBC}}}{d \log D_p} d \log D_p}, \quad (19)$$

203 which was used to depict the spectral variation of eBCMSD because eBCMSD did not always had an explicit modal pattern
 204 and the corresponding peak diameter was not always easy to be distinguished.

205 With the development of pollution, \bar{D}_p shifted apparently from around 400 nm to around 600 nm (Fig. 2a). $m_{\text{eBC,bulk}}$
 206 ($m_{\text{eBC,bulk,>700}}$) increased from less than 0.5 (0.15) $\mu\text{g m}^{-3}$ to as large as 2.5 (1.0) $\mu\text{g m}^{-3}$ by 5.0 (6.6) times. $\sigma_{\text{ab,bulk}}$ ($\sigma_{\text{ab,bulk,>700}}$)
 207 increased from less than 4 (1) Mm^{-1} to as large as 25 (10) Mm^{-1} by 6.3 (10.0) times. DRF_{eBC} ($\text{DRF}_{\text{eBC,>700}}$) increased from 1
 208 (0.2) W m^{-2} to as large as 4 (1) W m^{-2} by 4.0 (5.0) times. It could be seen that the variability of eBC_{>700} was significant. $f_{\text{m,>700}}$,
 209 $f_{\text{ab,>700}}$ and $f_{\text{DRF,>700}}$ increased from about 20 %, 20 % and 20 % to as large as 50 %, 50 % and 40 %, respectively (Fig. 2b),
 210 clearly showing important role of eBC_{>700} in BC mass, absorption as well as radiative effect.

211 **3.2 Equivalent black carbon mass size distribution**

212 **3.2.1 Overview**

213 The timeseries of eBCMSD in Changzhou and Beijing was shown in Fig. S1a and Fig. S1b1 – S1b4. eBCMSD was
 214 presented with normalized probability density function (pdf) to study general characteristics of eBCMSD. Figure 3a1 and 3a2

were the normalized pdf over the whole campaign of Changzhou and Beijing, respectively. It could be seen that eBCMSD in Changzhou was significantly different from that in Beijing. There were two modes in the median of eBCMSD in Changzhou, which peaked at around 240 nm and 1249 nm, respectively. Yu et al. (2010) found 3 modes in ECMSD, namely modes around 300 nm, 1 μm and 5 μm , and named the 3 modes as condensation mode, droplet mode and coarse mode, respectively. Following the nomenclature by Yu et al. (2010), the mode peaking at 240 nm and 1249 nm could be termed as condensation mode and droplet mode, respectively. In contrast, only condensation mode was identified in median eBCMSD in Beijing, which peaked at 427 nm. The variation of eBCMSD, defined as the difference between upper quartile and lower quartile, in Changzhou was overall smaller than that in Beijing. The variation of eBCMSD value in Changzhou (Beijing) ranged from 0.52 (0.54) $\mu\text{g m}^{-3}$ to 0.91 (1.73) $\mu\text{g m}^{-3}$ with average value of 0.75 (1.05) $\mu\text{g m}^{-3}$. The maximum upper quartile of eBCMSD in Changzhou was 1.58 $\mu\text{g m}^{-3}$. In comparison, the upper quartile of eBCMSD in Beijing could reached up to 2.14 $\mu\text{g m}^{-3}$, indicating the evolution of eBCMSD in Beijing was more drastic than that in Changzhou.

3.2.2 Evolution with respect to pollution level

In order to investigate the evolution of eBCMSD under different pollution stages, eBCMSD was grouped into 3 periods: (1) clean period in which $m_{\text{eBC,bulk}}$ was lower than 0.5 $\mu\text{g m}^{-3}$, (2) transitional period in which $m_{\text{eBC,bulk}}$ was greater than 0.5 $\mu\text{g m}^{-3}$ but lower than 1.0 $\mu\text{g m}^{-3}$, (3) polluted period in which $m_{\text{eBC,bulk}}$ was greater than 1.0 $\mu\text{g m}^{-3}$. Data from clean, transitional and polluted period accounted for 22.6 % (30.9 %), 51.3 % (31.9 %) and 26.0 % (37.2 %) of total data in Changzhou (Beijing), respectively, showing that Changzhou (Beijing) was dominated by transitional (polluted) period in this study.

In the clean period, there was no distinct difference in eBCMSD between Changzhou (Fig. 3b1) and Beijing (Fig. 3b2). Neither eBCMSD in Changzhou nor eBCMSD in Beijing exhibited obvious modal structure in the size range of measurement. The value of eBCMSD in both Changzhou and Beijing decreased with increasing D_p in general. For Changzhou (Beijing), the median of eBCMSD decreased from 0.87 (0.47) $\mu\text{g m}^{-3}$ at 200 nm to 0.26 (0.26) $\mu\text{g m}^{-3}$ at 1500 nm with average value of 0.42 (0.34) $\mu\text{g m}^{-3}$. The variation of eBCMSD in Changzhou (Beijing) was 0.24 (0.24) $\mu\text{g m}^{-3}$ ~ 0.47 (0.55) $\mu\text{g m}^{-3}$ with average value of 0.32 (0.35) $\mu\text{g m}^{-3}$, showing that the variation of eBCMSD in Changzhou was comparable to that in Beijing.

As polluted stage evolved to transitional period, the value of eBCMSD increased in both Changzhou (Fig. 3c1) and Beijing (Fig. 3c2) compared to that in clean period. The variation of eBCMSD in Changzhou (Beijing) reached 0.41 (0.44) $\mu\text{g m}^{-3}$ ~ 0.86 (0.86) $\mu\text{g m}^{-3}$ with average value of 0.53 (0.61) $\mu\text{g m}^{-3}$, about twice as much as that in clean period. It could be seen that the value of median and variation of eBCMSD in Changzhou were comparable to that in Beijing. However, the pattern of eBCMSD in Changzhou was obviously different from that in Beijing. The peak value of median eBCMSD located at 240 (347) nm in Changzhou (Beijing). Median eBCMSD in Changzhou exhibited two modes, namely condensation mode and droplet, with boundary at around 866 nm. In comparison, median eBCMSD in Beijing only had one mode, namely condensation mode. It should be noted that two modes (one mode) meant that there were two distinct groups (one group) of

247 BC-containing particles with respect to D_p , not the BC core size (D_{BC}). The difference in peak diameter of condensation mode
248 between Changzhou and Beijing was as large as 107 nm. Median eBCMSD at clean period was subtracted from that at
249 transitional period to study eBC mass increment at each D_p , as shown in Fig. S4a1. It could be clearly seen that mass increment
250 in Changzhou peaked at 289 nm and 1249 nm, contributing to condensation mode and droplet mode in eBCMSD, respectively.
251 In contrast, mass increment in Beijing only peaked at 385 nm, contributing to condensation mode in eBCMSD.

252 As the pollution stage came to polluted period, the value of eBCMSD increased drastically in both Changzhou (Fig. 3d1)
253 and Beijing (Fig. 3d2) compared to that in clean period. Both the median value and the variation of eBCMSD increased with
254 the development of pollution. The median eBCMSD increased to 0.88 (0.61) ~ 2.12 (2.45) $\mu\text{g m}^{-3}$ with average value of 1.49
255 (1.52) $\mu\text{g m}^{-3}$ in Changzhou (Beijing), about 4 times as much as the median eBCMSD in clean period. The variation of
256 eBCMSD in Changzhou (Beijing) reached 0.60 (0.73) ~ 1.11 (1.06) $\mu\text{g m}^{-3}$ with average value of 0.92 (0.94) $\mu\text{g m}^{-3}$, about 3
257 times as much as that in clean period. The difference in pattern of eBCMSD between Changzhou and Beijing became more
258 distinct. Median eBCMSD in Changzhou clearly exhibited a bimodal structure where the condensation mode and droplet
259 mode peaked at 289 nm and 1249 nm, respectively. Median eBCMSD in Beijing exhibited a unimodal structure where the
260 condensation mode peaked a 527 nm. As shown in Fig. S4b1, the peak of mass increment in Changzhou (Beijing) shifted
261 from 289 (385) nm to 347 (527) nm, varied by 58 (142) nm. The significant difference in the shift of peak indicated that aging
262 processes in regional background site was significantly different from that in urban site.

263 **3.2.3 Contribution of equivalent black carbon-containing particle larger than 700 nm to bulk equivalent black carbon** 264 **mass concentration**

265 The median (lower quartile ~ upper quartile) of $m_{\text{eBC,bulk}}$ was 0.73 (0.52 ~ 1.03) $\mu\text{g m}^{-3}$ in Changzhou and 0.79 (0.43 ~ 1.31)
266 $\mu\text{g m}^{-3}$ in Beijing (Fig. 4a1). The median of $m_{\text{eBC,bulk}}$ was comparable between Changzhou and Beijing. The variation of
267 $m_{\text{eBC,bulk}}$ in Changzhou, 0.51 $\mu\text{g m}^{-3}$, was smaller than that in Beijing, 0.88 $\mu\text{g m}^{-3}$. $m_{\text{eBC,bulk},>700}$ in Changzhou was overall
268 comparable to that in Beijing (Fig. 4a2). $m_{\text{eBC,bulk},>700}$ was 0.20 (0.13 ~ 0.32) $\mu\text{g m}^{-3}$ in Changzhou and 0.18 (0.10 ~ 0.33) μg
269 m^{-3} in Beijing. Therefore, eBC_{>700} was ubiquitous. Considering that the variation of $m_{\text{eBC,bulk},>700}$ in Changzhou, 0.19 $\mu\text{g m}^{-3}$,
270 was comparable to that in Beijing, 0.23 $\mu\text{g m}^{-3}$, the larger variation in $m_{\text{eBC,bulk}}$ in Beijing was mainly from eBC-containing
271 particles less than 700 nm. $f_{\text{m},>700}$ was 27.8 (20.9 ~ 36.5) % in Changzhou and 24.1 (17.5 ~ 34.2) % in Beijing (Fig. 4a3),
272 indicating that eBC_{>700} was overall one quarter of $m_{\text{eBC,bulk}}$. $f_{\text{m},>700}$ in Changzhou was slightly larger than that in Beijing, which
273 was contributed by droplet mode of eBCMSD in Changzhou. A summary table with respect to m_{eBC} was presented in Table 1.

274 The statistics of mass contribution of eBC_{>700} were studied with different pollution stages. As shown in Fig. 4a1, $m_{\text{eBC,bulk}}$
275 increased from 0.41 (0.33 ~ 0.45) $\mu\text{g m}^{-3}$ in clean period through 0.71 (0.58 ~ 0.83) $\mu\text{g m}^{-3}$ in transitional period to 1.33 (1.16
276 ~ 1.71) $\mu\text{g m}^{-3}$ in polluted period by 3.2 times in Changzhou and increased from 0.32 (0.22 ~ 0.41) $\mu\text{g m}^{-3}$ in clean period
277 through 0.73 (0.61 ~ 0.85) $\mu\text{g m}^{-3}$ in transitional period to 1.47 (1.21 ~ 1.82) $\mu\text{g m}^{-3}$ in polluted period by 4.6 times in Beijing.
278 As shown in Fig. 4a2, the change of $m_{\text{eBC,bulk},>700}$ with pollution level was substantial in both Changzhou and Beijing. For

279 Changzhou, $m_{\text{eBC,bulk},>700}$ increased from 0.11 (0.07 ~ 0.15) $\mu\text{g m}^{-3}$ in clean period to 0.20 (0.14 ~ 0.27) $\mu\text{g m}^{-3}$ in transition
280 period, and reached 0.40 (0.29 ~ 0.50) $\mu\text{g m}^{-3}$ in polluted period, increasing by as large as 3.6 times from clean period to
281 polluted period. For Beijing, $m_{\text{eBC,bulk},>700}$ increased from 0.07 (0.05 ~ 0.12) $\mu\text{g m}^{-3}$ in clean period to 0.17 (0.11 ~ 0.23) μg
282 m^{-3} in transition period, and reached 0.36 (0.25 ~ 0.52) $\mu\text{g m}^{-3}$ in polluted period, increasing by as large as 5.1 times from
283 clean period to polluted period. The change in $m_{\text{eBC,bulk}}$ and $m_{\text{eBC,bulk},>700}$ was overall consistent with the development of
284 pollution, leading to unobvious change in $f_{\text{m},>700}$ (Fig. 4a3). $f_{\text{m},>700}$ in Changzhou changed from 28.5 (20.3 ~ 36.0) % in
285 clean period through 28.4 (20.7 ~ 36.9) % in transitional period to 27.4 (22.6 ~ 36.2) % in polluted period. $f_{\text{m},>700}$ in Beijing
286 varied from 26.2 (18.4 ~ 36.8) % in clean period through 22.8 (16.3 ~ 32.3) % in transitional period to 23.8 (18.1 ~ 31.9) %
287 in polluted period.

288 3.2.4 Diurnal cycle

289 It could be seen clearly that the value of eBCMSD during daytime was overall lower than that during nighttime in both
290 Changzhou (Fig. 5a1) and Beijing (Fig. 5a2), indicating that eBCMSD was regulated by planetary boundary layer or
291 difference in surface emission source (Liu et al., 2019). For Changzhou (Beijing), eBCMSD from 10:00 to 18:00 (08:00 to
292 18:00) was obviously lower than that from 20:00 to 06:00 (20:00 to 06:00). Accordingly, $m_{\text{eBC,bulk}}$ in Changzhou reached minimum
293 of 0.56 (0.48 ~ 0.88) $\mu\text{g m}^{-3}$ at 12:00 and maximum of 0.97 (0.80 ~ 1.24) $\mu\text{g m}^{-3}$ at 21:00 (Fig. 5b1). $m_{\text{eBC,bulk}}$ in Beijing
294 reached minimum of 0.65 (0.42 ~ 1.02) $\mu\text{g m}^{-3}$ at 14:00 and maximum of 1.08 (0.55 ~ 1.52) $\mu\text{g m}^{-3}$ at 00:00, (Fig. 5b2). The
295 apparent diurnal cycle was found in the condensation mode of eBCMSD, which was mostly less than 700 nm. In contrast,
296 diurnal cycle was not obvious for eBCMSD larger than 700 nm for both Changzhou and Beijing. Consequently, neither
297 $m_{\text{eBC,bulk},>700}$ in Changzhou (Fig. 5c1) nor $m_{\text{eBC,bulk},>700}$ in Beijing (Fig. 5c2) exhibited obvious diurnal cycle. $m_{\text{eBC,bulk},>700}$ in
298 both Changzhou and Beijing fluctuated around 0.2 $\mu\text{g m}^{-3}$, consistent with Sect. 3.2.3. Combining the diurnal variation of
299 $m_{\text{eBC,bulk}}$ and $m_{\text{eBC,bulk},>700}$, $f_{\text{m},>700}$ was negatively correlated to $m_{\text{eBC,bulk}}$ according to Eq. (9) with higher value during the
300 daytime and lower value during the nighttime. $f_{\text{m},>700}$ reached maximum of 35.4 (26.6 ~ 41.1) % at 09:00 and reached
301 minimum of 23.6 (13.9 ~ 30.8) % at 21:00 in Changzhou (Fig. 5d1). $f_{\text{m},>700}$ reached maximum of 31.0 (20.8 ~ 36.9) % at
302 15:00 and reached minimum of 23.5 (16.1 ~ 27.8) % at 01:00 in Beijing (Fig. 5d2).

303 3.3 Size-resolved absorption coefficient

304 3.3.1 Overview

305 The general characteristics (timeseries) of $\sigma_{\text{ab,size-resolved}}$ in Changzhou and Beijing was shown in Fig. 3a3 (Fig. S2a) and Fig.
306 3a4 (Fig. S2b1 – S2b4), respectively. The median $\sigma_{\text{ab,size-resolved}}$ in both Changzhou and Beijing both exhibited unimodal
307 structure. For Changzhou (Beijing), $\sigma_{\text{ab,size-resolved}}$ had maximum value of 7.88 (10.59) Mm^{-1} at 416.1 (427.2) nm and minimum
308 value of 1.63 (2.90) Mm^{-1} at 1500 (1500) nm with average value of 5.39 (6.21) Mm^{-1} . The maximum value was 4.9 (3.7)
309 times as large as minimum value in Changzhou (Beijing), showing the significant dependence of absorption on particle size.
310 D_p which had higher median value of $\sigma_{\text{ab,size-resolved}}$ corresponded to larger variation on the whole. The variation of $\sigma_{\text{ab,size-}}$

resolved ranged from 2.25 (2.82) Mm^{-1} at 1500 (1500) nm to 7.43 (17.90) Mm^{-1} at 500 (527) nm with average value of 4.99 (8.97) Mm^{-1} in Changzhou (Beijing). The variation of $\sigma_{\text{ab,size-resolved}}$ was as large as the median value of $\sigma_{\text{ab,size-resolved}}$ in both Beijing and Changzhou, showing the large variability of BC absorption. The variation of $\sigma_{\text{ab,size-resolved}}$ in Beijing was overall 1.8 times as large as that in Changzhou, indicating that the evolution of $\sigma_{\text{ab,size-resolved}}$ in different sites could be significantly different.

3.3.2 Evolution with respect to pollution level

$\sigma_{\text{ab,size-resolved}}$ was grouped into 3 periods based on $m_{\text{eBC,bulk}}$ as described in Sect. 3.2.2. In clean period, the value of $\sigma_{\text{ab,size-resolved}}$ overall decreased with increasing D_p in both Changzhou (Fig. 3b3) and Beijing (Fig. 3b4), and the pattern of $\sigma_{\text{ab,size-resolved}}$ had no obvious modal structure. In Changzhou (Beijing), the value of $\sigma_{\text{ab,size-resolved}}$ decreased from 4.67 (3.43) Mm^{-1} at 200 (427) nm to 0.88 (1.80) Mm^{-1} at 1500 (1500) nm with average value of 2.95 (2.49) Mm^{-1} . The variation of $\sigma_{\text{ab,size-resolved}}$ in Changzhou (Beijing) ranged from 1.06 (1.57) Mm^{-1} to 2.72 (3.12) Mm^{-1} with average value of 2.04 (2.47) Mm^{-1} .

During the transitional period, the unimodal pattern could be identified in both Changzhou (Fig. 3c3) and Beijing (Fig. 3c4). Median $\sigma_{\text{ab,size-resolved}}$ peaked at 416 (427) nm with value of 7.80 (10.04) Mm^{-1} in Changzhou (Beijing). Median $\sigma_{\text{ab,size-resolved}}$ in clean period was subtracted from that in transitional period to study absorption increment at each D_p , as shown in Fig. S4a2. The increment of $\sigma_{\text{ab,size-resolved}}$ in Changzhou (Beijing) had maximum value of 3.94 (6.61) Mm^{-1} at 416 (427) nm and minimum value of 0.66 (1.15) Mm^{-1} at 1500 (1500) nm. The increment of absorption was most at around 420 nm and least at 1500 nm, showing the significant difference in the change of absorption at different D_p with the development of pollution. The maximum increment of absorption in Beijing was 1.7 times as large as that in Changzhou. Hence, the evolution of absorption could be different substantially in different locations. The variation of $\sigma_{\text{ab,size-resolved}}$ in Changzhou (Beijing) ranged from 1.94 (2.32) Mm^{-1} to 4.03 (6.43) Mm^{-1} with average value of 3.08 (4.45) Mm^{-1} , increasing by about 1.5 times compared to clean period.

In the polluted period, the unimodal pattern of $\sigma_{\text{ab,size-resolved}}$ was significant in both Changzhou (Fig. 3d3) and Beijing (Fig. 3d4). Median $\sigma_{\text{ab,size-resolved}}$ peaked at 416 (527) nm with value of 16.79 (25.85) Mm^{-1} and had minimum value of 2.85 (4.23) Mm^{-1} at 1500 (1500) nm in Changzhou (Beijing). Compared to transition period, peak diameter remained unchanged in Changzhou but increased by 100 nm in Beijing, indicating the evolution of $\sigma_{\text{ab,size-resolved}}$ with aging process was different between regional background site and typical urban site. The increment of absorption in Changzhou (Beijing) was most significant at 416 (527) nm with value of 12.93 (22.94) Mm^{-1} and least at 1500 (1500) nm with value of 1.97 (2.44) Mm^{-1} , as shown in Fig. S4b2. It could be seen that the diameter of increment in absorption remain unchanged in Changzhou and shifted by 100 nm in Beijing, indicating that absorption at different D_p varied differently at different locations with the deterioration of pollution. The variation of $\sigma_{\text{ab,size-resolved}}$ in Changzhou (Beijing) ranged from 2.19 (3.82) Mm^{-1} to 9.05 (15.61) Mm^{-1} with average value of 5.72 (8.22) Mm^{-1} , increasing by about 3 times compared to clean period, indicating that the variability of $\sigma_{\text{ab,size-resolved}}$ increased with the development of pollution.

3.3.3 Contribution of equivalent black carbon-containing particle larger than 700 nm to bulk absorption coefficient

It could be seen from the timeseries of $\sigma_{ab,size-resolved}$ in both Changzhou (Fig. S2a) and Beijing (Fig. S2b1 – S2b4) that absorption of eBC_{>700} was nonnegligible. $\sigma_{ab,bulk}$ was 4.93 (3.53 ~ 7.24) Mm⁻¹ in Changzhou and 6.37 (3.31 ~ 11.68) Mm⁻¹ in Beijing on the whole, as shown in Fig. 4b1. Both median and variation of $\sigma_{ab,bulk}$ in Changzhou were less than that in Beijing. $\sigma_{ab,bulk,>700}$ was 1.03 (0.62 ~ 1.59) Mm⁻¹ in Changzhou, accounting for 19.6 (15.8 ~ 24.6) % of $\sigma_{ab,bulk}$, and 1.47 (0.81 ~ 2.83) Mm⁻¹ in Beijing, accounting for 25.9 (19.6 ~ 33.7) % of $\sigma_{ab,bulk}$, respectively, as shown in Fig. 4b2 and Fig. 4b3. It could be clearly seen that eBC_{>700} contributed to substantial part of total absorption, and should be explicitly considered in BC radiative estimation. A summary table with respect to σ_{ab} was presented in Table 1.

With the aggravation of pollution, the change of $m_{eBC,bulk}$ in Changzhou was overall in agreement with that in Beijing (Fig. 4a1). However, the change of $\sigma_{ab,bulk}$ with the development of pollution was different between Changzhou and Beijing (Fig. 4b1). In the clean period, $\sigma_{ab,bulk}$ in Changzhou with value of 2.71 (2.30 ~ 3.28) Mm⁻¹ was comparable to that in Beijing with value of 2.47 (1.65 ~ 3.28) Mm⁻¹. In the transitional period, $\sigma_{ab,bulk}$ was 4.83 (4.04 ~ 6.02) Mm⁻¹ in Changzhou and 5.93 (4.72 ~ 7.33) Mm⁻¹ in Beijing. The deviation in $\sigma_{ab,bulk}$ was about 1 Mm⁻¹ between Changzhou and Beijing. In the polluted period, $\sigma_{ab,bulk}$ was 9.61 (7.99 ~ 11.93) Mm⁻¹ in Changzhou and 13.65 (10.94 ~ 17.59) Mm⁻¹ in Beijing. The deviation in $\sigma_{ab,bulk}$ came to 4 Mm⁻¹ between Changzhou and Beijing. It could be seen that with the development of pollution, the change of $\sigma_{ab,bulk}$ in Changzhou was less than that in Beijing. MAC_{bulk}, defined as the ratio of median $\sigma_{ab,bulk}$ to median $m_{eBC,bulk}$, changed from 6.61 (7.72) m² g⁻¹ through 6.80 (8.13) m² g⁻¹ to 7.23 (9.29) m² g⁻¹ in Changzhou (Beijing). The increase in MAC_{bulk} in both Changzhou and Beijing with the aggravation of pollution indicated the aging of BC. MAC_{bulk} in Changzhou was overall lower than that in Beijing and increased slower than that in Beijing with the development of pollution, indicating that the BC properties and aging process in Changzhou (regional background site) differentiate from that in Beijing (typical urban site).

$\sigma_{ab,bulk,>700}$ in both Changzhou and Beijing increased with the development of pollution, as shown in Fig. 4b2. $\sigma_{ab,bulk,>700}$ increased from 0.54 (0.62 ~ 1.59) Mm⁻¹ through 0.96 (0.72 ~ 1.32) Mm⁻¹ to 1.75 (1.53 ~ 2.36) Mm⁻¹ in Changzhou and increased from 0.63 (0.43 ~ 0.91) Mm⁻¹ through 1.36 (1.01 ~ 1.79) Mm⁻¹ to 3.45 (2.46 ~ 5.34) Mm⁻¹ in Beijing. $\sigma_{ab,bulk,>700}$ increased by 3.2 (5.5) times in Changzhou (Beijing). The relative increase of $\sigma_{ab,bulk,>700}$ was overall consistent with that of $\sigma_{ab,bulk}$ in both Changzhou and Beijing. As a result, there was no significant change in $f_{ab,>700}$ with the development of pollution (Fig. 4b3). $f_{ab,>700}$ varied from 19.8 (15.2 ~ 23.8) % through 19.3 (15.9 ~ 25.3) % to 19.6 (15.5 ~ 24.5) % in Changzhou and varied from 27.9 (20.7 ~ 36.4) % through 23.2 (17.8 ~ 30.7) % to 26.7 (20.4 ~ 34.7) % in Beijing. It could be seen that the increase of $\sigma_{ab,bulk,>700}$ in Changzhou was less than that in Beijing with the development of pollution. Specifically, $\sigma_{ab,bulk,>700}$ in Beijing was 2.0 times larger than that in Changzhou, showing that the change of $\sigma_{ab,bulk,>700}$ with the aggravation of pollution could be different significantly in different sites.

3.3.4 Diurnal cycle

$\sigma_{ab,size-resolved}$ exhibited clear diurnal cycle in both Changzhou (Fig. 5a3) and Beijing (Fig. 5a4) with lower value of $\sigma_{ab,size-resolved}$

375 resolved during daytime and higher value during nighttime. Accordingly, $\sigma_{ab,bulk}$ had minimum value of 3.51 (3.16 ~ 4.26) Mm^{-1}
 376 ¹ at 14:00 and maximum value of 7.20 (3.80 ~ 10.58) Mm^{-1} at 01:00 in Changzhou (Fig. 5b3). $\sigma_{ab,bulk}$ had minimum value of
 377 3.96 (2.97 ~ 9.10) Mm^{-1} at 14:00 and maximum value of 7.86 (4.04 ~ 13.19) Mm^{-1} at 00:00 in Beijing (Fig. 5b4), reflecting
 378 the regulation by planetary boundary layer. In contrast, neither $\sigma_{ab,bulk,>700}$ in Changzhou (Fig. 5c3) nor $\sigma_{ab,bulk,>700}$ in Beijing
 379 (Fig. 5c4) exhibited obvious diurnal cycle. Therefore, $f_{ab,>700}$, inversely proportional to $\sigma_{ab,bulk}$, had higher value during
 380 daytime and lower value during nighttime. For Changzhou, $f_{ab,>700}$ reached maximum at 09:00 with value of 25.3 (20.4 ~
 381 27.4) % and came to minimum at 21:00 with value of 16.6 (13.0 ~ 19.6) % (Fig. 5d3). For Beijing, $f_{ab,>700}$ reached maximum
 382 at 10:00 with value of 30.4 (21.1 ~ 36.3) % and came to minimum at 01:00 with value of 24.5 (17.2 ~ 28.1) % (Fig. 5d4).

383 3.4 Direct radiative forcing of equivalent black carbon

384 3.4.1 Overview

385 The timeseries of DRF_{eBC} in Changzhou and Beijing was shown in Fig. S3a1 and Fig. S3b1 – S3b4, respectively. It could
 386 be seen that DRF_{eBC} varied significantly in both Changzhou and Beijing. DRF_{eBC} was estimated to be 0.93 (0.70 ~ 1.39) W
 387 m^{-2} in Changzhou and 1.10 (0.65 ~ 2.00) W m^{-2} in Beijing, respectively (Fig. 4c1). The variation of DRF_{eBC} was as large as
 388 the median value of DRF_{eBC} , clearly indicating the large variability of BC radiative effect. DRF_{eBC} increased substantially
 389 with the aggravation of pollution (Fig. 4c1). DRF_{eBC} increased from 0.38 (0.38 ~ 0.38) W m^{-2} through 0.77 (0.70 ~ 0.98) W
 390 m^{-2} to 1.67 (1.29 ~ 2.07) W m^{-2} by 4.4 times in Changzhou and from 0.42 (0.33 ~ 0.66) W m^{-2} through 1.17 (0.79 ~ 1.45) W
 391 m^{-2} to 2.41 (1.68 ~ 2.86) W m^{-2} by 5.7 times in Beijing with the development of pollution. A summary table with respect to
 392 DRF_{eBC} was presented in Table 1.

393 3.4.2 Contribution of equivalent black carbon-containing particle larger than 700 nm to direct radiative forcing of 394 equivalent black carbon

395 $DRF_{eBC,>700}$ was estimated to be 0.19 (0.13 ~ 0.26) W m^{-2} in Changzhou and 0.20 (0.13 ~ 0.37) W m^{-2} in Beijing (Fig. 4c2),
 396 respectively, which accounted for 20.5 (18.4 ~ 22.2) % and 21.0 (16.3 ~ 26.1) % of DRF_{eBC} (Fig. 4c3), respectively. Therefore,
 397 $eBC_{>700}$ contributed to an important portion of BC radiative effect. With the aggravation of pollution, $DRF_{eBC,>700}$ increased
 398 substantially and was different regionally (Fig. 4c2), $DRF_{eBC,>700}$ increased from 0.10 (0.10 ~ 0.10) W m^{-2} through 0.17 (0.12
 399 ~ 0.26) W m^{-2} to 0.24 (0.22 ~ 0.30) W m^{-2} by 2.4 times in Changzhou and from 0.10 (0.08 ~ 0.12) W m^{-2} through 0.20 (0.17
 400 ~ 0.24) W m^{-2} to 0.47 (0.34 ~ 0.71) W m^{-2} by 4.7 times in Beijing. The characteristics of $f_{DRF,>700}$ with increasing pollution
 401 was complicated (Fig. 4c3). $f_{DRF,>700}$ varied from 25.0 (25.0 ~ 25.0) % through 21.1 (20.3 ~ 22.3) % to 17.6 (15.5 ~ 18.9) %
 402 in Changzhou, exhibiting a decreasing trend. However, $f_{DRF,>700}$ varied from 24.4 (17.4 ~ 27.7) % through 18.4 (15.4 ~ 24.5) %
 403 to 21.5 (19.1 ~ 26.9) % in Changzhou, without systematical change.

404 4 Conclusions

405 Black carbon (BC) mass size distribution (BCMSD) was an important factor influencing environmental and radiative effect
 406 of BC. However, current BCMSD measurements mainly focused on BC-containing particle less than 700 nm. The

characteristics of BC-containing particle greater than 700 nm ($BC_{>700}$) remained uncertain due to limit in technique. In this study, the characteristics of equivalent $BC_{>700}$ ($eBC_{>700}$) were measured and studied based on field measurements in eastern China.

Equivalent BCMSD ($eBCMSD$) was measured from 150 nm up to 1.5 μm with time resolution of 1 hour based on the method proposed by Zhao et al. (2022), where $eBCMSD$ was determined by an aerodynamic aerosol classifier (AAC) in tandem with an aethalometer (model AE33, AAC – AE33) and size-resolved particle number concentration was measured concurrently to model the influence of particle size on mass absorption cross section (Zhao et al., 2021). AAC – AE33 was applied to two field measurements in eastern China, namely Changzhou located in the Yangtze River Delta from May 17th to June 3rd in 2021 and Beijing located in the North China Plain from October 29th 2021 to January 26th 2022. Changzhou was a regional background site and Beijing was a typical urban site. The direct radiative forcing of eBC (DRF_{eBC}) was estimated by Santa Barbara DISORT (discrete ordinates radiative transfer) atmospheric radiative transfer (SBDART) model (Ricchiazzi et al., 1998).

$eBCMSD$ was different between Changzhou and Beijing. Campaign-averaged $eBCMSD$ in Changzhou exhibited two modes, peaking at 240 nm and 1249 nm, respectively. In contrast, campaign-averaged $eBCMSD$ in Beijing exhibited one mode, peaking at 427 nm. $eBC_{>700}$ was ubiquitous in both Changzhou and Beijing. The campaign-averaged mass, absorption as well as radiative contribution of $eBC_{>700}$ to bulk eBC mass concentration ($m_{eBC,bulk}$), bulk absorption coefficient ($\sigma_{ab,bulk}$), as well as DRF_{eBC} in Changzhou and Beijing were 27.8 (20.9 ~ 36.5) % and 24.1 (17.5 ~ 34.2) %, 19.6 (15.8 ~ 24.6) % and 25.9 (19.6 ~ 33.7) %, as well as 20.5 (18.4 ~ 22.2) % and 21.0 (16.3 ~ 26.1) %, respectively, manifesting the important role of $eBC_{>700}$ in environment and climate. Both $eBCMSD$ and size-resolved absorption coefficient ($\sigma_{ab,size-resolved}$) exhibited diurnal variation with lower value during the daytime and higher value during the nighttime in both Changzhou and Beijing.

With the aggravation of pollution, the evolution of $eBCMSD$ and $\sigma_{ab,size-resolved}$ in Changzhou was significantly different from that in Beijing. The peak diameter of $eBCMSD$ shifted from 240 (347) nm to 289 (527) nm in Changzhou (Beijing) and the peak diameter of $\sigma_{ab,size-resolved}$ shifted from 416 (427) nm to 416 (527) nm in Changzhou (Beijing), indicating the aging process in regional background site was distinct from that in urban site. Both the value of $eBCMSD$ and $\sigma_{ab,size-resolved}$ increased with the development of pollution in both Changzhou and Beijing. Accordingly, $m_{eBC,bulk}$, $\sigma_{ab,bulk}$ and DRF_{eBC} in Changzhou (Beijing) increased by 3.2 (4.6) times, 3.5 (5.5) times and 4.4 (5.7) times, respectively. $m_{eBC,bulk}$, $\sigma_{ab,bulk}$ and DRF_{eBC} of $eBC_{>700}$ in Changzhou (Beijing) increased by 3.6 (5.1) times, 3.2 (5.5) times and 2.4 (4.7) times, respectively, clearly showing the large variation of $eBC_{>700}$. Case study exhibited that contribution of $eBC_{>700}$ to $m_{eBC,bulk}$, $\sigma_{ab,bulk}$ and DRF_{eBC} could increase from 20 % to 50 %, from about 20 % to 50 % and from 20 % to 40 %, respectively. Therefore, $BC_{>700}$ is an important part of BC-containing particles and it was highly recommended to take $BC_{>700}$ into account in both BC field measurement and model evaluation of BC climate effect.

Code and data availability

439 The code and measurement data involved in this study are available upon request to the authors. The data involved in this study is
440 also available online at: https://pan.baidu.com/s/1IE2lyPg0vb8O_GPTl-dSog?pwd=pzi8.

441 **Author contribution**

442 CZ determined the main goal of this study. WZ carried experiments out and prepared the paper with contributions from all co-
443 authors.

444 **Competing interests**

445 The authors declare that they have no conflict of interest.

446 **References**

- 447 Artaxo, P., Fernandes, E. T., Martins, J. V., Yamasoe, M. A., Hobbs, P. V., Maenhaut, W., Longo, K. M., and Castanho, A.: Large-
448 scale aerosol source apportionment in Amazonia, *J. Geophys. Res.-Atmos.*, 103, 31837-31847, 10.1029/98jd02346, 1998.
- 449 Berner, A., Reischl, G., and Puxbaum, H.: Size distribution of traffic derived aerosols, *Sci. Total Environ.*, 36, 299-303,
450 10.1016/0048-9697(84)90280-8, 1984.
- 451 Bond, T. C.: Spectral dependence of visible light absorption by carbonaceous particles emitted from coal combustion, *Geophys. Res.*
452 *Let.*, 28, 4075-4078, 10.1029/2001gl013652, 2001.
- 453 Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H., and Klimont, Z.: A technology-based global inventory of black
454 and organic carbon emissions from combustion, *J. Geophys. Res.-Atmos.*, 109, 43, 10.1029/2003jd003697, 2004.
- 455 Bond, T. C., and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Science and*
456 *Technology*, 40, 27-67, 10.1080/02786820500421521, 2006.
- 457 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Karcher, B., Koch,
458 D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin,
459 N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo,
460 T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys.*
461 *Res.-Atmos.*, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- 462 Chakrabarty, R. K., Beres, N. D., Moosmuller, H., China, S., Mazzoleni, C., Dubey, M. K., Liu, L., and Mishchenko, M. I.: Soot
463 superaggregates from flaming wildfires and their direct radiative forcing, *Sci Rep*, 4, 8, 10.1038/srep05508, 2014.
- 464 Chow, J. C., Watson, J. G., Crow, D., Lowenthal, D. H., and Merrifield, T.: Comparison of IMPROVE and NIOSH Carbon
465 Measurements, *Aerosol Science and Technology*, 34, 23-34, 10.1080/02786820119073, 2001.
- 466 Drinovec, L., Mocnik, G., Zotter, P., Prevot, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Muller, T., Wiedensohler,
467 A., and Hansen, A. D. A.: The "dual-spot" Aethalometer: an improved measurement of aerosol black carbon with real-time loading
468 compensation, *Atmospheric Measurement Techniques*, 8, 1965-1979, 10.5194/amt-8-1965-2015, 2015.
- 469 Fuller, K. A., Malm, W. C., and Kreidenweis, S. M.: Effects of mixing on extinction by carbonaceous particles, *J. Geophys. Res.-*
470 *Atmos.*, 104, 15941-15954, 10.1029/1998jd100069, 1999.

471 Guo, Y. H.: Characteristics of size-segregated carbonaceous aerosols in the Beijing-Tianjin-Hebei region, *Environmental Science*
472 *and Pollution Research*, 23, 13918-13930, 10.1007/s11356-016-6538-z, 2016.

473 Hansen, A. D. A., Rosen, H., and Novakov, T.: The aethalometer - an instrument for the real-time measurement of optical-absorption
474 by aerosol-particles, *Sci. Total Environ.*, 36, 191-196, 10.1016/0048-9697(84)90265-1, 1984.

475 Johnson, T. J., Irwin, M., Symonds, J. P. R., Olfert, J. S., and Boies, A. M.: Measuring aerosol size distributions with the aerodynamic
476 aerosol classifier, *Aerosol Science and Technology*, 52, 655-665, 10.1080/02786826.2018.1440063, 2018.

477 Kuang, Y., Zhao, C. S., Tao, J. C., and Ma, N.: Diurnal variations of aerosol optical properties in the North China Plain and their
478 influences on the estimates of direct aerosol radiative effect, *Atmospheric Chemistry and Physics*, 15, 5761-5772, 10.5194/acp-15-
479 5761-2015, 2015.

480 Liu, D., Joshi, R., Wang, J., Yu, C., Allan, J. D., Coe, H., Flynn, M. J., Xie, C., Lee, J., Squires, F., Kotthaus, S., Grimmond, S., Ge,
481 X., Sun, Y., and Fu, P.: Contrasting physical properties of black carbon in urban Beijing between winter and summer, *Atmos. Chem.*
482 *Phys.*, 19, 6749-6769, 10.5194/acp-19-6749-2019, 2019.

483 Liu, D. T., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S. F., Williams, P. I., Ting,
484 Y. C., Haslett, S., Taylor, J. W., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe, H., and Allan, J. D.: Black-carbon absorption
485 enhancement in the atmosphere determined by particle mixing state, *Nature Geoscience*, 10, 184-U132, 10.1038/ngeo2901, 2017.

486 Liu, P. F., Zhao, C. S., Zhang, Q., Deng, Z. Z., Huang, M. Y., Ma, X. C., and Tie, X. X.: Aircraft study of aerosol vertical distributions
487 over Beijing and their optical properties, *Tellus Ser. B-Chem. Phys. Meteorol.*, 61, 756-767, 10.1111/j.1600-0889.2009.00440.x,
488 2009.

489 Ma, N., Zhao, C. S., Muller, T., Cheng, Y. F., Liu, P. F., Deng, Z. Z., Xu, W. Y., Ran, L., Nekat, B., van Pinxteren, D., Gnauk, T.,
490 Mueller, K., Herrmann, H., Yan, P., Zhou, X. J., and Wiedensohler, A.: A new method to determine the mixing state of light absorbing
491 carbonaceous using the measured aerosol optical properties and number size distributions, *Atmospheric Chemistry and Physics*, 12,
492 2381-2397, 10.5194/acp-12-2381-2012, 2012.

493 Matsui, H., Hamilton, D. S., and Mahowald, N. M.: Black carbon radiative effects highly sensitive to emitted particle size when
494 resolving mixing-state diversity, *Nature Communications*, 9, 10.1038/s41467-018-05635-1, 2018.

495 Moosmuller, H., Chakrabarty, R. K., and Arnott, W. P.: Aerosol light absorption and its measurement: A review, *Journal of*
496 *Quantitative Spectroscopy & Radiative Transfer*, 110, 844-878, 10.1016/j.jqsrt.2009.02.035, 2009.

497 Peng, J. F., Hu, M., Guo, S., Du, Z. F., Zheng, J., Shang, D. J., Zamora, M. L., Zeng, L. M., Shao, M., Wu, Y. S., Zheng, J., Wang,
498 Y., Glen, C. R., Collins, D. R., Molina, M. J., and Zhang, R. Y.: Markedly enhanced absorption and direct radiative forcing of black
499 carbon under polluted urban environments, *Proceedings of the National Academy of Sciences of the United States of America*, 113,
500 4266-4271, 10.1073/pnas.1602310113, 2016.

501 Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus
502 activity, *Atmospheric Chemistry and Physics*, 7, 1961-1971, 10.5194/acp-7-1961-2007, 2007.

503 Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S. M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N.,
504 Wehrli, C., Wiedensohler, A., and Zhang, X. Y.: Recommendations for reporting "black carbon" measurements, *Atmospheric*
505 *Chemistry and Physics*, 13, 8365-8379, 10.5194/acp-13-8365-2013, 2013.

506 Ramachandran, S., and Rajesh, T. A.: Black carbon aerosol mass concentrations over Ahmedabad, an urban location in western
507 India: Comparison with urban sites in Asia, Europe, Canada, and the United States, *J. Geophys. Res.-Atmos.*, 112, 19,
508 10.1029/2006jd007488, 2007.

509 Ricchiazzi, P., Yang, S. R., Gautier, C., and Sowle, D.: SBDART: A research and teaching software tool for plane-parallel radiative
510 transfer in the Earth's atmosphere, *Bulletin of the American Meteorological Society*, 79, 2101-2114, 10.1175/1520-
511 0477(1998)079<2101:Sarats>2.0.Co;2, 1998.

512 Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., Wilson, J. C., Reeves, J. M., Darbeheshti, M., Baumgardner,
513 D. G., Kok, G. L., Chung, S. H., Schulz, M., Hendricks, J., Lauer, A., Karcher, B., Slowik, J. G., Rosenlof, K. H., Thompson, T. L.,
514 Langford, A. O., Loewenstein, M., and Aikin, K. C.: Single-particle measurements of midlatitude black carbon and light-scattering
515 aerosols from the boundary layer to the lower stratosphere, *J. Geophys. Res.-Atmos.*, 111, 15, 10.1029/2006jd007076, 2006.

516 Schwarz, J. P., Spackman, J. R., Fahey, D. W., Gao, R. S., Lohmann, U., Stier, P., Watts, L. A., Thomson, D. S., Lack, D. A., Pfister,
517 L., Mahoney, M. J., Baumgardner, D., Wilson, J. C., and Reeves, J. M.: Coatings and their enhancement of black carbon light
518 absorption in the tropical atmosphere, *J. Geophys. Res.-Atmos.*, 113, 10, 10.1029/2007jd009042, 2008.

519 Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Berntsen, T., Collins, W. D., Fuzzi, S., Gallardo, L., Kiendler Scharr, A., Klimont, Z.,
520 Liao, H., Unger, N., and Zanis, P.: Short-Lived Climate Forcers. In *Climate Change 2021: The Physical Science Basis. Contribution*
521 *of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte,
522 V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K.,
523 Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press.,
524 Cambridge, United Kingdom and New York, NY, USA, 2021.

525 Tan, W. S., Zhao, G., Yu, Y. L., Li, C. C., Li, J., Kang, L., Zhu, T., and Zhao, C. S.: Method to retrieve cloud condensation nuclei
526 number concentrations using lidar measurements, *Atmospheric Measurement Techniques*, 12, 3825-3839, 10.5194/amt-12-3825-
527 2019, 2019.

528 Tavakoli, F., and Olfert, J. S.: An Instrument for the Classification of Aerosols by Particle Relaxation Time: Theoretical Models of
529 the Aerodynamic Aerosol Classifier, *Aerosol Science and Technology*, 47, 916-926, 10.1080/02786826.2013.802761, 2013.

530 Wang, H. L., An, J. L., Zhu, B., Shen, L. J., Duan, Q., and Shi, Y. Z.: Characteristics of Carbonaceous Aerosol in a Typical Industrial
531 City-Nanjing in Yangtze River Delta, China: Size Distributions, Seasonal Variations, and Sources, *Atmosphere*, 8, 14,
532 10.3390/atmos8040073, 2017.

533 Wang, J. D., Wang, S. X., Wang, J. P., Hua, Y., Liu, C., Cai, J., Xu, Q. C., Xu, X. T., Jiang, S. Y., Zheng, G. J., Jiang, J. K., Cai, R.
534 L., Zhou, W., Chen, G. Z., Jin, Y. Z., Zhang, Q., and Hao, J. M.: Significant Contribution of Coarse Black Carbon Particles to Light

535 Absorption in North China Plain, *Environ. Sci. Technol. Lett.*, 9, 134-139, 10.1021/acs.estlett.1c00953, 2022.

536 Wex, H., Neususs, C., Wendisch, M., Stratmann, F., Koziar, C., Keil, A., Wiedensohler, A., and Ebert, M.: Particle scattering,
537 backscattering, and absorption coefficients: An in situ closure and sensitivity study, *J. Geophys. Res.-Atmos.*, 107, 18,
538 10.1029/2000jd000234, 2002.

539 Yu, H., Wu, C., Wu, D., and Yu, J. Z.: Size distributions of elemental carbon and its contribution to light extinction in urban and
540 rural locations in the pearl river delta region, China, *Atmospheric Chemistry and Physics*, 10, 5107-5119, 10.5194/acp-10-5107-
541 2010, 2010.

542 Zhang, R. Y., Khalizov, A. F., Pagels, J., Zhang, D., Xue, H. X., and McMurry, P. H.: Variability in morphology, hygroscopicity, and
543 optical properties of soot aerosols during atmospheric processing, *Proceedings of the National Academy of Sciences of the United
544 States of America*, 105, 10291-10296, 10.1073/pnas.0804860105, 2008.

545 Zhao, G., Zhao, C. S., Kuang, Y., Bian, Y. X., Tao, J. C., Shen, C. Y., and Yu, Y. L.: Calculating the aerosol asymmetry factor based
546 on measurements from the humidified nephelometer system, *Atmospheric Chemistry and Physics*, 18, 9049-9060, 10.5194/acp-18-
547 9049-2018, 2018.

548 Zhao, G., Tao, J. C., Kuang, Y., Shen, C. Y., Yu, Y. L., and Zhao, C. S.: Role of black carbon mass size distribution in the direct
549 aerosol radiative forcing, *Atmospheric Chemistry and Physics*, 19, 13175-13188, 10.5194/acp-19-13175-2019, 2019.

550 Zhao, G., Yu, Y., Tian, P., Li, J., Guo, S., and Zhao, C.: Evaluation and Correction of the Ambient Particle Spectral Light Absorption
551 Measured Using a Filter-based Aethalometer, *Aerosol and Air Quality Research*, 20, 1833-1841, 10.4209/aaqr.2019.10.0500, 2020.

552 Zhao, W., Zhao, G., Li, Y., Guo, S., Ma, N., Tang, L., Zhang, Z., and Zhao, C.: New method to determine black carbon mass size
553 distribution, *Atmos. Meas. Tech.*, 15, 6807-6817, 10.5194/amt-15-6807-2022, 2022.

554 Zhao, W. L., Tan, W. S., Zhao, G., Shen, C. Y., Yu, Y. L., and Zhao, C. S.: Determination of equivalent black carbon mass
555 concentration from aerosol light absorption using variable mass absorption cross section, *Atmospheric Measurement Techniques*,
556 14, 1319-1331, 10.5194/amt-14-1319-2021, 2021.

557




Figure 1: Instrumental setup used in this study. Instruments used to measure $N_{\text{size-resolved}}$ was colored with red (green) for Changzhou (Beijing).






Figure 2: (a) eBCMSD from October 31st 2021 to November 6th 2021 in Beijing and (b) the corresponding $f_{m, > 700}$ (red solid line), $f_{ab, > 700}$ (green dashed line) as well as $f_{DRF, > 700}$ (black dotted line with triangle marker). The black solid line was \bar{D}_p .



570
 571 **Figure 3: Normalized pdf of eBCMSD measured in (a1 – d1) Changzhou and (a2 – d2) Beijing as well as $\sigma_{ab,size-resolved}$**
 572 **measured in (a3 – d3) Changzhou and (a4 – d4) Beijing. (a1 – a4), (b1 – b4), (c1 – c4) and (d1 – d4) were statistics over**
 573 **the whole campaign, clean period, transitional period and polluted period. Red solid line and red dashed lines were**
 574 **median and lower as well as upper quartiles.**



588
 589 **Figure 4: Box plots of (a1) $m_{eBC,bulk}$, (a2) $m_{eBC,bulk,>700}$, (a3) $f_{m,>700}$, (b1) $\sigma_{ab,bulk}$, (b2) $\sigma_{ab,bulk,>700}$, (b3) $f_{ab,>700}$, (c1) DRF_{eBC} ,**
 590 **(c2) $DRF_{eBC,>700}$ and (c3) $f_{DRF,>700}$ over the whole campaign (All), clean (C), transitional (T) as well as polluted (P) period,**
 591 **respectively. The box extended from the first quartile to the third quartile with a line at the median. The whiskers**
 592 **marked 5 % and 95 % percentile. The circle inside the box was the mean value. Statistics from Changzhou (Beijing)**
 593 **were colored red (green). The 95 percentile of $m_{eBC,bulk,>700}$ under polluted period for Beijing (a2) was $1.00 \mu g m^{-3}$. The**
 594 **95 percentile of $\sigma_{ab,bulk,>700}$ and that under polluted period for Beijing (b2) was 7.80 and 10.30 Mm^{-1} , respectively. The**
 595 **95 percentile of $DRF_{eBC,>700}$ under polluted period for Beijing (c2) was $1.41 W m^{-2}$.**
 596




Figure 5: Diurnal variation of (a1) eBCMSD, (b1) $m_{eBC,bulk}$, (c1) $m_{eBC,bulk,>700}$, (d1) $f_{m,>700}$ in Changzhou; (a2) eBCMSD, (b2) $m_{eBC,bulk}$, (c2) $m_{eBC,bulk,>700}$, (d2) $f_{m,>700}$ in Beijing; (a3) $\sigma_{ab,size-resolved}$, (b3) $\sigma_{ab,bulk}$, (c3) $\sigma_{ab,bulk,>700}$, (d3) $f_{ab,>700}$ in Changzhou and (a4) $\sigma_{ab,size-resolved}$, (b4) $\sigma_{ab,bulk}$, (c4) $\sigma_{ab,bulk,>700}$, (d4) $f_{ab,>700}$ in Beijing. Red solid line and green dashed lines were median and lower as well as upper quartiles.

Table 1: Summary of evolution of m_{eBC} , σ_{ab} and DRF_{eBC} .

	size range	clean		transition		polluted		average	
		Changzhou	Beijing	Changzhou	Beijing	Changzhou	Beijing	Changzhou	Beijing
m_{eBC}	bulk	0.41(0.33~0.45)	0.32(0.22~0.41)	0.71(0.58~0.83)	0.73(0.61~0.85)	1.33(1.16~1.71)	1.47(1.21~1.82)	0.73(0.52~1.03)	0.79(0.43~1.31)
	> 700 nm	0.11(0.07~0.15)	0.07(0.05~0.12)	0.20(0.14~0.27)	0.17(0.11~0.23)	0.40(0.29~0.50)	0.36(0.25~0.52)	0.20(0.13~0.32)	0.18(0.10~0.33)
σ_{ab}	bulk	2.71(2.30~3.28)	2.47(1.65~3.28)	4.83(4.04~6.02)	5.93(4.72~7.33)	9.61(7.99~11.93)	13.65(10.94~17.59)	4.93(3.53~7.24)	6.37(3.31~11.68)
	> 700 nm	0.54(0.62~1.59)	0.63(0.43~0.91)	0.96(0.72~1.32)	1.36(1.01~1.79)	1.75(1.53~2.36)	3.45(2.46~5.34)	1.03(0.62~1.59)	1.47(0.81~2.83)
DRF_{eBC}	bulk	0.38(0.38~0.38)	0.42(0.33~0.66)	0.77(0.70~0.98)	1.17(0.79~1.45)	1.67(1.29~2.07)	2.41(1.68~2.86)	0.93(0.70~1.39)	1.10(0.65~2.00)
	> 700 nm	0.10(0.10~0.10)	0.10(0.08~0.12)	0.17(0.12~0.26)	0.20(0.17~0.24)	0.24(0.22~0.30)	0.47(0.34~0.71)	0.19(0.13~0.26)	0.20(0.13~0.37)

616