

Supplements of

Roles of marine biota in the formation of atmospheric bioaerosols, cloud condensation nuclei, and ice-nucleating particles over the North Pacific Ocean, Bering Sea, and Arctic Ocean

5

Kaori Kawana¹, Fumikazu Taketani¹, Kazuhiko Matsumoto¹, Takuma Miyakawa¹, Yutaka Tobo², Yoko Iwamoto³, Akinori Ito⁴, Yugo Kanaya¹

¹ Earth Surface System Research Center, Research Institute for Global Change, Japan Agency for Marine-Earth
10 Science and Technology (JAMSTEC), Yokohama, Kanagawa, 2360001, Japan

² National Institute of Polar Research, Tachikawa, Tokyo, 1908518, Japan

³ Graduate School of Integrated Sciences for Life, Hiroshima University, Higashi Hiroshima, Hiroshima, 7390046,
Japan

⁴ Yokohama Institute for Earth Sciences, JAMSTEC, Yokohama, Kanagawa, 2360001, Japan

15 *Correspondence to:* Kaori Kawana (kawanak@jamstec.go.jp)




Figure S1. (a) Mass concentration and (b) mass fraction of chemical components in PM_{2.5} aerosol particles in each period. (c) Number concentrations and (d) fractions of types of fluorescent fine particles ($1 < D_p < 2.5 \mu\text{m}$), and (e) and (f) those of fluorescent coarse particles ($D_p > 2.5 \mu\text{m}$).




Figure S2. Time series of nutrient concentrations in seawater.




Figure S3. (a)–(d) Same as Fig. 4 but for fluorescent particles with diameter of $>2.5 \mu\text{m}$.





Figure S4. Scatterplots of nonfluorescent particles and bioindicators as (a) TEPs, (b) CSPs, and (c) Chl-*a*, and
 5 scatterplots of nonfluorescent particles and (d) WS, and (e) TEPs, (f) CSPs, and (g) Chl-*a*, multiplied with WS. Colored lines represent the orthogonal regression lines. Error bars represent one standard deviation.



5

Figure S5. (a)–(l) Same as Fig. 8 but with exclusion of the highest data point.