

The underappreciated role of transboundary pollution in future air quality and health improvements in China

Jun-Wei Xu¹, Jintai Lin^{1*}, Dan Tong², Lulu Chen¹

¹Laboratory for Climate and Ocean–Atmosphere Studies, Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, China

²Department of Earth System Science, Ministry of Education Key Laboratory for Earth System Modelling, Tsinghua University, Beijing, China

Correspondence: Jintai Lin (linjt@pku.edu.cn).

Abstract

Studies assessing the achievability of future air quality goals in China have focused on the role of reducing China’s domestic emissions, yet the influence of transboundary pollution of foreign origins has been largely underappreciated. Here, we assess the extent to which future changes in foreign transboundary pollution would affect the achievability of air quality goals in 2030 and 2060 for China. We find that in 2030, under the current-policy scenario in China, transboundary contributions to population-weighted PM_{2.5} in China would be reduced by 29% (1.2 μg m⁻³) as foreign countries transition from the fossil fuel-intensive to the low-carbon scenario. By 2060, the difference would be increased to 45% (1.8 μg m⁻³). Adopting the low-carbon instead of the fossil fuel-intensive pathway in foreign countries would avoid 10 million Chinese people from being exposed to PM_{2.5} concentrations above China’s Ambient Air Quality Standard (35 μg m⁻³) in 2030 and 5 million Chinese people from being exposed to PM_{2.5} concentrations above the World Health Organization Air Quality Guideline (5 μg m⁻³) in 2060. Meanwhile, China adopting the carbon-neutral pathway rather than its current pathway would also be helpful to reduce transboundary PM_{2.5} produced from the chemical interactions between foreign-transported and locally-emitted pollutants. In 2060, adopting a low-carbon pathway in China and foreign countries coincidentally would avoid 63% of transboundary pollution and 386,000 associated premature deaths in China, relative to adopting a fossil fuel-intensive pathway in both regions. Thus, the influence of transboundary pollution should be carefully considered when making future air quality expectations and pollution mitigation strategies.

1. Introduction

Long-term exposure to ambient fine particulate matter (PM_{2.5}, particulate matter smaller than 2.5 μm in aerodynamic diameter) is the largest environmental risk factor for human health, with an estimated 4.1 million attributable deaths worldwide (7.3% of the number of global deaths in 2019; Murray et al., 2020). Countries have taken diverse actions to improve air quality and public health, including setting ambitious future air quality and/or climate goals. The European Commission set the 2030 air quality goal as reducing the number of PM_{2.5}-attributable premature deaths by at least 55% compared with 2005 levels, equivalent to reducing PM_{2.5} concentrations to below 10 μg m⁻³ EU-wide (European Commission, 2022). The United States aims for a 50–52% reduction of greenhouse gas emissions relative to 2005 levels by 2030, which would avoid tens of thousands of premature deaths associated with PM_{2.5} as a co-benefit (Burtraw et al.,

Formatted: Indent: First line: 0.25 cm

Deleted:

Deleted: We also find that a

Deleted:

Deleted: We find that adopting the low-carbon instead of the fossil fuel-intensive pathway in foreign countries would avoid millions of Chinese people from being exposed to PM_{2.5} concentrations above China’s air quality standard level (35 μg m⁻³) in 2030 and the World Health Organization Air Quality Guideline (5 μg m⁻³) in 2060.

1 2021). However, the achievability of air quality goals or the extent of health co-benefits of
2 climate strategies is subject to large uncertainties as it is affected by a wide range of factors such
3 as the domestic and global socioeconomic development pathways, energy choices, and air
4 pollution control measures (Cheng et al., 2021b; O'Neill et al., 2020; Rao et al., 2017).

5 China is a key country to examine factors affecting the achievability of future air quality goals.
6 On the one hand, China suffers from serious air pollution and adverse health effects, with more
7 than a quarter of the world's total PM_{2.5}-associated premature deaths in 2015 estimated to occur
8 in China (Zhang et al., 2019). Despite remarkable achievements of the 5-year Clean Air Action
9 since 2013 (Zhang et al., 2019; Zheng et al., 2018a), annual mean PM_{2.5} concentration in China
10 still exceeded the newly revised World Health Organization (WHO) Air Quality Guideline
11 (WHO, 2021; AQG; 5 µg m⁻³) by 8 times by the end of the Clean Air Action in 2017. On the
12 other hand, China has set the most challenging air quality and climate goals among all the
13 developing countries in the world (Ascensão et al., 2018). In 2018, the Chinese government
14 proposed the roadmap for The Beautiful China Initiative (The State Council of the People's
15 Republic of China, 2016), which requires all cities to achieve the national ambient air quality
16 standards (NAAQS, 35 µg m⁻³) between 2030 and 2035 (The State Council of the People's
17 Republic of China, 2016). In 2019, China further announced an ambitious climate commitment
18 to achieve carbon neutrality by 2060. These policies have been regarded as interim steps towards
19 the WHO AQG. Therefore, investigating potential pathways and factors shaping the future air
20 quality in China could provide an excellent reference for other countries facing the dual
21 challenges of economic development and environmental protection.

22
23 A number of studies seek to answer whether the air quality goals in China can be achieved in
24 the future. These studies have been primarily focused on potential mitigation pathways to reduce
25 China's domestic emissions. Cheng et al. (2021b) found that the nation's current emission
26 control measures could reduce China's PM_{2.5} levels to below 30 µg m⁻³ by 2030, yet the benefits
27 of such measures would be mostly exhausted by then (Cheng et al., 2021b; Xing et al., 2020).
28 Xing et al. (2020) and Tang et al. (2022) suggested that the co-benefits of climate targets alone
29 (even the most ambitious target of 1.5 °C limit in global warming) were not able to help China
30 achieve the 35 µg m⁻³ target by 2035. Instead, Cheng et al. (2021b) proposed that a combination
31 of stringent clean air policies and ambitious climate targets (i.e., carbon neutrality by 2060 or a
32 global warming limit of 1.5 °C) could successfully reduce PM_{2.5} to below 35 µg m⁻³ by 2030 and
33 to below 10 µg m⁻³ (WHO interim target 4) by 2060. This would require a fundamental
34 transformation of China's economic-environmental development pathway by phasing out
35 polluting industries and moving towards renewable energy while implementing strict end-of-pipe
36 emission control.

37
38 However, these previous studies have largely neglected how the future changes in foreign
39 transboundary pollution would affect air quality in China, likely due to the perception that the
40 relatively short lifetime of PM_{2.5} (a few days) does not permit long-distance transport. This
41 negligence would put into question the confidence of their estimated achievability of air quality
42 goals in China. Due to large uncertainties in the future socioeconomic development pathways,
43 environmental commitments, financial supports and technology capabilities, future emissions
44 from China's neighboring countries might be highly variable, leading to different levels of
45 transboundary impacts to China's air quality. For example, emissions in South Asia, Central Asia
46 and Southeast Asia have been estimated to increase in the future by various projections

1 (International Energy Agency, 2021; Koplitz et al., 2017), due to their rapid-economic growth
2 and a lack of clear commitments on energy choices, climate actions and air pollution control
3 efforts. In this case, transboundary pollution from neighboring countries to China could
4 potentially become increasingly important to affect the achievability of air quality goals in
5 China. Alternatively, these surrounding countries may undergo a sustainable development
6 pathway, facilitated in part by external financial aids and technology supports, leading to
7 lowered transboundary pollution to China. Thus, the different prospects of transboundary
8 pollution could be a significant yet highly uncertain factor in the achievement of future air
9 quality goals in China.

10
11 In addition, the mechanism of transboundary pollution poses additional complexity to its
12 impacts on China's air quality. Foreign emissions can affect air quality in China through direct
13 transboundary transport in the atmosphere. Moreover, portions of foreign-transported pollution
14 can also interact chemically with China's locally emitted pollutants. Xu et al. (2023) found that
15 the transport and transformation of non-methane volatile organic compounds (NMVOCs) from
16 foreign countries could enhance the atmospheric oxidizing capacity and facilitate the oxidation
17 of Chinese nitrogen oxides (NO_x) to form nitric acid (HNO₃) and nitrate over North China
18 (referred to as eastern China in that study), leading to considerable foreign contributions to
19 China's air pollution.

20
21 The chemical interactions mean that the extent of transboundary pollution will depend on
22 emission changes in China as well. Therefore, the influence of transboundary pollution on
23 China's air quality in the future is a complex result of future emissions in China and in foreign
24 countries, along with their interactions. However, whether the changes in transboundary
25 pollution via direct transport and chemical interactions could affect the achievement of future air
26 quality goals in China and to what extent the influence could be remain poorly understood.

27
28 Here, we assess the potential influences that transboundary pollution could make on the
29 achievement of future air quality goals in China, considering the changes in direct pollution
30 transport and in the interactions between transported and China's locally emitted pollution. We
31 regard the air quality goal stated in The Beautiful China Initiative, which is that all cities have
32 annual mean PM_{2.5} concentrations below 35 μg m⁻³ between 2030–2035 (The State Council of
33 the People's Republic of China, 2016), as China's 2030 air quality goal. We also regard the
34 WHO AQG (annual mean PM_{2.5} below 5 μg m⁻³) as China's 2060 air quality goal. We then
35 discuss the likelihood of the achievement under currently proposed development pathways in
36 China and foreign countries. As detailed in Method, given the large uncertainties on future
37 emissions of a country (Rao et al., 2017), we consider three anthropogenic emission scenarios
38 (low: SSP119, medium: SSP245, and high: SSP370; O'Neill et al., 2014) for foreign countries
39 and two anthropogenic emission scenarios (current-policy and carbon-neutral; Cheng et al.,
40 2021b; Tong et al., 2020) for China (Table 1), so as to understand transboundary pollution in
41 China from the present (2015) to the future (2030 and 2060) under a wide range of plausible
42 futures. We do not consider the effects of physical climate change (e.g., temperature and
43 precipitation) on future transboundary pollution because Jiu et al. (2021) found that future PM_{2.5}
44 changes in China driven by anthropogenic emission reduction was 7 times more than the changes
45 due to meteorological fields by 2050 in both SSP126 and SSP585 scenarios. Other studies have

Formatted: Indent: First line: 0.59 ch

Deleted: (

Deleted: ,

Deleted: d

Deleted: eastern

Deleted: the

Deleted: of

Deleted: nitrate and PM2.5

Deleted: to eastern China

Formatted: English (CAN)

Deleted: , leading to additional transboundary effects (e.g., through formation of nitrate and ammonium; Xu et al., 2022).

Deleted: and

Deleted: (

Deleted: ,

Formatted: Subscript

1 ~~also indicted that the impact of climate change on future PM_{2.5} were~~ smaller than those of
2 anthropogenic emissions of pollutants (Hong et al., 2019; Jiang et al., 2013; Silva et al., 2017).
3 For each combination of foreign and Chinese anthropogenic emission scenarios, we simulate
4 PM_{2.5} concentrations at a 0.5° x 0.625° resolution using a chemical transport model (GEOS-
5 Chem; <http://www.geos-chem.org>) that can represent the complex pollutant emissions and
6 chemical reactions across a large spatial domain. Then, we correct the systematic bias in the
7 model simulated PM_{2.5} concentrations using a large set of ground-based observations of PM_{2.5} in
8 China; the same correction factor is applied to all present and future concentrations. Finally, we
9 quantify the transboundary impacts on Chinese PM_{2.5} under each emission scenario. We further
10 quantify the corresponding transboundary impacts on public health in China, measured by
11 premature deaths, using socio-demographic projections consistent with SSPs and the state-of-
12 the-art concentration–response relationships (the GEMM model; Burnett et al., 2018).

Deleted: and natural emissions of pollutants on future transboundary pollution, since their impacts on PM_{2.5}

Deleted: are

Formatted: Subscript

14 2. Method

16 In this study, we use a set of data and models to investigate future air quality and health burden
17 in China. Projected air pollutant emissions for foreign countries under SSP-RCP scenarios are
18 obtained from the International Institute for Applied Systems Analysis (IIASA; Rao et al., 2017;
19 Riahi et al., 2017) with updates on base year emissions and the harmonization year in this study.
20 Projected air pollutant emissions for China are developed by Tong et al. (2020) and Cheng et al.
21 (2021b). Ambient PM_{2.5} concentrations under each scenario are simulated by the GEOS-Chem
22 chemical transport model (<http://www.geos-chem.org>) and further corrected for systematic bias
23 by a suite of ground-based observations (Xu et al., 2023). PM_{2.5}-associated mortalities are
24 calculated using the Global Exposure Mortality Model (GEMM; Burnett et al., 2018).

25 2.1 Scenarios of future anthropogenic air pollutant emissions

27 Future pollutant emission outcome is a cumulative result of a range of variables including
28 socio-economic development, technological change, efficiency improvements and policies
29 directed at pollution control as well as alternative concerns including climate change, energy
30 access, and agricultural production (Rao et al., 2017). The Shared Socioeconomic Pathways
31 (SSP) includes 5 five distinctly different pathways about how the future might unfold in terms of
32 major socioeconomic, demographic, technological, lifestyle, policy and institutional trends
33 (O'Neill et al., 2014; van Vuuren et al., 2017). ~~We select three scenarios to represent low,~~
34 ~~medium and high emission cases:~~ SSP1 - sustainability, SSP2 - middle-of-the-road, SSP3 -
35 regional rivalry. An assumption about the degree of air pollution control (strong, medium or
36 weak) is included on top of the baseline pathway. Weak air pollution controls occur in SSP3,
37 with medium controls in SSP2 and strong controls in SSP1 (Turnock et al., 2020). However, SSP
38 scenarios do not include explicit climate policies (O'Neill et al., 2020). Instead, the
39 Representative Concentration Pathways (RCPs) generate climate projections targeting at a range
40 of climate forcing levels, such as 1.9 W m⁻² (1.5 °C warming), 4.5 W m⁻² (3 °C) and 7.0 W m⁻²
41 (4 °C) in 2100. Thus, the combination of SSP-RCP scenario framework depicts societal and
42 climate futures in parallel and explores plausible futures of human activities, the changing
43 climate and emissions (O'Neill et al., 2020).

Deleted:

Deleted: , SSP4 - inequality, SSP5 - fossil fuel development

Deleted: and SSP4

Deleted: and SSP5

1 Here, we use anthropogenic aerosol emissions projected under SSP-RCP scenarios for foreign
2 countries. We select 3 scenarios to represent low, middle and high air pollutant emissions in
3 plausible futures: SSP119 (a sustainable development pathway targeting a rise of the global
4 mean surface temperature below 1.5 °C from the pre-industrial levels by the end of the century),
5 SSP245 (a business-as-usual development pathway with 3 °C warming), SSP370 (a regional-
6 rivalry development pathway with 4 °C warming). Table 1 summarizes the scenario settings in
7 more detail.

8
9 The original SSP-RCP anthropogenic emissions future projection starts from 2015. Here, we
10 update the start year to 2019, which is the latest year that anthropogenic emissions from the
11 Community Emissions Data System (CEDS; Hoesly et al., 2018) are available. Because CEDS
12 historical emissions and SSP-RCP future emissions were developed upon different assumptions
13 and methods, it is important to harmonize the two datasets to ensure smooth transitions between
14 the two sets of emissions trajectories (Gidden et al., 2019). The update of the harmonization year
15 from 2015 to 2019 in this study could better represent the actual emission trajectory between
16 2015 and 2019. We also use the most recently developed CEDS emissions version 2
17 (<https://data.pnnl.gov/dataset/CEDS-4-21-21>) to harmonize the future SSP-RCP emission
18 projections as the new emissions can better represent the historical trend of pollutant emissions
19 ([https://github.com/JGCRI/CEDS/blob/master/documentation/Version_comparison_figures_v_2021_04_21_vs_v_2016_07_16\(CMIP6\).pdf](https://github.com/JGCRI/CEDS/blob/master/documentation/Version_comparison_figures_v_2021_04_21_vs_v_2016_07_16(CMIP6).pdf)). The final future SSP-RCP emissions for foreign
20 countries used in this study are presented in Fig. S1a.

22 Future scenarios of anthropogenic pollutant emissions for China were developed by Tong et al.
23 (2020) and Cheng et al. (2021b). We adopt these emissions instead of SSP-RCP emissions
24 because Tong et al. (2020) and Cheng et al. (2021b) developed the emissions specifically to meet
25 China's recent climate goals (i.e., 2030 carbon peak and 2060 carbon neutrality) and used
26 parameters (i.e., emission factors, energy use, etc.) to reflect the most up-to-date pollutant
27 control policies (i.e., clean air action since 2013) and technologies in China. Their emissions
28 have been suggested to better capture China's PM_{2.5} concentration decline during 2015–2019
29 than those driven by SSP-RCP emissions (Cheng et al., 2021a). We select two plausible
30 scenarios: the current-policy scenario and the carbon-neutral scenario. The current-policy
31 scenario seeks to achieve China's NDC pledges and the national PM_{2.5} air quality goal (i.e. 35 µg
32 m⁻³) by 2030, elucidating China's future air pollution mitigation pathway towards all the released
33 and determined upcoming clean air policies since 2015. The carbon-neutral scenario pursues
34 China's carbon-neutral commitment and the WHO's old PM_{2.5} guideline (10 µg m⁻³) by 2060. It
35 implements the best available end-of-pipe technologies and more stringent pollution control
36 policies than the current-policy scenario. Future anthropogenic pollutant emissions for China
37 under these scenarios are developed by firstly simulating China's future energy and
38 socioeconomic evolution using the Global Change Assessment Model (GCAM-China) and then
39 translating into pollutant emissions by the Dynamic Projection model for Emissions in China
40 (DPEC; Tong et al., 2020). More details are summarized in Table 1. The actual future air
41 pollutant emissions for China used in this study are presented in Fig. S1b.

42 2.2 Simulations of ambient PM_{2.5} concentrations

43

Deleted:

Deleted: future

Deleted: are harmonized to match anthropogenic emissions from the Community Emissions Data System (CEDS; Hoesly et al., 2018) for 2015, so that the resulting future trajectories provide a smooth transition from the historical emissions (Gidden et al., 2019).

Deleted: Here, we update the harmonization year from 2015 to the most recent year 2019 in CEDS historical emissions.

Deleted: are

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Deleted: .

Formatted: Font: (Asian) +Body Asian (DengXian)

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Formatted: Font: (Default) Times New Roman, (Asian) +Body Asian (DengXian), Not Bold

Deleted: Briefly,

Deleted: w

Formatted: Font: (Asian) +Body Asian (DengXian), Font color: Auto

1 We use the GEOS-Chem model to simulate PM_{2.5} concentrations in China and other Asian
2 countries under each emission scenario and year. Detailed descriptions of the model
3 configurations can be found in Xu et al. (2023). Briefly, we use the Flex-Grid capability of the
4 GEOS-Chem classic model v13.2.1 to simulate PM_{2.5} concentrations over Asia and its adjacent
5 regions (11° S–60° N, 30°–150° E; covering the entire Asia, eastern Africa and eastern Europe).
6 We use MERRA-2 assimilated meteorological data provided by the Global Modeling and
7 Assimilation Office (GMAO) at NASA Goddard Space Flight Center to drive the model. Our
8 simulation is conducted at a horizontal resolution of 0.5° latitude × 0.625° longitude with 47
9 vertical levels between the surface and ~ 0.01 hPa. Chemical boundary conditions are taken
10 from corresponding global simulations under each emission scenario in Table 2 at a resolution of
11 2° latitude × 2.5° longitude. The regeneration of boundary conditions under each emission
12 scenario could ensure the inclusion of pollution transported from countries both within and
13 outside the Flex-Grid domain as transboundary pollution to China. We spin up every simulation
14 for 1 month to remove the effects of initial conditions.

16 Anthropogenic emissions for the base year (2015) for China are taken from the Multi-
17 resolution Emission Inventory (MEIC) for 2015 (Zheng et al., 2018b), and for the rest of the
18 world are taken from the Community Emissions Data System (CEDS) version 2 for 2015
19 (<https://data.pnnl.gov/dataset/CEDS-4-21-21>). For future simulations, anthropogenic emissions
20 for China and foreign countries for each scenario are described above and are specified in Table
21 2. Other emissions are default in GEOS-Chem following Xu et al. (2023) and are fixed in all
22 present and future scenarios.

24 We conduct simulations for January, April, July and October, and treat the mean of the four
25 months as annual mean. Our meteorological fields are fixed to 2015 for all scenarios and years to
26 exclude the influence of climate on the results. More details of our model configurations can be
27 found in Table 2. We conduct two types of simulations: 1) baseline simulations (simulations with
28 “Base_” prefix in Table 2) that include complete anthropogenic emissions for both China and
29 foreign countries; 2) sensitivity simulations (simulations with “China_” prefix in Table 2) that
30 exclude anthropogenic emissions for foreign countries from the baseline simulation. Baseline
31 simulations calculate PM_{2.5} concentrations in China that are driven by both Chinese and foreign
32 emissions, while sensitivity simulations calculate PM_{2.5} concentrations in China that are driven
33 merely by China’s domestic emissions. The impacts of transboundary pollution on China’s PM_{2.5}
34 is calculated as the difference in China’s PM_{2.5} between a baseline simulation and a sensitivity
35 simulation in a specified year and under a specified Chinese emission scenario.

37 We correct our simulated PM_{2.5} concentrations under each scenario, based on a large set of
38 ground-based observations of PM_{2.5}, because our simulated PM_{2.5} concentrations are biased high
39 by roughly 15.7% (Fig. S2). Descriptions and data screening method of our ground-based
40 observations can be found in Xu et al. (2023). Further evaluations of PM_{2.5} composition
41 concentrations in China and PM_{2.5} concentrations in other Asian countries with ground-based
42 observations are presented in Xu et al. (2023). We correct simulated PM_{2.5} concentrations in
43 anthropogenic concentration-dominated grid cells (where anthropogenic emissions exceed
44 natural emissions) by observations located in anthropogenic concentration-dominated grid cells.
45 Similarly, we correct simulated concentrations in natural concentration-dominated grid cells
46 (where natural emissions exceed anthropogenic emissions) by observations located in natural

Formatted: Indent: First line: 0.25 cm

Deleted: (

Deleted: ,

Deleted: A number of previous studies have applied the GEOS-Chem to simulate PM_{2.5} concentrations and have shown consistency between observations and model results (Choi et al., 2019; Kopplitz et al., 2017; Venkataraman et al., 2018; Zhang et al., 2017).

Deleted: W

Deleted: area

Deleted: China, Southern Asia, Northern Asia and Central Asia

Deleted:

Deleted: The lowest vertical layer has a thickness of about 130 m. We regard the pollutant concentrations in this layer as “ground-level”. Detailed descriptions of the flex-grid setup can be found at <http://wiki.seas.harvard.edu/geos-chem/index.php/FlexGrid>. Our Flex-Grid domain extend the traditionally-defined nested Asia domain (11° S–55° N, 60°–150° E) in the model to better represent the transport of anthropogenic pollutants from Central Asia to China.

Our simulations are driven by assimilated meteorological data from MERRA-2 provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center. Convective transport in the model is computed from the convective mass fluxes in the meteorological archive as described by Wu et al. (2007). A non-local scheme is used to represent vertical mixing within the planetary boundary layer (PBL), as it accounts for different states of mixing based on the static instability (Lin and McElroy, 2010).

Formatted: Font: (Default) Times New Roman, Not Bold

Deleted: .

Deleted: ¶

GEOS-Chem simulates PM_{2.5} concentrations as the sum of sulfate (SO₄²⁻), nitrate (NO₃⁻), ammonium (NH₄⁺), organic [1]

Deleted:

Deleted: (

Deleted: ,

Deleted: Fine anthropogenic fugitive dust emissions from combustion and industrial sources for countries except China

Field Code Changed

Deleted: 2

Field Code Changed

Deleted: 2

Deleted:

Deleted: We correct PM_{2.5} concentrations by observations.

Deleted: grids

Deleted: grids

1 concentration-dominated grid cells. Natural and anthropogenic concentration-dominated grid
 2 cells in the model are shown in Fig. S3. To isolate anthropogenic concentration-dominated grid
 3 cells, we conduct a sensitivity that excludes natural emissions in China and in foreign countries.
 4 PM_{2.5} concentrations from this sensitivity simulation are referred to as natural PM_{2.5}
 5 concentrations. The difference between concentrations from the Base_2015 simulation (with
 6 complete natural and anthropogenic emissions) in Table 2 and the natural concentrations is the
 7 anthropogenic emission-contributed concentrations (anthropogenic concentrations). We scale
 8 anthropogenic concentrations by the average ratio of observed concentrations and simulated
 9 concentrations at each observation site that falls into an anthropogenic dominated grid cell.
 10 Concentrations before and after the correction for anthropogenic and all grid cells (including
 11 natural grid cells) are shown in Fig. S2. The overestimation in both anthropogenic and total
 12 PM_{2.5} concentrations is removed after the correction.

Deleted: in grids where PM_{2.5} contributed by anthropogenic emissions (anthropogenic grids) exceeds that of natural emissions (natural grids).

Formatted: Not Highlight

Deleted: pollution

14 We calculate sectoral contributions of foreign anthropogenic emissions to China's PM_{2.5}
 15 concentrations under China's current-policy and foreign countries' SSP370 scenario at a
 16 resolution of 2° x 2.5° for January in 2030 and 2060. Sectoral contributions are calculated by
 17 taking the difference of a simulation that includes one sector of SSP370 foreign anthropogenic
 18 emissions (agriculture, industry, energy, traffic, residential combustion, solvent use, waste
 19 burning) at a time and a simulation without foreign anthropogenic emissions
 20 ("China_current_2030" and "China_current_2060" runs in Table 2).

Deleted: grids

Deleted: 2

Deleted: PM_{2.5}

Deleted: overall

Deleted: d

22 We also conduct sensitivity simulations at a resolution of 2° x 2.5° to separate transboundary
 23 contributions to China's future PM_{2.5} concentrations from anthropogenic emissions within or
 24 outside the Flex-Grid domain of the model under the current-policy and SSP370 scenario.
 25 Contributions of foreign anthropogenic emissions from within the Flex-Grid domain are
 26 calculated as the difference of a simulation without foreign anthropogenic emissions outside the
 27 Flex-Grid domain and a simulation without foreign anthropogenic emissions over the globe.
 28 Contributions of foreign anthropogenic emissions from outside the Flex-Grid domain are
 29 calculated as the difference between the total foreign contributions and foreign contributions
 30 from Flex-Grid countries.

Formatted: Not Highlight

Formatted: Subscript

Deleted: (11° S–60° N, 30°–150° E, including the entire Asia, eastern Africa and eastern Europe)

31 **2.3 Health impact assessment**

Deleted: ¶

Formatted: English (US)

34 We use the GEMM model (Burnett et al., 2018) to estimate premature deaths attributable to
 35 ambient PM_{2.5} exposure for noncommunicable diseases (NCDs) and lower respiratory infections
 36 (LRIs) in China under each scenario. GEMM NCD+LRI calculates premature deaths associated
 37 with ambient PM_{2.5} exposure (M) for each population subgroup s (by age and gender) in grid cell
 38 g as:

Deleted:

$$40 M_{s,g}(C_g) = B_s \times AF_s(C_g) \times P_g \quad (1)$$

42 where B_s is the national baseline mortality rate of NCD+LRI for the exposed population
 43 subgroup s. AF_s(C_g) is the attributable fraction of NCD+LRI to PM_{2.5} exposure at level C_g for
 44 population subgroup s. P_g represents the total exposed population in grid cell g. In particular AF
 45 was calculated as AF = (RR - 1)/RR, where RR is the relative risk of NCD+LRI attributable to

1 ambient PM_{2.5} exposure. The dependence of RR of NCD+LRI on PM_{2.5} concentrations is
2 calculated as

3

$$4 \quad RR = e^{1+e^{\frac{\theta \times \ln(\frac{z}{\alpha} + 1)}{(-\frac{z-\mu}{v})}}}, \text{ where } z = \max(0, C_g - 2.4) \quad (2)$$

5 where θ , α , μ and v are fitted parameters of PM_{2.5}–mortality relationships. According to the
6 GEMM model, the RR of NCD+LRI is calculated by age for adults aged from 25 to greater than
7 85 years in 5-year intervals.

8 In this study, we use the national baseline mortality data by age group and disease type
9 from the Global Burden of Disease Results Tool 2017 version (GBD 2017; Institute for Health
10 Metrics and Evaluation, 2017). For future baseline mortality, we use the age-specific baseline
11 mortality rates projected by the International Futures (IFs) model v7.89 (Hughes et al., 2011).
12 Population and age structure data for China for 2015 and future years are obtained from the SSP
13 dataset (Samir and Lutz, 2017; Riahi et al., 2017), because China’s current-policy scenario is
14 built upon the SSP2 scenario and the carbon-neutral scenario is built upon the SSP1 scenario
15 (Cheng et al., 2021b; Tong et al., 2020). The gridded population data for China on a $0.5^\circ \times 0.5^\circ$
16 spatial resolution for 2015 and future years under each SSP scenario are developed by Huang et
17 al. (2019). Future cause-and-effect relationship between PM_{2.5} exposure and mortality follows
18 the PM_{2.5}-mortality hazard ratio function in the GEMM model, which has been widely used in
19 the calculation of PM_{2.5}-associated mortality in the future (Hong et al., 2019; Liu et al., 2021;
20 Yang et al., 2022).

Formatted: Subscript

Formatted: Font: (Default) Times New Roman, (Asian)
+Body Asian (DengXian), Not Bold

Formatted: Subscript

21 With the baseline mortality, the population data and the age-structure data, we calculate, grid
22 cell by grid cell, the age-specific baseline mortality rate under present and future scenarios. The
23 impacts of transboundary pollution on mortality in China is calculated as the difference between
24 mortality associated with PM_{2.5} simulated by the full anthropogenic emissions for China and
25 foreign countries, and mortality associated with PM_{2.5} simulated by excluding foreign emissions
26 from the full anthropogenic emissions.

27 3. Results

28

29 3.1 Achievability of future air quality goals in China.

30

31 In 2015, the national mean population-weighted PM_{2.5} over China is about $48 \mu\text{g m}^{-3}$ (Fig. 1)
32 after the observation-based correction, consistent with previous studies ($48\text{--}55 \mu\text{g m}^{-3}$) that used
33 various models (Burnett et al., 2018; Cheng et al., 2021a, 2021b; Tang et al., 2022; Zhang and
34 Cao, 2015). The correction reduces the overestimation in the model by roughly 16% (Fig. S2);
35 details about the correction approach are described in Method. From 2015 to 2030 and 2060,
36 there is a remarkable decreasing trend in China’s annual mean population-weighted PM_{2.5}
37 concentrations under plausible futures (Fig. 1). In 2030, achieving the $35 \mu\text{g m}^{-3}$ goal on a
38 national average level would be feasible even under the fossil fuel-intensive pathways in China
39 (current-policy) and foreign countries (SSP370), yet the most polluted provinces might not be
40 able to achieve the goal under such scenarios (upper whiskers in Fig. 1). In 2060, achieving the

1 WHO AQG goal of $5 \mu\text{g m}^{-3}$ would be highly unlikely under any emission pathway analyzed in
2 this study (Fig. 1).

3
4 Considering air quality goals at the city level in China ($35 \mu\text{g m}^{-3}$ in 2015 and 2030; $5 \mu\text{g m}^{-3}$ in
5 2060), the fraction of cities achieving air quality goals increases considerably as China and
6 foreign countries transition from the fossil fuel-intensive to the low-carbon pathway, yet the
7 achievement is not fully attainable in all cities. In 2015, only 30% of 365 cities in the national
8 $\text{PM}_{2.5}$ monitoring network has an annual mean population-weighted $\text{PM}_{2.5}$ below the $35 \mu\text{g m}^{-3}$
9 threshold (Fig. 2a). In 2030, the percentage increases considerably (Fig. 2b). Under the current-
10 policy emission pathway in China (the current clean air policies and Nationally Determined
11 Contribution pledges), roughly 65% of Chinese cities (average of the three foreign scenarios) is
12 able to achieve the goal, doubling the percentage in 2015. Under the carbon-neutral emission
13 pathway in China (stringent clean air policies and carbon neutrality commitments), the
14 percentage further increases to about 92% (average of the three foreign scenarios). However,
15 even the cleanest emission pathway in both China and foreign countries cannot allow all cities to
16 attain the $35 \mu\text{g m}^{-3}$ goal. In 2060 (Fig. 2c), the WHO AQG goal of $5 \mu\text{g m}^{-3}$ is not achievable
17 for the majority of cities ($\geq 75\%$) in China. Emission pathways adopted by foreign countries can
18 affect up to 6% of cities achieving the AQG goal, and the influence could be even larger over
19 border regions (as will be shown in Fig. 3).

20 21 3.2 Transboundary impacts on $\text{PM}_{2.5}$ concentrations in China.

22
23 In 2015, transboundary pollution contributes about $3.8 \mu\text{g m}^{-3}$ population-weighted $\text{PM}_{2.5}$ to
24 China (Fig. 3a), accounting for roughly 8% of the total population-weighted $\text{PM}_{2.5}$ (Fig. 4). In
25 the future, transboundary pollution becomes increasingly important in China, as the share of
26 transboundary pollution in China's total population-weighted $\text{PM}_{2.5}$ increases to 12%–22% in
27 2060 (Fig. 4).

28
29 For future $\text{PM}_{2.5}$ concentrations, the contribution of transboundary pollution to $\text{PM}_{2.5}$ in China
30 decreases as foreign countries and China undergo the low-carbon pathways (Fig. 3a). In 2030,
31 under the current-policy scenario in China, transboundary contributions to $\text{PM}_{2.5}$ in China would
32 be reduced by $1.2 \mu\text{g m}^{-3}$ (29%) as foreign countries transition from the fossil fuel-intensive
33 (SSP370) to the low-carbon (SSP119) scenario. By 2060, the difference would be increased to
34 $1.8 \mu\text{g m}^{-3}$ (45%). The transboundary pollution will also depend on Chinese domestic emissions,
35 because of their chemical interactions with foreign-transported pollution (Xu et al., 2023). In
36 2030, under the SSP370 scenario in foreign countries, transboundary contributions to $\text{PM}_{2.5}$ in
37 China could be reduced by $0.6 \mu\text{g m}^{-3}$ (14%) as China transitions from the current-policy to the
38 carbon-neutral scenario. In 2060, the $\text{PM}_{2.5}$ reduction could be increased to $1.8 \mu\text{g m}^{-3}$ (45%).

39
40 The direct atmospheric transport and the chemical interactions play different roles in
41 transboundary pollution over different regions in China. Over North China (outlined in Fig. 3c-
42 o), the influence of chemical interactions on transboundary pollution is prominent. Xu et al.
43 (2023) found that foreign-transported NMVOCs could enhance the atmospheric oxidizing
44 capacity and facilitated the oxidation of Chinese nitrogen oxides (NO_x) to form nitrate over
45 North China. This feature persists in the future. Due to chemical interactions, North China is the
46 region that is the most strongly affected by transboundary pollution in most scenarios. In 2030,

Deleted: (

Deleted: ,

Deleted:

Deleted: ,

Moved (insertion) [1]

1 transboundary pollution contributes 3 to 10 $\mu\text{g m}^{-3}$ $\text{PM}_{2.5}$ concentrations to North China, even
2 under China's carbon-neutral scenario (Fig. 3b and Fig. 3i-l), greatly increasing the difficulty for
3 this region to achieve the 35 $\mu\text{g m}^{-3}$ goal. Further reduction of transboundary pollution on North
4 China requires the reduction of not only foreign anthropogenic emissions but also China's
5 domestic emission. For example, in 2060, if China adopts the current-policy pathway (Fig. 3f-h),
6 achieving the WHO AQG goal of 5 $\mu\text{g m}^{-3}$ would be highly unlikely for the majority of North
7 China, as transboundary pollution alone would contribute roughly 3~10 $\mu\text{g m}^{-3}$ of $\text{PM}_{2.5}$
8 concentration (Fig. 3b and Fig. 3f-h). Alternatively, adopting the carbon-neutral rather than the
9 current-policy pathway in China reduces roughly 41%~58% of transboundary $\text{PM}_{2.5}$ over North
10 China in 2060 (Fig. 3b), making it possible for the region to achieve the 5 $\mu\text{g m}^{-3}$ goal. Thus, a
11 low emission pathway for China has a large co-benefit on reducing transboundary pollution
12 exerted upon its populous northern area by reducing the aforementioned chemical interactions.

13
14 The western border provinces of Yunnan and Xinjiang (denoted in Fig. 3c-o) are also
15 influenced substantially by transboundary pollution, with the magnitude of transboundary
16 pollution determined predominantly by foreign (but not Chinese) emission pathways. For
17 example, in 2060, under China's carbon-neutral emission pathway, transboundary $\text{PM}_{2.5}$ over
18 Yunnan decreases from 6 $\mu\text{g m}^{-3}$ to 3 $\mu\text{g m}^{-3}$ (a 50% reduction) as the foreign pathway switches
19 from fossil fuel-intensive (SSP370; Fig. 3m) to low-carbon (SSP119; Fig. 3o), reflecting the
20 considerable impact of Southern Asian pollution to China (Jiang et al., 2013). Over Xinjiang, the
21 transboundary pollution driven predominantly by direct atmospheric transport can reach 6 $\mu\text{g m}^{-3}$
22 in many scenarios (i.e., current-policy plus SSP370 for 2030 and 2060; and carbon-neutral plus
23 SSP245 for 2030), indicating the important influence of anthropogenic emissions from Central
24 Asia in the future that has hardly been investigated in previous studies.

25
26 As transboundary contributions are the largest under China's current-policy and foreign
27 countries' SSP370 emission pathways (Fig. 3a), we further investigate major emission sectors in
28 foreign countries that contribute to China's future $\text{PM}_{2.5}$ concentrations. We take January as an
29 example because Xu et al. (2023) found that transboundary contributions to China's $\text{PM}_{2.5}$ were
30 the largest in January. Fig. S4 reveals that agriculture, energy, industry, transportation and
31 residential combustion emissions in foreign countries are major sources of transboundary
32 pollution to China's national average future $\text{PM}_{2.5}$ concentrations in 2030 and 2060 January, and
33 that these sectors contribute roughly evenly (15%~23% for each source). There is pronounced
34 spatial heterogeneity in the source attribution. Solvent use emissions in foreign countries make
35 roughly 3 times larger contributions to North China (18% in 2030 and 2060) than to the national
36 average (6%~7% in 2030 and 2060). This is in line with the result of Xu et al. (2023) that the
37 NMVOCs are the primary drivers of transboundary $\text{PM}_{2.5}$ over North China. Residential
38 combustion in foreign countries contributes about 16% in 2030 and 10% 2060 more $\text{PM}_{2.5}$ to
39 Yunnan province than to the entire China, as Yunnan is mostly affected by emissions from South
40 Asia (Jiang et al., 2013) where residential combustion is intensive (McDuffie et al., 2020).

41
42 Transboundary pollution to China could arise from emissions of countries within and outside
43 the Flex-Grid domain of the model. Thus, we conduct a sensitivity simulation driven by China's
44 current-policy and foreign countries' SSP370 emissions in 2030 and 2060 as an example to
45 explore the relative importance of anthropogenic emissions from countries within and outside the
46 Flex-Grid domain for transboundary pollution to China. Fig. S5 shows that anthropogenic

Moved up [1]: In 2030, transboundary pollution contributes 3 to 10 $\mu\text{g m}^{-3}$ $\text{PM}_{2.5}$ concentrations even under China's carbon-neutral scenario (Fig. 3b and Fig. 3i-l), greatly increasing the difficulty for this region to achieve the 35 $\mu\text{g m}^{-3}$ goal.

Deleted: making China's domestic emission pathway an important driver to transboundary pollution and to the achievement of its air quality goals.

Deleted:

Deleted: I

Formatted: English (US)

Deleted: .

Deleted: emissions

Formatted: Subscript

Deleted: (

Deleted: ,

Formatted: Subscript

Deleted: from

Deleted: from

Deleted: (

Deleted: ,

Formatted: Subscript

Deleted: from

Formatted: Subscript

Deleted: high

Deleted: ¶

Deleted: from

Deleted: (11° S–60° N, 30°–150° E, including the entire Asia, eastern Africa and eastern Europe)

Deleted: W

Deleted: on

1 emissions from countries within the Flex-Grid domain dominate the transboundary pollution to
2 China in all seasons (>80%), and their contributions remain in 2030 and 2060. Contributions of
3 anthropogenic emissions from countries outside the Flex-Grid domain is larger in April and
4 October when westerly winds prevail and wet deposition is not as strong as in July (Jiang et al.,
5 2013; Leibensperger et al., 2011; Ni et al., 2018). North China is more influenced by emissions
6 outside the Flex-Grid domain (13% in April and 24% in October) than China on average is (7%
7 in April and 13% in October), indicating that North China is more influenced by emissions
8 outside Asia, such as Europe and North America. Leibensperger et al. (2011) have also found
9 that NO_x emission in the US could lead to an enhancement of PM_{2.5} over North China.

11 3.3 Health threats by transboundary pollution in China.

12 Figure 5 shows our estimated PM_{2.5}-associated premature deaths in 2015 and the future. Our
13 estimated PM_{2.5}-associated premature deaths in China in 2015 (2.03 million) is comparable with
14 other studies (2 to 2.4 million; Burnett et al., 2018; Geng et al., 2021; Tang et al., 2022). Our
15 estimated increasing trend of premature deaths in China from SSP370 to SSP119 is also
16 consistent in previous works (Tang et al., 2022; Yang et al., 2022), which is driven primarily by
17 population ageing (Fig. S6). Our estimated premature deaths in 2030 and 2060 under the current-
18 policy emission scenario (2.68 to 2.82 million for 2030; 1.8 to 2.05 million for 2060) is
19 substantially lower than those in Tang et al. (2022) (3.5 to 4 million for 2030; 6.5 to 7.5 million
20 for 2060). The difference is primarily because Tang et al. (2022) fixed the baseline mortality
21 rates in future years at the 2015 level, while our future baseline mortality rates from the IFs are
22 projected on the basis of income, education and technology advancement, and other factors
23 (Hughes et al., 2011).

24 The PM_{2.5} concentrations contributed by transboundary pollution can lead to an extra number
25 of people (i.e., excess population) exposed to PM_{2.5} concentrations above the targeted air quality
26 levels in China (35 µg m⁻³ in 2015 and 2030; and 5 µg m⁻³ in 2060), which may lead to potential
27 health threats. In the future, the number of excess population due to transboundary pollution
28 depends on both foreign and China's emission pathways (Fig. 6a). In 2030, with Chinese
29 emissions following the carbon-neutral scenario, adopting the low-carbon (SSP119) rather than
30 the fossil fuel-intensive pathway (SSP370) in foreign countries could avoid 10 million Chinese
31 people from being exposed to PM_{2.5} concentrations above 35 µg m⁻³. In 2060, 5 million people
32 could be avoided from being exposed to PM_{2.5} concentrations above 5 µg m⁻³ if the foreign
33 scenario switches from SSP370 to SSP119. These results indicate remarkable health benefits that
34 the low-carbon emission pathway in foreign countries would bring to China.

35
36 For a given future year and foreign emission scenario, the excess population due to
37 transboundary pollution tends to be larger when China adopts the carbon-neutral pathway than
38 when China adopts the current-policy pathway (Fig. 6a). This reflects the increasing influence of
39 transboundary pollution on air quality in China as China's overall PM_{2.5} concentrations drop
40 sharply towards the air quality goals under the carbon-neutral pathway (Fig. 1).

41
42 Another measure of health threat that transboundary pollution could exert upon China is PM_{2.5}-
43 related premature deaths. As shown in Fig. 6b, there is an obvious decreasing trend of
44 transboundary-contributed PM_{2.5}-related premature deaths in China as foreign countries and

Deleted: (

Deleted: ,

Formatted: Subscript

Formatted: Subscript

Formatted: English (US)

Formatted: Indent: First line: 0.25 cm, Don't adjust right indent when grid is defined, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Deleted: S3

1 China transition from the fossil fuel-intensive to the low-carbon pathway. In 2030, adopting the
2 low-carbon pathway (SSP119) in foreign countries would avoid 41% (178,000 under China's
3 carbon-neutral pathway) to 45% (207,000 under China's current-policy pathway) of premature
4 deaths that would occur under the fossil fuel-intensive pathway (SSP370) in foreign countries. In
5 2060, the avoidance would be as large as 76% (270,000 under China's current-policy pathway)
6 to 91% (63,000 under China's carbon-neutral pathway). In addition, China's low carbon
7 emission pathway could also bring considerable health benefits through reducing the chemical
8 interaction-related transboundary pollution and associated premature deaths in China. In 2030,
9 adopting the carbon-neutral pathway in China would avoid 99,000 (SSP245) to 211,000
10 (SSP119) people from transboundary pollution-associated premature deaths relative to adopting
11 the current-policy emission pathway. In 2060, the avoided deaths would be 76,000 (SSP119) to
12 283,000 (SSP370). These findings highlight the considerable health benefits to China if foreign
13 countries and China could adopt the low-carbon emission pathways coincidentally.

14 4. Discussion

15 This study reveals the increasingly important role that transboundary pollution would play in
16 the achievement of future air quality goals and the protection of public health in China. The
17 magnitude of transboundary pollution depends on both Chinese and foreign emissions, given the
18 direct pollution transport and the indirect impact through chemical interactions between
19 transported and China's locally emitted pollutants. Adopting the low-carbon (SSP119) instead of
20 the fossil fuel-intensive (SSP370) pathway in foreign countries would avoid millions of Chinese
21 people (Fig. 6a) from being exposed to PM_{2.5} concentrations above the targeted air quality levels
22 in 2030 (35 μg m⁻³) and 2060 (5 μg m⁻³), and would avoid 63,000~270,000 of transboundary
23 PM_{2.5}-associated mortality in China in 2060 (Fig. 6b). Adopting the carbon-neutral instead of
24 current-policy pathway in China would avoid 76,000~283,000 premature mortality associated
25 with transboundary pollution in 2060 (Fig. 6b). If China and foreign countries undergo the low-
26 carbon pathways coincidentally, transboundary pollution in China would be reduced by 63%
27 relative to adopting a fossil fuel-intensive emission pathway in both regions (Fig. 4a), and could
28 avoid 386,000 premature deaths in China (Fig. 6b). Cutting foreign emissions are particularly
29 effective at reducing transboundary pollution upon the western border provinces of Yunnan and
30 Xinjiang that are dominated by direct transport. Fully achieving the WHO AQG goal of 5 μg m⁻³
31 over the populous North China would be possible only when both China and foreign countries
32 adopt the low-carbon pathways (carbon-neutral and SSP119, respectively).

33 The importance of transboundary pollution is not confined in China. In the future, significant
34 emission changes are expected in many developing countries, affecting air quality locally and in
35 the downwind regions. These developing countries are often financially and/or technologically
36 less capable to control emissions by themselves. Thus, enhanced external aids would be essential
37 for these developing countries to undergo a low-carbon development in the future, which in turn
38 would benefit air quality and public health of the entire globe. External aids could be done
39 through global cooperation programs such as the Paris Agreement or through inter-regional
40 collaboration such as the Belt and Road Initiative.

41 Uncertainties arise from several factors in this study. The future development pathway of a
42 country is highly uncertain, leading to a wide spread of projected emission trajectories in the

Deleted: China, as a pioneer in climate mitigation and economic development in developing countries, could transfer emission control technologies and/or provide financial aid

Deleted: to neighboring countries

Deleted: Such aids could be deployed through

Deleted: or inter-regional

Deleted: and

Deleted: Promoting the environmental cooperation is particularly meaningful nowadays when the 2019 coronavirus (COVID-19) pandemic, the Russia-Ukraine conflict and the emergence of regional rivalry (e.g., the Sino-US trade war; Du et al., 2020) disrupt the global society and environment, threatening cooperation at all levels.

Deleted: These efforts could accelerate the low-carbon development in neighboring countries, which could in turn improve air quality and public health in China by reducing transboundary effects.

Deleted: to a certain degree

1 future. We thereby use a set of emission projections to represent the plausible range of future
2 emissions. The simulation of PM_{2.5} is subject to uncertainties in aerosol chemical and physical
3 processes, such as the wet deposition of nitrate (Luo et al., 2020) and the simplified secondary
4 organic aerosol formation scheme (Pai et al., 2020). Our correction to the simulated PM_{2.5}
5 concentrations using ground-based observations could reduce the uncertainty by 15% to 18%
6 (Fig. S2). Future population and age structure change are projected based on their historical
7 relationships with GDP and urbanization (O'Neill et al., 2020; Riahi et al., 2017). Thus, they
8 may introduce biases if the future development of global GDP and urbanization deviates from
9 the historical path (e.g., due to the emergence of anti-globalization (Dür et al., 2020) and regional
10 rivalry (O'Neill et al., 2014; van Vuuren et al., 2014). There are additional uncertainties from
11 PM_{2.5}-related death estimates due to the limited epidemiology evidence and statistical estimation
12 of the GEMM model, such as the influences of particulate species and size on health outcomes
13 (Burnett et al., 2018). We estimate the overall uncertainties of PM_{2.5}-related death in each
14 scenario as 95% CI in the main text. Besides, we do not consider potential influences of climate
15 change and the change of natural emissions on PM_{2.5} and transboundary pollution, yet their
16 influences are found small compared to the influence of anthropogenic emissions (Hong et al.,
17 2019; Jiang et al., 2013; Liu et al., 2021; Silva et al., 2017).

19 **Data availability**

20
21 The global SSP-RCP emissions data for 2015 and future scenarios, and the area-weighted
22 PM_{2.5} concentrations in China for 2015 and future scenarios are available upon request to the
23 corresponding author. All other data used in this study are publicly available and can be
24 downloaded from the following links. (1) China's future emission scenarios 2015–
25 2060: <http://www.meicmodel.org/dataset-dpec.html>. (2) Chinese future population data:
26 [https://www.scidb.cn/en/detail?dataSetId=73c1ddb79e54638bd0ca2a6bd48e3ff&dataSetType=](https://www.scidb.cn/en/detail?dataSetId=73c1ddb79e54638bd0ca2a6bd48e3ff&dataSetType=organization)
27 [organization](https://www.scidb.cn/en/detail?dataSetId=73c1ddb79e54638bd0ca2a6bd48e3ff&dataSetType=organization). (3) 2015 baseline mortality rate: <https://gbd2017.healthdata.org/gbd-search/>. (4)
28 Future baseline mortality rate projection: https://www.ifs.du.edu/IFs/frm_MortCohorts/.

30 **Code availability**

31
32 The GEOS-Chem model v13.2.1 source code used for PM_{2.5} concentration simulations is
33 available at: <http://www.geos-chem.org>. The SSP-RCP emission harmonization source code is
34 available at <http://software.ene.iiasa.ac.at/aneris/>. All computer codes generated during this study
35 are available from the corresponding authors upon reasonable request.

37 **Acknowledgements**

38
39 This work was supported by the National Natural Science Foundation of China (42075175), the
40 China Postdoctoral Science Foundation (2021M700191) and the Peking University Boya
41 Postdoctoral Fellowship.

43 **Author contributions**

44
45 J.L. led the study. J.X. and J.L. designed the study. J.X. performed the model simulations and
46 conducted the data analysis. D.T. provided China's future emissions data. L.C. provided SSP-

1 RCP emission harmonization and health impact assessment methods. J.X. wrote the manuscript
2 with inputs from J.L. All authors commented on the manuscript.
3

4
5 **Competing interests**
6

7 The authors declare that they have no conflict of interest.
8

9 **References**
10

11 WHO ((WHO), W. H. O.: WHO Global Air Quality Guidelines: Particulate Matter (PM2.5 and
12 PM10), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide, Geneva,, 2021.

13 Anon: Global Burden of Disease Study 2017, Seattle, WA, USA. [online] Available from:
14 http://www.healthdata.org/sites/default/files/files/policy_report/2019/GBD_2017_Booklet.pdf,
15 2017.

16 Ascensão, F., Fahrig, L., Clevenger, A. P., Corlett, R. T., Jaeger, J. A. G., Laurance, W. F. and
17 Pereira, H. M.: Environmental challenges for the Belt and Road Initiative, *Nat. Sustain.*, 1(5),
18 206–209, doi:10.1038/s41893-018-0059-3, 2018.

19 Burnett, R., Chen, H., Szyszkowicz, M., Fann, N., Hubbell, B., Pope, C. A., Apte, J. S., Brauer,
20 M., Cohen, A., Weichenthal, S., Coggins, J., Di, Q., Brunekreef, B., Frostad, J., Lim, S. S., Kan,
21 H., Walker, K. D., Thurston, G. D., Hayes, R. B., Lim, C. C., Turner, M. C., Jerrett, M.,
22 Krewski, D., Gapstur, S. M., Diver, W. R., Ostro, B., Goldberg, D., Crouse, D. L., Martin, R. V.,
23 Peters, P., Pinault, L., Tjepkema, M., Van Donkelaar, A., Villeneuve, P. J., Miller, A. B., Yin, P.,
24 Zhou, M., Wang, L., Janssen, N. A. H., Marra, M., Atkinson, R. W., Tsang, H., Thach, T. Q.,
25 Cannon, J. B., Allen, R. T., Hart, J. E., Laden, F., Cesaroni, G., Forastiere, F., Weinmayr, G.,
26 Jaensch, A., Nagel, G., Concin, H. and Spadaro, J. V.: Global estimates of mortality associated
27 with longterm exposure to outdoor fine particulate matter, *Proc. Natl. Acad. Sci. U. S. A.*,
28 115(38), 9592–9597, doi:10.1073/pnas.1803222115, 2018.

29 Cheng, J., Tong, D., Liu, Y., Yu, S., Yan, L., Zheng, B., Geng, G., He, K. and Zhang, Q.:
30 Comparison of Current and Future PM2.5 Air Quality in China Under CMIP6 and DPEC
31 Emission Scenarios, *Geophys. Res. Lett.*, 48(11), 1–11, doi:10.1029/2021GL093197, 2021a.

32 Cheng, J., Tong, D., Zhang, Q., Liu, Y., Lei, Y., Yan, G., Yan, L., Yu, S., Cui, R. Y., Clarke, L.,
33 Geng, G., Zheng, B., Zhang, X., Davis, S. J. and He, K.: Pathways of China's PM2.5 air quality
34 2015–2060 in the context of carbon neutrality, *Natl. Sci. Rev.*, (December 2020),
35 doi:10.1093/nsr/nwab078, 2021b.

36 Dallas Burtraw, Jhih-Shyang Shih, Maya Domeshek, Seth Villanueva, and K. F. L.: The
37 Distribution of Air Quality Health Benefits from Meeting US 2030 Climate Goals, Washington
38 D.C., 2022.

39 Dür, A., Eckhardt, J. and Poletti, A.: Global value chains, the anti-globalization backlash, and
40 EU trade policy: a research agenda, *J. Eur. Public Policy*, 27(6), 944–956,
41 doi:10.1080/13501763.2019.1619802, 2020.

42 Geng, G., Zheng, Y., Zhang, Q., Xue, T., Zhao, H., Tong, D., Zheng, B., Li, M., Liu, F., Hong,
43 C., He, K. and Davis, S. J.: Drivers of PM2.5 air pollution deaths in China 2002–2017, *Nat.*

- 1 Geosci., 14(9), 645–650, doi:10.1038/s41561-021-00792-3, 2021.
- 2 Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., Van Vuuren, D. P.,
3 Van Den Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O.,
4 Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp, A., Stehfest, E.
5 and Takahashi, K.: Global emissions pathways under different socioeconomic scenarios for use
6 in CMIP6: A dataset of harmonized emissions trajectories through the end of the century,
7 Geosci. Model Dev., 12(4), 1443–1475, doi:10.5194/gmd-12-1443-2019, 2019.
- 8 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert,
9 J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.,
10 Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R. and Zhang, Q.: Historical (1750–2014)
11 anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data
12 System (CEDS), Geosci. Model Dev., 11(1), 369–408, doi:10.5194/gmd-11-369-2018, 2018.
- 13 Hong, C., Zhang, Q., Zhang, Y., Davis, S. J., Tong, D., Zheng, Y., Liu, Z., Guan, D., He, K. and
14 Schellnhuber, H. J.: Impacts of climate change on future air quality and human health in China,
15 Proc. Natl. Acad. Sci. U. S. A., 116(35), 17193–17200, doi:10.1073/pnas.1812881116, 2019.
- 16 Huang, J., Qin, D., Jiang, T., Wang, Y., Feng, Z., Zhai, J., Cao, L., Chao, Q., Xu, X., Wang, G.
17 and Su, B.: Effect of Fertility Policy Changes on the Population Structure and Economy of
18 China: From the Perspective of the Shared Socioeconomic Pathways, Earth's Futur., 7(3), 250–
19 265, doi:https://doi.org/10.1029/2018EF000964, 2019.
- 20 Hughes, B. B., Kuhn, R., Peterson, C. M., Rothman, D. S., Solórzano, J. R., Mathers, C. D. and
21 Dickson, J. R.: Projections of global health outcomes from 2005 to 2060 using the International
22 Futures integrated forecasting model, Bull. World Health Organ., 89(7), 478–486,
23 doi:10.2471/BLT.10.083766, 2011.
- 24 IEA/OECD: IEA: World Energy Outlook, Int. Energy Agency [online] Available from:
25 <https://www.iea.org/reports/world-energy-outlook-2021> (Accessed 21 November 2022), 2021.
- 26 Jiang, H., Liao, H., Pye, H. O. T., Wu, S., Mickley, L. J., Seinfeld, J. H. and Zhang, X. Y.:
27 Projected effect of 2000-2050 changes in climate and emissions on aerosol levels in China and
28 associated transboundary transport, Atmos. Chem. Phys., 13(16), 7937–7960, doi:10.5194/acp-
29 13-7937-2013, 2013.
- 30 KC, S. and Lutz, W.: The human core of the shared socioeconomic pathways: Population
31 scenarios by age, sex and level of education for all countries to 2100, Glob. Environ. Chang., 42,
32 181–192, doi:https://doi.org/10.1016/j.gloenvcha.2014.06.004, 2017.
- 33 Koplitz, S. N., Jacob, D. J., Sulprizio, M. P., Myllyvirta, L. and Reid, C.: Burden of Disease
34 from Rising Coal-Fired Power Plant Emissions in Southeast Asia, Environ. Sci. Technol., 51(3),
35 1467–1476, doi:10.1021/acs.est.6b03731, 2017.
- 36 Leibensperger, E. M., Mickley, L. J., Jacob, D. J. and Barrett, S. R. H.: Intercontinental influence
37 of NO_x and CO emissions on particulate matter air quality, Atmos. Environ., 45(19), 3318–3324,
38 doi:10.1016/j.atmosenv.2011.02.023, 2011.
- 39 Liu, S., Xing, J., Wang, S., Ding, D., Cui, Y. and Hao, J.: Health Benefits of Emission Reduction
40 under 1.5 °C Pathways Far Outweigh Climate-Related Variations in China, Environ. Sci.
41 Technol., doi:10.1021/acs.est.1c01583, 2021.
- 42 Luo, G., Yu, F. and Moch, J. M.: Further improvement of wet process treatments in GEOS-

1 Chem v12.6.0: Impact on global distributions of aerosols and aerosol precursors, *Geosci. Model*
2 *Dev.*, 13(6), 2879–2903, doi:10.5194/gmd-13-2879-2020, 2020.

3 McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A.,
4 Zheng, B., Crippa, M., Brauer, M. and Martin, R. V.: A global anthropogenic emission inventory
5 of atmospheric pollutants from sector- And fuel-specific sources (1970-2017): An application of
6 the Community Emissions Data System (CEDS), *Earth Syst. Sci. Data*, 12(4), 3413–3442,
7 doi:10.5194/essd-12-3413-2020, 2020.

8 Murray, C. J. L., Aravkin, A. Y., Zheng, P., Abbafati, C., Abbas, K. M., Abbasi-Kangevari, M.,
9 Abd-Allah, F., Abdelalim, A., Abdollahi, M., Abdollahpour, I., Abegaz, K. H., Abolhassani, H.,
10 Aboyans, V., Abreu, L. G., Abrigo, M. R. M., Abualhasan, A., Abu-Raddad, L. J., Abushouk, A.
11 I., Adabi, M., Adekanmbi, V., Adeoye, A. M., Adetokunboh, O. O., Adham, D., Advani, S. M.,
12 Agarwal, G., Aghamir, S. M. K., Agrawal, A., Ahmad, T., Ahmadi, K., Ahmadi, M., Ahmadi, H.,
13 Ahmed, M. B., Akalu, T. Y., Akinyemi, R. O., Akinyemiju, T., Akombi, B., Akunna, C. J.,
14 Alahdab, F., Al-Aly, Z., Alam, K., Alam, S., Alam, T., Alanezi, F. M., Alanzi, T. M., Alemu, B.
15 wassihun, Alhabib, K. F., Ali, M., Ali, S., Alicandro, G., Alinia, C., Alipour, V., Alizade, H.,
16 Aljunid, S. M., Alla, F., Allebeck, P., Almasi-Hashiani, A., Al-Mekhlafi, H. M., Alonso, J.,
17 Altirkawi, K. A., Amini-Rarani, M., Amiri, F., Amugsi, D. A., Ancuceanu, R., Anderlini, D.,
18 Anderson, J. A., Andrei, C. L., Andrei, T., Angus, C., Anjomshoa, M., Ansari, F., Ansari-
19 Moghaddam, A., Antonazzo, I. C., Antonio, C. A. T., Antony, C. M., Antriyandarti, E., Anvari,
20 D., Anwer, R., Appiah, S. C. Y., Arabloo, J., Arab-Zozani, M., Ariani, F., Armoon, B., Årnlöv,
21 J., Arzani, A., Asadi-Aliabadi, M., Asadi-Pooya, A. A., Ashbaugh, C., Assmus, M., Atafar, Z.,
22 Atnafu, D. D., Atout, M. M. W., Ausloos, F., Ausloos, M., Ayala Quintanilla, B. P., Ayano, G.,
23 Ayanore, M. A., Azari, S., Azarian, G., Azene, Z. N., et al.: Global burden of 87 risk factors in
24 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease
25 Study 2019, *Lancet*, 396(10258), 1223–1249, doi:10.1016/S0140-6736(20)30752-2, 2020.

26 Ni, R., Lin, J., Yan, Y. and Lin, W.: Foreign and domestic contributions to springtime ozone
27 over China, *Atmos. Chem. Phys.*, 18(15), 11447–11469, doi:10.5194/acp-18-11447-2018, 2018.

28 O'Neill, B. C., Kriegler, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R. and
29 van Vuuren, D. P.: A new scenario framework for climate change research: The concept of
30 shared socioeconomic pathways, *Clim. Change*, 122(3), 387–400, doi:10.1007/s10584-013-
31 0905-2, 2014.

32 O'Neill, B. C., Carter, T. R., Ebi, K., Harrison, P. A., Kemp-Benedict, E., Kok, K., Kriegler, E.,
33 Preston, B. L., Riahi, K., Sillmann, J., van Ruijven, B. J., van Vuuren, D., Carlisle, D., Conde,
34 C., Fuglestedt, J., Green, C., Hasegawa, T., Leininger, J., Monteith, S. and Pichs-Madruga, R.:
35 Achievements and needs for the climate change scenario framework, *Nat. Clim. Chang.*, 10(12),
36 1074–1084, doi:10.1038/s41558-020-00952-0, 2020.

37 Of, T. S. C. of the P. R.: *The thirteenth Five-Year Plan.*, 2016.

38 Pai, S. J., Heald, C. L., Pierce, J. R., Farina, S. C., Marais, E. A., Jimenez, J. L., Campuzano-
39 Jost, P., Nault, B. A., Middlebrook, A. M., Coe, H., Shilling, J. E., Bahreini, R., Dingle, J. H. and
40 Vu, K.: An evaluation of global organic aerosol schemes using airborne observations, *Atmos.*
41 *Chem. Phys.*, 20(5), 2637–2665, doi:10.5194/acp-20-2637-2020, 2020.

42 Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K.,
43 Amann, M., Bodirsky, B. L., van Vuuren, D. P., Aleluia Reis, L., Calvin, K., Drouet, L., Fricko,
44 O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J.,

- 1 Luderer, G., Masui, T., Stehfest, E., Strefler, J., van der Sluis, S. and Tavoni, M.: Future air
2 pollution in the Shared Socio-economic Pathways, *Glob. Environ. Chang.*, 42, 346–358,
3 doi:<https://doi.org/10.1016/j.gloenvcha.2016.05.012>, 2017.
- 4 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N.,
5 Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M.,
6 Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F.,
7 Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J.,
8 Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L.,
9 Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M.,
10 Tabeau, A. and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and
11 greenhouse gas emissions implications: An overview, *Glob. Environ. Chang.*, 42, 153–168,
12 doi:[10.1016/j.gloenvcha.2016.05.009](https://doi.org/10.1016/j.gloenvcha.2016.05.009), 2017.
- 13 Shindell, D., Ru, M., Zhang, Y., Seltzer, K., Faluvegi, G., Nazarenko, L., Schmidt, G. A.,
14 Parsons, L., Challapalli, A., Yang, L. and Glick, A.: Temporal and spatial distribution of health,
15 labor, and crop benefits of climate change mitigation in the United States, *Proc. Natl. Acad. Sci.*,
16 118(46), e2104061118, doi:[10.1073/pnas.2104061118](https://doi.org/10.1073/pnas.2104061118), 2021.
- 17 Silva, R. A., West, J. J., Lamarque, J. F., Shindell, D. T., Collins, W. J., Faluvegi, G., Folberth,
18 G. A., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S. T., Sudo, K., Takemura, T.,
19 Bergmann, D., Cameron-Smith, P., Doherty, R. M., Josse, B., MacKenzie, I. A., Stevenson, D. S.
20 and Zeng, G.: Future global mortality from changes in air pollution attributable to climate
21 change, *Nat. Clim. Chang.*, 7(9), 647–651, doi:[10.1038/nclimate3354](https://doi.org/10.1038/nclimate3354), 2017.
- 22 Tang, R., Zhao, J., Liu, Y., Huang, X., Nielsen, C. P., Wang, H., Zhou, D. and Ding, A.: Air
23 quality and health co-benefits of China’s carbon dioxide emissions peaking before 2030, , 1–9,
24 doi:[10.1038/s41467-022-28672-3](https://doi.org/10.1038/s41467-022-28672-3), 2022.
- 25 Tong, D., Cheng, J., Liu, Y., Yu, S., Yan, L., Hong, C., Qin, Y., Zhao, H., Zheng, Y., Geng, G.,
26 Li, M., Liu, F., Zhang, Y., Zheng, B., Clarke, L. and Zhang, Q.: Dynamic projection of
27 anthropogenic emissions in China: Methodology and 2015–2050 emission pathways under a
28 range of socio-economic, climate policy, and pollution control scenarios, *Atmos. Chem. Phys.*,
29 20(9), 5729–5757, doi:[10.5194/acp-20-5729-2020](https://doi.org/10.5194/acp-20-5729-2020), 2020.
- 30 Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P.,
31 Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O’Connor, F. M.,
32 Olivié, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis, K.,
33 Wu, T. and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models,
34 *Atmos. Chem. Phys.*, 20(23), 14547–14579, doi:[10.5194/acp-20-14547-2020](https://doi.org/10.5194/acp-20-14547-2020), 2020.
- 35 van Vuuren, D. P., Kriegler, E., O’Neill, B. C., Ebi, K. L., Riahi, K., Carter, T. R., Edmonds, J.,
36 Hallegatte, S., Kram, T., Mathur, R. and Winkler, H.: A new scenario framework for Climate
37 Change Research: Scenario matrix architecture, *Clim. Change*, 122(3), 373–386,
38 doi:[10.1007/s10584-013-0906-1](https://doi.org/10.1007/s10584-013-0906-1), 2014.
- 39 van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., Doelman, J. C., van den Berg, M., Harmsen,
40 M., de Boer, H. S., Bouwman, L. F., Daioglou, V., Edelenbosch, O. Y., Girod, B., Kram, T.,
41 Lassaletta, L., Lucas, P. L., van Meijl, H., Müller, C., van Ruijven, B. J., van der Sluis, S. and
42 Tabeau, A.: Energy, land-use and greenhouse gas emissions trajectories under a green growth
43 paradigm, *Glob. Environ. Chang.*, 42, 237–250, doi:[10.1016/j.gloenvcha.2016.05.008](https://doi.org/10.1016/j.gloenvcha.2016.05.008), 2017.

1 Xing, J., Lu, X., Wang, S., Wang, T., Ding, D., Yu, S., Shindell, D., Ou, Y., Morawska, L., Li,
2 S., Ren, L., Zhang, Y., Loughlin, D., Zheng, H., Zhao, B., Liu, S., Smith, K. R. and Hao, J.: The
3 quest for improved air quality may push China to continue its CO₂reduction beyond the Paris
4 Commitment, *Proc. Natl. Acad. Sci. U. S. A.*, 117(47), 29535–29542,
5 doi:10.1073/pnas.2013297117, 2020.

6 Xu, J.-W., Lin, J., Luo, G., Adeniran, J. and Kong, H.: Foreign emissions exacerbate PM_{2.5}
7 pollution in China through nitrate chemistry, *Atmos. Chem. Phys.*, 23, 4149–4163,
8 doi:10.5194/acp-2022-646, 2023.

9 Yang, H., Huang, X., Westervelt, D. M., Horowitz, L. and Peng, W.: Socio-demographic factors
10 shaping the future global health burden from air pollution, *Nat. Sustain.*, doi:10.1038/s41893-
11 022-00976-8, 2022.

12 Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu,
13 W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L.,
14 Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B.,
15 Liu, Z., Yang, F., He, K. and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013
16 to 2017, *Proc. Natl. Acad. Sci.*, 116(49), 24463–24469, doi:10.1073/pnas.1907956116, 2019.


17 Zhang, Y.-L. and Cao, F.: Fine particulate matter (PM_{2.5}) in China at a city level, *Sci. Rep.*,
18 5(1), 14884, doi:10.1038/srep14884, 2015.

19 Zheng, B., Chevallier, F., Ciais, P., Yin, Y., Deeter, M. N., Worden, H. M., Wang, Y., Zhang, Q.
20 and He, K.: Rapid decline in carbon monoxide emissions and export from East Asia between
21 years 2005 and 2016, *Environ. Res. Lett.*, 13(4), 044007, doi:10.1088/1748-9326/aab2b3, 2018a.


22 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L.,
23 Zhang, Y., Zhao, H., Zheng, Y., He, K. and Zhang, Q.: Trends in China’s anthropogenic
24 emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18(19),
25 14095–14111, doi:10.5194/acp-18-14095-2018, 2018b.

26


27



1
2 **Figure 1.** Population-weighted PM_{2.5} concentrations over China. Box-and-whisker plots
3 represent 5th, 25th, 75th, and 95th percentiles of provincial population-weighted PM_{2.5}
4 concentrations in 2015, 2030 and 2060. Lines in the middle of each box represent the national
5 mean population-weighted PM_{2.5} concentrations. Future emission scenarios in China are labeled
6 as text at the top and in foreign countries are represented by colors according to the legend.
7



8
9
10 **Figure 2.** Percent of cities achieving air quality goals in China. **(a)** Percent of cities in China
11 with an annual mean population-weighted PM_{2.5} concentration below 35 µg m⁻³ in 2015. **(b-c)**
12 Percent of cities with an annual mean population-weighted PM_{2.5} concentration achieving the 35
13 µg m⁻³ goal in 2030 (b) and the 5 µg m⁻³ goal in 2060 (c). PM_{2.5} concentrations are simulated
14 under different future emission scenarios in China (current-policy and carbon-neutral) and
15 foreign countries (represented by colors following the legend).
16
17



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

Figure 3. Contributions of transboundary pollution to PM_{2.5} concentrations over China. **(a)** Transboundary contributions to national annual mean population-weighted PM_{2.5} in China in 2015, 2030 and 2060. Future scenarios are estimated by different emission scenarios in China represented by light (carbon-neutral scenario) and dark shadings (current-policy scenario), along with different emission scenarios in other countries (SSP370, SSP245, SSP119) represented by colors according to the legend. Text on top of each bar represents the transboundary-contributed population-weighted PM_{2.5} under China's current-policy emission scenarios. Text in the light shading of each bar represents the transboundary-contributed population-weighted PM_{2.5} under China's carbon-neutral emission scenarios. **(b)** Same as a, but for North China. **(c-o)** Spatial distributions of transboundary-contributed annual mean PM_{2.5} concentrations over China in 2030 and 2060 under different emission scenarios in China and in other countries. YN represents Yunnan province. XJ represents Xinjiang province. North China is outlined by thick black lines.

Moved (insertion) [2]

Moved up [2]: North China is outlined by thick black lines.




Figure 4. Fractional transboundary contributions to PM_{2.5} concentrations over China. The percentage fraction of transboundary-contributed PM_{2.5} in China's total population-weighted PM_{2.5} in 2015, 2030 and 2060. Future anthropogenic emission scenarios are represented by different colors and markers following the legend.





Figure 5. PM_{2.5}-associated premature deaths in China. Total PM_{2.5}-associated premature deaths in China for 2015, 2030 and 2060 under each emission scenario in China (denoted as text at the top) and in other countries (represented by colors in the legend). Numbers denote the estimated deaths in each scenario. Error bars represent the 95% confidence interval of the RR function.



1
2
3 **Figure 6.** Potential health threats associated with transboundary pollution in China. **(a)**
4 Population exposed to an annual mean population-weighted PM_{2.5} concentration above the goals
5 (35 μg m⁻³ for 2015 and 2030; 5 μg m⁻³ for 2060) due to transboundary contributions of PM_{2.5} in
6 China under different emission scenarios in China (denoted as text at the top) and in other
7 countries (represented by colors in the legend). **(b)** Transboundary-contributed
8 premature mortality in China under different emission scenarios in China (denoted as text at the
9 top) and in other countries (represented by colors in the legend). Error bars represent the 95%
10 confidence interval of the RR function.
11

1 **Table 1.** Description of future scenario settings

Scenarios	Socioeconomic pathway	Climate target	Pollution control strength	PM _{2.5} emission level	Key features
Foreign					
SSP119	Sustainability	1.9 W m ⁻² (1.5 °C)	Strong	Low	Strong economic growth via sustainable pathway. Incomes increase substantially and inequality within and between countries is greatly decreased. Significantly lower demand for energy- and resource-intensive agricultural commodities. Effective pollution controls result in substantial reductions in air pollutant emissions.
SSP245	Middle-of-the-road	4.5 W m ⁻² (3 °C)	Medium	Medium	An intermediate case between SSP1 and SSP3.
SSP370	Regional rivalry	7.0 W m ⁻² (~4 °C)	Weak	High	High inequity between countries. GDP growth is low and concentrated in current high-income nations, while population growth is focused in low- and middle- income countries. Energy system is coal-intensive. The implementation of pollution controls is delayed and less effective.
China					
Current-policy	Middle-of-the-road (SSP2)	4.5 W m ⁻² (3 °C)	Medium	Medium	Achieve China's Nationally Determined Contribution (NDC) pledges and the national PM _{2.5} air quality standard (i.e. 35 µg m ⁻³) by 2030, elucidating China's future air pollution mitigation pathway towards all the released and determined upcoming clean air policies since 2015.
Carbon-neutral	Sustainability (SSP1)	Net-zero CO ₂ emissions in 2060	Strong	Low	Pursue China's carbon-neutral commitment and the WHO's old PM _{2.5} guideline (10 µg m ⁻³) by 2060. It implements the best available end-of-pipe technologies and more stringent pollution control policies than the current-policy.

2
3
4
5
6

1 **Table 2.** Simulation configurations

Simulation type	Anthropogenic Emissions		Emission year	Met field year
	China	Foreign countries		
Base_2015 China_2015	MEIC MEIC	CEDS None	2015	2015
Base_current_SSP119_2030 Base_current_SSP245_2030 Base_current_SSP370_2030 China_current_2030	Current-policy Current-policy Current-policy Current-policy	SSP119 SSP245 SSP370 None	2030	
Base_neutral_SSP119_2030 Base_neutral_SSP245_2030 Base_neutral_SSP370_2030 China_neutral_2030	Carbon-neutral Carbon-neutral Carbon-neutral Carbon-neutral	SSP119 SSP245 SSP370 None		
Base_current_SSP119_2060 Base_current_SSP245_2060 Base_current_SSP370_2060 China_current_2060	Current-policy Current-policy Current-policy Current-policy	SSP119 SSP245 SSP370 None	2060	
Base_neutral_SSP119_2060 Base_neutral_SSP245_2060 Base_neutral_SSP370_2060 China_neutral_2060	Carbon-neutral Carbon-neutral Carbon-neutral Carbon-neutral	SSP119 SSP245 SSP370 None		

2
3

1 **Supplementary Information for**
2 **The underappreciated role of transboundary pollution in future air quality and health**
3 **improvements in China**

4
5 Jun-Wei Xu, Jintai Lin*, Dan Tong, Lulu Chen

6
7 ¹Laboratory for Climate and Ocean–Atmosphere Studies, Department of Atmospheric and
8 Oceanic Sciences, School of Physics, Peking University, Beijing, China


9 ²Department of Earth System Science, Ministry of Education Key Laboratory for Earth System
10 Modelling, Tsinghua University, Beijing, China

11
12 Correspondence: Jintai Lin (linjt@pku.edu.cn).


13 This Supplemental Information document presents the additional data to support the primary
14 document.

15

16




1
2 **Figure S1.** Future emissions of PM_{2.5}-related pollutants in China and in other countries. **(a)**
3 Emissions for foreign countries within our Flex-Grid simulation domain (11° S–60° N, 30°–150°
4 E) for future years are projected under SSP-RCP scenarios, with updates on base year emissions
5 and the harmonization year in this study. Colors represent the emissions of different pollutants
6 following the legend. **(b)** Emissions for China for future years are projected with the current-
7 policy scenario and the carbon-neutral scenario.
8




1
2
3
4
5
6
7
8
9
10
11
12

Figure S2. Correction and evaluation of simulated $PM_{2.5}$ over China. **(a)** Comparison of the observed and unscaled simulated annual mean concentrations $PM_{2.5}$ at collocated grids for 2015. Both the observed and the simulated $PM_{2.5}$ were in grids where anthropogenic $PM_{2.5}$ exceeds natural $PM_{2.5}$. r refers to the correlation coefficient. NMB refers to normalized mean bias. N indicates the number of grids shown in the figure. **(b)** Same as (a), but for scaled simulated $PM_{2.5}$ concentrations to remove the systematic bias. **(c)** Same as (a), but for all grids (including both anthropogenic pollution-dominated and natural pollution-dominated grids). **(d)** Same as (b), but for all grids.



1
2 **Figure S3.** Anthropogenic emission-dominated grid cells (red) and natural emission-dominated
3 grid cells (blue) at a $0.5^\circ \times 0.625^\circ$ resolution in China. Circles indicate locations of $PM_{2.5}$
4 observations used in this study.
5
6
7
8
9
10




1
2 **Figure S4.** Simulated sectoral contributions of foreign anthropogenic emissions to PM_{2.5}
3 concentrations over the entire China, North China and Yunnan province for January in 2030
4 (top) and in 2060 (bottom) under the current-policy scenario in China and SSP370 scenario in
5 foreign countries. Sectors include agriculture (AGR), energy (ENE), industry (IND),
6 transportation (TRA), residential combustion (RES), solvent use (SLV), and waste burning
7 (WST). The simulation is conducted at a resolution of 2° x 2.5°.
8



1
2
3
4
5
6
7
8

Figure S5. Fractional contributions of foreign anthropogenic emissions from countries within and outside the Flex-Grid domain (11° S–60° N, 30°–150° E, including the entire Asia, eastern Africa and eastern Europe) to PM_{2.5} concentrations over China, North China and Yunnan province in 2030 (top) and in 2060 (bottom) under the current-policy scenario in China and SSP370 scenario in foreign countries. The simulation is conducted at a resolution of 2° x 2.5°.



9
10
11
12
13
14

Figure S6. Future population and age structure in China. Total populations in China in 2015 and future years projected under different SSP scenarios (SSP1 corresponds to China’s carbon-neutral scenario; SSP2 corresponds to China’s current-policy scenario). Colors represent different age groups according to the legend. White numbers indicate the fraction of population above the age of 65 in the total population.

Page 6: [1] Deleted

Microsoft Office User

5/30/23 10:14:00 AM

▼.....
Page 6: [2] Deleted

Microsoft Office User

5/30/23 10:33:00 AM

▼.....