

Photo-Induced Shrinking of Aqueous Glycine Aerosol Droplets

Shinnosuke Ishizuka,^{a,b,c,†,*} Oliver Reich,^{a,†} Grégory David,^a and Ruth Signorell^{a,*}

^a Laboratory of Physical Chemistry, ETH Zurich, Vladimir-Prelog-Weg 2, CH-8093, Zurich, Switzerland

^b Institute of Advanced Research, Nagoya University, Nagoya 046-8601, Japan

^c Institute of Space-Earth Environmental Research, Nagoya University, Nagoya 046-8601, Japan

ABSTRACT: Due to their small size, micrometer and submicron sized solution droplets can respond differently to physical and chemical processes compared with extended bulk material. Using optically trapped micrometer sized aqueous glycine droplets, we demonstrate a photo-induced degradation of glycine upon irradiation with visible light, even though molecular glycine does not absorb light in the near UV/vis range to any significant extent. This reaction is observed as photo-induced shrinking of the droplet, which we characterize by analyzing the elastic light scattering and the Raman spectrum of the droplet over the course of the reaction. We find the volume to shrink with a constant rate over the major part of the shrinking process. This indicates the presence of a rate limiting photocatalyst, which we attribute to mesoscopic glycine clusters in the droplet solution. Our findings relate to previous reports about enhanced absorption and fluorescence rates of amino acid solutions. However, to the best of our knowledge, this is the first experimental evidence of a photochemical pathway facilitated by mesoscopic clusters. Light interaction with such mesoscopic photoactive molecular aggregates might be more important for aerosol photochemistry than previously anticipated.

* rsignorell@ethz.ch, ishizuka.shinnosuke.a3@f.mail.nagoya-u.ac.jp

† S.I. and O.R. contributed equally to this work.

1 1. INTRODUCTION

2 Aerosols, dispersions of solid and liquid particles in a gas, are ubiquitous in Earth's atmosphere and as such play
3 an important role for many atmospheric processes (Boucher et al., 2013; PöSchl and Shiraiwa, 2015). Size, chem-
4 ical composition, viscosity and thermodynamic phase of aerosol particles respond to their environment including
5 the surrounding gas species, temperature, humidity (Bones et al., 2012; Zieger et al., 2017; Tang et al., 1997;
6 Swietlicki et al., 2008) and light irradiation (PöSchl and Shiraiwa, 2015; Cremer et al., 2016; Walser et al., 2007;
7 Corral Arroyo et al., 2022). Particular attention has been paid to their chemistry distinct from that in the bulk, a
8 phenomenon possibly arising from high surface to volume ratio of aerosol particles and the accessibility to highly
9 supersaturated states (Altaf et al., 2016; Kucinski et al., 2019; Bzdek and Reid, 2017). Various chemical reactions
10 have been shown to be accelerated in microdroplets (Cremer et al., 2016; Lee et al., 2015; Girod et al., 2011),
11 with some reactions being exclusive to the droplet phase (Lee et al., 2019). These unique reactors can act as
12 medium for the birth, growth and degradation of atmospherically relevant particles (Ruiz-Lopez et al., 2020), and
13 be utilized for organic synthesis (Bain et al., 2017). Chemical processes in prebiotic aerosols have also been
14 proposed as potential mechanisms for the origin of life (Tervahattu et al., 2004). However, molecular processes
15 leading to the anomalous chemistry in micrometer and submicron aerosol particles are still largely unknown.
16 Although some processes may be ascribed to the discontinuous and asymmetric intermolecular interactions at the
17 particle surface (Ruiz-Lopez et al., 2020), the microphysical origins behind many of the aforementioned particle
18 specific phenomena are still not adequately explored.

19
20 Glycine is an amino acid that acts as precursor to proteins and fulfills a number of other biological functions
21 (Arnstein, 1954; Hall, 1998; Jackson, 1991). With its small size and simple structure, glycine often serves as a
22 proxy for other amino acids and physiologically relevant molecules, and as such has been studied extensively in
23 the past. It is generally accepted that molecular glycine does not absorb light in the near UV/vis range, similar to
24 other amino acids (Bhat and Dharmaparakash, 2002). However, it has been shown that their optical properties
25 change when glycine molecules arrange themselves into mesoscopic clusters by forming hydrogen bond networks
26 (Terpugov et al., 2021). Furthermore, these formations respond to light irradiation in non-trivial ways which can
27 be exploited to induce long range order inside the glycine solution (Alexander and Camp, 2019; Sugiyama et al.,
28 2012; Zaccaro et al., 2001; Garetz et al., 2002; Urquidi et al., 2022) on a scale of up to millimeters (Yuyama et
29 al., 2010). While these interactions have the potential to change the optical properties of glycine ensembles sig-
30 nificantly and to enable new photochemical reaction pathways, there has been little experimental evidence for
31 such reactions so far.

32
33 In this work, we study the response of aqueous glycine droplets to irradiation by visible light. We observe the
34 shrinking of optically trapped micrometer sized glycine droplets, which can be unambiguously attributed to the
35 exposure to the trapping laser with wavelength 532 nm. To the best of our knowledge, this interaction has not
36 been reported before. We characterize it here with particular focus on the shrinking rate and its dependence on the

37 light intensity. To explain our results, we discuss possible reaction schemes based on the available experimental
38 data. Although further data is needed to elucidate the exact photochemical pathways of the observed reaction,
39 these findings demonstrate the existence of a photochemical reaction for molecules which previously have been
40 considered photochemically inert at visible wavelengths.

41

42

43 2. METHODS

44 Dual beam optical traps are widely used to confine and isolate single particles (Ashkin, 1997; Gong et al., 2018;
45 Čižmár et al., 2005; Esat et al., 2018; Reich et al., 2020). The counter-propagating tweezers (CPT) setup for
46 trapping aqueous glycine droplets is shown in Fig. 1 and consists of a continuous green laser beam (Novanta
47 Photonics Opus 532 6W), which is expanded and then split into two beams of equal power. These two beams are
48 aligned counter-propagating on a single axis and focused into the trapping cell, where a single droplet is trapped
49 between the two focii.

50

51 The droplets are generated from 1.0 M or 2.0 M aqueous solutions of glycine (purity 100%, HPLC certificate,
52 Sigma-Aldrich G7126) using a commercial atomizer (TSI 3076) with pressurized, humidified nitrogen gas (purity
53 5.0). We performed additional elementary and HPLC analyses of the glycine and the glycine solutions. The results
54 showed no indication of any trace impurities absorbing in the visible range. On this basis, we conclude that our
55 samples do not contain any contaminants that could act as photosensitizer for photochemical reactions (see be-
56 low). A system of copper tubings directs the spray of particles into the trapping cell, where the droplets agglom-
57 erate at the designated trapping position. The humidification of the nitrogen flow is necessary to ensure that the
58 droplets reach the trapping position in the liquid state. The trapping cell is filled with nitrogen gas (purity 5.0) and
59 a steady nitrogen flow formed by combining wet and dry nitrogen with adjustable flow ratios is used to control
60 the relative humidity (RH) in the cell. For the experiments reported here, the RH is set at 77 ± 3 % well above the
61 efflorescence RH of glycine at approximately 55 % (Chan et al., 2005). At this RH, the droplet solution is super-
62 saturated with an estimated glycine concentration of 60 % in mass (Chan et al., 2005), corresponding to approxi-
63 mately 5 M. Temperature and RH inside the trapping cell are monitored by a sensor (Sensirion SHT35) placed a
64 few millimeters away from the trapping position. When the agglomerated particle reaches a size of approximately
65 2-3 μ m in radius, the remainder of the droplets in the cell are flushed out with nitrogen for 20-30 minutes to ensure
66 that only the trapped droplet remains in the cell. After flushing, the power of the trapping laser is kept constant
67 until the end of the measurement.

68


69 The particle shrinking is monitored by imaging the polarization resolved two-dimensional angular optical scatter-
70 ing (polarization resolved TAOS) of the particle (Parmentier et al., 2022), as shown in Fig. 2. The TAOS image
71 is obtained by collecting the elastically scattered light of the trapping beams under a scattering angle of $90 \pm 24^\circ$
72 with an objective (Mitutoyo 20x NA 0.42). The parallel and perpendicular polarization components with respect
73 to the scattering plane (TAOS PPol and TAOS SPol) are separated using a polarization beam splitter and recorded
74 with separate CMOS cameras (Thorlabs DCC1545M). The scattering intensity for each polarization is calculated
75 from the average of the respective TAOS image and recorded over time. At specific times, the shrinking spherical
76 particle reaches a size at which it is in resonance with the light of the trapping beams, which corresponds to a Mie
77 resonance (Bohren and Huffman, 2008). From the comparison of the recorded evolution of the polarization re-
78 solved scattering intensity to simulations using Mie theory, the size of the particle can be determined at the specific

79 times. Fig. 3 shows an example of such a TAOS analysis. The values of the size at the discrete points in time,
80 obtained from the times where the particles experience a Mie resonance, can then be interpolated with high accu-
81 racy to obtain the full size evolution of the particle over the course of the measurement.

82

83 The molecular composition of the particle is monitored by continuous recording of Raman spectra (David et al.,
84 2020) during the shrinking process. To this end, the light scattered by the particle is collected under a scattering
85 angle of $90 \pm 24^\circ$ by a second objective and fiber coupled into a low noise, high sensitivity spectrograph (Andor
86 KY-328i-A). The inelastically scattered light is analyzed in the range 540-680 nm which corresponds to Raman
87 shifts of 280-4100 cm^{-1} . This range contains in particular the O-H symmetric stretching mode of water ($\nu_2\text{-H}_2\text{O}$,
88 2700-3750 cm^{-1}) as well as several vibrational modes of glycine, which we exploit for the characterization of the
89 molecular composition (see later data for an example).


90



91

92 **Figure 1** Counter-propagating tweezers setup. The laser beam is first expanded by a factor of 5 and then split into
93 two beams by the polarizing beam splitter (PBS). The half-waveplate ($\lambda/2$) rotates the polarization to 45° with
94 respect to the axes of the PBS to ensure equal power splitting between the two beams. The beams are aligned on
95 a single axis and focused into the trapping cell using two aspherical lenses (AL). An optical isolator introduced
96 at the start of the beam path prevents unwanted optical feedback into the laser head.

97



98

99

100

101


102

103

104

105

Figure 2 Setup for TAOS imaging and Raman spectroscopy. The trapping beams run perpendicular to the figure plane. The scattered light of the trapping beams is collected horizontally and vertically at a scattering angle of $90 \pm 24^\circ$. The vertical beam is split into parallel and perpendicular polarized light with respect to the scattering plane and the respective beam is loosely focused on a CMOS camera (P TAOS and S TAOS respectively). The horizontal beam is filtered for the spectral range of 540 – 680 nm and fiber coupled into a low noise, high sensitivity spectrometer (Spect. Raman) for measurement of the Raman spectrum.




106

107 **Figure 3** Analysis of the two-dimensional angular optical scattering spectrum (TAOS). **a)** Experimental polari-
 108 zation resolved total TAOS signal over time. **b)** Zoom on a specific time interval for clarity **c)** Simulated polari-
 109 zation resolved total TAOS signal as function of particle radius in the specific time interval. Arrows indicate the
 110 peak assignment based on the similarities between the peak shapes.

111

112 **3. RESULTS AND DISCUSSION**


113 The shrinking of the aqueous glycine droplets over time is shown in Fig. 4 for three representative examples. The
114 droplets are trapped using different laser powers, which affects the rate at which their volume is decreasing. For
115 all laser powers, the volume is observed to shrink linearly with time over a large portion of the shrinking process.
116 After a certain point (‘point of acceleration’), when the droplet has lost 60 – 80 % of its volume, the shrinking
117 suddenly accelerates, only to continue again approximately linearly with time, albeit at a higher rate.



118

119 **Figure 4** Volume shrinking of glycine droplets as a function of time. **a)** Droplet trapped at 300 mW nominal laser
120 power **b)** droplet trapped at 1000 mW nominal laser power **c)** droplet trapped at 3000 mW nominal laser power.
121 At higher power, the shrinking proceeds faster (visible by the larger slopes). Solid and dashed lines indicate the
122 best linear fit before and after the acceleration of the shrinking observed at approximately **a)** 280 h, **b)** 59 h and
123 **c)** 53 h.

124 To quantify the dependence of the shrinking rate on the light intensity incident on the particle, a linear fit is
125 performed on the data before and after the point of acceleration. Since between different experiments, the align-
126 ment of the optical trap, and hence the focusing of the laser light on the particle, is subject to temporal mechanical
127 drifts, the nominal laser power used for trapping of the droplets is not an optimal indicator for the incident inten-
128 sity. Instead, we use the intensity of the scattered light as a measure that is proportional to the incident light
129 intensity. The intensity of the scattered light is obtained from the average signal of the TAOS PPol and SPol
130 images during the time of the droplet shrinking. To ensure consistency between the different experiments, the
131 average of the light intensity is taken over the same volume interval of $[65.4, 51.0] \mu\text{m}^3$ (radius $[2.5, 2.1] \mu\text{m}$) for
132 all droplets. This interval corresponds to the interval for which we obtained data for most droplets. Fig. 5 shows
133 the resulting initial shrinking rates as function of light intensity. The shrinking rate is observed to be proportional
134 to the light intensity, confirming that the shrinking is induced by the trapping laser at a wavelength of 532 nm.



135


136 **Figure 5** Fitted shrinking rate as function of light intensity. The solid line indicates the best linear fit through
137 the origin.

138 To gain further insight into the shrinking mechanism, we analyze the evolution of the Raman spectra over the
139 course of the shrinking process. A representative example of such an evolution is shown in Fig. 6. For Raman
140 spectra of single spherical particles, so called morphology dependent resonances, or whispering gallery modes
141 (WGMs) (Oraevsky, 2002), are superimposed on the molecular signals. To separate the WGMs from the molecular
142 signal of interest, the Raman spectra are normalized and stacked in chronological order from left to right, as shown
143 in Fig. 6a. The WGMs show up as a manifold of thin slanted lines bending towards lower wavenumbers with
144 increasing time as the droplet shrinks. Molecular band positions on the other hand are independent of particle
145 size, and are identified as horizontal lines in the evolution of the Raman spectra.

146

147 From Fig. 6a it is evident that the molecular Raman signal remains qualitatively the same, indicating that no
148 significant change in the molecular composition takes place in the droplet over the course of the shrinking process.
149 This behavior is observed for all investigated droplets. Since the particle loses the major part of its volume during

150 the shrinking, this implies that glycine is removed from the droplet as a consequence of chemical reaction (*vide*
 151 *infra*). As the water vapor pressure of the droplet is given by the surrounding RH of 77 % and therefore has to
 152 remain constant, the removal of glycine from the droplet must be accompanied by the evaporation of water in
 153 order to maintain the equilibrium glycine concentration. A quantitative analysis of the Raman spectrum (Fig. 6b)
 154 reveals that the spectral intensity of glycine modes with respect to the O-H stretching mode of water remains
 155 constant, confirming a constant concentration of glycine molecules during the shrinking process. The measure-
 156 ment ends when the particle becomes too small for stable trapping, and therefore leaves the optical trap.
 157



158
 159 **Figure 6** Temporal evolution of Raman signal during droplet shrinking. **a)** Normalized Raman spectra stacked
 160 chronologically from left to right. The molecular Raman bands are visible as horizontal straight white lines on the
 161 dark background. The finer, slanted and curved lines correspond to whispering gallery modes. After approxi-
 162 mately 49 h, the particle leaves the optical trap as it reaches a size that is too small for trapping, and only back-
 163 ground is recorded. The band remaining afterwards at 2323 cm^{-1} corresponds to the nitrogen gas in the trapping
 164 cell. **b)** Ratio of the molecular glycine bands centered at 1064 cm^{-1} and 1440 cm^{-1} to the water band at around
 165 3400 cm^{-1} . Peaks and dips in the graphs correspond to spectra where a whispering gallery mode is superimposed
 166 on the glycine signal and the water signal, respectively, and are not relevant for the molecular composition. The
 167 dashed horizontal lines are a guide to the eye. In this example, the acceleration of the droplet shrinking is observed
 168 at $t_{\text{acc}} = 46$ h. From this point in time onwards until the particle is lost, the faster shrinking leads to more frequent
 169 WGMs perceived as an apparent increase of noise in the data.

170 Aqueous droplets which contain a stable non-volatile solute do not shrink over time when they are optically
 171 trapped (see Supplementary Section S1 for an example). The observation of droplet shrinking in the case of dis-
 172 solved glycine proves that glycine does not remain stable inside the droplet solution. Furthermore, the linear
 173 dependence of the shrinking rate on the light intensity (Fig. 5) proves that the observed shrinking is induced by
 174 the laser light. In particular, the volume shrinking rate increases from 0.18 to 2.25 $\mu\text{m}^3 \text{h}^{-1}$ when the light intensity

175 is increased by the same relative amount, thus spanning over one order of magnitude. This observation is intri-
176 guing as aqueous glycine is not known to absorb light in the visible range, similar to other amino acids (Bhat and
177 Dharmaparakash, 2002). In addition, the linear dependence between shrinking rate and light intensity rules out the
178 possibility of multiphoton absorption, which might otherwise populate energetically excited states of glycine to
179 induce chemical reactions. The fact that the observed shrinking is driven by the incident light indicates a photo-
180 chemical reaction or photothermal processes occurring in the aqueous glycine droplet. As is shown in the follow-
181 ing however, photothermal processes can be ruled out by our experimental data.

182


183 The observation of a constant shrinking rate over a large portion of the experiments (Fig. 4) provides another
184 important piece of information, as this allows us to rule out some of the possible shrinking mechanisms. For
185 instance, assuming some residual absorption of light at 532 nm by the droplet solution, one might argue that the
186 shrinking is a consequence of enhanced evaporation due to the droplet heating by the laser. The evaporation of
187 glycine from the droplet can be approximated by the Hertz-Knudsen equation:

$$188 \quad \frac{dN}{dt} = S \cdot \frac{\alpha p}{\sqrt{2\pi MRT}} \quad (1)$$

189 where $\frac{dN}{dt}$ is the molar evaporation rate of glycine, S is the droplet's surface, p is the partial pressure and M is the
190 molar mass of glycine, R the gas constant, T the temperature and α a heuristic sticking coefficient with values
191 between 0 and 1. It is evident from Eq. (1) that the rate of shrinking by evaporation should scale with the surface
192 area of the droplet, and that in this case a deceleration of the shrinking should be observed over time, contrary to
193 the experimental data. Moreover, any heating by laser light absorption would at most lead to a very small temper-
194 ature rise due to the efficient cooling by the surrounding gas (see Supplementary Section S2 for an upper estimate
195 of the temperature increase). In addition to these arguments, if evaporation were significant, some evaporation
196 should still be observable even at low light intensity, where the heating of the droplet is negligible and hence any
197 deviations from room temperature can be neglected. As seen from Fig. 5 however, there is no shrinking observable
198 for low light intensities. We can therefore rule out evaporation as the dominant shrinking mechanism.

199

200 The observation of a constant shrinking rate also excludes photochemical reactions in which molecular glycine
201 directly absorbs photons:





203 where Gly(aq) is a solvated glycine molecule, $h\nu$ is the energy of the incoming photon and P is the reaction prod-
204 uct that is removed from the droplet (aq) into the surrounding gas phase (g) afterwards. As mentioned above,
205 molecular glycine is considered non-absorbing at 532 nm. However, if we nevertheless assume that molecular
206 glycine could be very weakly absorbing as in Eq. (2), the photon density inside the droplet would be constant over
207 the course of the photochemical reaction. Earlier studies (Parmentier et al., 2021; Corral Arroyo et al., 2022) show
208 that no significant size dependence of the optical confinement is observed for weakly absorbing particles in the
209 micrometer- and submicrometer-size ranges. Hence, the first step in Eq. (2) would be pseudo-first order, for which

210 the reaction rate is proportional to the concentration of glycine molecules in the droplet. Therefore, one would
211 expect a decrease in the observed shrinking rate over time, which contradicts the experimental observation.

212

213 The examples above illustrate that any mechanism, in which glycine molecules directly absorb incoming photons,
214 cannot explain the constant shrinking rates. This implies that a more intricate reaction must take place in the
215 droplet. We suggest the following simplified scheme with an additional reaction partner M:



218 where M^* denotes a photoexcited state of M, and P is the reaction product of glycine in the presence of this pho-
219 toexcited species. This scheme represents a mediated reaction in which the reaction partner M is activated by light
220 absorption, and then reacts with glycine and returns back to the ground state. M shows the characteristics of a
221 photosensitizer (Corral Arroyo et al., 2018; George et al., 2015; Wang et al., 2020; Rapf and Vaida, 2016), which
222 is not consumed during the reaction, and therefore the amount of M remains constant in the droplet. We further
223 assume that the light absorption of the photosensitizer M is the rate limiting step, i.e., that Eq. (3) proceeds much
224 slower than Eq. (4). This is equivalent to requiring that M is only weakly absorbing, or that the concentration of
225 M in the droplet is low. Since the amount of M in the rate limiting step (Eq. (3)) remains constant during the
226 shrinking process, so does M^* (quasistationary), resulting in a constant rate of degradation of Gly (Eq. (4)). As-
227 suming that the absolute concentration of M is much smaller than that of Gly at all times, and therefore has no
228 relevant effect on the equilibrium water vapor pressure of the droplet, the volume shrinking rate is predicted to be
229 constant in accordance with the experimental data.

230

231 Although the proposed mechanism in Eq. (3) and (4) concurs with the observed constant shrinking rates, it does
232 not yet specify the nature of the photosensitizer and its reaction with glycine. We first discuss potential candidates
233 for the photosensitizer. Contamination during the preparation of the different aqueous glycine solutions used in
234 this study was minimized by using pure substances (glycine purity $\geq 99\%$, water resistivity $18.2 \text{ M}\Omega\cdot\text{s}$). No
235 correlation between the shrinking rates in Fig. 5 and the age of the solution at the time of the measurements was
236 observed, which indicates that there is no accumulation of photoactive contaminants in the solution after the
237 preparation. We therefore argue that contamination is not the origin of the photosensitizer.

238

239 It is evident from the previous discussion that the photosensitizer must possess an absorption band at 532 nm,
240 which is not the case for single solvated glycine molecules. The optical properties of molecules may change
241 however when forming intermolecular bonds, e.g. leading to an enhancement of the absorption and fluorescence
242 in the case of protein aggregates (Homchaudhuri and Swaminathan, 2004; Shukla et al., 2004; Chan et al., 2013;
243 Pinotsi et al., 2013) and amino acid clusters (Chen et al., 2018). For glycine solutions in particular, observations
244 of light absorption in the near UV/vis range have been attributed to the presence of mesoscopic clusters (Jawor-

245 Baczynska et al., 2013; Zimbitas et al., 2019), specifically due to the formation of hydrogen bonds between indi-
246 vidual molecules (Terpugov et al., 2021).

247


248 Mesoscopic clusters occur naturally in both undersaturated and supersaturated glycine solutions, though a kinetic
249 barrier may have to be overcome for their formation (Jawor-Baczynska et al., 2013).). A similar barrier needs to
250 be overcome to dissolve these clusters. Hence the clusters can remain kinetically stable even if the number of
251 glycine monomers decreases. The average size of the mesoscopic clusters, typically of the order of 100 nm for
252 bulk solutions, depends not only on the monomer concentration but also on the history of the sample, indicating
253 that the clusters do not necessarily remain in thermodynamic equilibrium after formation (Jawor-Baczynska et
254 al., 2013). Based on these observations, we propose that the photo-induced droplet shrinking is mediated by light
255 absorption of mesoclusters in the glycine solution. These mesoclusters remain kinetically stable during the droplet
256 shrinking, despite the decrease of the absolute number of glycine monomers (i. e. constant glycine monomer
257 concentration). Hence, these clusters act as stable photosensitizers inside the droplets, in accordance with the
258 observed constant shrinking rate. The observation of a constant initial shrinking rate thus allows us to narrow
259 down the possible reaction mechanisms at work in the droplets. Alternative schemes might be conceivable, such
260 as the existence of several reaction partners for glycine. However, there is no further experimental evidence in
261 favor of more complex alternatives to the simple mechanism proposed (Eq. (3) and (4)). Furthermore, as pointed
262 out above, mesoscopic glycine clusters match the requisite characteristics of the proposed photosensitizer, and are
263 therefore likely candidates. To pursue the argument, let us further discuss the role of the mesoscopic clusters in
264 the observed acceleration of the droplet shrinking.

265

266 The acceleration of the droplet shrinking proceeds relatively promptly at the point of acceleration when the parti-
267 cle has lost a typical amount of 75 % of its volume (Fig. 4). Assuming that the total number of mesoclusters
268 remains approximately constant (M in Eqs.(3) and (4)), their concentration has increased by an approximate factor
269 of 4 at this point. The sudden nature of the shrinking acceleration hints at a phase transition inside the particle. In
270 particular, the increase in nanocluster concentration may trigger the separation of a dense cluster phase inside the
271 droplets as part of a liquid-liquid phase transition. While a definite conclusion has to await more detailed micro-
272 scopic investigations, we present the following arguments in favor of this explanation. Mesoscopic clusters in
273 aqueous solutions are known to interact with focused light irradiation by assembling in the focal point of the light
274 beam due to the optical force that acts on the individual clusters (Sugiyama et al., 2012). This mechanism is the
275 basis of laser-induced phase transitions(Alexander and Camp, 2019), in the particular case of glycine both for
276 liquid-liquid phase separation (Sugiyama et al., 2012; Yuyama et al., 2010) and solid crystal nucleation (Sugiyama
277 et al., 2012; Alexander and Camp, 2019; Yuyama et al., 2010; Zaccaro et al., 2001; Garetz et al., 2002). It should
278 be noted at this point that the spot size in the focus of our optical trap is slightly larger (5 μm) than the typical
279 droplet size, and that therefore, it might appear unlikely that the electromagnetic field gradient is strong enough
280 to induce cluster aggregation in our case. However, micrometer sized droplets exhibit a large variance in the
281 spatial distribution of the internal light field, owing to the nanofocusing effect (Cremer et al., 2016; Corral Arroyo

282 et al., 2022), which can provide the field gradients necessary for aggregation. If in the case of our trapped droplets,
283 the acceleration is due to a phase transition, it would likely be assisted by the light irradiance. One would therefore
284 expect a dependence of the observed shrinking acceleration on the incident light intensity. Fig. 7 shows the meas-
285 ured acceleration ratio, that is, the ratio between the shrinking rate after to before the point of acceleration, as a
286 function of light intensity. From this data it is evident that the ratio increases with higher light intensity, which
287 agrees with our explanation of a light-induced phase transition. Since in this scheme, the clusters are expected to
288 aggregate in the regions of high light intensity after reaching a critical concentration, this would lead to a larger
289 absorption rate and thus a larger subsequent reaction rate, in agreement with the observation. Further studies will
290 be necessary to provide conclusive evidence for this explanation.

291



292

293 **Figure 7** Acceleration of droplet shrinking versus light intensity. The solid line represents the best linear fit
294 through the measurement data.

295 More data will also be required to understand the specifics of the interaction between the photosensitizer and the
296 solvated glycine molecules in Eq. (4). Here, we can only provide a qualitative discussion based on the available
297 data. The Raman spectra (Fig. 6) show no detectable change in the molecular composition, even after the droplet
298 has lost the major part of its volume during the shrinking process. Since this observation rules out the accumula-
299 tion of reaction products in the droplets over time, the reaction products must be small, volatile compounds that
300 quickly evaporate into the surrounding gas phase. Known degradation mechanisms of glycine and other amino
301 acids in aqueous solution proceed via reaction with radical species, in particular solvated electrons e_{solv}^- and hy-
302 droxyl $\cdot\text{OH}$ radicals (Moenig et al., 1985; Garrison, 1964, 1972), which form as part of a photosensitized reaction
303 with chromophoric organic matter in water (Lundeen et al., 2014; Sun et al., 2018; Mopper and Zika, 1987).
304 Currently, our data does not allow us to distinguish between different degradation pathways.

305

306 4. CONCLUSIONS

307 We have demonstrated that single micrometer sized aqueous glycine droplets respond to the illumination with
308 laser light of 532 nm by shrinking, despite the fact that molecular glycine does not absorb in the near UV/vis

309 range. Most remarkably, the volume shrinking rate remained constant over the major part of the shrinking process.
310 This indicates a photo-induced decay of glycine molecules in the presence of a rate limiting catalyst, or photosen-
311 sitizer, and the subsequent evaporation of small, volatile reaction products. Based on the available literature data,
312 we propose that intrinsic mesoscopic clusters of glycine molecules formed by hydrogen bonding in the aqueous
313 solution are the most plausible candidates for this photosensitizer. The presence of mesoscopic glycine clusters
314 would also explain the sudden acceleration of the shrinking rate occurring at a volume loss of ~75 %. Because of
315 its dependence on the light intensity, we attribute this sudden rate change to arise from the interaction of the
316 mesoclusters with the incident light, possibly initiating a light-induced phase transition.

317

318 This study provides yet another example of the non-trivial interactions of light with aqueous glycine solutions
319 (Alexander and Camp, 2019; Sugiyama et al., 2012; Zaccaro et al., 2001; Garetz et al., 2002; Yuyama et al., 2010),
320 which facilitate previously undiscovered reaction pathways - interactions that are likely not exclusive to glycine.
321 Light harvesting by and light interaction with such mesoscopic photosensitizers in aerosol droplets might also
322 have played a role in the formation of more complex organic molecules under prebiotic conditions. Further in-
323 vestigations are needed to shed light on the specifics of the observed phenomena, and to yield new insight into
324 the underlying reaction mechanisms, which remain elusive in part. Studying solutions in micrometer sized drop-
325 lets (attoliter volumes) using high laser powers offers the advantage of much higher sensitivity to photo-induced
326 reactions than typically achievable with bulk solutions.

327

328 **Acknowledgements**

329 This project was supported by the Japan Society for the Promotion of Science (Oversea Research Fellowship and
330 Grant-in-aid for Scientific Research 1186520; S.I.), by Tokai Pathways to Global Excellence, part of MEXT
331 Strategic Professional Development Program for Young Researchers (S.I.), by the Swiss National Science Foun-
332 dation (SNSF project number 200020_200306), and by ETH Zürich. We are very grateful to D. Stapfer and M.
333 Steger from our workshops for technical support, and to D. Zindel for the chemical analysis of our samples and
334 helpful discussions.

335

336 **Data Repository**

337 The data that support our findings are deposited on the ETH Research Collection under <https://doi.org/xx>.

338

- Alexander, A. J. and Camp, P. J.: Non-photochemical laser-induced nucleation, *The Journal of chemical physics*, 150, 040901, 2019.
- Altaf, M. B., Zuend, A., and Freedman, M. A.: Role of nucleation mechanism on the size dependent morphology of organic aerosol, *Chem. Commun.*, 52, 9220-9223, 2016.
- Arnstein, H.: The metabolism of glycine, in: *Adv. Protein Chem.*, Elsevier, 1-91, 1954.
- Ashkin, A.: Optical trapping and manipulation of neutral particles using lasers, *Proceedings of the National Academy of Sciences*, 94, 4853-4860, 1997.
- Bain, R. M., Sathyamoorthi, S., and Zare, R. N.: "On - droplet" chemistry: the cycloaddition of diethyl azodicarboxylate and quadricyclane, *Angew. Chem.*, 129, 15279-15283, 2017.
- Bhat, M. N. and Dharmaprakash, S.: Growth of nonlinear optical γ -glycine crystals, *J. Cryst. Growth*, 236, 376-380, 2002.
- Bohren, C. F. and Huffman, D. R.: *Absorption and scattering of light by small particles*, John Wiley & Sons 2008.
- Bones, D. L., Reid, J. P., Lienhard, D. M., and Krieger, U. K.: Comparing the mechanism of water condensation and evaporation in glassy aerosol, *Proceedings of the National Academy of Sciences*, 109, 11613-11618, 2012.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., and Lohmann, U.: Clouds and aerosols, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 571-657, 2013.
- Bzdek, B. R. and Reid, J. P.: Perspective: Aerosol microphysics: From molecules to the chemical physics of aerosols, *The Journal of Chemical Physics*, 147, 220901, 2017.
- Chan, F. T., Schierle, G. S. K., Kumita, J. R., Bertocini, C. W., Dobson, C. M., and Kaminski, C. F.: Protein amyloids develop an intrinsic fluorescence signature during aggregation, *Analyst*, 138, 2156-2162, 2013.
- Chan, M. N., Choi, M. Y., Ng, N. L., and Chan, C. K.: Hygroscopicity of water-soluble organic compounds in atmospheric aerosols: Amino acids and biomass burning derived organic species, *Environ. Sci. Technol.*, 39, 1555-1562, 2005.
- Chen, X., Luo, W., Ma, H., Peng, Q., Yuan, W. Z., and Zhang, Y.: Prevalent intrinsic emission from nonaromatic amino acids and poly (amino acids), *Science China Chemistry*, 61, 351-359, 2018.
- Čižmár, T., Garcés-Chávez, V., Dholakia, K., and Zemánek, P.: Optical conveyor belt for delivery of submicron objects, *Appl. Phys. Lett.*, 86, 174101, 2005.
- Corral Arroyo, P., David, G., Alpert, P. A., Parmentier, E. A., Ammann, M., and Signorell, R.: Amplification of light within aerosol particles accelerates in-particle photochemistry, *Science*, 376, 293-296, 2022.
- Corral Arroyo, P., Bartels-Rausch, T., Alpert, P. A., Dumas, S. p., Perrier, S. b., George, C., and Ammann, M.: Particle-phase photosensitized radical production and aerosol aging, *Environ. Sci. Technol.*, 52, 7680-7688, 2018.
- Cremer, J. W., Thaler, K. M., Haisch, C., and Signorell, R.: Photoacoustics of single laser-trapped nanodroplets for the direct observation of nanofocusing in aerosol photokinetics, *Nature communications*, 7, 1-7, 2016.
- David, G., Parmentier, E. A., Taurino, I., and Signorell, R.: Tracing the composition of single e-cigarette aerosol droplets in situ by laser-trapping and Raman scattering, *Scientific reports*, 10, 1-8, 2020.
- Esat, K., David, G., Poulkas, T., Shein, M., and Signorell, R.: Phase transition dynamics of single optically trapped aqueous potassium carbonate particles, *Physical Chemistry Chemical Physics*, 20, 11598-11607, 2018.
- Garetz, B. A., Matic, J., and Myerson, A. S.: Polarization switching of crystal structure in the nonphotochemical light-induced nucleation of supersaturated aqueous glycine solutions, *Phys. Rev. Lett.*, 89, 175501, 2002.
- Garrison, W. M.: Actions of ionizing radiations on nitrogen compounds in aqueous media, *Radiation Research Supplement*, 4, 158-174, 1964.
- Garrison, W. M.: *Radiation-induced reactions of amino acids and peptides*, Univ. of California, Berkeley, 1972.
- George, C., Ammann, M., D'Anna, B., Donaldson, D., and Nizkorodov, S. A.: Heterogeneous photochemistry in the atmosphere, *Chem. Rev.*, 115, 4218-4258, 2015.
- Girod, M., Moyano, E., Campbell, D. I., and Cooks, R. G.: Accelerated bimolecular reactions in microdroplets studied by desorption electrospray ionization mass spectrometry, *Chemical Science*, 2, 501-510, 2011.
- Gong, Z., Pan, Y.-L., Videen, G., and Wang, C.: Optical trapping and manipulation of single particles in air: Principles, technical details, and applications, *J. Quant. Spectrosc. Radiat. Transfer*, 214, 94-119, 2018.
- Hall, J. C.: Glycine, *Journal of Parenteral and Enteral Nutrition*, 22, 393-398, 1998.
- Homchaudhuri, L. and Swaminathan, R.: Near ultraviolet absorption arising from lysine residues in close proximity: a probe to monitor protein unfolding and aggregation in lysine-rich proteins, *Bull. Chem. Soc. Jpn.*, 77, 765-769, 2004.
- Jackson, A. A.: The glycine story, *Eur J Clin Nutr*, 45, 59-65, 1991.
- Jawor-Baczynska, A., Moore, B. D., Lee, H. S., McCormick, A. V., and Sefcik, J.: Population and size distribution of solute-rich mesospecies within mesostructured aqueous amino acid solutions, *Faraday Discuss.*, 167, 425-440, 2013.
- Kucinski, T. M., Dawson, J. N., and Freedman, M. A.: Size-Dependent Liquid-Liquid Phase Separation in Atmospherically Relevant Complex Systems, *The Journal of Physical Chemistry Letters*, 10, 6915-6920, 2019.
- Lee, J. K., Banerjee, S., Nam, H. G., and Zare, R. N.: Acceleration of reaction in charged microdroplets, *Q. Rev. Biophys.*, 48, 437-444, 2015.
- Lee, J. K., Samanta, D., Nam, H. G., and Zare, R. N.: Micrometer-sized water droplets induce spontaneous reduction, *Journal of the American Chemical Society*, 141, 10585-10589, 2019.
- Lundeen, R. A., Janssen, E. M.-L., Chu, C., and McNeill, K.: Environmental photochemistry of amino acids, peptides and proteins, *CHIMIA International Journal for Chemistry*, 68, 812-817, 2014.
- Moenig, J., Chapman, R., and Asmus, K. D.: Effect of the protonation state of the amino group on the $\cdot\text{OH}$ radical induced decarboxylation of amino acids in aqueous solution, *The Journal of Physical Chemistry*, 89, 3139-3144, 1985.
- Mopper, K. and Zika, R. G.: *Natural photosensitizers in sea water: riboflavin and its breakdown products*, in, ACS Publications, 1987.
- Oravsky, A. N.: Whispering-gallery waves, *Quantum electronics*, 32, 377, 2002.

404 Parmentier, E. A., Corral Arroyo, P., Gruseck, R., Ban, L., David, G., and Signorell, R.: Charge Effects on the Photodegradation of Single
405 Optically Trapped Oleic Acid Aerosol Droplets, *The Journal of Physical Chemistry A*, 126, 4456-4464, 2022.
406 Parmentier, E. A., David, G., Arroyo, P. C., Bibawi, S., Esat, K., and Signorell, R.: Photochemistry of single optically trapped oleic acid
407 droplets, *J. Aerosol Sci*, 151, 105660, 2021.
408 Pinotsi, D., Buell, A. K., Dobson, C. M., Kaminski Schierle, G. S., and Kaminski, C. F.: A label - free, quantitative assay of amyloid fibril
409 growth based on intrinsic fluorescence, *ChemBioChem*, 14, 846-850, 2013.
410 Pöschl, U. and Shiraiwa, M.: Multiphase chemistry at the atmosphere-biosphere interface influencing climate and public health in the
411 anthropocene, *Chem. Rev.*, 115, 4440-4475, 2015.
412 Rapf, R. J. and Vaida, V.: Sunlight as an energetic driver in the synthesis of molecules necessary for life, *Physical Chemistry Chemical*
413 *Physics*, 18, 20067-20084, 2016.
414 Reich, O., David, G., Esat, K., and Signorell, R.: Weighing picogram aerosol droplets with an optical balance, *Communications Physics*, 3,
415 223, 10.1038/s42005-020-00496-x, 2020.
416 Ruiz-Lopez, M. F., Francisco, J. S., Martins-Costa, M. T., and Anglada, J. M.: Molecular reactions at aqueous interfaces, *Nature Reviews*
417 *Chemistry*, 4, 459-475, 2020.
418 Shukla, A., Mukherjee, S., Sharma, S., Agrawal, V., Kishan, K. R., and Guptasarma, P.: A novel UV laser-induced visible blue radiation
419 from protein crystals and aggregates: scattering artifacts or fluorescence transitions of peptide electrons delocalized through hydrogen
420 bonding?, *Arch. Biochem. Biophys.*, 428, 144-153, 2004.
421 Sugiyama, T., Yuyama, K.-i., and Masuhara, H.: Laser trapping chemistry: from polymer assembly to amino acid crystallization, *Acc. Chem.*
422 *Res.*, 45, 1946-1954, 2012.
423 Sun, Z., Zhang, C., Xing, L., Zhou, Q., Dong, W., and Hoffmann, M. R.: UV/nitritotriacetic acid process as a novel strategy for efficient
424 photoreductive degradation of perfluorooctanesulfonate, *Environ. Sci. Technol.*, 52, 2953-2962, 2018.
425 Swietlicki, E., Hansson, H. C., Hämeri, K., Svenningsson, B., Massling, A., McFiggans, G., McMurry, P. H., Petäjä, T., Tunved, P., Gysel,
426 M., Topping, D., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of
427 submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments—a review, *Tellus B: Chemical*
428 *and Physical Meteorology*, 60, 432-469, 10.1111/j.1600-0889.2008.00350.x, 2008.
429 Tang, I. N., Tridico, A. C., and Fung, K. H.: Thermodynamic and optical properties of sea salt aerosols, *Journal of Geophysical Research:*
430 *Atmospheres*, 102, 23269-23275, 10.1029/97jd01806, 1997.
431 Terpugov, E. L., Kondratyev, M. S., and Degtyareva, O. V.: Light-induced effects in glycine aqueous solution studied by Fourier transform
432 infrared-emission spectroscopy and ultraviolet-visible spectroscopy, *J. Biomol. Struct. Dyn.*, 39, 108-117, 2021.
433 Tervahattu, H., Tuck, A., and Vaida, V.: Chemistry in prebiotic aerosols: a mechanism for the origin of life, in: *Origins*, Springer, 153-165,
434 2004.
435 Urquidi, O., Brazard, J., LeMessurier, N., Simine, L., and Adachi, T. B.: In situ optical spectroscopy of crystallization: One crystal nucleation
436 at a time, *Proceedings of the National Academy of Sciences*, 119, e2122990119, 2022.
437 Walser, M. L., Park, J., Gomez, A. L., Russell, A. R., and Nizkorodov, S. A.: Photochemical aging of secondary organic aerosol particles
438 generated from the oxidation of d-limonene, *The Journal of Physical Chemistry A*, 111, 1907-1913, 2007.
439 Wang, X., Gemayel, R., Hayeck, N., Perrier, S., Charbonnel, N., Xu, C., Chen, H., Zhu, C., Zhang, L., and Wang, L.: Atmospheric
440 photosensitization: a new pathway for sulfate formation, *Environ. Sci. Technol.*, 54, 3114-3120, 2020.
441 Yuyama, K.-i., Sugiyama, T., and Masuhara, H.: Millimeter-scale dense liquid droplet formation and crystallization in glycine solution
442 induced by photon pressure, *The Journal of Physical Chemistry Letters*, 1, 1321-1325, 2010.
443 Zaccaro, J., Matic, J., Myerson, A. S., and Garetz, B. A.: Nonphotochemical, laser-induced nucleation of supersaturated aqueous glycine
444 produces unexpected γ -polymorph, *Crystal Growth & Design*, 1, 5-8, 2001.
445 Zieger, P., Väisänen, O., Corbin, J. C., Partridge, D. G., Bastelberger, S., Mousavi-Fard, M., Rosati, B., Gysel, M., Krieger, U. K., Leck, C.,
446 Nenes, A., Riipinen, I., Virtanen, A., and Salter, M. E.: Revising the hygroscopicity of inorganic sea salt particles, *Nature Communications*,
447 8, 15883, 10.1038/ncomms15883, 2017.
448 Zimbitas, G., Jawor-Baczynska, A., Vesga, M. J., Javid, N., Moore, B. D., Parkinson, J., and Sefcik, J.: Investigation of molecular and
449 mesoscale clusters in undersaturated glycine aqueous solutions, *Colloids Surf. Physicochem. Eng. Aspects*, 579, 123633, 2019.

450