



Impacts of maritime shipping on air pollution along the U.S. East Coast

Maryam Golbazi¹ and Cristina Archer¹

¹Center for Research in Wind (CRew), University of Delaware, 221 Academy Street, Newark, DE 19716, USA

Correspondence: Maryam Golbazi, Research Application Laboratory, National Center for Atmospheric Research, Boulder, CO 80301, USA (mgolbazi@ucar.edu)

Abstract.

Air pollution is considered a leading threat to human health in the U.S. and worldwide. An important source of air pollution in coastal areas is the globally increasing maritime shipping traffic. In this study, we take a high-resolution modeling approach to investigate the impacts of ship emissions on concentrations of various atmospheric pollutants, under the meteorological conditions and emissions of the year 2018. We utilize the Comprehensive Air Quality Model with extensions (CAMx) to simulate transport, diffusion, and chemical reactions, and the Weather Research and Forecasting (WRF) model to provide the meteorological inputs. We focus on four criteria pollutants – fine particulate matter with a diameter smaller than 2.5 μm (PM_{2.5}), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), and ozone (O₃) – as well as nitrogen oxide (NO), and we calculate their concentrations in the presence and absence of the ship emissions along the U.S. East Coast, particularly in the proximity of major ports.

We find that ship emissions increase the PM_{2.5} concentrations over the ocean and sparse areas inland. The 98-th percentile of the 24-hour average PM_{2.5} concentrations (the “design value” used by the U.S. Environmental Protection Agency) increased by up to 3.2 $\mu\text{g m}^{-3}$ in some coastal areas. In addition, ships contribute significantly to SO₂ concentrations, up to 95% over the Atlantic and up to 90% over land in coastal states, which represents a ~45 ppb increase in the SO₂ design values in some states. The 98-th percentile of the hourly NO₂ concentrations also increased by up to 15 ppb at the major ports and along the shore. In addition, we find that the impact of shipping emissions on O₃ concentrations is not uniform, meaning that ships affect ozone pollution in both positive and negative ways. Over the ocean, O₃ concentrations were significantly higher in the presence of ships, but in major coastal cities O₃ concentrations decreased in the presence of ships. Our simulation results show that ships emit significant amounts of fresh NO in the atmosphere, which then helps scavenge O₃ in VOC-limited areas, such as major ports. By contrast, over the ocean (NO_x-limited regime), enhanced NO_x concentrations due to ships contribute to the formation of O₃ and therefore enhance O₃ concentrations. Overall, due to the dominant southwesterly wind direction in the region, the impacts of ships on air pollutants mainly remain offshore. However, in coastal states near major ports, the impacts are significantly important.



1 Introduction

25 Globally, it is estimated that, in 2019, ambient air pollution, particularly particulate matter (PM) and ozone (O₃), was responsible for 4.5 million premature deaths worldwide (Fuller et al., 2022). This ranked air pollution as a leading risk factor in the Global Burden of Disease Study by the Institute for Health Metrics and Evaluation in 2019 (Murray and Lopez, 1996). Meanwhile, ship traffic is globally increasing and is becoming an important source of air pollution, especially in coastal areas (Corbett and Fischbeck, 1997; Eyring et al., 2010; Schnurr and Walker, 2019). Sea transport accounts for 80% of goods transported
30 worldwide (Schnurr and Walker, 2019), while recent studies estimate demand growth of almost 40% for seaborne trade by 2050 (Serra and Fancello, 2020). Marine vessels are important sources of air pollutants, emitting sulfur oxides (SO_x), nitrogen oxides (NO_x = NO + NO₂), particulate matter (PM), carbon monoxide (CO), volatile organic compounds (VOC), and carbon dioxide (CO₂) (Corbett et al., 2009; Eyring et al., 2007; Eyring, 2008; Smith et al., 2015). Low-grade marine fuel oil contains 3,500 times more sulfur than road diesel (Wan et al., 2016). Studies report fuel consumption of ocean-going ships between 200-290
35 million metric tons for the year 2000 (Corbett and Köhler, 2003; Endresen et al., 2007). Ships are responsible for about 15% of all global anthropogenic NO_x emissions and 4–9% of sulfur dioxide SO₂ emissions. In addition, oceangoing ships are estimated to emit 1.2–1.6 million metric tons (Tg) of PM annually (Corbett et al., 2009; Eyring et al., 2010; Viana et al., 2014).

About 70% of ship emissions occur within 400 km of the shore (Endresen et al., 1999, 2003). Thus, ships can be a major source of pollution in coastal areas and can impact human health. For instance, ship emissions in East Asia have caused
40 14,500–37,500 premature deaths in 2013, the amount of which had doubled since 2005 (Liu et al., 2016). Similarly, particulates emitted from ships cause 60,000 cardiopulmonary and lung cancer deaths each year worldwide (Corbett et al., 2009). Studies from different parts of the world like China show that shipping emissions increased the annual averaged PM_{2.5} concentrations in the eastern coastal regions up to 5.2 μg m⁻³, which was carried 900 km inland (Lv et al., 2018). In Europe, although the increase in PM_{2.5} concentrations by ships is found to be small, their relative contribution is large because of the low background
45 PM_{2.5} concentrations (Viana et al., 2009; Aksoyoglu et al., 2016).

Fine particulate matter (PM_{2.5}) is a harmful air pollutant that consists of microscopic particles with a diameter smaller than 2.5 μm. These particles can penetrate human lungs and even the bloodstream and cause serious health problems (U.S. Environmental Protection Agency (EPA), 2020b). NO_x are a group of highly reactive gases; although seven compounds are technically part of the NO_x family (NO, NO₂, nitrous oxide N₂O, dinitrogen dioxide N₂O₂, dinitrogen trioxide N₂O₃, dinitrogen
50 tetroxide N₂O₄, dinitrogen pentoxide N₂O₅), the most abundant are NO and NO₂, but only NO₂ is actually regulated in the U.S. NO₂ is harmful to humans by irritating the human respiratory system and to the environment by creating acid rain (U.S. Environmental Protection Agency (EPA), 2020a; Lin and McElroy, 2011); it is also a precursor to tropospheric ozone (O₃) formation, which has further negative impacts on human health (, EPA). Similarly, short-term exposure to SO₂ can harm the human respiratory system. These four pollutants – PM_{2.5}, SO₂, NO₂, and O₃ – are both primary (i.e., they can be directly emitted
55 into the atmosphere) and secondary (i.e., they can also form after chemical reactions in the atmosphere) pollutants. Here, we will focus on these four pollutants which are among the seven “criteria” pollutants that are regulated at the federal level by the



U.S. Environmental Protection Agency (EPA) via the National Ambient Air Quality Standards (NAAQS)(U.S. Environmental Protection Agency (EPA), 2022b).

60 Due to the complex nature of the atmosphere and its processes, such as chemical reactions, transport, and diffusion, high concentrations of these pollutants are not necessarily found where their emissions are highest. Therefore, although the ship emissions are released in marine environments, the atmospheric conditions can play an essential role in transporting those pollutants, some of which are precursors for the formation of secondary EPA-regulated pollutants, like O₃.

Here, we explore the impacts of ocean-going ship emissions on the air quality along the U.S. East Coast by utilizing the Comprehensive Air Quality Model with extensions (CAMx) for our simulations. For the first time, we use the most recent high
65 spatial (4 km) and temporal (hourly) resolution ship emission data from the EPA's National Emission Inventory (NEI). We also include the ship stack height to consider at what vertical layer the emissions are emitted into the atmosphere, to be able to account for stability and atmospheric impacts on the pollutants. We investigate the pollution concentrations in a control scenario based on the shipping activities in the year 2018. Then, we conduct another simulation for a hypothetical condition where we eliminate the ship emissions altogether while keeping everything else the same. The difference between the two scenarios gives
70 insights into the net contribution of the ships to air pollution.

Ozone pollution is one of the main focuses of this study. The rate of ozone production can be limited by the concentration of either VOC or NO_x and depends on the relative sources of hydroxyl radical (OH) and NO_x (Finlayson-Pitts and Pitts Jr, 1993). When the rate of OH production is greater than the rate of NO_x production, the rate of ozone production is NO_x-limited. In this situation, ozone concentrations are sensitive to NO_x emissions rather than VOC concentrations. In contrast, when the
75 rate of OH production is less than the rate of NO_x production, ozone production is VOC-limited. In this case, ozone is most effectively reduced by lowering VOC concentrations. NO_x is generally higher where human mobility and transportation are higher (Archer et al., 2020) and While O₃ is generally NO_x-limited in rural areas and downwind suburban areas, in urban areas with high population and high traffic emissions O₃ is often VOC-limited (Seinfeld and Pandis, 1998). Motor vehicles are among the major sources of ozone pollution in the region through their NO_x and VOC emissions (Niemeier et al., 2006; Yao et al.,
80 2015; Zhang et al., 2014). However, the impact of ocean ship emissions along the East Coast of the United States is lacking in the literature and we fill this gap in this study. In locations that exceed the EPA ozone standards by only 2-3 ppb, like the small state of Delaware (Moghani et al., 2018), the ship contribution could be of even higher importance.

2 Methods

2.1 Setup of the WRF-CAMx modeling system

85 We take a modeling approach to explore the pollution concentration across the study domain. The models used in this study are the Weather Research and Forecasting (WRF) model, version 4.3, and the Comprehensive Air quality Model with extensions (CAMx) version 7.1 with the Carbon Bond version 6 revision 5 (CB6r5) chemical mechanism. WRF is developed at the National Center for Atmospheric Research (NCAR) (Skamarock et al., 2019) and is one of the most widely used numerical weather prediction models. CAMx is a modular, Eulerian, 3-dimensional photochemical air quality model (Ramboll Environment



90 and Health, 2020) that simulates the emission, production, advection, diffusion, chemical transformation, and removal of
atmospheric pollutants at regional scales and is among the few that are recommended by the EPA for regulatory purposes (U.S.
Environmental Protection Agency (EPA), 2022a). We use the WRF-CAMx modeling system to conduct simulations of two
separate scenarios, based on the exact same setup and inputs: the first scenario includes the ship emissions (WithShips) while in
the second hypothetical scenario we remove the ship emissions altogether (NoShips). The difference in pollution levels between
95 the two cases provides the net contribution of ship emissions to regional air quality.

CAMx requires input data to characterize meteorology and chemistry, initial and boundary conditions for all the modeling
domains, and other environmental conditions such as the photolysis rates. Meteorology is an essential factor in the formation of
many secondary pollutants, both directly and indirectly. Atmospheric stability plays a significant role in determining pollutant
faith (Arya et al., 1999). In CAMx, the plume rise calculations for point sources including the CMV emissions depend on
100 meteorological conditions and atmospheric stability to determine what vertical layer the emissions are emitted in. In the
summertime, various atmospheric stabilities have been found to be dominant over the Atlantic ocean depending on different
locations (Golbazi et al., 2022; Golbazi and Archer, 2019; Archer et al., 2016). We use the WRF model to provide meteorological
inputs to CAMx. The publicly available WRF-CAMx data processing program (Ramboll Environment and Health, 2020) is used
to generate CAMx meteorological input files from WRF output files. Details on the WRF model setup are provided in Table 1. Our
105 period of study is the summer of 2018, selected to reflect the most recent emission inventory available (discussed next in Section
2.2). Photolysis rate inputs to CAMx were calculated using the Tropospheric Ultraviolet and Visible (TUV) radiative transfer
and photolysis model (<https://www2.acom.ucar.edu/modeling/tropospheric-ultraviolet-and-visible-tuv-radiation-model>).



Table 1. Details of the WRF-CAMx model setup.

Simulation period	1 June – 30 August, 2018
Horizontal grid resolution	4 km
Vertical layers	35
Spin-up time	48 hours
WRF version 4.3	
Initial/boundary conditions	NAM reanalysis, 6-hourly, 12-km resolution
LSM	Noah-modified 21-category IGBP-MODIS
PBL Scheme	MYNN2
Shortwave radiation	RRTMG shortwave
Longwave radiation	RRTMG scheme
SST update	NASA-JPL 1km resolution data
Grid size	400 × 400 grid cells
CAMx version 7.1	
Chemistry	Carbon bond 6 revision 5
Meteorological inputs	WRF model v4.3
Emission data	EPA/NEI 2018
CMV emissions	Inline point sources
Initial/boundary conditions	EPA 2018
Grid size	315 × 300 grid cells

The domain of this study covers the East Coast of the United States (Figure 1) and includes major cities and highly populated regions. Furthermore, it contains several major ports, which are found to experience high shipping traffic. The meteorological files have 400 × 400 horizontal grid points and 35 vertical layers covering the entire CAMx domain, which consists of 315 × 300 grid points, the same as the emission files. Details about the model configuration are discussed in Table 1. Both the WRF and CAMx models have a 4-km horizontal resolution, the same as the emission inventory, in order to avoid spatial interpolation of gridded emissions data. To minimize the impacts of the initial conditions on modeling results, we consider at least 48 hours of spin-up time for both models. Furthermore, as the areas of interest are far from the boundaries, the effects of boundary conditions on modeling results are expected to be minimal.

2.2 Emission data

For emission inputs, we use the most recent emission inventory developed by the EPA, which includes the period May 1st, 2018 to October 1st, 2018 (U.S. Environmental Protection Agency (EPA), 2017) at 4-km horizontal grid resolution and hourly temporal resolution. The emissions are distributed on a 315 × 300 grid, which covers the entire East Coast of the U.S. (Figure 1),




Figure 1. The study domain with 315×300 grid cells and 4-km horizontal grid resolution.

120 with 35 layers vertically. Emissions are treated in two basic ways within CAMx: gridded 2D emissions that are released into each
grid cell of the modeling domain near the surface (i.e., “area sources”, such as traffic or residential heating) and stack-specific
“point sources”, where each stack is assigned unique coordinates and parameters (i.e., smokestacks or ship chimneys). For inline
point source emissions, CAMx computes the plume rise using stack parameters and the hourly emissions for each emission
sector.

125 In the 2018 NEI data, the gridded 2D emissions are merged, meaning that they are provided as one set of surface emissions
that include all emission sectors. On the other hand, the elevated point sources are provided for each potential sector separately.
For ship emissions, we use the emission data for the Commercial Marine Vessels (CMV) sector, which includes Category 1, 2
(small engine), and 3 (large engine) ships. CAMx computes the time-varying buoyant plume rise using stack parameters and
the hourly emissions for each emissions sector, including CMV. Unlike previous EPA data sets, the CMV emissions in 2018 are
130 at a one-hour temporal resolution. The initial and boundary conditions for this study are also provided by the EPA.



3 Results

We study the impacts of ship emissions on the concentrations of four criteria pollutants: PM_{2.5}, SO₂, NO₂, and O₃. We calculate the time-average concentrations over different time periods depending on the pollutant as follows, to match the EPA national standards: one hour for SO₂ and NO₂; 8 hours for O₃; and 24 hours for PM_{2.5}. We analyze every pollutant from two perspectives:
135 1) from a regulatory perspective, thus calculating the statistics that are as close as possible to the EPA design value for each pollutant (Table 2), and 2) from a worst-case perspective, thus calculating the maximum contribution of ships to each pollutant over the entire three-month study period.

Table 2. Design values for criteria pollutants (U.S. Environmental Protection Agency (EPA), 2022b). For attainment purposes, the design values should be calculated by taking the average of the various percentiles over the past three years; since only one year was simulated in this study, the 3-year average could not be calculated and therefore only the actual percentiles were used.

Pollutant	Design value	Threshold for attainment
O ₃	4-th highest 8-hr averaged daily maximum	70 ppb
SO ₂	99-th percentile 1-hr daily maximum	75 ppb
NO ₂	98-th percentile 1-hr daily maximum	100 ppb
PM _{2.5}	98-th percentile 24-hr average	35 µg m ⁻³

To calculate the maximum contribution, we first find the differences between the two cases (WithShips minus NoShips) at every grid cell, averaged over the relevant time interval, which depends on the pollutant (Table 2); then, we find the maximum
140 difference through the 3 months at every grid cell as follows:

$$\max(\Delta P_{i,j}) = \max_{t \in [1 \dots n]} (P_{i,j}^{WithShips}(t) - P_{i,j}^{NoShips}(t)) \quad (1)$$

where n is the number of data on the 1-, 8-, or 24-hr averaged pollutant P concentration values (the exact time averaging window depends on the pollutant, see Table 2) over the 3-month period of study, $P^{WithShips}$ and $P^{NoShips}$ are pollutant P concentrations with and without the ships, respectively, and i and j correspond to the model grid cell indices.

145 Although the maximum contribution from Eq. 1 is not valuable in terms of reaching or maintaining the EPA attainment for states, it is essential to understand the importance of maritime shipping on air quality, physically and statistically. A summary of the design values defined here for each pollutant to represent the EPA standards and the threshold for attainment are presented in Table 2. The defined design values follow the same criteria as defined by the EPA (U.S. Environmental Protection Agency (EPA), 2022b) but only for the time period of this study. For the remainder of the article, we will assume that the design values
150 defined in Table 2 are serving the purpose of analyzing the pollution from a regulatory perspective and are the same as the EPA standards for those pollutants for the time period of this study.




We find that the concentration of carbon monoxide (CO) remains unchanged in presence of ships (not shown), suggesting that the CMV sector has minimal impact on the CO concentrations in the region. As such, we do not discuss the CO results in the rest of this study.

155 3.1 Fine particulate matter (PM_{2.5})

The PM_{2.5} species used in this study are those included in the CAMx model output (Ramboll Environment and Health, 2020). The EPA requires that the 3-year average of the 98-th percentile of the daily mean PM_{2.5} concentrations should not exceed 35 µg m⁻³. Here, we calculated the 98-th percentile of the 24-h averaged PM_{2.5} concentrations at every grid cell during the simulation period in both scenarios, WithShips, and NoShips.

160 We find that PM_{2.5} levels stayed below 35 µg m⁻³ across most of the domain and that only two locations, i.e., Manhattan, NY, and Easton, PA (Figure 2a), crossed the 35 µg m⁻³ maximum allowed concentrations and therefore were in non-attainment based on the design value defined in this study in Table 2. From a policy perspective, the CMV sector increases PM_{2.5} levels up to 3.2 µg m⁻³ in Manhattan, NY, and up to 2 µg m⁻³ elsewhere (Figure 2c). This is while the percent contribution to PM_{2.5} concentrations remains below 27% across the domain (Figure 2d). In a worst-case scenario, however, the maximum contribution of the ships to PM_{2.5} concentrations within the 3 months is significantly high across the domain but due to the dominant southwesterly wind direction in the region (Golbazi et al., 2022), it mostly remains over the Atlantic Ocean (Figure 2b). The maximum impact on PM_{2.5} during the 3 months reaches as high as 8 µg m⁻³. Across the domain, the highest impacts are found offshore of MD and VA and in the Chesapeake Bay, DE. Over the land, the highest impacts are in Manhattan, NY, CT, and coastal MA.





170 3.2 Sulfur dioxide (SO₂)

The SO₂ design value is defined as the 99-th percentile daily maximum SO₂ concentrations in the simulation period, which should not exceed 75 ppb (Table 2). Here we calculated the 99-th percentile of daily maximum SO₂ concentrations at every grid cell over the simulation period in the two scenarios, i.e., WithShips and NoShips. Then, we subtracted these two cases from one other (WithShips minus NoShips) to obtain the net effect of the maritime shipping sector.

175 Our results show that ships have a significantly high impact on SO₂ concentrations, up to 95% and 90% over the Atlantic Ocean and inland, respectively (Figure 3d). This suggests that the CMV sector is one of the highest contributors to SO₂ levels in the region. The increase in the SO₂ design value by ships remains mainly offshore and around the major shipping routes (Figure 3c). However, it reaches the interior of land in major ports and some parts of the coastal states. Over the simulation period, the contribution of the ships to the 99-th percentile daily SO₂ maxima is up to 45 ppb, with the highest impact in Baltimore,
180 Maryland (MD), and Norfolk, Virginia (VA), and parts of New Jersey and Long Island (Figure 3c). We note that, however, the SO₂ design value in the region remained below 75 ppb in all states (Figure 3a). It is worth mentioning that the locations with the highest SO₂ concentrations are the ones highly impacted by the ships.

In addition, we calculated the 3-month maximum contributions of the ships to SO₂ concentrations, which indicates the worst-case scenario at every grid cell (figure 3b). Although the increase in SO₂ design value was mainly offshore, the maximum
185 contribution of the ships to SO₂ showed a different pattern, with a maximum increase of up to ~185 ppb at a few grid cells around Norfolk, VA. We note that an occasional and short-term spike of high concentrations of SO₂, as we report here for Norfolk, is not necessarily associated with a strong health impact.




Figure 3. SO₂ concentration results: a) 99-th percentile of daily 1-hour maximum SO₂ concentrations with ships (WithShips); b) maximum contribution of the ships during the simulation period (Eq. 1); c) differences between the 99-th percentile daily maximum between the two scenarios (WithShips minus NoShips); and d) percent contribution of the ships to the 99-th percentile of the daily SO₂ maximum.



3.3 Nitrogen dioxide (NO₂)

NO₂ is a precursor to tropospheric ozone (O₃) formation, which has further negative impacts on human health (, EPA). NO₂ is generally directly emitted into the atmosphere from emission sources, including ships. We find that ships cause a significant increase in the 98-th percentile of daily maximum NO₂ concentrations, up to 34 ppb, but only at a few locations along the coast and in coastal states with major ports (Figure 4c), suggesting that, except for states with large ports, ships do not significantly impact the state compliance with the EPA standards. However, in NC, VA, DE, NY, and CT the shipping impacts reach beyond 15 ppb from a regulatory standpoint. Among these states, while NC remains in attainment with regulations (below 100 ppb), it experiences up to 80% contribution from the ships to its NO₂ concentrations. On the other hand, NY is the only state in the study domain that exceeds the 100 ppb standard for NO₂ concentrations, and shipping contribution to its non-attainment is 20-25%.

The maximum contribution of the ships to NO₂ concentrations, which is illustrated in Figure 4b, shows that, in a worst-case scenario, ships contributed to up to ~50 ppb of NO₂ along the coast and 75 ppb over the Atlantic, which is significantly high compared to 100 ppb standard. The 3-month highest impact happens near the major ports and shipping routes but stretches to the land and over the ocean.

3.4 Ozone (O₃)

Tropospheric ozone is formed by both naturally occurring and anthropogenic sources. Ozone is not emitted directly into the air, but, in the presence of sunlight, it is created by its precursors: VOC and NO_x. The rate of ozone production can be limited by either VOCs or NO_x. As a result, a specific location can be either VOC-limited or NO_x-limited. The rate of production/destruction of O₃ in the atmosphere is different in either of these regimes. We will further discuss this matter later in this section.

We use 8-hour average ozone concentrations for our analysis to maintain consistency with the EPA standards (U.S. Environmental Protection Agency (EPA), 2022b). We calculated the 8-hr averaged ozone values by averaging consecutive eight hours of O₃ outputs at each hour of the day and storing it at the start hour (Cohen et al., 1999). For instance, O₃ at 11:00 in a day indicates the time-averaged O₃ concentrations between hours 11:00 and 19:00 in that day. Hereafter, we will refer to the 8-hr averaged O₃ concentrations simply as O₃ concentrations.

Ambient ozone concentrations are directly affected by temperature, solar radiation, wind speed, and other meteorological factors. Since O₃ production is a photochemical reaction, its peak concentrations are found during the daytime when tropospheric ultraviolet radiation in the atmosphere is highest. Since the focus of this study is the daily high episodes of O₃ that are associated with adverse health impacts, we limit our analysis of the maximum impacts to only daytime hours. To select the daytime hours, we considered the 8-hr averaged O₃ daily profiles in 10 different locations along the coast from which 4 selected locations are shown in Figure 5. The lowest concentrations of O₃ are associated with hours 00:00 to 08:00 UTC (20:00 to 04:00 local time). We eliminated these hours from our analysis to only focus on concentrations during the high episodes. Therefore, we select the hours with peak O₃ concentrations during the 24-h period i.e., 09:00 to 23:00 UTC (05:00 to 19:00 local time).




Figure 4. NO_2 concentration results: a) 98-th percentile of 1-hour daily maximum NO_2 concentrations with ships (WithShips); b) maximum contribution of the ships during the simulation period (Eq. 1); c) differences between the 98-th percentile of the 1-hour daily maximum between the two scenarios (WithShips minus NoShips); and d) percent contribution of the ships to the 98-th percentile of the daily NO_2 maximum.



220 We find that although ship emissions contribute to O_3 enhancement in the region, they reduce O_3 at some urban locations. We detect a significant increase or decrease in O_3 concentrations in presence of the ships, depending on whether the location was NO_x or VOC limited.




Figure 5. Eight-hour average O_3 concentration time series at four different locations along the East Coast from June 1st to August 30th, 2018: a) Atlantic City, NJ; b) Manhattan, NY; c) Cape Cod, MA; and d) Providence, RI. The thin lines represent the daily time series and the thick lines are the 3-month average profiles with red lines for WithShips and blue lines for the NoShips simulations.

The maximum contribution of the ships to the O_3 levels over the entire 3-month period is illustrated in Figure 6c as a worst-case scenario. Over the ocean, the maximum increase is large, up to 8.6 ppb. However, the pollution increase remained primarily offshore and did not significantly impact the coastal areas. O_3 increased by 4-5 ppb at most in parts of North Carolina, Baltimore, MD, parts of CT, and MA. Otherwise, the maximum increase over the land was up to 3.5 ppb. It's important to note that the maximum impact is not necessarily at the time when high O_3 episodes (from a regulatory perspective) are found. Despite the O_3 increase over the Atlantic, at the shore where the major ports are built, the maximum contribution of the ships is negative, suggesting that O_3 was destructed in presence of the ships. This is due to the complex nature of atmospheric chemistry, where the fresh NO emissions from the ships scavenge O_3 and reduce its concentrations in urban VOC-limited areas.

While Figure 6b is important to understand the worst-case scenario of the shipping impact on O_3 pollution in the region, it does not help to measure the impact of these changes on state compliance with EPA standards. This is because a high impact on O_3 in the worst-case scenario may not correspond to the time of the day when O_3 daily maxima occurred. Therefore, to study the impacts of ship emissions from a policy perspective, it is necessary to explore the impacts on the O_3 design value at every grid cell. For O_3 , the EPA defines this standard as the 4-th highest 8-hr averaged O_3 daily maxima, averaged over a 3-year period




Figure 6. O₃ concentration results: a) 4-th highest 8-hour daily maximum O₃ concentrations with ships (WithShips); b) maximum contribution of the ships during the 3-month simulation period (Eq. 1); c) differences between the 4-th highest 8-hour daily maximum between the two scenarios (WithShips minus NoShips); and d) percent contribution of the ships to the 4-th highest daily O₃ maximum.



which should not exceed 70 ppb (U.S. Environmental Protection Agency (EPA), 2022b). Since we do not have data for three consecutive years, we focus on the 4-th highest daily maximum in our study period, the summer of 2018. Hereafter, we will refer to this value as O₃ design value in this study. We assume that this value represents the 4th highest daily maximum in the year 2018 since the highest O₃ episodes are expected to occur during the summer period. Thus, the regions with higher than 70
240 ppb O₃ concentrations in 6a are most susceptible to being in non-attainment with the EPA standards and therefore the impacts of the ships are of higher significance in those regions. Out of all states in the domain, NY, NJ, and MD are the only states that exceed the 70 ppb standard and are likely to be in non-attainment. All other states stay in attainment with the standards defined in Table 2 in either scenario.

From a policy perspective, O₃ design values increased in presence of the ships by up to 3.5 over the Atlantic Ocean. However,
245 we find a reduction (up to 6.5 ppb) in O₃ concentrations at major ports along the East Coast (Figure 6c), where fresh NO is emitted by the ships into the atmosphere in VOC-limited regions (Figure 7). In most parts, the major impact of the ships remains offshore away from the coastal areas. However, in some regions in MA, RI, CT, ME, VA, NC, and MD ships contribute to O₃ increase at the coast from which, only MD is likely to be in non-attainment. The highest increase in O₃ design value inland is found in VA and NC and is up to 2.5 ppb, while we note that in NY, the 4-th highest daily maximum is decreased by 4 ppb in
250 presence of the ships for the reasons discussed later in this section. However, the decrease in O₃ values remains in Manhattan, NY, and is not associated with the parts of Long Island (NY) where O₃ exceeds the standards.

In the atmosphere, the formation or destruction of ozone depends on the concentrations of both NO_x and VOC and the ratio between them (*VOC/NO_x*). Transportation usually is associated with high NO_x emissions, therefore O₃ is generally NO_x-limited in rural areas and VOC-limited in urban areas, with low and high traffic densities, respectively.

255 In the VOC-limited regions, high concentrations of freshly emitted NO locally scavenge O₃ and lead to the formation of NO₂. Close to the emission sources, this titration process can be considered an ozone sink. In addition, high NO₂ concentrations deflect the initial oxidation step of VOC by forming other products such as nitric acid (HNO₃), which slows down the formation of O₃ (National Research Council, 1992; Beck et al., 1998). Because of these reactions, an increase in NO leads to a decrease in O₃ at VOC-limited regions.

260 In CAMx, the VOC-limited regime is defined when the rate of change of hydrogen peroxide (H₂O₂) is lower compared to the rate of change of HNO₃. A higher $\Delta\text{H}_2\text{O}_2/\Delta\text{HNO}_3$ ratio indicates a NO_x-limited regime, while a lower $\Delta\text{H}_2\text{O}_2/\Delta\text{HNO}_3$ ratio corresponds to a VOC-limited regime. There are other indicators for determining the NO_x/VOC-limited regimes that are discussed in the literature (Li et al., 2022). However, here we use the $\Delta\text{H}_2\text{O}_2/\Delta\text{HNO}_3$ ratio as is used in the CAMx model (Ramboll Environment and Health, 2020).

265 To understand the formation/destruction of ozone in presence of the ships in our study domain, we calculated the frequency of the VOC-limited regime based on the ratio at every grid cell. Figure 7a illustrates the percentage of the times that a VOC-limited regime occurred at every cell, which is the highest along the coast and in major cities. This indicates that O₃ may be affected by titration when ship emissions are present. We find that the NO concentrations increase along the coast where we detect a decrease in O₃ concentrations (Figure 7b) and a VOC-limited regime (Figure 7c).



270 This finding does not necessarily mean that ships help create better air quality since a reduction in O_3 is due to a significant increase in other important air pollutants i.e., NO_x concentrations.




Figure 7. VOC- versus NO_x -limited regime determination: a) percentage of the times when $H_2O_2/HNO_3 < 0.35$ as determined in CAMx model, which is an indication of a VOC-limited regime; b) maximum contribution of ships to O_3 pollution (Figure 6c); and c) same as in b) but for NO.

4 Conclusions

Ships emit significant amounts of pollutants within 400 km of the shores. Here, we studied the ocean-going ship emissions on the air quality of the U.S. East Coast. We utilized the WRF-CAMx modeling system to simulate the pollution concentrations in the presence and absence of shipping activities along the East Coast and at the major ports. We used the WRF model to provide the meteorological inputs for the CAMx air quality model for the year 2018, on which the most recent EPA/NEI emission inventory is based. We particularly focused on $PM_{2.5}$, SO_2 , NO_2 , and O_3 . Overall, we studied the outcomes on every pollutant from two perspectives: 1) from the EPA perspective concerning the national concentration standards for each specific pollutant, and 2) the maximum contribution of ships to that pollutant over the 3 months.

280 We find that shipping increases the $PM_{2.5}$ concentrations across the domain. the 98-th percentile daily average $PM_{2.5}$ levels increased by $3.2 \mu g m^{-3}$ over the ocean and in some coastal areas. However, in a worst-case scenario, ships contribute up to approximately $8.0 \mu g m^{-3}$ to $PM_{2.5}$ concentrations, only over the Atlantic off the coast of MD, and VA. In addition, we find that ships have a significantly high impact, up to 95% and 90%, on the SO_2 concentrations over the Atlantic and inland, respectively. This suggests that the CMV sector is one of the highest contributors to SO_2 levels in the region. The shipping contribution to SO_2 levels was up to 45 ppb over coastal regions. Ship emissions also impacted the NO_2 design value by up

285



to 34 ppb. In addition, our simulation results show that the impact of ship emissions on O₃ concentrations is not uniform, meaning that maritime shipping affects ozone pollution in both positive and negative ways. Although over the ocean O₃ concentrations increase significantly in the presence of ships (up to 8.6 ppb), in coastal areas with major cities and major ports O₃ concentrations decrease by up to ~6.5 ppb. To understand the reasons behind the O₃ reduction in the presence of ships, we
290 analyzed the $\Delta\text{H}_2\text{O}_2/\Delta\text{HNO}_3$ ratio in the region, which is used to determine NO_x- or VOC-limited ozone production, as well as changes to NO concentrations, since they play a significant role in O₃ formation and destruction. We found that ships emit significant amounts of fresh NO in the atmosphere, which then helps scavenge O₃ in VOC-limited regimes. As a result, with higher NO concentrations in the atmosphere produced by ship emissions, O₃ is destructed in major cities and urban areas. By contrast, over the ocean (a NO_x-limited regime), excessive NO_x emissions due to the ships contribute to the formation of O₃
295 and therefore an enhancement in O₃ concentrations. It is important to note that the destruction of O₃ by ship emissions in major cities does not necessarily mean that the ships create better air quality because a decrease in O₃ is a consequence of a significant increase in other pollutants, like NO. Overall, the majority of the time, due to the dominant southwesterly wind direction in the region, the impacts on different pollutants remained spatially confined offshore. However, in some coastal areas near the major ports, the impacts were significant.

300 **Data availability**

The data for the model setup is available in the GitHub repository at:
<https://github.com/golbazimaryam/ShippingEmissionsAndAirQuality>

Author contribution

M. Golbazi contributed to the design of the study, preparing the manuscript, setting up and carrying out simulations, and
305 analysis. C.L. Archer contributed to the design of the project, preparing the manuscript and editing, as well as data analysis.

Competing interests

"The authors declare that they have no conflict of interest."

Acknowledgements. *

Partial funding for this research came from the University of Delaware (UD) Graduate College Doctoral Fellowship and from the Delaware
310 Natural Resources and Environmental Control (DNREC, award no. 18A00378). The simulations were conducted on the UD Caviness and NCAR Cheyenne high-performance computer clusters.



References

- Aksoyoglu, S., Baltensperger, U., and Prévôt, A. S.: Contribution of ship emissions to the concentration and deposition of air pollutants in Europe, *Atmospheric Chemistry and Physics*, 16, 1895–1906, 2016.
- 315 Archer, C. L., Colle, B. A., Veron, D. L., Veron, F., and Sienkiewicz, M. J.: On the predominance of unstable atmospheric conditions in the marine boundary layer offshore of the US northeastern coast, *Journal of Geophysical Research: Atmospheres*, 121, 8869–8885, 2016.
- Archer, C. L., Cervone, G., Golbazi, M., Al Fahel, N., and Hultquist, C.: Changes in air quality and human mobility in the USA during the COVID-19 pandemic, *Bulletin of Atmospheric Science and Technology*, 1, 491–514, 2020.
- Arya, S. P. et al.: *Air pollution meteorology and dispersion*, vol. 310, Oxford University Press New York, 1999.
- 320 Beck, J., Krzyzanowski, M., and Koffi, B.: Tropospheric Ozone in the European Union - The consolidated report, Tech. rep., EEA Topic Report, 1998.
- Cohen, J., Fitz-Simons, T., and Wayland, M.: Guideline on data handling conventions for the PM NAAQS, Tech. rep., Environmental Protection Agency, Office of Air Quality Planning and . . . , 1999.
- Corbett, J. J. and Fischbeck, P.: Emissions from ships, *Science*, 278, 823–824, 1997.
- 325 Corbett, J. J. and Köhler, H. W.: Updated emissions from ocean shipping., *Journal of Geophysical Research*, 108, 2003.
- Corbett, J. J., Winebrake, J. J., Green, E. H., Kasibhatla, P., Eyring, V., and Lauer, A.: Mortality from Ship Emissions: A Global Assessment, *Science and Technology*, 2009.
- Endresen, O., Sorgard, E., Sundet, J. K., Dalsoren, S. B., Isaksen, I. S. A., Berglen, T. F., and Gravir, G.: Global nitrogen and sulfur inventories for oceangoing ships., *Journal of Geophysical Research: Atmospheres*, 104, 3457–3470, 1999.
- 330 Endresen, O., Sorgard, E., Sundet, J., Dalsoren, S., Isaksen, I., Berglen, T., and Gravir, G.: Emission from international sea transportation and environmental impact., *Journal of Geophysical Research*, 108, 2003.
- Endresen, O., Sorgard, E., Behrens, H. L., Brett, P. O., and Isaksen, I. S. A.: A historical reconstruction of ships fuel consumption and emissions., *Journal of Geophysical Research*, 112, 2007.
- (EPA), U. E. P. A.: Ground-level ozone pollution, <https://www.epa.gov/no2-pollution/basic-information-about-no2#WhatisNO2>, 2020.
- 335 Eyring, V.: Updated Study on Greenhouse Gas Emissions from Ships: Climate Impacts, Tech. rep., 2008.
- Eyring, V., Corbett, J. J., Lee, D. S., and Winebrake, J. J.: Brief summary of the impact of ship emissions on atmospheric composition, climate, and human health, Tech. rep., Health and Environment sub-group of the International Maritime Organization, 2007.
- Eyring, V., Isaksen, I. S. A., Berntsen, T., Collins, W. J., Corbett, J. J., Endresen, O., Grainger, R. G., Moldanova, J., Schlager, H., and Stevenson, D. S.: Transport impacts on atmosphere and climate: Shipping, *Atmospheric Environment*, 44, 4735–4771, 2010.
- 340 Finlayson-Pitts, B. and Pitts Jr, J.: Atmospheric chemistry of tropospheric ozone formation: scientific and regulatory implications, *Air & Waste*, 43, 1091–1100, 1993.
- Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., Caravanos, J., Chiles, T., Cohen, A., Corra, L., et al.: Pollution and health: a progress update, *The Lancet Planetary Health*, 2022.
- Golbazi, M. and Archer, C. L.: Methods to Estimate Surface Roughness Length for Offshore Wind Energy, *Advances in Meteorology*, 2019.
- 345 Golbazi, M., Archer, C. L., and Alessandrini, S.: Surface impacts of large offshore wind farms, *Environmental Research Letters*, 17, 064 021, 2022.



- Li, X., Qin, M., Li, L., Gong, K., Shen, H., Li, J., and Hu, J.: Examining the implications of photochemical indicators for O₃–NO_x–VOC sensitivity and control strategies: A case study in the Yangtze River Delta (YRD), China, *Atmospheric Chemistry and Physics*, 22, 14 799–14 811, 2022.
- 350 Lin, J.-T. and McElroy, M. B.: Detection from space of a reduction in anthropogenic emissions of nitrogen oxides during the Chinese economic downturn, *Atmospheric Chemistry and Physics*, 2011.
- Liu, H., Fu, M., Jin, X., Shang, Y., Shindell, D., Faluvegi, G., Shindell, C., and He, K.: Health and climate impacts of ocean-going vessels in East Asia, *Nature climate change*, 6, 1037–1041, 2016.
- Lv, Z., Liu, H., Ying, Q., Fu, M., Meng, Z., Wang, Y., Wei, W., Gong, H., and He, K.: Impacts of shipping emissions on PM 2.5 pollution in
355 China, *Atmospheric Chemistry and Physics*, 18, 15 811–15 824, 2018.
- Moghani, M., Archer, C. L., and Mirzakhali, A.: The importance of transport to ozone pollution in the U.S. Mid-Atlantic, *Atmospheric Environment*, 191, 420–431, 2018.
- Murray, C. J. L. and Lopez, A. D.: Evidence-based health policy — Lessons from the Global Burden of Disease Study, *Science*, 274, 740–743, 1996.
- 360 National Research Council: Rethinking the ozone problem in urban and regional air pollution, National Academies Press, Washington, DC, <https://doi.org/10.17226/1889>, 1992.
- Niemeier, U., Granier, C., Kornblueh, L., Walters, S., and Brasseur, G.: Global impact of road traffic on atmospheric chemical composition and on ozone climate forcing., *Journal of Geophysical Research: Atmospheres*, 2006.
- Ramboll Environment and Health: COMPREHENSIVE AIR QUALITY MODEL WITH EXTENSIONS Version 7.10 – User’s guide, Tech.
365 rep., https://camx-wp.azurewebsites.net/Files/CAMxUsersGuide_v7.10.pdf, 2020.
- Schnurr, R. E. and Walker, T. R.: Marine Transportation and Energy Use, in: Reference Module in Earth Systems and Environmental Sciences, Elsevier, 2019.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics*, John Wiley and sons, 1998.
- Serra, P. and Fancello, G.: Towards the IMO’s GHG goals: A critical overview of the perspectives and challenges of the main options for
370 decarbonizing international shipping, *Sustainability*, 12, 3220, 2020.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., et al.: A description of the advanced research WRF model version 4, Tech. rep., National Center for Atmospheric Research, Boulder, CO, USA, 2019.
- Smith, T., Jalkanen, J., Anderson, B., Corbett, J., Faber, J., Hanayama, S., O’Keeffe, E., Parker, S., Johansson, L., Aldous, L., et al.: Third
375 IMO greenhouse gas study 2014, Tech. rep., 2015.
- U.S. Environmental Protection Agency (EPA): PROFILE OF VERSION 1 OF THE 2014 NATIONAL EMISSIONS INVENTORY, Tech. rep., Office of Air Quality Planning and Standards, https://www.epa.gov/sites/production/files/2017-04/documents/2014neiv1_profile_final_april182017.pdf, Accessed 06/17/2020, 2017.
- U.S. Environmental Protection Agency (EPA): Nitrogen Dioxide (NO₂) Pollution, <https://www.epa.gov/ground-level-ozone-pollution/ground-level-ozone-basics>, accessed 07/28/2020, 2020a.
- 380 U.S. Environmental Protection Agency (EPA): Particulate Matter (PM) Pollution, <https://www.epa.gov/pm-pollution/particulate-matter-pm-basics#PM>, accessed 06/10/2020, 2020b.
- U.S. Environmental Protection Agency (EPA): Photochemical Air Quality Modeling, Support Center for Regulatory Atmospheric Modeling (SCRAM), <https://www.epa.gov/scram/photochemical-air-quality-modeling>, accessed on 12/01/2022, 2022a.



- 385 U.S. Environmental Protection Agency (EPA): NAAQS Table, <https://www.epa.gov/criteria-air-pollutants/naaqs-table>, accessed 04/04/2022,
2022b.
- Viana, M., Amato, F., Alastuey, A., Querol, X., Moreno, T., Garcia Dos Santos, S., Hecce, M. D., and Fernández-Patier, R.: Chemical tracers
of particulate emissions from commercial shipping, *Environmental science & technology*, 43, 7472–7477, 2009.
- Viana, M., Hammingh, P., Colette, A., Querol, X., Degraeuwe, B., Vlieger, I., and Aardenne, J.: Impact of maritime transport emissions on
390 coastal air quality in Europe, *Atmospheric Environment*, 2014.
- Wan, Z., Zhu, M., Chen, S., and Sperling, D.: Pollution: Three steps to a green shipping industry, *Nature*, 530, 275–277, 2016.
- Yao, Z., Wu, B., Shen, X., Cao, X., Jiang, X., Ye, Y., and He, K.: On-road emission characteristics of VOCs from rural vehicles and their
ozone formation potential in Beijing, China., *Atmospheric Environment*, 2015.
- Zhang, L., Jacob, D. J., Yue, X., Downey, N. V., Wood, D. A., and Blewitt, D.: Sources contributing to background surface ozone in the US
395 Intermountain West., *Atmospheric Chemistry and Physics*, 2014.